


## CORRECTION

# Correction: Operative treatment of fragility fractures of the pelvis is connected with lower mortality. A single institution experience

Pol Maria Rommens, Mehdi Boudissa, Sven Krämer, Miha Kisilak, Alexander Hofmann, Daniel Wagner

There is an error in the caption of [Table 7](#) Please see the complete, correct [Table 7](#) caption here.



## OPEN ACCESS

**Citation:** Rommens PM, Boudissa M, Krämer S, Kisilak M, Hofmann A, Wagner D (2021) Correction: Operative treatment of fragility fractures of the pelvis is connected with lower mortality. A single institution experience. PLoS ONE 16(9): e0258076. <https://doi.org/10.1371/journal.pone.0258076>

**Published:** September 27, 2021

**Copyright:** © 2021 Rommens et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 7. Demographics, LoS, mortality, situation at discharge and at follow up of FFP type III and FFP type IV patients, depending on type of treatment.

Type of treatment	All patients	non-operative	operative	p-value
Number of patients (n, %)	238 (100)	186 (78.2)	52 (21.8)	
Age (median)	82	82.5	79	<b>0.006</b>
Women (n, %)	204 (85.7)	159 (85.5)	45 (86.5)	
Men (n, %)	34 (14.3)	27 (14.5)	7 (13.5)	0.848
Patients with comorbidities (n, %)	223 (93.7)	174 (93.5)	49 (94.2)	0.787
Patients with two or more comorbidities (n, %)	131 (55.0)	104 (55.9)	27 (51.9)	0.609
Median length of hospital stay (days)	10	9	17	<b>&lt;0.001</b>
Minimum	0	0	3	
Maximum	92	28	92	
IQR	7–14.8	6–12	14.8–25	
General in-hospital complications (n, %)	53 (22.3)	39 (19.8)	14 (27.5)	0.362
Surgery-related complications (n, %)	n.a.	n.a.	12 (23.1)	
Surgical revisions (n, %)			9 (17.3)	
<b>Mobility at discharge (n, %)</b>				
Ward	143 (60.1)	121 (65.1)	22 (42.3)	
Room	43 (18.1)	35 (18.8)	8 (15.4)	
Transfer	41 (17.2)	23 (12.4)	18 (34.6)	
Bedridden	4 (1.7)	3 (1.6)	1 (1.9)	
Not documented	5 (2.1)	2 (1.1)	3 (5.8)	
Death in hospital	2 (0.8)	2 (1.1)	0 (0.0)	
<b>Mobility at discharge (n, %)</b>				
Walker	186 (80.5)	156 (85.7)	30 (61.2)	
Non-walker	45 (19.5)	26 (14.3)	19 (38.8)	<b>&lt;0.001</b>
<b>Destination at discharge (n, %)</b>				
Home	107 (45.0)	88 (47.3)	19 (36.5)	
Geriatrics	67 (28.1)	56 (30.1)	11 (21.2)	
Rehabilitation	49 (20.6)	31 (16.7)	18 (34.6)	
Hospital	8 (3.4)	5 (2.7)	3 (5.8)	<b>0.022</b>
Not documented	5 (2.1)	4 (2.2)	1 (1.9)	
Death in hospital	2 (0.8)	2 (1.1)	0 (0.0)	
<b>Destination at discharge (n, %)</b>				
Independent	167 (57.4)	88 (48.8)	19 (37.3)	
Dependent	124 (42.6)	92 (51.2)	32 (62.7)	0.141
<b>Mobility at discharge (n, %)</b>				
Walking	186 (80.5)	156 (85.7)	30 (61.2)	
Non walking	45 (19.5)	26 (14.3)	19 (38.8)	<b>&lt;0.001</b>
Mortality in hospital (%)	1.1	1.1	0.0	0.453
One-year mortality (%)	14.2	15.9	8.0	0.154
Two-year mortality (%)	21.6	25.3	8.0	
Five-year mortality (%)	59.1	67.1	33.2	
Overall mortality				<b>0.001</b>
Mortality before follow up (n, %)	87 (36.6)	76 (40.9)	11 (21.2)	
Lost to follow up (n, %)	16 (6.7)	10 (5.4)	6 (11.5)	
Follow up (n, %)	135 (56.7)	100 (53.8)	35 (67.3)	
Follow up of surviving patients (n, %)	135 (89.4)	100 (90.9)	35 (85.4)	
Median follow up time (weeks)	164	117	119.7	
<b>Patients with PCS and MCS (n, %)</b>	67/135 (49.6)	51/100 (51.0)	16/35 (45.7)	

(Continued)

Table 7. (Continued)

Type of treatment	All patients	non-operative	operative	p-value
Median SF-8 physical (PCS)	32.21	33.41	29.39	0.734
Min SF-8 physical	16.34	16.34	17.34	
Max SF-8 physical	62.51	62.51	56.68	
Median SF-8 mental (MCS)	54.41	53.55	55.42	0.938
Min SF-8 mental	19.16	19.16	19.83	
Max SF-8 mental	68.44	68.44	65.17	
<b>Patients with PMS (n, %)</b>	134/135 (99.3)	99/100 (99.0)	35/35 (100.0)	
Median PMS	5	5	6	0.285
Min PMS	0	0	0	
Max PMS	9	9	9	
<b>Patients with NRS (n, %)</b>	133/135 (98.5)	98/100 (98.0)	35/35 (100.0)	
Median NRS	3	1	4	<b>0.024</b>
Min NRS	0	0	0	
Max NRS	10	10	9	

P-values below 0.05 are shown in bold.

<https://doi.org/10.1371/journal.pone.0258076.t001>

## Reference

1. Rommens PM, Boudissa M, Krämer S, Kisilak M, Hofmann A, Wagner D (2021) Operative treatment of fragility fractures of the pelvis is connected with lower mortality. A single institution experience. PLoS ONE 16(7): e0253408. <https://doi.org/10.1371/journal.pone.0253408> PMID: 34242230