



Predictive value of frailty in the mortality of hospitalized patients with COVID-19: a systematic review and meta-analysis

Yupei Zou^{1,2}, Maonan Han³, Jiarong Wang³, Jichun Zhao³, Huatian Gan^{1,2}, Yi Yang^{2,3}

¹The Center of Gerontology and Geriatrics, West China Hospital, Sichuan University, Chengdu, China; ²National Clinical Research Center for Geriatrics, West China Hospital, Sichuan University, Chengdu, China; ³Department of Vascular Surgery, West China Hospital, Sichuan University, Chengdu, China

Contributions: (I) Conception and design: Y Yang; (II) Administrative support: H Gan; (III) Provision of study materials or patients: Y Zou, M Han; (IV) Collection and assembly of data: Y Zou, M Han; (V) Data analysis and interpretation: Y Zou, M Han; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Huatian Gan, MD. The Center of Gerontology and Geriatrics, West China Hospital, Sichuan University, 37 Guo Xue Alley, Chengdu 610041, China. Email: ganhuatian123@163.com; Yi Yang, MD. Department of Vascular Surgery, West China Hospital, Sichuan University, 37 Guo Xue Alley, Chengdu 610041, China. Email: yangy@wchscu.cn.

Background: The present study aimed to analyze the impact of frailty on mortality risk among hospitalized patients with coronavirus disease 2019 (COVID-19).

Methods: Literature searches were conducted using the MEDLINE, Embase, and Cochrane databases for articles reporting the association between frailty and mortality in hospitalized patients with COVID-19. The quality of the included studies was assessed using the Newcastle-Ottawa scale (NOS). A random-effects meta-analysis was performed to calculate the pooled effects.

Results: A total of 21 studies with 26,652 hospitalized patients were included. Sixteen studies used the Clinical Frailty Score (CFS), and five used other frailty assessment tools. The pooled estimates of frailty in hospitalized patients with COVID-19 were 51.4% [95% confidence interval (CI): 39.9–62.9%]. In the CFS group, frail patients experienced a higher rate of short-term mortality than non-frail patients [odds ratio (OR) =3.0; 95% CI: 2.3–3.9; $I^2=72.7\%$; $P<0.001$]. In the other tools group, frail patients had a significantly increased short-term mortality risk compared with non-frail patients (OR =2.4; 95% CI: 1.4–4.1; $P=0.001$). Overall, a higher short-term mortality risk was observed for frail patients than non-frail patients (OR =2.8; 95% CI: 2.3–3.5; $P<0.001$). In older adults, frail patients had a higher rate of short-term mortality than non-frail patients (OR =2.3; 95% CI: 1.8–2.9; $P<0.001$).

Conclusions: Compared to non-frail hospitalized patients with COVID-19, frail patients suffered a higher risk of all-cause mortality, and this result was also found in the older adult group.

Keywords: Meta-analysis; frailty; mortality; coronavirus disease 2019 (COVID-19)

Submitted Nov 22, 2021. Accepted for publication Feb 14, 2022.

doi: 10.21037/atm-22-274

View this article at: <https://dx.doi.org/10.21037/atm-22-274>

Introduction

Since the first reported case of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in China at the end of 2019, coronavirus disease 2019 (COVID-19) has rapidly become a pandemic causing unprecedented health and economic challenges (1). The majority of

COVID-19 patients are asymptomatic or have mild symptoms. However, some individuals develop severe complications leading to death. According to the World Health Organization (WHO) (2), there have been more than four million deaths from COVID-19 across 223 countries as of September 1, 2021. Studies have shown that COVID-19 patients who are elderly and have pre-

existing comorbidities, such as obesity, hypertension, diabetes, chronic renal insufficiency, or coronary heart disease, have worse clinical outcomes (3).

Frailty is a multidimensional geriatric syndrome characterized by impaired stress tolerance due to a decline in physiological reserves and functioning of different organs and is associated with poor health outcomes, including mortality, admissions to nursing homes, and falls. Frailty is a common syndrome in older individuals (4), with a prevalence rate of 41.3% in those aged over 65 and 65.2% in those aged over 80 (5). Nevertheless, aging and frailty are not synonymous (6). Compared with aging, a diagnosis of frailty arises from a comprehensive geriatric assessment that reflects the overall health status of individuals. Several recent studies have explored the association between frailty and the clinical outcomes of patients infected by SARS-CoV-2, but the results remain inconclusive (7-10). Frailty was associated with being female and having comorbidities in patients with COVID-19, and the level of white blood cell count, D-dimer, and C-reactive protein were higher in frail patients (11,12). Most studies have demonstrated that frailty in hospitalized COVID-19 patients is a risk factor for short-term outcomes, such as in-hospital death and intensive care unit (ICU) admission (7,8). However, some studies have found that frailty is not correlated with mortality (9,10). Therefore, it is necessary to synthesize all available evidence to understand whether frailty is a predictor of mortality so as to inform evidence-based decisions in clinical practice. This study provides a systematic review and meta-analysis to compare the mortality risk of frail *vs.* non-frail hospitalized patients with COVID-19. We present the following article in accordance with the MOOSE reporting checklist (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-274/rc>).

Methods

Search strategy

MEDLINE, Embase, and the Cochrane Library databases were comprehensively searched for relevant studies up to September 11, 2021. The search algorithms consisted of Medical Subject Headings (MeSH) and keywords as follows: “frail” or “frailty” (MeSH) and (“COVID-19” OR “Coronavirus Infection Disease 2019” OR “SARS-CoV-2” OR “Coronavirus Infection”) and (“mortality” or “deceased” or “death”). Additionally, we reviewed the reference lists of eligible articles to identify potential gray literature. The

complete search strategy is available in [Tables S1,S2](#).

Eligibility criteria

Two reviewers (PZ and MH) blindly and independently selected the eligible studies if they met the following inclusion criteria: (I) studies that reported the associations between frailty and mortality in hospitalized patients with COVID-19; (II) studies that reported short-term mortality outcomes, *i.e.*, in-hospital mortality, 30-day mortality, or 60-day mortality. Studies were excluded if they met the following criteria: (I) review articles or letters; (II) studies with less than 20 participants; (III) studies with insufficient data; (IV) non-English language studies. Any differences of opinion between the reviewers regarding an article’s qualification for inclusion were resolved by discussion.

Data collection

The data were extracted by two reviewers (PZ and MH) independently, and any differences of opinion were resolved by discussion. Data extracted from the articles included the following: name of the first author, study year, institution(s), number of participating centers, study design, enrollment period, the number of patients, age, gender, frailty assessment tools, and prevalence of frailty. The primary outcome was short-term mortality, defined as a composite outcome of in-hospital mortality, 30-day mortality, or 60-day mortality. The secondary outcome was ICU admission.

Quality assessment

The methodological quality of the included studies was assessed using the Newcastle-Ottawa scale (NOS) (13). Two reviewers (PZ and MH) independently assessed the methodological quality of the included studies. The results of the quality assessment are reported in [Table S3](#).

Statistical analysis

The effect size (ES) (the exact number of frail patients and mortality) was used to assess the association between frailty and mortality risk in patients with COVID-19. The pooled odds ratio (OR) with 95% confidence intervals (CIs) were calculated using a random-effects model to compare mortality and ICU admission rates between frail and non-frail patients from the data of the included




Figure 1 PRISMA flowchart of the study selection process. PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

studies. Heterogeneity between studies was tested using the I^2 statistic. The potential publication bias was evaluated quantitatively with Egger's test and assessed qualitatively with funnel plots. Sensitivity analysis was carried out to assess the heterogeneity and robustness of pooled outcomes. We performed subgroup analyses based on the frailty assessment tools. We also demonstrated the outcome estimates for older patients (age ≥ 65 years). All statistical analyses were carried out using STATA statistical software, version 15.1 (StataCorp., TX, USA). In all analyses, a P value less than 0.05 was considered statistically significant.

Results

Study selection

After removing duplicate publications, a total of 371 articles were identified from the electronic database searches. After screening the titles and abstracts, 315 irrelevant articles were removed. Full-text articles were carefully reviewed,

and an additional 35 studies were excluded. Eventually, 21 studies meeting the inclusion and exclusion criteria were included in this meta-analysis (7-10,14-29). A PRISMA flow chart of the article selection is shown in *Figure 1*.

Study characteristics

Twenty-one included studies, composed of 26,652 hospitalized patients with COVID-19, reported the association between frailty and mortality risk. The principal characteristics of the included studies are reported in *Table 1*. Over half of the studies included older adults (age ≥ 65 years), and 50.1% (13,187/26,303) of patients were female. Six frailty assessment tools were used, including the Clinical Frailty Score (CFS) (n=16) (7-9,11,16-25), the Hospital Frailty Risk Score (HFRS) (n=1) (26), the Geriatric Nutritional Risk Index (GRNI) (n=1) (27), the modified Frailty Index (mFI) (n=1) (28), the Multidimensional Prognostic Index (MPI) (n=1) (29), and the Frail Non-Disabled Survey (FIND) (n=1) (10). The prevalence of

Table 1 Characteristics of the studies included in the meta-analysis

Study	Design	Region	Sample	Consecutive	Female	Age, year ^a	Age limitation, year-old	Prevalence of frailty	Frailty criteria	Outcome measures
Poco 2021	RCS	Brazil	711	1	306	66±11	≥50	25.5% (181/711)	CFS	In-hospital mortality
Owen 2021	RCS	UK	206	NM	NM	NM	≥65	53.9% (110/206)	CFS	30-day mortality
Mendes 2020	RCS	Switzerland	235	NM	133	67 [61–74]	≥65	78.7% (185/235)	CFS	In-hospital mortality
Chinnadurai 2020	RCS	UK	215	1	85	[36–88]	NM	51.2% (110/215)	CFS	In-hospital mortality
Aw 2020	RCS	UK	664	1	305	NM	≥65	70.8% (470/664)	CFS	In-hospital mortality
Labenz 2020	RCS	Germany	42	NM	13	63 [55–71]	NM	14.2% (6/42)	CFS	In-hospital mortality
Hewitt 2020	RCS	UK and Italy	1,559	NM	661	69 [35–85]	NM	51.3% (799/1,559)	CFS	In-hospital mortality
De Smet 2020	RCS	Belgium	81	1	48	NM	≥65	79.0% (64/81)	CFS	In-hospital mortality
Aliberti 2021	RCS	Brazil	1,830	1	769	68.7±10.1	≥50	27.0% (494/1,830)	CFS	30-day mortality
Andrés-Esteban 2021	RCS	Spain	254	NM	96	62±7.2	≥65	17.3% (44/254)	CFS	In-hospital mortality
Dres 2021	PCS	France, Switzerland, and Belgium	1,085	NM	326	68.9±10.9	≥70	9.1% (99/1,085)	CFS	30-day mortality
Maniero 2021	RCS	UK	124	1	61	73 [48–88]	≥65	77.8% (89/124)	CFS	In-hospital mortality
Lozano-Montoya 2021	RCS	Spain	300	1	188	70.3±9.2	≥75	72.0% (216/300)	CFS	In-hospital mortality
Ramos-Rincon 2021	RCS	Spain	290	NM	119	66±13	≥18	60.1% (54/90)	CFS	In-hospital mortality
Burns 2020	RCS	UK	28	NM	13	63.5±10.4	NM	60.7% (17/28)	CFS	In-hospital mortality
Tehrani 2021	RCS	Sweden	143	0	NM	68.2±11.4	NM	49.7% (71/143)	CFS	60-day mortality
Kundi 2020	RCS	Turkey	18,234	NM	9736	74.1±7.4	≥65	68.1% (12420/18,234)	HFRS	In-hospital mortality
Recinella 2020	RCS	Bologna	109	1	55	NM	≥65	72.5% (79/109)	GRNI	In-hospital mortality
Fumagalli 2021	RCS	Italy	221	1	87	82 [78–86]	≥75	72.5% (79/221)	mFI	In-hospital mortality
Pilotto 2021	PCS	Italy	227	1	134	80.5 [65–99]	≥65	73.6% (167/227)	MPI	In-hospital mortality
Steinmeyer 2020	RCS	France	94	NM	52	85.5±7.5	≥65	87.2% (82/94)	FIND	In-hospital mortality

^a, data are expressed as mean ± standard deviation or median [range] or mean or [range]. RCS, retrospective cohort study; PCS, prospective cohort study; CFS, Clinical Frailty Score; NM, not mentioned; HFRS, Hospital Frailty Risk Score; GNRI, Geriatric Nutritional Risk Index; MFI, modified Frailty Index; MPI, Multidimensional Prognostic Index; FIND, Frail Non-Disabled Survey.




Figure 2 Pooled proportion of frailty in hospitalized patients with COVID-19. COVID-19, coronavirus disease 2019; ES, effect size; CI, confidence interval; CFS, Clinical Frailty Score.

frailty ranged from 9.1–87.2%, with a summary estimate rate of 51.4% (95% CI: 39.9–62.9%; $I^2=99.49\%$; *Figure 2*). The result of Egger's test indicated no apparent publication bias ($t=1.89$; $P=0.07$; *Figure S1*). The sensitive analysis showed that there was no substantial change on the overall pooled OR (*Figure S2*).

Association between frailty and short-term mortality

The short-term mortality outcome measures were in-hospital mortality in 17 studies, 30-day mortality in three studies, and 60-day mortality in one study. We performed a subgroup analysis based on the frailty assessment tools used. In the CFS group, frail patients experienced a higher rate of short-term mortality than non-frail patients (OR =3.0; 95% CI: 2.3–3.9; $I^2=72.7\%$; $P<0.001$; *Figure 3*). In the other tools group (HFRS, GRNI, mFI, MPI, and FIND), frail patients had a significantly higher short-term mortality risk than non-frail patients (OR =2.4; 95% CI: 1.4–4.1; $I^2=63.8\%$;

$P=0.001$). Overall, a higher rate of short-term mortality was observed for frail patients compared with non-frail patients (OR =2.8; 95% CI: 2.3–3.5; $I^2=78.0\%$; $P<0.001$).

The subgroup analysis of older adults (age ≥ 65 years) revealed that frail patients with COVID-19 in the CFS group had a 2.34-fold higher risk of short-term mortality compared with non-frail patients (OR =2.3; 95% CI: 1.7–3.3; $I^2=62.5\%$; $P<0.001$; *Figure 4*). In the other tools group, frail COVID-19 patients experienced a higher rate of short-term mortality than non-frail patients (OR =2.4; 95% CI: 1.4–4.1; $I^2=63.8\%$; $P=0.001$). The overall results revealed that frail patients over 65 years old had significantly increased short-term mortality compared with non-frail patients (OR =2.3; 95% CI: 1.8–2.9; $I^2=60.9\%$; $P<0.001$).

Association between frailty and ICU admission

Five studies investigated the association between frailty and ICU admission. Three studies used the CFS, one used the




Figure 3 Forest plots of outcomes for short-term mortality in frail vs. non-frail COVID-19 patients. COVID-19, coronavirus disease 2019; OR, odds ratio; CI, confidence interval; CFS, Clinical Frailty Score.

HFRS, and one used the MPI. In the studies using the CFS, the results showed that frail patients had a lower risk of ICU admission than non-frail patients (OR =0.2; 95% CI: 0.1–0.5; $I^2=0.0\%$; $P<0.001$; **Figure S3**). In contrast, in the other tools group, frail COVID-19 patients demonstrated a higher risk of ICU admission than non-frail patients (OR =2.0; 95% CI: 1.8–2.1; $I^2=0.0\%$; $P<0.001$). The overall results revealed no significant difference in ICU admission rates between frail and non-frail patients (OR =0.5; 95% CI: 0.1–2.1; $I^2=89.4\%$; $P=0.37$).

Discussion

This systematic review and meta-analysis included 21 studies with 26,652 hospitalized COVID-19 patients and aimed to assess the influence of frailty on clinical outcomes. We pooled the prevalence of frailty and found that over half of COVID-19 patients were frail. CFS was the most commonly used frailty assessment in the studies

of hospitalized COVID-19 patients. Frail patients had a higher rate of short-term mortality than non-frail patients, which was also evident in older adults. However, the results from studies investigating ICU admissions that used CFS and other frailty assessment tools were conflicting.

Frailty is a complex age-related clinical state or syndrome characterized by increasing susceptibility to stress, with a functional decline across multiple physiological systems (30,31). COVID-19 caused by SARS-CoV-2 has created a global pandemic and challenged healthcare systems worldwide (1,32). COVID-19 is a systemic disease that can rapidly cause hypoxemic respiratory symptoms and even acute respiratory distress syndrome (ARDS) (33). It is worth mentioning that, compared with computed tomography, lung ultrasound is a non-invasive bedside and low cost technique with high diagnostic accuracy. Lung ultrasound might play an important role in early diagnosis, monitoring of COVID-19 pneumonia, especially in the COVID-19 patients with frailty, which might reduce the pressure




Figure 4 Forest plots of outcomes for short-term mortality in frail vs. non-frail COVID-19 patients ≥ 65 years old. COVID-19, coronavirus disease 2019; OR, odds ratio; CI, confidence interval; CFS, Clinical Frailty Score.

on the patients' body and handling time, and decrease the risk of SARS-CoV-2 transmission (34,35). Besides, based on the blood dissemination of hemophagocytic lymphohistiocytosis, COVID-19 can have secondary effects on extrapulmonary organs, including the nervous system, heart, intestine, kidney, and liver, resulting in multiple organ system dysfunction syndrome (36-38). As a consequence, patients infected with COVID-19 have an increased vulnerability to frailty. Therefore, to improve the prognosis of patients with COVID-19, identifying whether they are frail and need more intensive care is of utmost importance.

There are several well-validated frailty assessment tools used for screening and quantifying frailty status. Many studies across different diseases have revealed that frailty is associated with a variety of poor health outcomes, such as falls, dementia, mortality, hospitalization, and poor quality of life (39-41). As such, it is important to evaluate frailty status comprehensively to optimize outcomes. In this study,

we found that frailty was a predictor of all-cause mortality in hospitalized COVID-19 patients, similar to the results of previous systematic reviews. There may be several reasons for this result. First, compared with non-frail patients, patients in a frail condition are characterized by reduced physiological reserves, chronic malnutrition, and a loss of resistance to stressors due to accumulated deficits, all of which increase the risk of a poor outcome for patients infected by SARS-CoV-2 (42,43). Second, frail patients tend to develop a pro-inflammatory status with an elevated level of inflammatory markers in cells and tissues. A leading cause of severe COVID-19 is the cytokine storm syndrome, associated with various cytokines such as IFN α , IL-6, and IL-1, which are elevated in frail patients (36,37). This can lead to immunologic derangement and increase the risk of extrapulmonary organ dysfunction. Third, most frail patients are older adults, and high COVID-19-associated mortality occurs among the older population. This may be due to the high prevalence of comorbidities, such as

hypertension, diabetes mellitus, chronic heart disease, and dementia in the geriatric population, which have been identified as risk factors for mortality in patients ≥ 65 years old with COVID-19 (44).

Based on the pooled analysis, we found contradictory results in the association between frailty and ICU admission. According to the results of studies using CFS, frail patients had a lower risk of ICU admission than non-frail patients (OR =0.2; 95% CI: 0.1–0.5). However, among the studies using other tools, frail patients experienced a higher risk of ICU admission than non-frail patients (OR =2.0; 95% CI: 1.8–2.1). Dumitrascu *et al.* (45) reported that eight studies showed an association between frailty as assessed by the CFS and ICU admission (OR =0.24; 95% CI: 0.08–0.71). This discrepancy in results might be caused by the sampling error between studies using the CFS *vs.* other assessment tools and the shortage of medical resources in the early stages of the COVID-19 pandemic. The National Institute for Health and Care Excellence (NICE) guidelines (46) suggest that the CFS should be used to assess whether COVID-19 patients need more intensive care. Based on previous research and the current meta-analysis, further studies are required to confirm this suggestion.

Several limitations of this meta-analysis need to be considered. Firstly, during the study selection, a number of studies reporting the association between frailty and mortality in COVID-19 hospitalized patients were excluded because of non-standard cut-off values for frailty groupings. Furthermore, the effect values of these studies were reported as either ORs or risk ratios. To maximize the pooling of the overall estimates and avoid inflated risk estimates by converting ORs into risk ratios, several studies were excluded where the data could not be extracted. Moreover, most studies were retrospective in design. Therefore, the pooled outcomes were inevitably affected by selection bias. Secondly, because most studies used the CFS as the frailty assessment tool, we conducted a pooled analysis combining the results from other tools used. The data were too limited to allow for further subgroup analysis, which may have decreased the reliability of the results in our meta-analysis.

Conclusions

The results of this systematic review and meta-analysis indicated that frail patients suffered from a higher risk of all-cause mortality than non-frail hospitalized COVID-19 patients, and this result was also found in the older adult

group. Therefore, an assessment of frailty could assist clinicians in identifying hospitalized COVID-19 patients who require more intensive treatment and care.

Acknowledgments

Funding: This study was supported by the Science and Technology Department of Sichuan Province (Grant No. 2019JY0346) and the National Clinical Research Center for Geriatrics, West China Hospital, Sichuan University (Grant No. Z2018A12).

Footnote

Reporting Checklist: The authors have completed the MOOSE reporting checklist. Available at <https://atm.amegroups.com/article/view/10.21037/atm-22-274/rc>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-274/coif>). The authors have no conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Jefferson T, Spencer EA, Brassey J, et al. Transmission of Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) from pre and asymptomatic infected individuals: a systematic review. *Clin Microbiol Infect* 2021. [Epub ahead of print]. doi: 10.1016/j.cmi.2021.10.015.
2. COVID-19 coronavirus pandemic. Available online: <https://www.worldometers.info/coronavirus/>

3. Gacche RN, Gacche RA, Chen J, et al. Predictors of morbidity and mortality in COVID-19. *Eur Rev Med Pharmacol Sci* 2021;25:1684-707.
4. Mitnitski A, Rockwood K. The rate of aging: the rate of deficit accumulation does not change over the adult life span. *Biogerontology* 2016;17:199-204.
5. Carneiro JA, Cardoso RR, Durães MS, et al. Frailty in the elderly: prevalence and associated factors. *Rev Bras Enferm* 2017;70:747-52.
6. Rockwood K, Song X, MacKnight C, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ* 2005;173:489-95.
7. Hewitt J, Carter B, Vilches-Moraga A, et al. The effect of frailty on survival in patients with COVID-19 (COPE): a multicentre, European, observational cohort study. *Lancet Public Health* 2020;5:e444-51.
8. Ramos-Rincon JM, Moreno-Perez O, Pinargote-Celorio H, et al. Clinical Frailty Score vs Hospital Frailty Risk Score for predicting mortality and other adverse outcome in hospitalised patients with COVID-19: Spanish case series. *Int J Clin Pract* 2021;75:e14599.
9. Owen RK, Conroy SP, Taub N, et al. Comparing associations between frailty and mortality in hospitalised older adults with or without COVID-19 infection: a retrospective observational study using electronic health records. *Age Ageing* 2021;50:307-16.
10. Steinmeyer Z, Vienne-Noyes S, Bernard M, et al. Acute Care of Older Patients with COVID-19: Clinical Characteristics and Outcomes. *Geriatrics (Basel)* 2020;5:65.
11. Poco PCE, Aliberti MJR, Dias MB, et al. Divergent: Age, Frailty, and Atypical Presentations of COVID-19 in Hospitalized Patients. *J Gerontol A Biol Sci Med Sci* 2021;76:e46-51.
12. Ma Y, Hou L, Yang X, et al. The association between frailty and severe disease among COVID-19 patients aged over 60 years in China: a prospective cohort study. *BMC Med* 2020;18:274.
13. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010;25:603-5.
14. Mendes A, Serratrice C, Herrmann FR, et al. Predictors of In-Hospital Mortality in Older Patients With COVID-19: The COVIDAge Study. *J Am Med Dir Assoc* 2020;21:1546-54.e3.
15. Chinnadurai R, Ogedengbe O, Agarwal P, et al. Older age and frailty are the chief predictors of mortality in COVID-19 patients admitted to an acute medical unit in a secondary care setting- a cohort study. *BMC Geriatr* 2020;20:409.
16. Aw D, Woodrow L, Ogliari G, et al. Association of frailty with mortality in older inpatients with Covid-19: a cohort study. *Age Ageing* 2020;49:915-22.
17. Labenz C, Kremer WM, Schattenberg JM, et al. Clinical Frailty Scale for risk stratification in patients with SARS-CoV-2 infection. *J Investig Med* 2020;68:1199-202.
18. De Smet R, Mellaerts B, Vandewinckele H, et al. Frailty and Mortality in Hospitalized Older Adults With COVID-19: Retrospective Observational Study. *J Am Med Dir Assoc* 2020;21:928-32.e1.
19. Andrés-Esteban EM, Quintana-Díaz M, Ramírez-Cervantes KL, et al. Outcomes of hospitalized patients with COVID-19 according to level of frailty. *PeerJ* 2021;9:e11260.
20. Aliberti MJR, Szejf C, Avelino-Silva VI, et al. COVID-19 is not over and age is not enough: Using frailty for prognostication in hospitalized patients. *J Am Geriatr Soc* 2021;69:1116-27.
21. Dres M, Hajage D, Lebbah S, et al. Characteristics, management, and prognosis of elderly patients with COVID-19 admitted in the ICU during the first wave: insights from the COVID-ICU study : Prognosis of COVID-19 elderly critically ill patients in the ICU. *Ann Intensive Care* 2021;11:77.
22. Maniero C, Patel D, Pavithran A, et al. A retrospective cohort study of risk factors and outcomes in older patients admitted to an inner-city geriatric unit in London during first peak of COVID-19 pandemic. *Ir J Med Sci* 2021. [Epub ahead of print]. doi: 10.1007/s11845-021-02679-z.
23. Lozano-Montoya I, Quezada-Feijoo M, Jaramillo-Hidalgo J, et al. Mortality risk factors in a Spanish cohort of oldest-old patients hospitalized with COVID-19 in an acute geriatric unit: the OCTA-COVID study. *Eur Geriatr Med* 2021;12:1169-80.
24. Burns GP, Lane ND, Tedd HM, et al. Improved survival following ward-based non-invasive pressure support for severe hypoxia in a cohort of frail patients with COVID-19: retrospective analysis from a UK teaching hospital. *BMJ Open Respir Res* 2020;7:e000621.
25. Tehrani S, Killander A, Åstrand P, et al. Risk factors for death in adult COVID-19 patients: Frailty predicts fatal outcome in older patients. *Int J Infect Dis* 2021;102:415-21.
26. Kundi H, Çetin EHÖ, Canpolat U, et al. The role of Frailty on Adverse Outcomes Among Older Patients with COVID-19. *J Infect* 2020;81:944-51.
27. Recinella G, Marasco G, Serafini G, et al. Prognostic role

- of nutritional status in elderly patients hospitalized for COVID-19: a monocentric study. *Aging Clin Exp Res* 2020;32:2695-701.
28. Fumagalli C, Ungar A, Rozzini R, et al. Predicting Mortality Risk in Older Hospitalized Persons With COVID-19: A Comparison of the COVID-19 Mortality Risk Score with Frailty and Disability. *J Am Med Dir Assoc* 2021;22:1588-92.e1.
 29. Pilotto A, Azzini M, Cella A, et al. The multidimensional prognostic index (MPI) for the prognostic stratification of older inpatients with COVID-19: A multicenter prospective observational cohort study. *Arch Gerontol Geriatr* 2021;95:104415.
 30. Clegg A, Young J, Iliffe S, et al. Frailty in elderly people. *Lancet* 2013;381:752-62.
 31. Kojima G. Frailty as a predictor of hospitalisation among community-dwelling older people: a systematic review and meta-analysis. *J Epidemiol Community Health* 2016;70:722-9.
 32. Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun* 2020;109:102433.
 33. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020;8:475-81.
 34. Allinovi M, Parise A, Giacalone M, et al. Lung Ultrasound May Support Diagnosis and Monitoring of COVID-19 Pneumonia. *Ultrasound Med Biol* 2020;46:2908-17.
 35. Smith MJ, Hayward SA, Innes SM, et al. Point-of-care lung ultrasound in patients with COVID-19 - a narrative review. *Anaesthesia* 2020;75:1096-104.
 36. Mehta P, McAuley DF, Brown M, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;395:1033-4.
 37. Kim JS, Lee JY, Yang JW, et al. Immunopathogenesis and treatment of cytokine storm in COVID-19. *Theranostics* 2021;11:316-29.
 38. Ferrucci L, Fabbri E. Inflammageing: chronic inflammation in ageing, cardiovascular disease, and frailty. *Nat Rev Cardiol* 2018;15:505-22.
 39. Qiu Y, Li G, Wang X, et al. Prevalence of cognitive frailty among community-dwelling older adults: A systematic review and meta-analysis. *Int J Nurs Stud* 2022;125:104112.
 40. Brown PJ, Ciarleglio A, Roose SP, et al. Frailty and Depression in Late Life: A High-Risk Comorbidity with Distinctive Clinical Presentation and Poor Antidepressant Response. *J Gerontol A Biol Sci Med Sci* 2021. [Epub ahead of print]. doi: 10.1093/gerona/glab338.
 41. Gracie TJ, Caufield-Noll C, Wang NY, et al. The Association of Preoperative Frailty and Postoperative Delirium: A Meta-analysis. *Anesth Analg* 2021;133:314-23.
 42. Moraes MB, Avgerinou C, Fukushima FB, et al. Nutritional interventions for the management of frailty in older adults: systematic review and meta-analysis of randomized clinical trials. *Nutr Rev* 2021;79:889-913.
 43. El Assar M, Angulo J, Carnicero JA, et al. Association between telomere length, frailty and death in older adults. *Geroscience* 2021;43:1015-27.
 44. Kojima G, Iliffe S, Walters K. Frailty index as a predictor of mortality: a systematic review and meta-analysis. *Age Ageing* 2018;47:193-200.
 45. Dumitrascu F, Branje KE, Hladkovicz ES, et al. Association of frailty with outcomes in individuals with COVID-19: A living review and meta-analysis. *J Am Geriatr Soc* 2021;69:2419-29.
 46. National Institute for Health and Care Excellence. COVID-19 Rapid Guideline: Critical Care in Adults. London: National Institute for Health and Care Excellence, 2020. Available online: <https://www.nice.org.uk/guidance/ng159>
- (English Language Editor: D. Fitzgerald)

Cite this article as: Zou Y, Han M, Wang J, Zhao J, Gan H, Yang Y. Predictive value of frailty in the mortality of hospitalized patients with COVID-19: a systematic review and meta-analysis. *Ann Transl Med* 2022;10(4):166. doi: 10.21037/atm-22-274