

Erosive Hand Osteoarthritis is Associated with Subclinical Atherosclerosis and Endothelial Dysfunction

Athanasios Koutroumpas¹, Athanasios Giannoukas², Elias Zintzaras³,
Ekaterini Exarchou¹, Aris Baliakos¹, Konstantinos Makaritsis¹, Lazaros I. Sakkas^{1, 4}

¹Department of Rheumatology, Faculty of Medicine, School of Health Sciences, University of Thessaly, Greece; ²Department of Vascular Surgery, Faculty of Medicine, School of Health Sciences, University of Thessaly, Greece; ³Department of Biomathematics, Faculty of Medicine, School of Health Sciences, University of Thessaly, Greece; ⁴Center for Molecular Medicine, Old Dominion University, Norfolk, VA, USA

ABSTRACT

Chronic inflammatory disorders have been associated with accelerated atherosclerosis and increased cardiovascular (CV) risk. Recent evidence suggests that erosive hand osteoarthritis (EOA) has considerable inflammation; therefore, we examined the presence of subclinical atherosclerosis and endothelial dysfunction in EOA. Twenty-four patients with EOA and 24 age- and sex-matched healthy individuals without clinical OA were included in the study. No subject had a history of CV disease. Intima-media thickness (IMT) and atheromatous plaques in the common carotid and common femoral arteries were measured by Doppler ultrasonography. The endothelium-dependent, flow-mediated dilatation (FMD) and endothelium-independent, sublingual glyceryl trinitrate (NTG)-induced dilatation (NMD) of the brachial artery were assessed. The EOA patients had significantly elevated systolic and diastolic blood pressure ($p < 0.001$ for both). The 10-year risk of general CV disease, as predicted with the Framingham Risk Score, was similar in patients and controls ($p = 0.18$). IMT of both common carotid and common femoral artery were increased in EOA ($p = 0.01$ and $p < 0.01$, respectively), but the frequency of atherosclerotic plaques was not increased. There was no difference in FMD and NMD between the two groups, but the difference between FMD and NMD was increased in EOA. In conclusion, this small controlled study showed an association between EOA and subclinical atherosclerosis that cannot be fully attributed to traditional CV risk factors, as assessed by the Framingham score. These results suggest that chronic, low-grade inflammation is implicated in atherosclerosis in EOA. (*Int J Biomed Sci* 2013; 9 (4): 217-223)

Keywords: atherosclerosis; Doppler ultrasonography; dilatation; endothelium; erosive osteoarthritis; Framingham risk score

Corresponding author: Prof. L. I. Sakkas, Department of Rheumatology, Faculty of Medicine, University of Thessaly, 41110, Biopolis, Larissa, Greece. Tel: +302413502813; Fax: +302413501016; E-mail: lsakkas@med.uth.gr.

Note: Financial support from ELKE, University of Thessaly.

Received May 30, 2013; **Accepted** June 20, 2013

Copyright: © 2013 Athanasios Koutroumpas et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.5/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

INTRODUCTION

Erosive hand osteoarthritis (EOA) is a subset of hand osteoarthritis (HOA) which is defined by erosions in interphalangeal (IP) joints on plain radiographs and occurs primarily in postmenopausal women (1). Traditionally, osteoarthritis (OA) has been considered a degenerative disorder of the aging population. However, synovial inflammation has been frequently described in early and advanced knee or hip OA (2, 3) and inflammation is increasingly recognized as an important factor in the pathophysiology of this disease (4, 5). Newer imaging modalities, such as ultrasound and MRI, revealed frequent local inflammation in HOA (6, 7). The inflammatory component is more evident in EOA (1, 8, 9). Systemic low-grade inflammation, as exemplified by elevated serum C-reactive protein, has been reported in EOA (10).

A growing body of evidence supports that endothelial dysfunction and atherosclerosis are significantly more prevalent in patients with chronic inflammatory disorders, compared to age-matched controls (11). This has been convincingly demonstrated in rheumatoid arthritis, a prototype of chronic inflammatory diseases, which is associated with increased cardiovascular (CV) risk (12). Therefore, we sought to investigate the occurrence of atherosclerosis and endothelial dysfunction in patients with EOA compared to normal controls.

METHODS

Selection and description of participants

Patients participating in the study were followed at the Rheumatology outpatient clinic of the University General Hospital of Larissa. All fulfilled the 1990 American College of Rheumatology criteria for the classification of HOA (13). In addition, they exhibited IP joint central area erosions in the form of “gull-wing” or “saw-teeth” on plain radiography and classified as having EOA. A second group of age- and sex-matched individuals served as healthy controls, with no hand pain or clinical signs of hand osteoarthritis, such as Heberden’s and Bouchard’s nodes, or 1st carpometacarpal joint squaring. In addition, they had no history or clinical signs and symptoms of osteoarthritis in other joints. All participants, in both groups, had no history of CV disease, such as an acute coronary syndrome, stroke or peripheral arterial disease. History of diabetes mellitus (DM), hypertension (HT), hyperlipidemia (HL) and smoking, and any relevant medication were recorded.

The study protocol was approved by the Institutional Scientific and Ethical Committee, all participants gave a written informed consent, and the study was conducted according to good clinical practice.

Technical information

Ultrasonographic (US) examination was performed by a single trained and senior vascular surgeon (GA). Common carotid and common femoral artery intima-media thickness (IMT), were measured using high resolution B-mode ultrasonography, using a 12 L, 5 Hz linear transducer (GE Logic equipment), according to recently published recommendations (14). Measurements were conducted twice for both the right and the left artery, and the mean values were recorded. IMT was used as a marker of preclinical atherosclerosis, as common carotid artery IMT is an early atherosclerotic index and a strong predictor of cardiovascular end-points, such as stroke and coronary events (15, 16).

The presence and type of atheromatous plaques was recorded for both sites. A plaque was defined as a focal structure encroaching into the lumen, of at least 0.5 mm, or 50% of the surrounding IMT. Alternatively, a plaque should demonstrate thickness >1.5 mm, measured from the media-adventitia to the media-lumen interface (14). Carotid plaques were characterized as type 1-5, depending on the echogenicity and homogeneity of the plaque (17). This characterization has clinical implications, as type 1 and 2 plaques carry a high risk for complications, type 3 and 4 plaques carry a low risk, and type 5 calcified plaques carry an undetermined risk (18).

Two definitions were used for subclinical atherosclerosis: plaque formation and a common carotid artery IMT >1 mm. The cut-off of 1 mm was used because this level is associated with coronary artery disease and increased CV risk (18). As common carotid artery IMT is a better surrogate for CV risk than common femoral artery IMT, risk analyses were performed using carotid artery measurements.

An US “biopsy score” (USBS) has been assessed in all subjects (19). This is a non-invasive procedure for the detection and grading of early carotid and femoral artery atherosclerosis. Subjects were graded into A-F classes, according to vessel wall characteristics: (A) normal (intima media and adventitia clearly separated), 0 points; (B) interface disruption, 2 points; (C) intima-media granulation, 4 points; (D) plaque without hemodynamic disturbance, 6 points; (E) asymptomatic plaque causing stenosis, 8 points; (F) plaque causing stenosis and symptoms, 10 points. The

points for the two carotid and two femoral arteries were added, giving a numerical score ranging from 0 to 40. Classes D-E are associated with higher risk of ischemic cardiovascular or cerebrovascular events and higher risk of mortality than classes A-C (20).

Endothelial function was assessed by measuring flow-mediated endothelium-dependent dilatation (FMD). The end-diastolic diameter of both brachial arteries was measured at rest and within 15 sec after release of a sphygmomanometer cuff inflated at a pressure of at least 50 mmHg above the subject's systolic blood pressure (BP) for 3 minutes (21). Similar measurements were conducted 5 minutes after administration of 0.4 mg of glyceryl trinitrate (NTG) sublingually for the assessment of endothelium-independent dilatation (NMD) (21). Vasodilatation was expressed as the difference between the two measurements. FMD reflects endothelial dysfunction and is mediated by the release of nitric oxide from arterial endothelial cells. It is predictive of CV events, and is related to both the presence and extent of coronary atherosclerosis (22).

As the vasodilator potential is not the same across individuals, NMD can be used as a means of measuring the maximum obtainable vascular wall reaction to a vasodilatory stimulus (23). Therefore, we calculated the difference between NMD and FMD and compared the means between the two groups of individuals. In this way, FMD is opposed not just to the baseline brachial artery diameter, but also to the maximum dilatation obtained by administration of NTG. Apart from baseline diameter, dilatation is influenced by hypertension and vasoactive anti-hypertensive drugs. All differences were expressed as numerical values (in mm), rather than percentage of change from baseline value. Expressing the vasodilator effect as the percentage of the baseline value is more suitable for measurements in the same individual before and after an intervention (23).

All assessments were conducted in a quiet, dark room and subjects were examined in the supine position. Patients remained at the supine position for at least 10 minutes before the examination.

Serum total cholesterol, high density lipoprotein (HDL), low density lipoprotein (LDL) and triglycerides were measured by standard methods. The cut-off values for total cholesterol, LDL and HDL are in accordance with the Third Report of the National Cholesterol Education Program (NCEP) (24).

The Framingham 10-year risk of CVD was estimated according to published method (25). This algorithm incorporates age, sex, systolic blood pressure, smoking, blood

pressure treatment and LDL and HDL levels and provides an estimate of the 10-year risk of clinical CVD. The Framingham risk score provides the risk that is attributed to traditional CVD risk factors.

Statistics

Qualitative variables were compared with the Fischer's exact test and the chi-square test, as appropriate. Comparisons between means were carried out with the Student's t-test and the Mann-Whitney U, as appropriate. Statistical analysis has been performed using the SPSS, version 13, software (SPSS Inc. Chicago, USA).

For each group the risk of preclinical atherosclerosis was separately assessed with regard to age, sex, history of hypertension, hyperlipidaemia, diabetes and smoking, total cholesterol (whether >240 or not), LDL (whether >160 or not), HDL (whether <50 or not), systolic blood pressure (whether > or <135 mmHg, which is the upper normal limit, according to international guidelines) and diastolic blood pressure (whether > or <90 mmHg), and regular use of vasoactive and hypolipidaemic agents (β -blockers, calcium channel blockers, angiotensin converting enzyme inhibitors, angiotensin II receptor antagonists, diuretics and statins), with the Fisher's exact test and the Mann-Whitney U test.

RESULTS

Demographics and lipids

The two groups were comparable for all demographic variables. No significant difference was found in the frequency of diabetes mellitus, hypertension, hyperlipidaemia and smoking status between EOA patients and controls. Similarly, no difference was found in the rate of prescription of vasoactive and hypolipidemic medications between the two groups (Table 1). Fourteen of the EOA patients (58%) were receiving hydroxychloroquine (HCQ).

Serum lipid concentrations were available for 23 of the 24 subjects in the EOA group and all subjects in the control group. There was a non-significant trend towards increased serum lipid concentrations for total cholesterol, LDL and triglycerides in the EOA group (Table 2). EOA patients had significantly elevated systolic (160 ± 32.9 mmHg [mean \pm SD]) and diastolic (99.9 ± 20.4 mmHg) blood pressure (BP) at the time of examination, compared to the controls (systolic BP = 143.4 ± 32.1 mmHg, diastolic BP = 83.42 ± 11.9 mmHg) ($p < 0.001$ for both) (Table 2).

The Framingham 10-year risk of CVD was calculated for 23 of the 24 participants in the EOA group, as 1 patient

lacked serum lipid determinations. Patients with EOA had a non-significant trend towards an increased Framingham risk score (mean=23.9% 10-year risk of CVD), compared to the control group (mean=17.3% 10-year risk of CVD, $p=0.18$) (Table 3).

Ultrasonography

EOA was associated with a marginally significant increase in the risk of IMT>1mm (OR=3.33, 95% CI 1.02-10.9, $p=0.043$) (Table 4). Common carotid artery IMT was significantly elevated in the EOA group, compared to the control group ($p=0.012$). Similarly, common femoral artery IMT was elevated in the EOA compared to the control group ($p<0.01$) (Table 3). Fifteen EOA patients (62.5%) had carotid and/or femoral artery atherosclerotic plaques, identified by high-resolution B-mode ultrasonography,

compared to 13 controls (54.2%) with atherosclerotic plaques, but this difference was not statistically significant ($p=0.77$). For each type of plaque, the difference between the two groups was not statistically significant (data not shown). One patient in the EOA group had occluded internal carotid artery. All other plaques were not hemodynamically significant. EOA was not associated with an increased risk of plaque, compared to controls.

The total USBS was higher in the EOA group compared to the controls ($p=0.03$) (Table 3). As classes D-F are associated with higher risk of ischemic events, we compared the number of subjects in each group that had at least one class D-F plaque. A non-significant trend in favor of the EOA group was found (Table 3). For each group the risk of subclinical atherosclerosis did not correlate with any demographic characteristic, lipids, blood pressure, or

Table 1. Demographic characteristics of patients with EOA and normal controls

Parameter	EOA (n=24)	Controls (n=24)	<i>p</i>
Age, mean (SD)	62.5 (6.6)	60.7 (5.8)	0.33
Sex (Female)	22	22	0.99
History-Hypertension, n (%)	16 (66.6)	12 (50)	0.38
Hyperlipidemia, n (%)	7 (29.2)	13 (54.2)	0.14
Diabetes mellitus, n (%)	0 (0)	2 (8.3)	0.49
Smoking, n (%)	3 (12.5)	4 (16.7)	0.99
Treatment-statin, n (%)	4 (16.7)	8 (33.3)	0.32
ACE inhibitor, n (%)	6 (25)	3 (12.5)	0.46
Beta blocker, n (%)	2 (8.3)	6 (25)	0.244
AT II receptor blockers, n (%)	5 (20.8)	3 (12.5)	0.46
Diuretics, n (%)	6 (25)	5 (20.8)	0.74
Hydroxychloroquine, n (%)	14 (58.3)	0 (0)	<0.0001

SD, standard deviation; ACE, angiotensin converting enzyme; AT II, Angiotensin II; NS, non-significant.

Table 2. Serum lipid profiles and blood pressure measurements in patients with EOA and normal controls

Variable	EOA (n=24)	Controls (n=24)	<i>P</i>
Cholesterol (mg/dL), mean (SD)	237.6 (49.5)	230 (35.4)	0.55
HDL (mg/dL), mean (SD)	59.4 (14.3)	59.1 (13.1)	0.94
LDL (mg/dL), mean (SD)	157.5 (48.8)	147 (33.3)	0.37
Triglycerides (mg/dL), mean (SD)	141.8 (75.2)	124.4 (56)	0.63
Systolic BP (mmHg), mean (SD)	160 (32.9)	143.4 (32.1)	<0.001
Diastolic BP (mmHg), mean (SD)	99.9 (20.4)	83.42 (11.9)	<0.001

Table 3. Intima media thickness and flow-mediated and nitroglycerine-mediated vasodilatation in patients with EOA and normal controls

Variable	EOA (n=24)	Controls (n=24)	P
Framingham risk score (10-year CVD rate, mean, SD)	23.9 (14.7)	18.8 (11.2)	0.18
Intima-media thickness, common carotid artery (mm), mean (SD)	0.91 (0.17)	0.82(0.19)	0.012
Intima-media thickness, common femoral artery (mm), mean (SD)	0.7 (0.19)	0.6 (0.2)	<0.01
Ultrasound Biopsy Score, mean (SD)	14 (3.3)	12 (2.8)	0.03
Ultrasound Biopsy Score, no of patients with at least one Class D-F plaque (%)	15 (62.5)	13 (54.2)	0.77
Δ Brachial artery diameter- reactive hyperemia (flow-mediated vasodilatation-FMD) (mm), mean (SD)	0.36 (0.39)	0.36 (0.44)	0.92
Δ Brachial artery diameter after glyceryl Trinitrate administration (NMD) (mm), mean (SD)	0.62 (0.49)	0.45 (0.48)	0.057
Δ Brachial artery diamener FMD-NMD	0.348 (0.2)	0.1 (0.49)	0.026

Table 4. Risk of subclinical atherosclerosis in EOA patients and normal controls

Lesion	EOA (n=24)	Controls (n=24)	OR
IMT>1mm, n (%)	16 (66.7)	9 (37.5)	OR=3.33 (95% CI 1.02-10.9) <i>p</i> =0.043 (chi-square test) <i>P</i> =0.08 (Fischer's exact test)
Plaque, n (%)	15 (62.5)	13 (54.2)	OR=1.41 (95% CI 0.46-4.46) <i>P</i> =0.77

medications, thus no effect modifiers were found. Therefore, no multivariate analysis was required

Flow-mediated and nitroglycerine induced vasodilatation

Twenty-two individuals in both groups received glyceryl trinitrate for the assessment of endothelium-independent vasodilatation. Four individuals did not receive glyceryl trinitrate because of aortic valve stenosis (one individual) and hypotension (3 individuals). No difference was found between the two groups in both flow-mediated, endothelium-dependent vasodilatation and nitroglycerine-induced, endothelium-independent vasodilatation. However, the difference between FMD and NMD was significantly greater in EOA patients compared to healthy controls (*p*=0.026) (Table 3). Neither NMD nor FMD was correlated with systolic or diastolic BP.

DISCUSSION

In this controlled study we found evidence of subclinical atherosclerosis and endothelial dysfunction in EOA. This is in agreement with a recent Rotterdam population-based study in persons aged >55 years that investigated

carotid atherosclerosis and found an association with distal IP OA in women (26). Similarly, a population-based study in Iceland found hand OA in elderly women to be associated with carotid and coronary atherosclerosis (27). In the Framingham Heart study, an association was found between spinal facet joint OA and abdominal aortic calcifications, a surrogate for CV disease, after adjustment for epidemiologic factors associated with traditional CV risk factors (28). Also, patients with generalized OA had significantly increased popliteal artery wall thickness, assessed by MRI (29). However, other studies have not found association of hand OA with arterial stiffness (30) or abdominal aortic calcification (31).

Previous studies showed increased incidence of CV events and increased mortality in patients with OA compared to the general population (32-34). The enhanced CV risk in OA could be, at least in part, explained by the high prevalence of co-morbidities and traditional CV risk factors among older subjects with OA. Indeed, in the large population of the third National Health and Nutrition Examination Survey, 40% of the OA patients had hypertension, 11% had diabetes and 32% had high total cholesterol, compared with 25%, 6% and 24%, respectively, of the US

general population without arthritis (35). The Ulm study, in OA patients undergoing hip or knee surgery, showed a positive correlation of hypercholesterolaemia with widespread joint involvement (36). In our study, EOA did not exhibit a significant increase in the 10-year risk of CVD, as predicted by the Framingham Risk Score, which is based on the contribution of traditional CV risk factors.

We found an increased intima-media thickness (sub-clinical atherosclerosis) in patients with EOA that could not be attributed to traditional CV risk factors, as assessed by the 10-year Framingham risk score. Moreover, patients with EOA exhibited evidence of endothelial dysfunction, as estimated by the difference between FMD and NMD, that could not be attributed to differences in systolic or diastolic blood pressure.

The link between OA and atherosclerosis could be explained by shared pathogenetic mechanisms. Inflammation appears to play a role in OA and EOA in particular (1, 5, 8-10). Chronic inflammation also promotes endothelial cell activation and endothelial dysfunction (11). Inflammatory mediators, such as tumor necrosis factor- α (TNF- α), interleukin (IL)-6 and IL-1, up-regulated in OA (5), participate in the pathophysiology of atherosclerosis (37, 38). C-reactive protein, elevated in EOA (10), is associated with atherosclerosis and is predictive of future ischemic events (39, 40). Osteoprotegerin (OPG), part of the Receptor Activator Nuclear Factor κ -B (RANK)/RANK ligand system, inhibits osteoclastogenesis, and thus bone resorption, is up-regulated in hand OA (41). Elevated serum OPG levels have been associated with coronary artery disease and stroke and have been proposed as an independent risk factor for CV disease and associated mortality (42). Adiponectin may be a link between EOA and atherosclerosis. High adiponectin levels are associated with a decreased risk of radiographic progression in hand OA (43). Low levels of adiponectin are associated with obesity and CV disease. Adiponectin reduces lipid accumulation in macrophage foam cells. Macrophages transduced with adiponectin gene exhibit decreased uptake of oxidized low-density lipoprotein and increased HDL-mediated cholesterol efflux (44). However, the precise molecular basis of both disorders is still evading us.

A limitation of our study is the small sample size, and therefore, our results should be confirmed in a larger study. However, US examinations were performed by a single, experienced assessor, ruling out inter-observer variations. In addition, 14 out of 24 EOA patients (58.3%) were receiving HCQ for a mean period of 23.5 months. HCQ might have improved vascular function in our study

population, since it has been shown to exert a mild protective effect against atherosclerosis in patients with systemic lupus erythematosus (45). This means that the difference between the two populations would have been more significant, in the absence of HCQ.

In conclusion, in this small, cross-sectional, controlled study, we found evidence of subclinical atherosclerosis and endothelial dysfunction in patients with EOA.

CONFLICT OF INTEREST

The authors declare that no conflicting interests exist.

REFERENCES

1. Punzi L, Ramonda R, Sfriso P. Erosive osteoarthritis. *Best Pract. Res. Clin. Rheumatol.* 2004; 18: 739.
2. Smith MD, Triantafyllou S, Parker A, et al. Synovial membrane inflammation and cytokine production in patients with early osteoarthritis. *J. Rheumatol.* 1997; 24: 365.
3. Sakkas LI, Scanzello CR, Katsetos CD, et al. T cells and T-cell cytokine transcripts in the synovial membrane in patients with osteoarthritis. *Clin. Diagn. Lab. Immunol.* 1998; 5: 430.
4. Sakkas L, Platsoucas C. The role of T-cells in the pathogenesis of osteoarthritis. *Arthritis. Rheum.* 2007; 56: 409.
5. Kapoor M, Martel-Pelletier J, Lajeunesse D, et al. Role of proinflammatory cytokines in the pathophysiology of osteoarthritis. *Nat. Rev. Rheumatol.* 2011; 7: 33.
6. Tan AL, Grainger A, Tanner S, et al. High-resolution magnetic resonance imaging for the assessment of hand osteoarthritis. *Arthritis. Rheum.* 2005; 52: 2355.
7. Keen H, Wakefield R, Grainger A, et al. An ultrasonographic study of osteoarthritis of the hand: synovitis and its relationship to structural pathology and symptoms. *Arthritis. Rheum.* 2008; 59: 1756.
8. Vlychou M, Koutroumpas A, Malizos K, Sakkas LI. Ultrasonographic evidence of inflammation is frequent in hands of patients with erosive osteoarthritis. *Osteoarthritis. Cartilage.* 2009; 17: 1283.
9. Wittoek R, Carron P, Verbruggen G. Structural and inflammatory sonographic findings in erosive and non-erosive osteoarthritis of the interphalangeal joints. *Ann. Rheum. Dis.* 2010; 69: 2173.
10. Punzi L, Ramonda R, Oliviero F, et al. Value of C reactive protein in the assessment of erosive osteoarthritis of the hand. *Ann. Rheum. Dis.* 2005; 64: 955.
11. Kaplan MJ. Management of cardiovascular disease risk in chronic inflammatory disorders. *Nat. Rev. Rheumatol.* 2009; 5: 208.
12. Kitas GD, Gabriel SE. Cardiovascular risk in rheumatoid arthritis: state of the art and future perspectives. *Ann. Rheum. Dis.* 2011; 70: 8.
13. Altman R, Alarcón G, Appelrouth D, et al. The American College of Rheumatology criteria for the classification and reporting of osteoarthritis of the hand. *Arthritis. Rheum.* 1990; 33: 1601.
14. Touboul PJ, Hennerici MG, Meairs S, et al. Mannheim carotid intima-media thickness consensus (2004-2006). An update on behalf of the Advisory Board of the 3rd and 4th Watching the Risk Symposium, 13th and 15th European Stroke Conferences, Mannheim, Germany, 2004, and Brussels, Belgium, 2006. *Cerebrovasc. Dis.* 2007; 23: 75.
15. O'Leary DH, Polak JF, Kronmal RA, et al. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults. *N. Engl. J. Med.* 1999; 340: 14.

16. Lorenz MW, Markus HS, Bots ML, *et al.* Prediction of clinical cardiovascular events with carotid intima-media thickness: a systematic review and meta-analysis. *Circulation.* 2007; 115: 459.
17. Geroulakos G, Ramaswami G, Nicolaides A, *et al.* Characterization of symptomatic and asymptomatic carotid plaques using high-resolution real-time ultrasonography. *Br. J. Surg.* 1993; 80: 1274.
18. Komerovsky R, Desideri A. Carotid ultrasound assessment of patients with coronary artery disease: a useful index for risk stratification. *Vasc. Health Risk Manag.* 2005; 1: 131.
19. Belcaro G, Laurora G, Cesarone MR, *et al.* Noninvasive ultrasonic biopsy: evaluation of early arteriosclerotic lesions progression in normal asymptomatic, hyperlipidemic, and diabetic subjects. *Angiology.* 1993; 44: 93.
20. Belcaro G, Barsotti A, Nicolaides AN. "Ultrasonic biopsy"--a non-invasive screening technique to evaluate the cardiovascular risk and to follow up the progression and the regression of arteriosclerosis. *Vasa.* 1991; 20: 40.
21. Corretti MC, Anderson TJ, Benjamin EJ, *et al.* Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J. Am. Coll. Cardiol.* 2002; 39: 257.
22. Kuvin JT, Karas RH. Clinical utility of endothelial function testing: ready for prime time? *Circulation.* 2003; 107: 3243.
23. Moens AL, Goovaerts I, Claeys MJ, Vrints CJ. Flow-mediated vasodilation: a diagnostic instrument, or an experimental tool? *Chest.* 2005; 127: 2254.
24. Expert panel on Detection, Evaluation and Treatment of High Blood Pressure in Adults (Adult treatment panel III), Executive summary, National Cholesterol Education Program, National Heart, Lung and Blood Institute, National Institutes of Health, NIH publication No. 01-3670, May 2011.
25. D'Agostino RB Sr, Vasan RS, Pencina MJ, *et al.* General cardiovascular risk profile for use in primary care: the Framingham Heart Study. *Circulation.* 2008; 117: 743.
26. Hoeven TA, Kavousi M, Clockaerts S, *et al.* Association of atherosclerosis with presence and progression of osteoarthritis: the Rotterdam Study. *Ann. Rheum. Dis.* 2012 (In Press).
27. Jonsson H, Helgadóttir GP, Aspelund T, *et al.* Hand osteoarthritis in older women is associated with carotid and coronary atherosclerosis: the AGES Reykjavik study. *Ann. Rheum. Dis.* 2009; 68: 1696.
28. Suri P, Katz JN, Rainville J, *et al.* Vascular disease is associated with facet joint osteoarthritis. *Osteoarthritis Cartilage.* 2010; 18: 1127.
29. Kornaat PR, Sharma R, van der Geest RJ, *et al.* Positive association between increased popliteal artery vessel wall thickness and generalized osteoarthritis: is OA also part of the metabolic syndrome? *Skeletal Radiol.* 2009; 38: 1147.
30. Saleh AS, Najjar SS, Muller DC, *et al.* Arterial stiffness and hand osteoarthritis: a novel relationship? *Osteoarthritis Cartilage.* 2007; 15: 357.
31. Karasik D, Kiel DP, Kiely DK, *et al.* Abdominal aortic calcification and exostoses at the hand and lumbar spine: the Framingham Study. *Calcif. Tissue Int.* 2006; 78: 1.
32. Cerhan JR, Wallace RB, el-Khoury GY, *et al.* Decreased survival with increasing prevalence of full-body, radiographically defined osteoarthritis in women. *Am. J. Epidemiol.* 1995; 141: 225.
33. Haara MM, Mannonen P, Kroger H, *et al.* Osteoarthritis of finger joints in Finns aged 30 or over: prevalence, determinants, and association with mortality. *Ann. Rheum. Dis.* 2003; 61: 151.
34. Nuesch E, Dieppe P, Reichenbach S, *et al.* All cause and disease specific mortality in patients with knee or hip osteoarthritis: population based cohort study. *BMJ.* 2011; 342: d1165.
35. Singh G, Miller JD, Lee FH, *et al.* Prevalence of Cardiovascular Disease Risk Factors Among US Adults With Self-Reported Osteoarthritis: Data from the Third National Health and Nutrition Examination Survey. *Am. J. Manag. Care.* 2002; Suppl 8: S383.
36. Stürmer T, Sun Y, Sauerland S, *et al.* Serum cholesterol and osteoarthritis. The baseline examination of the Ulm Osteoarthritis Study. *J. Rheumatol.* 1998; 25: 1827.
37. Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N. Engl. J. Med.* 2005; 352: 1685.
38. Kleemann R, Zadelaar S, Kooistra T. Cytokines and atherosclerosis: a comprehensive review of studies in mice. *Cardiovasc Res.* 2008; 79: 360.
39. Cao JJ, Arnold AM, Manolio TA, *et al.* Association of carotid artery intima-media thickness, plaques, and C-reactive protein with future cardiovascular disease and all-cause mortality: the Cardiovascular Health Study. *Circulation.* 2007; 116: 32.
40. Haverkate E, Thompson SG, Pyke S, *et al.* Production of C-reactive protein and risk of coronary events in stable and unstable angina. *Lancet.* 2007; 349: 462.
41. Pansulaia I, Kalichman L, Kobylansky E. Association between radiographic hand osteoarthritis and RANKL, OPG and inflammatory markers. *Osteoarthritis Cartilage.* 2010; 18: 1448.
42. Kiechl S, Schett G, Wenning G, *et al.* Osteoprotegerin is a risk factor for progressive atherosclerosis and cardiovascular disease. *Circulation.* 2004; 109: 2175.
43. Yusuf E, Ioan-Facsinay A, Bijsterbosch J, *et al.* Association between leptin, adiponectin and resistin and long-term progression of hand osteoarthritis. *Ann. Rheum. Dis.* 2011; 70: 1282.
44. Tian L, Luo N, Klein RL, *et al.* Adiponectin reduces lipid accumulation in macrophage foam cells. *Atherosclerosis.* 2009; 202: 152.
45. Roman MJ, Shanker B-A, Davis A, *et al.* Prevalence and Correlates of Accelerated in Systemic Lupus Erythematosus. *N. Engl. J. Med.* 2003; 349: 2399.