

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 4-[[7-(Trifluoromethyl)quinolin-4-yl]-amino]benzenesulfonamide-ethanol-methanol (1/0.47/0.53)

Mostafa M. Ghorab,<sup>a</sup> Mansour S. Al-Said,<sup>a</sup> Abdullah A. Al-Mishari,<sup>a</sup> Ching Kheng Quah<sup>b,‡</sup> and Hoong-Kun Fun<sup>b,\*§</sup><sup>a</sup>Medicinal, Aromatic and Poisonous Plants Research Center (MAPPRC), College of Pharmacy, King Saud University, PO Box 2457, Riyadh 11451, Saudi Arabia, and<sup>b</sup>X-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia

Correspondence e-mail: hkfun@usm.my


Received 28 June 2012; accepted 29 June 2012

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å; disorder in main residue;  $R$  factor = 0.045;  $wR$  factor = 0.129; data-to-parameter ratio = 9.2.

In the title compound,  $\text{C}_{16}\text{H}_{12}\text{F}_3\text{N}_3\text{O}_2\text{S}\cdot 0.47\text{C}_2\text{H}_5\text{OH}\cdot 0.53\text{CH}_3\text{OH}$ , the quinoline ring system is approximately planar, with a maximum deviation of 0.035 (3) Å, and makes a dihedral angle of 52.67 (9)° with the benzene ring. The F atoms of the  $-\text{CF}_3$  group are disordered over two orientations, with refined site occupancies of 0.56 (2) and 0.44 (2). A single solvate site is occupied at random by ethanol or methanol, with refined site occupancies of 0.470 (6) and 0.530 (6), respectively. In the crystal, molecules are linked *via*  $\text{N}-\text{H}\cdots\text{O}$ ,  $\text{N}-\text{H}\cdots\text{N}$ ,  $\text{O}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds, thereby forming sheets lying parallel to (010).

## Related literature

For background to the biological and pharmacological activity of quinolines, see: Ghorab *et al.* (2011, 2012).



‡ Thomson Reuters ResearcherID: A-5525-2009.

§ Thomson Reuters ResearcherID: A-3561-2009.

## Experimental

## Crystal data

 $\text{C}_{16}\text{H}_{12}\text{F}_3\text{N}_3\text{O}_2\text{S}\cdot 0.47\text{C}_2\text{H}_5\text{OH}\cdot 0.53\text{CH}_3\text{O}$  $M_r = 405.98$ Triclinic,  $P\bar{1}$  $a = 8.6037$  (1) Å $b = 9.3146$  (2) Å $c = 11.4590$  (2) Å $\alpha = 92.463$  (1)° $\beta = 91.544$  (1)° $\gamma = 92.969$  (1)° $V = 915.85$  (3) Å<sup>3</sup> $Z = 2$ Cu  $K\alpha$  radiation $\mu = 2.07$  mm<sup>-1</sup> $T = 296$  K $0.83 \times 0.43 \times 0.11$  mm

## Data collection

Bruker SMART APEXII CCD diffractometer

Absorption correction: multi-scan (SADABS; Bruker, 2009)

 $T_{\min} = 0.279$ ,  $T_{\max} = 0.801$ 

9433 measured reflections

2845 independent reflections

2631 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.026$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.045$  $wR(F^2) = 0.129$  $S = 1.05$ 

2845 reflections

309 parameters

2 restraints

H atoms treated by a mixture of independent and constrained refinement

 $\Delta\rho_{\text{max}} = 0.34$  e Å<sup>-3</sup> $\Delta\rho_{\text{min}} = -0.41$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N2}-\text{H1N2}\cdots\text{O3}$	0.87	2.21	3.016 (6)	153
$\text{N3}-\text{H2N3}\cdots\text{N1}^{\text{i}}$	0.85 (3)	2.08 (3)	2.924 (3)	169 (3)
$\text{N3}-\text{H1N3}\cdots\text{O1}^{\text{ii}}$	0.87 (3)	2.26 (3)	3.107 (3)	163 (3)
$\text{O3}-\text{H1O3}\cdots\text{O1}^{\text{iii}}$	0.96	2.49	3.387 (6)	155
$\text{O3}-\text{H1O3}\cdots\text{O2}^{\text{iii}}$	0.96	2.59	3.425 (6)	146
$\text{C5}-\text{H5A}\cdots\text{O1}^{\text{iii}}$	0.93	2.50	3.343 (3)	151
$\text{C16}-\text{H16A}\cdots\text{O3}$	0.93	2.51	3.287 (6)	141

Symmetry codes: (i)  $-x + 1, -y + 2, -z + 2$ ; (ii)  $-x, -y + 2, -z + 1$ ; (iii)  $-x + 1, -y + 2, -z + 1$ .

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

The authors are grateful for the sponsorship of the Research Center, College of Pharmacy and the Deanship of Scientific Research, King Saud University, Riyadh, Saudi Arabia. HKF and CKQ also thank Universiti Sains Malaysia (USM) for the Research University Grant (No. 1001/PFIZIK/811160).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB6882).

## References

- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ghorab, M. M., Ragab, F. A., Heiba, H. I., El-Gazzar, M. G. & El-Gazzar, M. G. (2011). *Acta Pharm.* **61**, 415–425.
- Ghorab, M. M., Ragab, F. A., Heiba, H. I., Youssef, H. A. & El-Gazzar, M. G. (2012). *Med. Chem. Res.* **21**, 1376–1383.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supplementary materials

*Acta Cryst.* (2012). E68, o2396 [doi:10.1107/S1600536812029698]

**4-[[7-(Trifluoromethyl)quinolin-4-yl]amino]benzenesulfonamide–ethanol–methanol (1/0.47/0.53)**

**Mostafa M. Ghorab, Mansour S. Al-Said, Abdullah A. Al-Mishari, Ching Kheng Quah and Hoong-Kun Fun**

**Comment**

As a continuation of our efforts towards synthesizing biologically active heterocyclic compounds (Ghorab *et al.*, 2011, 2012), we have prepared the title quinoline carrying a sulfonamide moiety to evaluate its anticancer activity, which will be reported later.

In the title molecule, Fig. 1, the quinoline ring system (N1/C1-C9) is nearly planar with a maximum deviation of 0.035 (3) Å at atom C1 and it makes a dihedral angle of 52.67 (9)° with the benzene ring (C11-C16). The F atoms (F1/F2/F3) are each disordered over two positions with refined site-occupancies of 0.56 (2) and 0.44 (2). A single solvate site is occupied at random by ethanol or methanol with refined site-occupancies of 0.470 (6) and 0.530 (6) respectively.

In the crystal (Fig.2), molecules are linked *via* N2–H1N2···O3, N3–H1N3···O1, N3–H2N3···N1, O3–H1O3···O1, O3–H1O3···O2, C5–H5A···O1 and C16–H16A···O3 hydrogen bonds (Table 1) forming two-dimensional networks parallel to (010).

**Experimental**


A mixture of 4-chloro-7-trifluoromethylquinoline (0.01 mole) and sulfanilamide (0.01 mole) in absolute ethanol (30 ml) was refluxed for 8h. The solid obtained was recrystallized from ethanol to give the title compound. Colourless plates were obtained by slow evaporation from a methanol/ethanol solvent mixture at room temperature.

**Refinement**


Atoms H1N3 and H2N3 were located in a difference Fourier map and refined freely with N–H = 0.86 (3) and 0.88 (3) Å. Atom H1N2 was located in a difference Fourier map and was refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.3 U_{\text{eq}}(\text{N})$  [N–H = 0.8727]. The remaining hydrogen atoms were positioned geometrically [C–H = 0.93 or 0.96 Å, O–H = 0.95 Å] and were refined using a riding model, with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  or  $1.5 U_{\text{eq}}(\text{C}, \text{O})$ . Trifluoro atoms (F1/F2/F3) are disordered over two positions with refined site-occupancies of 0.56 (2) and 0.44 (2). A single solvate site is occupied at random by ethanol or methanol with refined site-occupancies of 0.470 (6) and 0.530 (6) respectively.

**Computing details**

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

**Figure 1**

The molecular structure of the title compound showing 30% probability displacement ellipsoids for non-H atoms. Both major and minor components of disorder are shown.

**Figure 2**

The crystal structure of the title compound, viewed along the *a* axis. H atoms not involved in hydrogen bonds (dashed lines) have been omitted for clarity. Only the major disorder component is shown.

4-[[7-(Trifluoromethyl)quinolin-4-yl]amino]benzenesulfonamide–ethanol– methanol (1/0.47/0.53)

Crystal data

$C_{16}H_{12}F_3N_3O_2S \cdot 0.47C_2H_6O \cdot 0.53CH_4O$   
 $M_r = 405.98$   
 Triclinic,  $P\bar{1}$   
 Hall symbol: -P 1  
 $a = 8.6037$  (1) Å  
 $b = 9.3146$  (2) Å  
 $c = 11.4590$  (2) Å  
 $\alpha = 92.463$  (1)°  
 $\beta = 91.544$  (1)°  
 $\gamma = 92.969$  (1)°  
 $V = 915.85$  (3) Å<sup>3</sup>

$Z = 2$   
 $F(000) = 420$   
 $D_x = 1.472$  Mg m<sup>-3</sup>  
 Cu  $K\alpha$  radiation,  $\lambda = 1.54178$  Å  
 Cell parameters from 2059 reflections  
 $\theta = 3.9$ – $70.4$ °  
 $\mu = 2.07$  mm<sup>-1</sup>  
 $T = 296$  K  
 Plate, colorless  
 $0.83 \times 0.43 \times 0.11$  mm

Data collection

Bruker SMART APEXII CCD  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 $\varphi$  and  $\omega$  scans  
 Absorption correction: multi-scan  
 (SADABS; Bruker, 2009)  
 $T_{\min} = 0.279$ ,  $T_{\max} = 0.801$

9433 measured reflections  
 2845 independent reflections  
 2631 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$   
 $\theta_{\text{max}} = 63.0$ °,  $\theta_{\text{min}} = 3.9$ °  
 $h = -9 \rightarrow 9$   
 $k = -10 \rightarrow 10$   
 $l = -12 \rightarrow 10$

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.045$   
 $wR(F^2) = 0.129$   
 $S = 1.05$   
 2845 reflections  
 309 parameters  
 2 restraints  
 Primary atom site location: structure-invariant  
 direct methods  
 Secondary atom site location: difference Fourier  
 map

Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0745P)^2 + 0.2909P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.001$   
 $\Delta\rho_{\text{max}} = 0.34$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.41$  e Å<sup>-3</sup>  
 Extinction correction: SHELXL,  
 $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$   
 Extinction coefficient: 0.0031 (7)

Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional R-factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.13678 (6)	1.19048 (6)	0.59873 (4)	0.0525 (2)	
F1A	1.3524 (19)	0.588 (2)	1.0571 (10)	0.172 (6)	0.56 (2)

F2A	1.2936 (12)	0.4416 (8)	0.9214 (12)	0.116 (4)	0.56 (2)
F3A	1.4169 (11)	0.630 (2)	0.887 (2)	0.165 (5)	0.56 (2)
F1B	1.321 (2)	0.470 (2)	0.8836 (18)	0.154 (7)	0.44 (2)
F2B	1.2955 (16)	0.5212 (19)	1.0590 (12)	0.137 (5)	0.44 (2)
F3B	1.4139 (11)	0.6569 (10)	0.961 (2)	0.118 (4)	0.44 (2)
N1	0.8884 (2)	0.8734 (2)	1.08416 (15)	0.0558 (5)	
N2	0.6928 (3)	0.8826 (3)	0.74398 (17)	0.0723 (6)	
H1N2	0.7094	0.8068	0.6996	0.094*	
N3	-0.0014 (2)	1.1225 (3)	0.67275 (19)	0.0607 (5)	
O1	0.0977 (2)	1.15432 (19)	0.47846 (13)	0.0640 (4)	
O2	0.1626 (2)	1.33775 (18)	0.63484 (16)	0.0724 (5)	
C1	0.7685 (3)	0.9498 (3)	1.05917 (19)	0.0577 (6)	
H1A	0.7267	1.0026	1.1203	0.069*	
C2	0.6992 (2)	0.9581 (3)	0.94910 (19)	0.0545 (5)	
H2A	0.6143	1.0145	0.9388	0.065*	
C3	0.7565 (2)	0.8827 (2)	0.85519 (18)	0.0512 (5)	
C4	0.8899 (2)	0.8004 (2)	0.87623 (18)	0.0499 (5)	
C5	0.9674 (3)	0.7244 (3)	0.7879 (2)	0.0648 (6)	
H5A	0.9300	0.7250	0.7110	0.078*	
C6	1.0946 (3)	0.6508 (3)	0.8123 (2)	0.0680 (7)	
H6A	1.1438	0.6016	0.7527	0.082*	
C7	1.1524 (3)	0.6487 (3)	0.9279 (2)	0.0606 (6)	
C8	1.0800 (3)	0.7194 (2)	1.0154 (2)	0.0564 (5)	
H8A	1.1178	0.7155	1.0919	0.068*	
C9	0.9493 (2)	0.7981 (2)	0.99220 (18)	0.0489 (5)	
C10	1.3000 (3)	0.5751 (3)	0.9526 (3)	0.0769 (8)	
C11	0.5640 (3)	0.9600 (3)	0.70824 (19)	0.0596 (6)	
C12	0.5560 (3)	1.1052 (3)	0.7346 (2)	0.0616 (6)	
H12A	0.6380	1.1554	0.7756	0.074*	
C13	0.4265 (3)	1.1765 (3)	0.7003 (2)	0.0578 (6)	
H13A	0.4199	1.2738	0.7203	0.069*	
C14	0.3073 (2)	1.1028 (2)	0.63654 (18)	0.0504 (5)	
C15	0.3168 (3)	0.9587 (3)	0.6061 (2)	0.0671 (7)	
H15A	0.2377	0.9102	0.5609	0.081*	
C16	0.4443 (3)	0.8871 (3)	0.6433 (2)	0.0718 (7)	
H16A	0.4499	0.7893	0.6246	0.086*	
H2N3	0.020 (3)	1.130 (3)	0.746 (3)	0.061 (7)*	
H1N3	-0.034 (3)	1.037 (3)	0.645 (3)	0.075 (9)*	
O3	0.6404 (7)	0.5947 (6)	0.6144 (5)	0.120 (2)	0.530 (6)
H1O3	0.7187	0.6422	0.5695	0.181*	0.530 (6)
C17	0.7032 (11)	0.4787 (8)	0.6295 (13)	0.148 (4)	0.530 (6)
H17A	0.7984	0.4949	0.6749	0.223*	0.530 (6)
H17B	0.6311	0.4193	0.6712	0.223*	0.530 (6)
H17C	0.7243	0.4313	0.5560	0.223*	0.530 (6)
C18	0.3657 (7)	0.5101 (10)	0.4490 (6)	0.084 (2)	0.470 (6)
H18A	0.4198	0.5162	0.3770	0.100*	0.470 (6)
H18B	0.3355	0.4103	0.4562	0.100*	0.470 (6)
C19	0.2268 (12)	0.5869 (11)	0.4363 (12)	0.131 (4)	0.470 (6)
H19A	0.1698	0.5435	0.3695	0.196*	0.470 (6)

H19B	0.1649	0.5786	0.5043	0.196*	0.470 (6)
H19C	0.2508	0.6867	0.4235	0.196*	0.470 (6)
O4	0.4659 (8)	0.5410 (5)	0.5282 (6)	0.130 (3)	0.470 (6)
H1O4	0.5659	0.5073	0.5128	0.195*	0.470 (6)

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0516 (3)	0.0669 (4)	0.0399 (3)	0.0193 (2)	-0.0105 (2)	0.0026 (2)
F1A	0.138 (8)	0.260 (12)	0.120 (7)	0.136 (8)	-0.081 (6)	-0.078 (7)
F2A	0.087 (3)	0.065 (3)	0.196 (12)	0.027 (3)	-0.023 (4)	0.014 (5)
F3A	0.059 (3)	0.210 (10)	0.243 (12)	0.052 (4)	0.031 (5)	0.124 (9)
F1B	0.152 (11)	0.182 (13)	0.128 (8)	0.127 (10)	-0.068 (7)	-0.080 (8)
F2B	0.103 (6)	0.187 (9)	0.132 (9)	0.069 (6)	-0.014 (5)	0.083 (9)
F3B	0.044 (3)	0.105 (4)	0.205 (12)	0.005 (3)	-0.022 (6)	0.029 (6)
N1	0.0571 (11)	0.0718 (11)	0.0393 (10)	0.0096 (9)	-0.0032 (8)	0.0045 (8)
N2	0.0639 (12)	0.1111 (17)	0.0443 (11)	0.0455 (12)	-0.0132 (9)	-0.0094 (10)
N3	0.0517 (11)	0.0880 (16)	0.0435 (12)	0.0193 (10)	-0.0071 (8)	0.0019 (10)
O1	0.0681 (10)	0.0856 (11)	0.0397 (9)	0.0222 (8)	-0.0117 (7)	0.0052 (7)
O2	0.0794 (12)	0.0654 (10)	0.0728 (11)	0.0213 (8)	-0.0194 (9)	-0.0012 (8)
C1	0.0546 (13)	0.0764 (14)	0.0431 (12)	0.0127 (10)	0.0047 (9)	0.0018 (10)
C2	0.0429 (11)	0.0743 (14)	0.0478 (12)	0.0140 (9)	0.0029 (8)	0.0068 (10)
C3	0.0403 (10)	0.0718 (13)	0.0424 (11)	0.0126 (9)	-0.0032 (8)	0.0047 (9)
C4	0.0434 (11)	0.0641 (12)	0.0428 (11)	0.0090 (9)	-0.0035 (8)	0.0037 (9)
C5	0.0629 (14)	0.0883 (17)	0.0447 (13)	0.0271 (12)	-0.0072 (10)	-0.0008 (11)
C6	0.0639 (15)	0.0850 (16)	0.0571 (14)	0.0314 (12)	-0.0046 (11)	-0.0039 (11)
C7	0.0502 (12)	0.0637 (13)	0.0683 (15)	0.0145 (10)	-0.0126 (10)	0.0030 (11)
C8	0.0536 (12)	0.0642 (13)	0.0510 (13)	0.0072 (10)	-0.0156 (10)	0.0052 (10)
C9	0.0448 (11)	0.0595 (11)	0.0423 (11)	0.0032 (9)	-0.0060 (8)	0.0065 (8)
C10	0.0619 (17)	0.0797 (18)	0.090 (2)	0.0252 (14)	-0.0209 (14)	0.0023 (15)
C11	0.0482 (12)	0.0918 (17)	0.0405 (12)	0.0249 (11)	-0.0059 (9)	0.0021 (10)
C12	0.0479 (12)	0.0813 (16)	0.0554 (14)	0.0065 (10)	-0.0125 (10)	0.0054 (11)
C13	0.0528 (12)	0.0680 (13)	0.0529 (13)	0.0087 (10)	-0.0089 (9)	0.0061 (10)
C14	0.0463 (11)	0.0670 (13)	0.0391 (11)	0.0158 (9)	-0.0059 (8)	0.0046 (9)
C15	0.0599 (14)	0.0769 (16)	0.0637 (15)	0.0210 (11)	-0.0241 (11)	-0.0112 (12)
C16	0.0725 (16)	0.0784 (16)	0.0649 (16)	0.0330 (13)	-0.0241 (12)	-0.0127 (12)
O3	0.144 (5)	0.093 (3)	0.119 (4)	0.004 (3)	-0.041 (3)	-0.016 (3)
C17	0.111 (8)	0.139 (10)	0.191 (13)	-0.020 (7)	-0.039 (8)	0.017 (8)
C18	0.068 (4)	0.124 (7)	0.065 (4)	0.050 (4)	0.002 (3)	0.026 (4)
C19	0.104 (7)	0.090 (6)	0.203 (12)	0.024 (5)	0.006 (7)	0.034 (6)
O4	0.160 (6)	0.071 (3)	0.159 (7)	0.004 (3)	0.039 (5)	-0.022 (3)

*Geometric parameters (Å, °)*

S1—O2	1.4202 (18)	C7—C10	1.500 (3)
S1—O1	1.4314 (16)	C8—C9	1.399 (3)
S1—N3	1.599 (2)	C8—H8A	0.9300
S1—C14	1.768 (2)	C11—C12	1.379 (4)
F1A—C10	1.266 (8)	C11—C16	1.385 (4)

F2A—C10	1.277 (8)	C12—C13	1.383 (3)
F3A—C10	1.362 (9)	C12—H12A	0.9300
F1B—C10	1.258 (10)	C13—C14	1.379 (3)
F2B—C10	1.339 (10)	C13—H13A	0.9300
F3B—C10	1.209 (9)	C14—C15	1.379 (3)
N1—C1	1.316 (3)	C15—C16	1.381 (3)
N1—C9	1.371 (3)	C15—H15A	0.9300
N2—C3	1.373 (3)	C16—H16A	0.9300
N2—C11	1.414 (3)	O3—C17	1.2497 (11)
N2—H1N2	0.8727	O3—H1O3	0.9600
N3—H2N3	0.86 (3)	C17—H17A	0.9600
N3—H1N3	0.88 (3)	C17—H17B	0.9600
C1—C2	1.387 (3)	C17—H17C	0.9600
C1—H1A	0.9300	C18—O4	1.2485 (11)
C2—C3	1.376 (3)	C18—C19	1.432 (11)
C2—H2A	0.9300	C18—O4 <sup>i</sup>	1.567 (11)
C3—C4	1.434 (3)	C18—H18A	0.9600
C4—C5	1.412 (3)	C18—H18B	0.9600
C4—C9	1.412 (3)	C19—H19A	0.9600
C5—C6	1.350 (3)	C19—H19B	0.9600
C5—H5A	0.9300	C19—H19C	0.9600
C6—C7	1.404 (3)	O4—O4 <sup>i</sup>	1.171 (11)
C6—H6A	0.9300	O4—C18 <sup>i</sup>	1.567 (11)
C7—C8	1.359 (4)	O4—H1O4	0.9500
O2—S1—O1	118.81 (11)	F3B—C10—C7	113.3 (5)
O2—S1—N3	108.50 (13)	F1B—C10—C7	114.0 (6)
O1—S1—N3	106.49 (12)	F1A—C10—C7	115.9 (4)
O2—S1—C14	107.14 (11)	F2A—C10—C7	114.0 (5)
O1—S1—C14	108.12 (10)	F2B—C10—C7	109.6 (5)
N3—S1—C14	107.28 (10)	F3A—C10—C7	110.3 (4)
C1—N1—C9	116.18 (18)	C12—C11—C16	119.6 (2)
C3—N2—C11	126.10 (19)	C12—C11—N2	121.9 (2)
C3—N2—H1N2	115.2	C16—C11—N2	118.5 (2)
C11—N2—H1N2	114.7	C11—C12—C13	120.2 (2)
S1—N3—H2N3	111.6 (17)	C11—C12—H12A	119.9
S1—N3—H1N3	112 (2)	C13—C12—H12A	119.9
H2N3—N3—H1N3	116 (3)	C14—C13—C12	119.8 (2)
N1—C1—C2	125.4 (2)	C14—C13—H13A	120.1
N1—C1—H1A	117.3	C12—C13—H13A	120.1
C2—C1—H1A	117.3	C13—C14—C15	120.4 (2)
C3—C2—C1	119.9 (2)	C13—C14—S1	120.16 (17)
C3—C2—H2A	120.0	C15—C14—S1	119.35 (17)
C1—C2—H2A	120.0	C14—C15—C16	119.6 (2)
N2—C3—C2	123.6 (2)	C14—C15—H15A	120.2
N2—C3—C4	119.04 (19)	C16—C15—H15A	120.2
C2—C3—C4	117.38 (19)	C15—C16—C11	120.3 (2)
C5—C4—C9	118.1 (2)	C15—C16—H16A	119.8
C5—C4—C3	124.07 (19)	C11—C16—H16A	119.8

C9—C4—C3	117.80 (19)	C17—O3—H1O3	99.6
C6—C5—C4	121.6 (2)	O3—C17—H17A	110.6
C6—C5—H5A	119.2	O3—C17—H17B	106.9
C4—C5—H5A	119.2	H17A—C17—H17B	109.5
C5—C6—C7	119.9 (2)	O3—C17—H17C	110.9
C5—C6—H6A	120.1	H17A—C17—H17C	109.5
C7—C6—H6A	120.1	H17B—C17—H17C	109.5
C8—C7—C6	120.2 (2)	O4—C18—C19	122.6 (9)
C8—C7—C10	120.4 (2)	O4—C18—O4 <sup>i</sup>	47.5 (5)
C6—C7—C10	119.3 (2)	C19—C18—O4 <sup>i</sup>	167.5 (9)
C7—C8—C9	121.1 (2)	O4—C18—H18A	105.7
C7—C8—H8A	119.5	C19—C18—H18A	107.2
C9—C8—H8A	119.5	O4 <sup>i</sup> —C18—H18A	72.6
N1—C9—C8	117.50 (19)	O4—C18—H18B	106.1
N1—C9—C4	123.32 (19)	C19—C18—H18B	107.9
C8—C9—C4	119.1 (2)	O4 <sup>i</sup> —C18—H18B	83.8
F3B—C10—F1B	111.2 (9)	H18A—C18—H18B	106.3
F3B—C10—F1A	69.1 (7)	C18—C19—H19A	106.8
F1B—C10—F1A	124.4 (7)	C18—C19—H19B	110.5
F3B—C10—F2A	127.6 (7)	H19A—C19—H19B	109.5
F1A—C10—F2A	107.9 (8)	C18—C19—H19C	111.1
F3B—C10—F2B	102.5 (7)	H19A—C19—H19C	109.5
F1B—C10—F2B	105.4 (9)	H19B—C19—H19C	109.5
F2A—C10—F2B	81.8 (8)	O4 <sup>i</sup> —O4—C18	80.7 (6)
F1B—C10—F3A	78.6 (8)	O4 <sup>i</sup> —O4—C18 <sup>i</sup>	51.8 (4)
F1A—C10—F3A	105.2 (7)	C18—O4—C18 <sup>i</sup>	132.5 (5)
F2A—C10—F3A	102.4 (9)	C18—O4—H1O4	114.4
F2B—C10—F3A	133.7 (6)		
C9—N1—C1—C2	0.9 (3)	C8—C7—C10—F1A	-2.1 (15)
N1—C1—C2—C3	-0.2 (4)	C6—C7—C10—F1A	174.3 (14)
C11—N2—C3—C2	1.4 (4)	C8—C7—C10—F2A	124.1 (7)
C11—N2—C3—C4	-178.6 (2)	C6—C7—C10—F2A	-59.5 (7)
C1—C2—C3—N2	178.6 (2)	C8—C7—C10—F2B	34.5 (10)
C1—C2—C3—C4	-1.4 (3)	C6—C7—C10—F2B	-149.1 (10)
N2—C3—C4—C5	3.9 (4)	C8—C7—C10—F3A	-121.4 (13)
C2—C3—C4—C5	-176.1 (2)	C6—C7—C10—F3A	55.0 (13)
N2—C3—C4—C9	-177.6 (2)	C3—N2—C11—C12	50.6 (4)
C2—C3—C4—C9	2.3 (3)	C3—N2—C11—C16	-130.4 (3)
C9—C4—C5—C6	0.2 (4)	C16—C11—C12—C13	2.5 (4)
C3—C4—C5—C6	178.7 (2)	N2—C11—C12—C13	-178.6 (2)
C4—C5—C6—C7	0.1 (4)	C11—C12—C13—C14	-2.0 (4)
C5—C6—C7—C8	0.6 (4)	C12—C13—C14—C15	-0.4 (4)
C5—C6—C7—C10	-175.8 (3)	C12—C13—C14—S1	176.73 (17)
C6—C7—C8—C9	-1.5 (4)	O2—S1—C14—C13	5.9 (2)
C10—C7—C8—C9	174.9 (2)	O1—S1—C14—C13	135.11 (19)
C1—N1—C9—C8	178.0 (2)	N3—S1—C14—C13	-110.4 (2)
C1—N1—C9—C4	0.2 (3)	O2—S1—C14—C15	-176.9 (2)
C7—C8—C9—N1	-176.2 (2)	O1—S1—C14—C15	-47.8 (2)



C7—C8—C9—C4	1.7 (3)	N3—S1—C14—C15	66.7 (2)
C5—C4—C9—N1	176.8 (2)	C13—C14—C15—C16	2.2 (4)
C3—C4—C9—N1	-1.8 (3)	S1—C14—C15—C16	-175.0 (2)
C5—C4—C9—C8	-1.0 (3)	C14—C15—C16—C11	-1.6 (4)
C3—C4—C9—C8	-179.6 (2)	C12—C11—C16—C15	-0.7 (4)
C8—C7—C10—F3B	-79.2 (13)	N2—C11—C16—C15	-179.7 (2)
C6—C7—C10—F3B	97.2 (13)	C19—C18—O4—O4 <sup>i</sup>	-170.3 (9)
C8—C7—C10—F1B	152.4 (15)	C19—C18—O4—C18 <sup>i</sup>	-170.3 (9)
C6—C7—C10—F1B	-31.2 (16)	O4 <sup>i</sup> —C18—O4—C18 <sup>i</sup>	-0.003 (1)

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N2—H1N2...O3	0.87	2.21	3.016 (6)	153
N3—H2N3...N1 <sup>ii</sup>	0.85 (3)	2.08 (3)	2.924 (3)	169 (3)
N3—H1N3...O1 <sup>iii</sup>	0.87 (3)	2.26 (3)	3.107 (3)	163 (3)
O3—H1O3...O1 <sup>iv</sup>	0.96	2.49	3.387 (6)	155
O3—H1O3...O2 <sup>iv</sup>	0.96	2.59	3.425 (6)	146
C5—H5A...O1 <sup>iv</sup>	0.93	2.50	3.343 (3)	151
C16—H16A...O3	0.93	2.51	3.287 (6)	141

Symmetry codes: (ii)  $-x+1, -y+2, -z+2$ ; (iii)  $-x, -y+2, -z+1$ ; (iv)  $-x+1, -y+2, -z+1$ .