

## Article

# Opposed elevational variation in prevalence and intensity of endoparasites and their vectors in a lizard

Lola ÁLVAREZ-RUIZ<sup>a</sup>, Rodrigo MEGÍA-PALMA<sup>b,\*</sup>, Senda REGUERA<sup>c</sup>,  
Santiago RUIZ<sup>d,e</sup>, Francisco J. ZAMORA-CAMACHO<sup>f</sup>, Jordi FIGUEROLA<sup>g,e</sup>, and  
Gregorio MORENO-RUEDA<sup>h</sup>

<sup>a</sup>Centro de Investigaciones sobre Desertificación-CSIC, Ctra. Náquera Km. 4.5, Moncada, Valencia, E-46113, Spain,

<sup>b</sup>Departamento de Ecología Evolutiva, Museo Nacional de Ciencias Naturales-CSIC, c/José Gutiérrez Abascal 2, Madrid, E-28006, Spain, <sup>c</sup>Unit of Ethology and Animal Welfare, Faculty of Veterinary, University CEU Cardenal Herrera, Alfara del Patriarca, (Valencia), E-46113, Spain, <sup>d</sup>Servicio de Control de Mosquitos, Diputación Provincial de Huelva, Centro de Investigación Biomédica en Red, Epidemiología y Salud Pública (CIBERESP), Huelva, Spain,

<sup>e</sup>CIBER de Epidemiología y Salud Pública (CIBERESP), Spain, <sup>f</sup>Departamento de Biogeografía y Cambio Global, Museo Nacional de Ciencias Naturales-CSIC, C/José Gutiérrez Abascal 2, Madrid, E-28006, Spain, <sup>g</sup>Departamento de Ecología de Humedales, Estación Biológica de Doñana-CSIC, c/Américo Vespucio s/n, Sevilla, E-41092, Spain, and <sup>h</sup>Departamento de Zoología, Facultad de Ciencias, Universidad de Granada, Granada, E-18071, Spain

\*Address correspondence to Rodrigo Megía-Palma. E-mail: rodrigo.megia@gmail.com

Handling editor: David Bierbach

Received on 10 August 2017; accepted on 8 January 2018

## Abstract

Studying the causes of parasite geographic distribution is relevant to understand ecological and evolutionary processes that affect host populations as well as for species conservation. Temperature is one of the most important environmental variables affecting parasite distribution, as raising temperatures positively affect development, reproduction, and rate of transmission of both endo- and ectoparasites. In this context, it is generally accepted that, in mountains, parasite abundance decreases with elevation. However, empirical evidence on this topic is limited. In the present study, we analyzed the elevational variation of hemoparasites and ectoparasites of a lizard, *Psammotromus algirus*, along a 2,200-m elevational gradient in Sierra Nevada (SE Spain). As predicted, ectoparasite (mites, ticks, mosquitoes, and sandflies) abundance decreased with elevation. However, hemoparasite prevalence and intensity in the lizard augmented with altitude, showing a pattern contrary to their vectors (mites). We suggest that tolerance to hemoparasites may increase with elevation as a consequence of lizards at high altitudes taking advantage of increased body condition and food availability, and reduced oxidative stress. Moreover, lizards could have been selected for higher resistance against hemoparasites at lowlands (where higher rates of replication are expected), thus reducing hemoparasite prevalence and load. Our findings imply that, in a scenario of climate warming, populations of lizards at high elevation may face increased abundance of ectoparasites, accompanied with strong negative effects.

**Key words:** elevation, immune system, parasite distribution, *Psammotromus algirus*, reptile

Parasites take resources from hosts, which is detrimental to their hosts because it reduces host fitness (reviewed in Schmid-Hempel 2011). Hence, parasites constitute an important selective pressure, with a preponderant role in host evolution (Dawkins 1990), particularly the evolution of antiparasitic strategies such as the immune system (Wakelin and Apanius 1997; Schmid-Hempel 2011; Owen and Hawley 2014). In fact, parasitism has a remarkable importance in sexual selection processes (Hamilton and Zuk 1982), with males usually having a higher parasite prevalence and load relative to females (Klein 2000). Moreover, parasites may increase risk of extinction in small or threatened host populations (Ebert et al. 2000), affecting community structure and ecosystem functioning (Hudson et al. 2002).

Predictive models of climatic change suggest that distribution shifts of wild populations (such as shifts into higher elevations in the mountains) sometimes will result in novel contact with other populations and transmission of parasites between these populations might have negative effects on naive new hosts (Rózsa et al. 2015). In this context, understanding how parasite–host dynamics vary geographically is important to comprehend ecological and evolutionary processes affecting host populations as well as for species conservation (Poisot et al. 2017). Spatiotemporal distribution of ectoparasites depends not only on environmental variables, particularly temperature (e.g., Lindsay and Birley 1996; Estrada-Peña 2001; Ekner et al. 2011), but also water availability (especially for mosquitoes; Hernández et al. 2017) and other environmental factors (Illera et al. 2017). Vector-borne pathogens usually increase their development and replication rates as well as their period of infectivity with increased temperature (reviewed in Altizer et al. 2013). Consequently, under a climate warming scenario, variations in parasite distribution and dynamics are to be expected, leading to new parasite–host encounters (Garamszegi 2011; Martínez and Merino 2011; Rózsa et al. 2015). In fact, parasite abundance in Europe has increased in the last years (Møller et al. 2013). The consequences of these changes are still poorly known, but they can exert a great impact on host populations. For example, the introduction of *Plasmodium relictum* (protozoa causing avian malaria) along with invasive birds in Hawaii has been associated with several local extinctions of endemic Hawaiian birds, which had no evolved defences against this parasite (Atkinson and LaPointe 2009).

In this sense, elevational gradients constitute an excellent experimental scenario to study geographical distribution of parasites, their vectors, and hosts along an environmental cline (see Körner 2007; Keller et al. 2013). It is generally accepted that parasite abundance decreases with elevation (Badyaev 1997). However, little evidence supports this assumption. The prevalence of parasitic mites *Riccardoella limacum* on the land snail *Arianta arbustorum* is nil at high elevations (Baur and Baur 2005). Similarly, the load of feather mites (ectosymbionts whose parasitic role remains controversial) on birds diminishes with altitude (Meléndez et al. 2014). However, the prevalence of parasitic mites on lizards has been reported to decrease (Carothers and Jaksic 2001) or increase (Spoecker 1967) with elevation, depending on the system studied. Regarding hemoparasites, the prevalence of avian malaria parasites has been reported to decrease with elevation (Zamora-Vilchis et al. 2012; Niebuhr et al. 2016). Nevertheless, in a recent study, Illera et al. (2017) found that elevational patterns of avian hemosporidian prevalence are complex, depending on the genus considered or even the mountain studied.

The objective of the present study was to examine the elevational variation in prevalence and intensity of hemoparasites and ectoparasites (acari and dipterans) in a Mediterranean lizard, *Psammmodromus algirus*. Some mosquito species may feed on lizard blood (Alcaide et al. 2009), whereas acari (mites and ticks) also act as transmitters

for hemoparasites (Svahn 1975; Ekner et al. 2011; Haklová-Kočíková et al. 2014). As temperature decreases with altitude, we expect a correlated decrease in ectoparasites as well, resulting in a concomitant elevational decrease in the prevalence of vector-borne hemoparasites. In addition, since hemoparasites that infect lizards increase their multiplication rate with host body temperature (Oppliger et al. 1996), and lizards maintain high body temperatures for even shorter durations as elevation increases (due to prolonged hibernation periods and reduced daily activity; Zamora-Camacho et al. 2013, 2016b), we predict that the intensity of hemoparasite infections will also decrease with elevation.

## Material and Methods

### Sampling

We sampled *P. algirus* lizards during their activity period (May–July), at 6 different elevations (300, 700, 1,200, 1,700, 2,200, and 2,500 m a.s.l.) along a 2,200-m altitudinal gradient in the Sierra Nevada range slopes in the southeast of Spain (3°38'00"W, 36°54'58"N). To facilitate statistical tests of elevational trends, sampling points were ultimately grouped into 3 categories: low (300 and 700 m a.s.l.), mid (1,200 and 1,700 m a.s.l.), and high (2,200 and 2,500 m a.s.l.). Sampling was conducted within the typical habitat of *P. algirus*: areas with shrubs ranging from 40 to 100 cm height and relatively high vegetation cover (Díaz and Carrascal 1991). Care was taken in selecting locations with similar habitat structure (more details in Zamora-Camacho et al. 2013). In alpine zones (above 2,000 m asl), precipitations fall as snow during the cold period, usually covering the soil from November to April. During the lizard activity period, average daily temperatures are below 18 °C (Mean ± SE; 2, 200 m asl: 16.54 ± 0.64 °C; 2,500 m asl: 17.22 ± 0.78 °C). In mid mountain, precipitations typically fall in the form of rain, with snowfalls restricted to winter. Average daily temperatures during lizard activity period are ~20 °C (1,200 m asl: 20.87 ± 0.60 °C; 1,700 m asl: 19.08 ± 0.61 °C). Lastly, at lowlands, precipitations are in the form of rain, snowfalls occurring only rarely, and temperature during lizard activity period is >20 °C (300 m asl: 25.02 ± 0.59 °C; 700 m asl: 23.01 ± 0.58 °C) (more details in Zamora-Camacho et al. 2013, 2016b).

In 2013, we sampled 56 males (*N* per elevation: low = 20, mid = 18, high = 18) and in 2016, we sampled 100 males (low = 43, mid = 41, high = 16) and 48 females (low = 17, mid = 15, high = 16). Every lizard was weighed with a digital balance to the nearest centigram and the snout to vent length (SVL) was measured with a ruler to the nearest millimeter. We only considered adult lizards for the study, which were identified as such according to SVL (details in Reguera et al. 2014a). We calculated a body condition index (BCI) as the residuals of the regression of body mass on SVL (log-transformed; Vervust et al. 2008). Lizards were collected by hand or using a pole with a loop of string with a slipknot that tightens around the neck of the lizards. These widely used techniques allow catching lizards unharmed. In 2013, and after measurements were taken, lizards were carried to the lab in cotton bags to be used in additional data collection (see Reguera et al. 2015), and released at the spot where they were caught 1 week after capture. In 2016, once samples and measurements were taken, lizards were released at exactly the same spot where they had been collected from.

### Quantification of parasites

During the breeding seasons of 2013 and 2016, we collected blood samples (<10 µL) from the ventral region of the base of the tail by

poking the lizards with sterile needles (25G) and recovering the blood with microcapillary tubes ( $75 \times 1.1$  mm, Na-heparinized). Before poking the tail, we cleaned the area with 96% ethanol. In males, we always avoided the area of the hemipenes poking the tail around 2–3 cm away from the cloaca. We pressed the wound after blood extraction to stop bleeding quickly. We performed thin blood smears that were air dried and fixed with methanol. Smears were stained for 40 min with Giemsa diluted 1: 10 in buffer, pH 7.2 (Megía-Palma et al. 2016a, 2016b). Two different authors (L.Á.-R. in 2013 and R.M.-P. in 2016) evaluated the presence of hemoparasites and counted them in a total of 10,000 blood cells per sample at  $\times 1,000$  magnification in an area with homogeneous distribution of red blood cells (Merino and Potti 1995) with a microscope BX41TF (Olympus, Tokyo, Japan). Quantification of parasitemia by observation of blood smears under the microscope has frequently been used in parasitological studies to compare variation in infection among individual hosts (e.g. Amo et al. 2005; Molnár et al. 2013; Megía-Palma et al. 2016a). The genus of the parasite was identified according to morphological characteristics of the parasites observed and the effect of these parasites on nucleus and level of cell membrane distortion in the host cell (Telford 2008). Prevalence was calculated as the percentage of individuals that presented at least 1 parasite (Rózsa et al. 2000). Intensity of infection is expressed as the mean ( $\pm$  standard error) of protozoa per 10,000 erythrocytes (Majláthová et al. 2010; Megía-Palma et al. 2016b).

In 2016, we examined the variation of ectoparasites in the geographic gradient. For this purpose, right after capturing the lizards, we checked all the lizard body for mites. Detectability of mites is low; hence, the same person (R.M.-P.) used a magnifying glass ( $\times 10$ ) to count in the field and one-by-one the number of mites beneath the scales in the base of the tail of the lizards, where mites were usually found. Some mites were collected and examined in the laboratory for identification following Moraza et al. (2009), and the presence of hemoparasites in mites was examined by performing smears of the mite body. Mean intensity was calculated as the total number of mites divided by the total number of infected individuals (Rózsa et al. 2000).

Additionally, we sampled hematophagous dipterans in 2015, by placing BG-sentinel mosquito traps (Biogents AG, Regensburg, Germany) along the elevational gradient (Roiz et al. 2015) (more details in Annex A in Supplementary Material). Moreover, given that ectoparasites may provoke immune response (both cellular and humoral) in hosts (Saino et al. 1998; Huyghe et al. 2010; Owen et al. 2010; Rynkiewicz et al. 2013), we measured, in 2010, the cellular immune response to phytohemagglutinin (PHA-P, Sigma Aldrich, L-8754) in 123 adult lizards. PHA is an innocuous protein that provokes an immune response mediated by T-cells (Martin et al. 2006; Brown et al. 2011) (more details in Annex B, in Supplementary Material).

### Statistical analyses

To analyze the distribution of endoparasites (i.e., intensity and prevalence of hemoparasites) in relation to elevation, we performed generalized linear models (GLMs) with Statistica 8.0 (StatSoft 2007). The analyses of the altitudinal variation in hemoparasite load (intensity) were fitted with a Poisson distribution linked to a log function. The variables “sex,” “elevation,” and their interaction were included as factors in the analysis. Meanwhile, the GLM performed to study prevalence of hemoparasites along elevation was fitted with a binomial distribution (absence/presence) linked to a logit function. Besides blood parasites, in 2016, we also analyzed

elevational variation in mite load. Intensity of mites was fitted by a Poisson distribution in GLMs with elevation, sex, and their interaction as factors. We did not explore the distribution of mite prevalence along the elevational gradient because it was close to 100% in the sample (see “Results” section).

The relationship between the parasite load and the SVL of the lizards along the elevational gradient was analyzed setting parasite load as dependent variable, with a GLM analysis fitted with a Poisson distribution linked to a log function. Sex was included as a factor, and SVL was included as an independent variable. However, as SVL is strongly related to elevation in this system (Zamora-Camacho et al. 2014), we performed separate analyses for each elevation to avoid the confounding effect of the altitude on SVL. Additionally, we tested for the variation in BCI with elevation by using a linear model. To control for differences in BCI between sexes and years, we included sex and year as cofactors (3 levels: males from 2013, males from 2016, and females from 2016). In addition, we analyzed the relationship between BCI and parasite load along the elevational gradient. To this end, we set parasite load (either hemoparasites or mites) as the dependent variable in two separate GLM analyses. Mite and hemoparasites loads were fitted with a Poisson distribution linked to a log function. Sex was included as a factor in both analyses, while year was only included in the analysis of hemoparasites (mites were only surveyed in 2016). We also included the interaction between elevation and BCI to explore whether the relationship between parasite load and BCI was consistent along the elevational gradient.


## Results

### Distribution of hemoparasites along the elevational gradient


We found hemoparasites of the genera *Karyolysus* (Apicomplexa: Adeleorina) and *Schellackia* (Apicomplexa: Eimeriorina) in the erythrocytes of the lizards (Supplementary Figure S1). However, the presence of the hemococcidia *Schellackia* was low with only 5.4% of the sample infected, with an intensity of infection of  $3.6 \pm 1.8$  parasites per 10,000 cells, range = 1–16. The prevalence of hemoparasites of the genus *Karyolysus* was 51% (104/204). Specifically, in 2013, 57% (32/56) of the males were infected, with a mean intensity  $\pm$  standard deviation (SD) of infection per 10,000 cells of  $59.3 \pm 11.1$  parasites, range = 1–310. In 2016, 47% (47/100) of the males ( $85.2 \pm 14.5$ , 1–479) and 50% (24/48) of the females ( $51.7 \pm 10.5$ , 1–148) were infected by parasites of the genus *Karyolysus*. The analysis of prevalence of this hemoparasite revealed a significant increase at higher elevations (GLM: Wald Stat. = 10.3,  $P = 0.005$ ; Figure 1A,B). Although there was no significant difference in prevalence between the sexes (Wald Stat. = 0.13,  $P = 0.7$ ), the intensity of infection in 10,000 cells was significantly explained by the interaction sex  $\times$  elevation (Wald Stat. = 166.9,  $P < 0.0001$ ; Figure 2), with higher infection intensities in males than in females at higher elevations (Table 1).

### Distribution of ectoparasites along the elevational gradient

We found two groups of ectoparasites on the body of the lizards: ticks (Acari: Ixodidae) and mites (Acari: Macronyssidae; Supplementary Figure S2A,B). However, only 5.4% of lizards carried ticks with a mean  $\pm$  standard error intensity of infestation per individual of  $4.1 \pm 2.1$ , range = 1–18. Regarding mite prevalence,



**Figure 1.** Prevalence of hemoparasites in *P. algirus* lizards sampled at 3 elevations in (A) 2013 (only males) and (B) 2016 (males and females) in Sierra Nevada (SE Spain).



**Figure 2.** Intensity of hemoparasites infection in *P. algirus* lizards in Sierra Nevada (SE Spain). Mean  $\pm$  SD intensity per 10,000 cells of hemoparasites of genus *Karyolysus* in males and females along the elevational gradient.

98% (98/100) of the males and 92% (49/53) of the females were infected by mites. Mite load decreased with altitude (Wald Stat. = 1603.1,  $P < 0.0001$ ) and the interaction sex  $\times$  elevation significantly explained the variation in mite load along the elevational gradient (Wald Stat. = 42.8,  $P < 0.0001$ ; Figure 3). Males had higher mite load than females in all the localities (Wald Stat. = 1708.9,  $P < 0.0001$ ; Table 1), but differences were more accentuated in low-elevation sites (Figure 3). We found different stages of the hemoparasite *Karyolysus* in body smears of mites (genus *Ophionyssus*) that

were infecting the lizards (Supplementary Figure S2C,D), thus suggesting these mites can act as transmitters of hemoparasites. Regarding dipterans, we only captured mosquitoes (Culicidae) and sandflies (Phlebotomidae) at low and mid elevation, with no dipteran caught at high elevation (more details in Annex A in Supplementary Material).

### Relationships between parasite infection and lizard traits

Sex and SVL were related to hemoparasite load in all elevations (Supplementary Table S1). Larger lizards had higher hemoparasite loads at lower and higher elevations, but the opposite was observed at mid elevations. However, infected lizards were larger than uninfected ones ( $F_{1, 141} = 11.90$ ,  $P < 0.001$ ) (mean SVL  $\pm$  SD of infected individuals =  $68.46 \pm 5.53$ ,  $n = 104$ ; uninfected individuals =  $62.72 \pm 5.29$ ,  $n = 100$ ). This significant result was independent of altitude or sex ( $F_{2, 141} = 0.90$ ,  $P = 0.42$ ; Supplementary Figure S3). Variation in mite load was also explained by sex and SVL at all elevations (Supplementary Table S1). In this case too, the relationship between mite load and SVL was inconsistent at different elevations. At the lowest elevations, where there were high mite loads, larger lizards had higher mite loads, whereas at mid and high elevations, larger lizards had lower mite loads.

Body condition tended to increase with elevation ( $F_{2, 193} = 2.53$ ,  $P = 0.08$ ; effect of sex-year:  $F_{2, 193} = 9.28$ ,  $P < 0.001$ ; Supplementary Figure S4). Intensity of hemoparasites tended to covary with body condition (Wald Stat. = 2.95,  $P = 0.08$ ), but this relationship differed inconsistently with elevation (interaction elevation  $\times$  body condition: Wald Stat. = 469.85,  $P < 0.001$ ; effect of elevation: Wald Stat. = 3614.80,  $P < 0.001$ ; Supplementary Figure S5). Males sampled in 2013 and 2016 had similar hemoparasite loads, and these were significantly higher than the hemoparasite loads of females in 2016 (Wald Stat. = 668.03,  $P < 0.001$ ). However, regarding mites (only in 2016), lizards in better body condition consistently harboured more mites at every elevation (Wald Stat. = 489.20,  $P < 0.001$ ; Supplementary Figure S6). Lastly, cellular immune response did not significantly vary with elevation (Annex B in Supplementary Material).

### Discussion


Although it is generally accepted that parasitism decreases with elevation, our findings suggest that patterns of elevational variation of parasitism may be complex. While arthropod ectoparasite (mites, ticks, mosquitoes, and sandflies) abundance decreased with elevation, hemoparasite prevalence and intensity in the lizard *P. algirus* augmented with altitude in Sierra Nevada (Spain), showing a pattern contrary to vector abundance and the expected impacts of temperature on hemoparasites.

Mites were the main ectoparasite on *P. algirus* lizards in Sierra Nevada. Although mite prevalence was almost 100% along the altitudinal gradient, mite abundance decreased with elevation. Similarly, Carothers and Jaksic (2001) reported decreased abundance of parasitic mites with elevation in *Liolaemus* lizards in the Andes. Arthropod cycles are very sensitive to temperature (e.g., Lindsay and Birley 1996; Dawson et al. 2005; Raffel et al. 2006), and ectoparasites are exposed to environmental conditions. Hence, as mites ascend in elevation, they are exposed to lower temperatures and reduced daily and annual time available to complete their life cycles. Long periods of lizard hibernation likely contribute

**Table 1.** Intensity of infection by endo- and ectoparasites of the lizard *P. algirus* along an elevational gradient

2013	Elevation	Mean intensity	2016	Elevation	Mean intensity
Hemoparasites	High	♂ 69 ± 18.2 (6–310)	Hemoparasites	High	♂ 114.8 ± 28.3 (2–429) ♀ 69.3 ± 14.2 (1–148)
	Mid	♂ 48.8 ± 11.7 (1–121)		Mid	♂ 75.7 ± 24.3 (7–479) ♀ 22.2 ± 12.3 (3–94)
	Low	♂ 43.4 ± 19.9 (20–123)		Low	♂ 67.2 ± 17.7 (1–174) ♀ 22 ± 16 (6–38)
			Mites	High	♂ 77.5 ± 14.1 (18–223) ♀ 33.8 ± 8.8 (3–111)
				Mid	♂ 119.6 ± 20.1 (3–490) ♀ 38 ± 11 (1–120)
				Low	♂ 222 ± 24.2 (53–745) ♀ 82.4 ± 14.7 (2–242)

Shown are mean ± standard error and range (minimum–maximum) of intensity of infection of hemoparasites (genus *Karyolysus*) in 2013, and hemoparasites and mites in 2016, along the elevational gradient.



**Figure 3.** Mite load (mean ± SD) in *P. algirus* lizards sampled along an elevational gradient in Sierra Nevada (SE Spain).

to increasing winter parasite mortality (Comas and Ribas 2015). Consequently, a reduction of ectoparasite abundance with elevation is consistent with our expectations.

However, contrary to our predictions, and the pattern found for ectoparasites, prevalence and intensity of hemoparasites increased with elevation. This finding contrasts with other studies reporting a decrease in hemoparasite infection with elevation in birds (Zamora-Vilchis et al. 2012; Niebuhr et al., 2016). However, an alpine population of the Iberian rock lizard, *Iberolacerta cyreni* (formerly *Lacerta monticola*), at similar high altitudes as studied here (1,900–2,200 m a.s.l.), also showed high prevalence of hemoparasites (77%; Amo et al. 2004). Moreover, a recent study in the Sierra Nevada revealed that the elevational pattern in avian malaria hemoparasites depends on the genus considered, because *Plasmodium* prevalence decreased with elevation, while *Haemoproteus* prevalence increased, and *Leucocytozoon* prevalence remained constant with elevation (Illera et al. 2017).

*Ophionyssus* mites are the known vectors of hemoparasites of the genus *Karyolysus* in lizard hosts (Svahn 1975; Telford 2008; Haklová-Kočiková et al. 2014) and we found different stages of this hemoparasite in body smears of the *Ophionyssus* mites which were infecting the lizards. Thus, hemoparasites and their main vector showed an opposed elevational pattern. Van Riper et al. (1986) already reported that the abundance of vectors did not explain the intensity of hemoparasite infection in Hawaiian birds. In fact, the hematophagous mites are present all along the elevational gradient

in Sierra Nevada, although with decreasing abundances. This puzzling result might be explained by different reasons such as elevational variation in lizards’ age (older individuals typically accumulate more parasites with time; Comas et al. 2014; Meléndez et al. 2014; Dudek et al. 2016), population density (e.g., Brown and Brown 1986), host susceptibility to infection, or host tolerance to parasites.

Mean SVL (a proxy of age in lizards) significantly increased with elevation, but the observed relationship between hemoparasite load and SVL was inconsistent along the elevational gradient. Moreover, average age of the lizards in our study system, measured by means of skeletochronology (Comas et al. 2016), is similar at the highest and the lowest elevations (Comas et al. In preparation). Thus, the elevational pattern of hemoparasite load is not explained by elevational differences in lizard age. Regarding density, no clear relation between density and parasitism has been reported in lizards (Sorci et al. 1997). Although the minimal lizard density in our study system is found at low elevation, the maximal density is at mid elevation (Zamora-Camacho et al. 2013), which is not consistent with the pattern of hemoparasite variation reported here. Lizards could be more susceptible to hemoparasites at high elevations, given that highland lizards invest more resources in colour signals (Reguera 2015), and other studies have reported that lizards with more colorful colorations suffer from increased infection by hemoparasites (Megía-Palma et al. 2016a, 2016b).

On the other hand, lizards could show higher tolerance (*sensu* Svensson and Råberg 2010; Råberg 2014) to hemoparasites at high elevations. Hosts in better condition may harbor larger intensities of parasites (Comas et al. 2014; Podmokła et al. 2014; Bonneaud et al. 2017). In our study system, we found that body condition tended (albeit not significantly) to improve with elevation, and lizards in better condition tend to harbor higher loads of hemoparasites, although the trend was inconsistent with elevation. This may be driven by increased food availability at higher elevation (Moreno-Rueda et al. 2017). Moreover, lizards suffer less oxidative damage at high altitudes (Reguera et al. 2014b, 2015), which could improve tolerance to parasites (Costantini 2014). Furthermore, lizards are darker at higher elevations (Reguera et al. 2014a), and higher melanization has been related to better tolerance to infections (Ducrest et al. 2008).

However, lowland habitats allow more time available for parasite development and are much strongly influenced by human disturbances (pesticides, pollution, habitat fragmentation, human

presence, etc.), which might decrease the lizards' capacity to tolerate hemoparasites (Giraudeau et al. 2014; also see Marcogliese 2005; Hudson et al. 2006). This might increase the cost of the infection in lowland lizards, increasing lizard mortality (see Atkinson et al. 2001), not only as a direct consequence of parasites but also as a consequence of increased risk of predation for lizards fighting the infection (Zamora-Camacho et al. 2015). Consequently, it would lead to a positive selection for parasite resistance (Sagonas et al. 2016), which might have resulted in both the observed low proportion of infected individuals and the low parasite intensities (Råberg 2014). We have not found elevational differences in cellular immune response (also see Reguera et al. 2015), although humoral immune response (more related to endoparasites) has not been examined. However, in a previous study, we found that lowland lizards responded to an antigen with hypothermia, while highland lizards did not (Zamora-Camacho et al. 2016a). Hypothermia in lowland lizards may be a way to reduce parasite replication rate (see Oppliger et al. 1996), while highland lizards would not need hypothermia to control for hemoparasites as they are more tolerant.

It is worthy to point out that we found sexual differences in mite and hemoparasite loads, and male lizards consistently having the highest ecto- and endoparasite loads at all localities. Increased parasite loads in males is a frequent finding (e.g., Amo et al. 2004; 2005; Václav et al. 2007; Dudek et al. 2016) that suggests a marked sexual dimorphism in response to infections, likely influenced by the higher levels of steroid hormones (testosterone) in males. Testosterone is immunosuppressive (Mondal and Rai 1999; Belliure et al. 2004), which explains sexual dimorphism in immune defence (Møller et al. 1998; Klein 2000, 2004; Schmid-Hempel 2011; Foo et al. 2016), thus increasing the susceptibility to parasites (Salvador et al. 1996). We only examined one component of the immune response and this might explain why we could not detect a sexual dimorphism in cellular immune response.

Reptiles are especially sensitive to global change due to their limited dispersal ability and their thermal physiology (Sinervo et al. 2010). Ectotherms could be more affected by global warming than endotherms since their survival and fitness are strongly linked to ambient temperatures (Chamaillé-James et al. 2006; Moreno-Rueda and Pleguezuelos 2007; Zamora-Camacho et al. 2010). In this sense, climate warming might result in increased hemoparasite replication rate as a consequence of expanded activity periods in highland lizards (see Moreno-Rueda et al. 2009), and might also lead to a higher abundance of vectors. In such a case, populations of lizards at high elevation could face parasite types or parasite loads for which they are not locally adapted (see e.g., Van Riper et al. 1986; Oppliger et al. 1999; Morgan et al. 2005). Thus, in a climate warming scenario, the pathogen impact could drive the decline of highland host populations.

## Acknowledgments

We thank the personnel from the Espacio Natural de Sierra Nevada for their constant support. Elena Melero, Laureano González-Granda and Sergio García Mesa participated in work during 2013. J.A. Hernández-Agüero and A. Jorge photographed the mite larvae with a SEM microscopy operating at the Museo Nacional de Ciencias Naturales-CSIC. Comments by Mar Comas, David Bierbach, and two anonymous referees improved the manuscript. The study was carried out with the permits for animal experimentation and for working with wild fauna, from Junta de Andalucía and Sierra Nevada National Park (references GMN/GyB/JMIF, ENSN/JSG/JEGT/MCF, ENSN/JSG/BRL/MCF, SGMN/GyB/JMIF, and SSA/SI/MD/ps).

## Funding

This work was funded by the European Union the Spanish government (Ministerio de Ciencia e Innovación and Ministerio de Economía y Ciencia (FEDER-MINECO), projects CGL2009-13185, CGL2014-55969-P, CGL2015-65055-P, and CGL2015-67789. F.J.Z.-C (AP2009-3505) and S.R. (AP2009-1325) were supported by two predoctoral grants from the Ministerio de Ciencia e Innovación (FPU program). F.J.Z.-C was partially supported by a Ramón Areces Foundation postdoctoral fellowship and by a Juan de la Cierva contract.

## Supplementary Material

Supplementary material can be found at <https://academic.oup.com/cz>.

## References

- Alcaide M, Rico C, Ruiz S, Soriguer R, Muñoz J et al., 2009. Disentangling vector-borne transmission networks: a universal DNA barcoding method to identify vertebrate hosts from arthropod bloodmeals. *PLoS ONE* 4:e7092.
- Altizer S, Ostfeld R, Johnson P, Kutz S, Harvell C, 2013. Climate change and infectious diseases: from evidence to a predictive framework. *Science* 341: 514–519.
- Amo L, López P, Martín J, 2004. Prevalence and intensity of haemogregarinid blood parasites in a population of the Iberian rock lizard, *Lacerta monticola*. *Parasitol Res* 94:290–293.
- Amo L, López P, Martín J, 2005. Prevalence and intensity of haemogregarine blood parasites and their mite vectors in the common wall lizard *Podarcis muralis*. *Parasitol Res* 96:378–381.
- Atkinson CT, LaPointe DA, 2009. Introduced avian diseases, climate change, and the future of Hawaiian honeycreepers. *J Avian Med Surg* 23:53–63.
- Atkinson CT, Lease JK, Drake BM, Shema NP, 2001. Pathogenicity, serological responses, and diagnosis of experimental and natural malarial infections in native Hawaiian thrushes. *Condor* 103:209–218.
- Badyaev AV, 1997. Altitudinal variation in sexual dimorphism: a new pattern and alternative hypotheses. *Behav Ecol* 8:675–690.
- Baur A, Baur B, 2005. Interpopulation variation in the prevalence and intensity of parasitic mite infection in the land snail *Arianta arbustorum*. *Invertebr Biol* 124:194–201.
- Belliure J, Smith L, Sorci G, 2004. Effect of testosterone on T cell-mediated immunity in two species of Mediterranean lacertid lizards. *J Exp Zool Part A* 301:411–418.
- Bonneaud C, Sepil I, Wilfert L, Calsbeek R, 2017. *Plasmodium* infections in natural populations of *Anolis sagrei* reflect tolerance rather than susceptibility. *Integr Comp Biol* 57:352–361.
- Brown CR, Brown MB, 1986. Ectoparasitism as a cost of coloniality in cliff swallows *Hirundo pyrrhonota*. *Ecology* 67:1206–1218.
- Brown GP, Shilton CM, Shine R, 2011. Measuring amphibian immunocompetence: validation of the phytohemagglutinin skin-swelling assay in the cane toad *Rhinella marina*. *Methods Ecol Evol* 2:341–348.
- Carothers JH, Jaksic FM, 2001. Parasite loads and altitudinal distribution of *Liolaemus* lizards in the central Chilean Andes. *Rev Chil Hist Nat* 74: 691–686.
- Chamaillé-James S, Massot M, Aragón P, Clobert J, 2006. Global warming and positive fitness response in mountain populations of common lizards *Lacerta vivipara*. *Glob Change Biol* 12:392–402.
- Comas M, Ribas A, 2015. Why are the prevalence and diversity of helminths in the endemic Pyrenean brook newt *Calotriton asper* (Amphibia, Salamandridae) so low? *J Helminthol* 89:175–181.
- Comas M, Ribas A, Milazzo C, Sperone E, Tripepi S, 2014. High levels of prevalence related to age and body condition: host-parasite interactions in a water frog *Pelophylax kl. hispanicus*. *Acta Herpetol* 9:25–31.
- Comas M, Reguera S, Zamora-Camacho FJ, Salvadó H, Moreno-Rueda G, 2016. Comparison of the effectiveness of phalanges vs. humeri and femurs to estimate lizard age with skeletochronology. *Anim Biodiv Conserv* 39: 237–240.

- Costantini D, 2014. *Oxidative Stress and Hormesis in Evolutionary Ecology and Physiology: A Marriage between Mechanistic and Evolutionary Approaches*. Berlin/Heidelberg: Springer.
- Dawkins R, 1990. Parasites, desiderata lists and the paradox of the organism. *Parasitology* 100:563–573.
- Dawson RD, Hillen KK, Whitworth TL, 2005. Effects of experimental variation in temperature on larval densities of parasitic *Protocalliphora* (Diptera: Calliphoridae) in nests of tree swallows (Passeriformes: Hirundinidae). *Environ Entomol* 34:563–568.
- Díaz JA, Carrascal LM, 1991. Regional distribution of a Mediterranean lizard: influence of habitat cues and prey abundance. *J Biogeogr* 18:291–297.
- Ducrest A-L, Keller L, Roulin A, 2008. Pleiotropy in the melanocortin system, coloration and behavioural syndromes. *Trends Ecol Evol* 23:502–510.
- Dudek K, Skórka P, Sajkowska ZA, Ekner-Grzyb A, Dudek M et al., 2016. Distribution pattern and number of ticks on lizards. *Ticks Tick-Borne Dis* 7: 172–179.
- Ebert D, Lipsitch M, Mangin KL, 2000. The effect of parasites on host population density and extinction: experimental epidemiology with *Daphnia* and six microparasites. *Am Nat* 156:459–477.
- Ekner A, Dudek K, Sajkowska Z, Majláthová V, Majláth I et al., 2011. Anaplasmataceae and *Borrelia burgdorferi* sensu lato in the sand lizard *Lacerta agilis* and co-infection of these bacteria in hosted *Ixodes ricinus* ticks. *Parasite Vectors* 4:182.
- Estrada-Peña A, 2001. Distribution, abundance, and habitat preferences of *Ixodes ricinus* (Acari: Ixodidae) in Northern Spain. *J Med Entomol* 38: 361–370.
- Foo YZ, Nakagawa S, Rhodes G, Simmons LW, 2016. The effects of sex hormones on immune function: a meta-analysis. *Biol Rev* 92:551–571.
- Garamszegi LZ, 2011. Climate change increases the risk of malaria in birds. *Global Change Biol* 17:1751–1759.
- Giraudeau M, Mousel M, Earl S, McGraw K, 2014. Parasites in the city: degree of urbanization predicts poxvirus and coccidian infections in house finches (*Haemorrhous mexicanus*). *PLoS ONE* 9:e86747.
- Haklová-Kočíková B, Hižňanová A, Majláth I, Račka K, Harris DJ et al., 2014. Morphological and molecular characterization of *Karyolysus* – a neglected but common parasite infecting some European lizards. *Parasite Vector* 7:555.
- Hamilton WD, Zuk M, 1982. Heritable true fitness and bright birds: a role for parasites? *Science* 218:384–387.
- Hernández MÁ, Rojo MÁ, Campos F, Gutiérrez-Corchero F, Moreno-Rueda G, 2017. Haemosporidian prevalence in Southern Grey Shrike *Lanius meridionalis* nestlings: impact on body condition and geographic distribution in the Iberian Peninsula. *Bird Study* 64:362–373.
- Hudson PJ, Dobson AP, Lafferty KD, 2006. Is a healthy ecosystem one that is rich in parasites? *Trends Ecol Evol* 21:381–385.
- Hudson PJ, Rizzoli A, Grenfell BT, Heesterbeek J, Dobson AP, 2002. *Ecology of Wildlife Diseases*. Oxford: Oxford University Press.
- Huyghe K, Van Oystaeyen A, Pasmans F, Tadić Z, Vanhooydonck B et al., 2010. Seasonal changes in parasite load and a cellular immune response in a colour polymorphic lizard. *Oecologia* 163:867–874.
- Illera JC, López G, García-Padilla L, Moreno Á, 2017. Factors governing the prevalence and richness of avian haemosporidian communities within and between temperate mountains. *PLoS ONE* 12:e0184587.
- Keller I, Alexander J, Holderegger R, Edwards P, 2013. Widespread phenotypic and genetic divergence along altitudinal gradients in animals. *J Evol Biol* 26:2527–2543.
- Klein SL, 2000. The effects of hormones on sex differences in infections: from genes to behavior. *Neurosci Biobehav Rev* 24:627–638.
- Klein S, 2004. Hormonal and immunological mechanisms mediating sex differences in parasite infection. *Parasite Immunol* 26:247–264.
- Körner C, 2007. The use of ‘altitude’ in ecological research. *Trends Ecol Evol* 22:569–574.
- Lindsay SW, Birley MH, 1996. Climate change and malaria transmission. *Ann Trop Med Parasitol* 90:573–588.
- Majláthová V, Majláth I, Haklová B, Hromada M, Ekner A et al., 2010. Blood parasites in two co-existing species of lizards (*Zootoca vivipara* and *Lacerta agilis*). *Parasitol Res* 107:1121–1127.
- Marcogliese DJ, 2005. Parasites of the superorganism: are they indicators of ecosystem health? *Int J Parasitol* 35:705–716.
- Martin LB, Han P, Lewittes J, Kuhlman JR, Klasing KC et al., 2006. Phytohemagglutinin-induced skin swelling in birds: histological support for a classic immunoeological technique. *Funct Ecol* 20:290–299.
- Martínez J, Merino S, 2011. Host-parasite interactions under extreme climatic conditions. *Curr Zool* 57:390–405.
- Megía-Palma R, Martínez J, Merino S, 2016a. A structural colour ornament correlates positively with parasite load and body condition in an insular lizard species. *Sci Nat* 103:1–10.
- Megía-Palma R, Martínez J, Merino S, 2016b. Structural- and carotenoid-based throat color patches in males of *Lacerta schreiberi* reflect different parasitic diseases. *Behav Ecol Sociobiol* 70:2017–2025.
- Meléndez L, Laiolo P, Mironov S, García M, Magaña O et al., 2014. Climate-driven variation in the intensity of a host-symbiont animal interaction along a broad elevation gradient. *PLoS ONE* 9:e101942.
- Merino S, Potti J, 1995. High prevalence of hematozoa in nestlings of a passerine species, the pied flycatcher *Ficedula hypoleuca*. *Auk* 112:1041–1043.
- Møller AP, Sorci G, Erritzøe J, 1998. Sexual dimorphism in immune defense. *Am Nat* 152:605–619.
- Møller AP, Merino S, Soler JJ, Antonov A, Badás EP et al., 2013. Assessing the effects of climate on host-parasite interactions: a comparative study of European birds and their parasites. *PLoS ONE* 8:e82886.
- Molnár O, Bajér K, Mészáros B, Török J, Herczeg G, 2013. Negative correlation between nuptial throat colour and blood parasite load in male European green lizards supports the Hamilton-Zuk hypothesis. *Naturwissenschaften* 100:551–558.
- Mondal S, Rai U, 1999. Sexual dimorphism in phagocytic activity of wall lizard’s splenic macrophages and its control by sex steroids. *Gen Comp Endocr* 116:291–298.
- Moraza ML, Irwin NR, Godinho R, Baird SJE, de Bellocq JG, 2009. A new species of *Ophionyssus* Mégnin (Acari: Mesostigmata: Macronyssidae) parasitic on *Lacerta schreiberi* Bedriaga (Reptilia: Lacertidae) from the Iberian Peninsula, and a world key to species. *Zootaxa* 2007:58–68.
- Moreno-Rueda G, Pleguezuelos JM, 2007. Long-term and short-term effects of temperature on snake detectability in the wild: A case study with *Malpolon monspessulanus*. *Herpetol J* 17:204–207.
- Moreno-Rueda G, Pleguezuelos JM, Alaminos E, 2009. Climate warming and activity period extension in the Mediterranean snake *Malpolon monspessulanus*. *Clim Change* 92:235–242.
- Moreno-Rueda G, Melero E, Reguera S, Zamora-Camacho FJ, Álvarez-Benito I, 2017. Prey availability, prey selection, and trophic niche width in the lizard *Psammotromus algerus* along an elevational gradient. *Curr Zool*. <https://doi.org/10.1093/cz/zox077>.
- Morgan AD, Gandon S, Buckling A, 2005. The effect of migration on local adaptation in a coevolving host-parasite system. *Nature* 437:253–256.
- Niebuhr CN, Poulin R, Tompkins DM, 2016. Is avian malaria playing a role in native bird declines in New Zealand? Testing hypotheses along an elevational gradient. *PLoS ONE* 11:e0165918.
- Oppliger A, Celerier ML, Clobert J, 1996. Physiological and behaviour changes in common lizards parasitized by haemogregarines. *Parasitology* 113:433–438.
- Oppliger A, Vernet R, Baez M, 1999. Parasite local maladaptation in the Canarian lizard *Gallotia galloti* (Reptilia: Lacertidae) parasitized by haemogregarian blood parasite. *J Evolution Biol* 12:951–955.
- Owen JP, Hawley D, 2014. Host-parasite interactions. In: Malagoli D, Ottaviani E, editors. *Eco-Immunology, Evolutionary Aspects and Future Perspectives*. Dordrecht: Springer, 73–92.
- Owen JP, Nelson AC, Clayton DH, 2010. Ecological immunology of bird-ectoparasite systems. *Trends Ecol Evol* 26:530–539.
- Podmokla E, Dubiec A, Drobniak SM, Arct A, Gustafsson L et al., 2014. Avian malaria is associated with increased reproductive investment in the blue tit. *J Avian Biol* 45:219–224.
- Poisot T, Guévenoux-Julien C, Fortin M-J, Gravel D, Legendre P, 2017. Hosts, parasites and their interactions respond to different climatic variables. *Global Ecol Biogeogr* 26:942–951.
- Råberg L, 2014. How to live with the enemy: understanding tolerance to parasites. *PLoS Biol* 12:e1001989.

- Reguera S, 2015. Adaptive color variation along an elevational gradient: the case of the Mediterranean lizard *Psammotromus algirus* [PhD Thesis]: Universidad de Granada.
- Reguera S, Zamora-Camacho FJ, Moreno-Rueda G, 2014a. The lizard *Psammotromus algirus* (Squamata: Lacertidae) is darker at high altitudes. *Biol J Linn Soc* 112:132–141.
- Reguera S, Zamora-Camacho FJ, Trenzado CE, Sanz A, Moreno-Rueda G, 2014b. Oxidative stress decreases with elevation in the lizard *Psammotromus algirus*. *Comp Biochem Phys A* 172:52–56.
- Reguera S, Zamora-Camacho FJ, Melero E, García-Mesa S, Trenzado CE et al., 2015. Ultraviolet radiation does not increase oxidative stress in the lizard *Psammotromus algirus* along an elevational gradient. *Comp Biochem Phys A* 183:20–26.
- Roiz D, Ruiz S, Soríguier R, Figuerola J, 2015. Landscape effects on the presence, abundance and diversity of mosquitoes in Mediterranean wetlands. *PLoS ONE* 10:e0128112.
- Rózsa L, Reicsigel J, Majoros G, 2000. Quantifying parasites in samples of hosts. *J Parasitol* 86:228–232.
- Rózsa L, Tryjanowski P, Vas Z, 2015. Under the changing climate: how shifting geographic distribution and sexual selection shape parasite diversification. In: Morand S, Krasnov B, Littlewood T, editors. *Parasite Diversity and Diversification: Evolutionary Ecology Meets Phylogenetics*. Cambridge (UK): Cambridge University Press, 58–76.
- Rynkiewicz EC, Hawlena H, Durden LA, Hastriter MW, Demas GE et al., 2013. Associations between innate immune function and ectoparasites in wild rodent hosts. *Parasitol Res* 112:1763–1770.
- Sagonas K, Rota IA, Tsitsilonis OE, Pafilis P, Valakos ED, 2016. Infection risk dictates immunological divergence among populations in a Mediterranean lizard. *J Evol Biol* 29:1680–1688.
- Saino N, Calza S, Möller AP, 1998. Effects of a dipteran ectoparasite on immune response and growth trade-offs in barn swallows *Hirundo rustica* nestlings. *Oikos* 81:217–228.
- Salvador A, Veiga JP, Martín J, López P, Abelenda M et al., 1996. The cost of producing a sexual signal: testosterone increases the susceptibility of male lizards to ectoparasitic infestation. *Behav Ecol* 7:145–150.
- Schmid-Hempel P, 2011. *Evolutionary Parasitology: The Integrated Study of Infections, Immunology, Ecology, and Genetics*. New York: Oxford University Press.
- Sinervo B, Méndez-de-la-Cruz F, Miles DB, Heulin B, Bastiaans B et al., 2010. Erosion of lizard diversity by climate change and altered thermal niches. *Science* 328:894–899.
- Sorci G, de Fraipont M, Clobert J, 1997. Host density and ectoparasite avoidance in the common lizard *Lacerta vivipara*. *Oecologia* 111:183–188.
- Spoecker PD, 1967. Ectoparasites of a Mojave desert population of the lizard *Uta stansburiana stejnegeri* Schmidt. *Am Midl Nat* 77:539–542.
- StatSoft, Inc. 2007. *STATISTICA* (data analysis software system), version 8.0. www.statsoft.com.
- Svahn K, 1975. Blood parasites of the genus *Karyolysus* (Coccidia, Adeleidae) in Scandinavian lizards: description and life cycle. *Norw J Zool* 23:277–295.
- Svensson EI, Råberg L, 2010. Resistance and tolerance in animal enemy-victim coevolution. *Trends Ecol Evol* 25:267–274.
- Telford SR, Jr, 2008. *Hemoparasites of the Reptilia: Color Atlas and Text*. Boca Raton (FL): CRC Press.
- Václav R, Prokop P, Fekiac V, 2007. Expression of breeding coloration in European green lizards (*Lacerta viridis*): variation with morphology and tick infestation. *Can J Zool* 85:1199–1206.
- Van Riper C, III, van Riper SG, Goff ML, Laird M, 1986. Epizootiology and ecological significance of malaria in Hawaiian land birds. *Ecol Monogr* 56:327–344.
- Vervust B, Lailvaux SP, Grbac I, Van Damme R, 2008. Do morphological condition indices predict locomotor performance in the lizard *Podarcis sicula*? *Acta Oecol* 34:244–251.
- Wakelin D, Apanius V, 1997. Immune defence: genetic control. In: Clayton DH, Moore J, editors. *Host-Parasite Evolution: General Principles and Avian Models*. New York: Oxford University Press, 30–58.
- Zamora-Camacho FJ, Moreno-Rueda G, Pleguezuelos JM, 2010. Long- and short-term impact of temperature on snake detection in the wild: further evidence from the snake *Hemorrhois hippocrepis*. *Acta Herpetol* 5:143–150.
- Zamora-Camacho FJ, Reguera S, Moreno-Rueda G, Pleguezuelos JM, 2013. Patterns of seasonal activity in a Mediterranean lizard along a 2200 m altitudinal gradient. *J Therm Biol* 38:64–69.
- Zamora-Camacho FJ, Reguera S, Moreno-Rueda G, 2014. Bergmann's Rule rules body size in an ectotherm: heat conservation in a lizard along a 2200-metre elevational gradient. *J Evol Biol* 27:2820–2828.
- Zamora-Camacho FJ, Reguera S, Rubiño-Hispán MV, Moreno-Rueda G, 2015. Eliciting an immune response reduces sprint speed in a lizard. *Behav Ecol* 26:115–120.
- Zamora-Camacho FJ, Reguera S, Moreno-Rueda G, 2016a. Elevational variation in body-temperature response to immune challenge in a lizard. *PeerJ* 4:e1972.
- Zamora-Camacho FJ, Reguera S, Moreno-Rueda G, 2016b. Thermoregulation in the lizard *Psammotromus algirus* along a 2200-m elevational gradient in Sierra Nevada (Spain). *Int J Biometeorol* 60:687–697.
- Zamora-Vilchis I, Williams SE, Johnson CN, 2012. Environmental temperature affects prevalence of blood parasites of birds on an elevation gradient: implications for disease in a warming climate. *PLoS ONE* 7:e39208.