



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Two-pore channels as master regulators of membrane trafficking and endocytic well-being

Kristin Vassileva<sup>1,2</sup>, Mark Marsh<sup>1</sup> and Sandip Patel<sup>2</sup>



Two-pore channels (TPCs) are a ubiquitous class of  $\text{Ca}^{2+}$ - and  $\text{Na}^{+}$ -permeable ion channels expressed within the endo-lysosomal system. They have emerged as central regulators of a wide array of physiological processes intimately linked to information processing. In this short review, we highlight how molecular and chemical strategies have uncovered multiple roles for TPCs in regulating various aspects of endo-lysosomal trafficking associated with disease. We summarise advances in the identification of new small molecules to pharmacologically target TPCs for medical benefit. Lastly, we discuss possible underpinning molecular mechanism(s) that translate TPC-mediated ionic fluxes to function.

## Addresses

<sup>1</sup> MRC Laboratory for Molecular Cell Biology, University College London, London, UK

<sup>2</sup> Department of Cell and Developmental Biology, University College London, London, UK

Corresponding author:

Current Opinion in Physiology 2020, 17:163–168

This review comes from a themed issue on **Calcium signaling**

Edited by **Indu S Ambudkar** and **Aldebaran M Hofer**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 14th August 2020

<https://doi.org/10.1016/j.cophys.2020.08.002>

2468-8673/© 2020 Published by Elsevier Ltd.

## Introduction

TPCs are evolutionarily ancient cation channels that reside on acidic organelles [1,2]. Humans possess two isoforms - TPC1 and TPC2, that are ubiquitously expressed through the endo-lysosomal system and associated structures such as lysosome-related organelles [3–5]. The endo-lysosomal system is a ‘super highway’ that connects the cell to its extracellular environment through a highly dynamic system of vesicles that communicate not only with each other [6] but also with other organelles such as the Golgi apparatus and the endoplasmic reticulum (ER) [7]. TPCs are thought to play key roles in regulating aspects of this communication through their ability to modulate cation flux. However, there has been some confusion regarding the activating stimuli and permeability properties of TPCs. On the one hand, TPCs are evidently activated by the  $\text{Ca}^{2+}$  mobilising messenger

NAADP, which has long been known to mobilise acidic stores of  $\text{Ca}^{2+}$  [3,4,8]. On the other hand, evidence suggests TPCs are  $\text{Na}^{+}$  channels gated directly by the endo-lysosomal phosphoinositide  $\text{PI}(3,5)\text{P}_2$  [9,10]. A recent study has clarified this issue by showing that TPCs are highly unusual in switching their ion selectivity in an agonist-dependent manner [11<sup>••</sup>]. Thus, in the presence of NAADP, TPCs behave as non-selective  $\text{Ca}^{2+}$ -permeable channels, whereas in the presence of  $\text{PI}(3,5)\text{P}_2$  they are  $\text{Na}^{+}$ -selective – a switch that can be phenocopied by novel cell permeable small molecules [11<sup>••</sup>]. The fact that both  $\text{Ca}^{2+}$  and  $\text{PI}(3,5)\text{P}_2$  are implicated in endo-lysosomal trafficking highlights the potential of TPCs as key regulators of information flow [12]. Here, we summarise recent work implicating TPCs in endo-lysosomal trafficking in both health and disease.

## TPCs and endo-lysosomal morphology

Form and function are inextricably linked, thus, changes in endo-lysosomal morphology can signal lysosomal malaise. This is well exemplified by the biomarker potential of an increase in the relative volume of acidic compartments in lysosomal storage diseases [13]. Overexpression of TPC2, but not TPC1, causes enlargement and clustering of endocytic organelles in *Xenopus* oocytes and is associated with aberrant trafficking of pigment granules [14]. Notably, the effects were reversed by the fast  $\text{Ca}^{2+}$  chelator, BAPTA, but not the slower  $\text{Ca}^{2+}$  chelator, EGTA, suggesting a role for local  $\text{Ca}^{2+}$  fluxes. Here, differences in the location of the channels appears to underlie isoform specificity, that is, the phenotype can be recapitulated by a chimeric channel in which the N-terminus of TPC1 is replaced with the N-terminus of TPC2 that contains a lysosomal targeting sequence [14,15].

Aberrant morphology of the endo-lysosomal system is also a feature of human cells overexpressing TPC2, and fibroblasts derived from people with Parkinson’s disease caused by the G2019S mutation in the Parkinson’s related protein, LRRK2 [16,17]. The latter adds to the growing body of evidence pointing to Parkinson’s being a lysosomal disease [18]. Importantly, this particular phenotype could be reversed by molecular silencing of TPC2 or NAADP antagonism [17]. As with studies in *Xenopus*, there was a selective dependence on TPC2 and local  $\text{Ca}^{2+}$  signals [17].

Collectively, these studies suggest that increased TPC2 activity, be it forced or pathological, has profound effects

on the structure of endocytic organelles which, in turn, is likely to impact lysosomal functions.

### TPCs and the vesicular trafficking of receptors and protein toxins

Cell homeostasis is dependent on the ongoing, bidirectional flow of information between the plasma membrane and the cell interior. This includes pathways that mediate the trafficking of internalised cell surface receptors. In this context, EGF receptors and LDL receptors have been shown to accumulate in late endosomes in TPC2-deficient cells [19]. Mis-trafficking of the latter is likely to account for the non-alcoholic fatty liver disease-like symptoms observed in TPC2 knockout mice fed a cholesterol-rich diet [19]. Knockout of TPC2 also modestly delays the trafficking of PDGF receptors [20]. In this study, the effects were isoform-specific as knockout of TPC1 had little effect. Additionally, knockdown of either TPC1 or TPC2 results in accumulation of integrins in early endosomes [21]. This effect likely contributes to the reduced adhesion and migration of cancer cell lines and metastatic tumour formation *in vivo* [21].

Receptor endocytosis is often hijacked by protein toxins and other pathogens to gain entry into cells. TPC1, but not TPC2, knockout impeded trafficking of cholera toxin to the Golgi apparatus [20]. This effect is reminiscent of overexpression studies of sea urchin TPC1 which also perturbed cholera toxin trafficking [16]. However, in an independent study, knockout of TPC1 had only modest effects on cholera toxin-induced cyclic AMP production [22]. TPC1 knockout reduced the trafficking and downstream effects of a number of so called 'short trip' bacterial toxins, such as anthrax and diphtheria toxins, that enter cells through early endocytic compartments [22].

In summary, multiple studies show that TPCs regulate the transit and biological activity of a number of receptors and cargoes that enter cells through receptor-mediated endocytosis, often in an isoform-selective manner (Figure 1). It remains to be seen how extensively this applies to other endocytic cargoes.

### TPCs and the trafficking of viruses

In recent years, TPCs have emerged as novel host factors supporting infection by clinically relevant viral pathogens. Sakurai *et al.* demonstrated that disrupting TPC function, molecularly or pharmacologically, prevented the cellular entry of Ebola virus (EBOV) [23\*\*]. Knock down or knockout of either TPC isoform, as well as overexpression of a dominant negative TPC2 pore mutant inhibited EBOV escape from the endosomal network. Similarly, the NAADP antagonist Ned-19 and a range of voltage-gated Ca<sup>2+</sup> channel blockers inhibited EBOV infection *in vitro*. Of the latter, the plant-derived alkaloid tetrandrine appeared to be the most potent inhibitor of EBOV entry with half-maximal inhibitory

concentration (IC<sub>50</sub>) in the nanomolar range. Notably, tetrandrine is a potent blocker of TPCs and showed therapeutic efficacy against EBOV in an *in vivo* mouse model [23\*\*].

More recent work by Marchant and colleagues [24] suggested that the intracellular trafficking and subsequent infection of another enveloped RNA virus - MERS-CoV, is also reliant on TPC activity. Loss of function analyses revealed that TPC knock down impairs MERS-CoV pseudovirus translocation through the endocytic system. Furthermore, overexpression of TPC1 or TPC2 also led to significant reduction in translocation events. The observed inhibitory effects were strongly influenced by the correct subcellular distribution of the channels as overexpression of a functionally active TPC2 re-routed to the plasma membrane had no notable effect on MERS-CoV pseudovirus infectivity. Importantly, knock down and overexpression of a distinct lysosomal Ca<sup>2+</sup> channel – TRPML1, exerted no inhibitory effects, thus attesting to specificity. A number of bisbenzylisoquinoline alkaloids related to tetrandrine, which were demonstrated to inhibit NAADP-evoked Ca<sup>2+</sup> release in intact cells, also potently impaired pseudovirus progression through the endosomal network [24]. Tetrandrine also blocks SARS-CoV-2 pseudo-virus infection [25] pointing to a general requirement for TPCs in Coronavirus infection.


TPCs were also found to regulate the endo-lysosomal escape of the HIV-1 encoded Tat peptide [26]. Following secretion from infected cells, Tat enters uninfected cells via receptor-mediated endocytosis and transits to the nucleus to enhance gene transcription. Khan *et al.* demonstrated that drug-induced blockade or molecular silencing of TPCs but not TRPML1 prevents Tat from escaping endocytic compartments [26]. Further investigations using more biologically relevant systems are needed to determine whether blockade of TPCs has therapeutic potential against HIV-1.

Collectively, TPCs have emerged as common host factors exploited by a number of non-related viral pathogens (Figure 1).

### TPCs and non-vesicular traffic

A hallmark of NAADP-mediated Ca<sup>2+</sup> signalling is its global nature, resulting from amplification of local Ca<sup>2+</sup> signals deriving from acidic Ca<sup>2+</sup> stores by ER Ca<sup>2+</sup> channels [27]. It has long been proposed that such inter-organellar communication occurs at membrane contact sites between the acidic organelles and the ER [28]. Both endosome-ER and lysosome-ER contacts are readily identifiable by EM and abundant [29,30]. Importantly, a recent study has provided evidence that local 'constitutive' NAADP-mediated Ca<sup>2+</sup> signals may act to strengthen contacts between late endosomes and the ER [31\*\*]. Previously, such contacts have been shown to

Figure 1



Schematic illustrating distinct TPC-dependent membrane trafficking events. TPCs have been implicated in the trafficking of internalised proteins including cell-surface receptors and integrins through early, late and recycling endosomes, and strengthening of membrane contact sites between endosomes and the endoplasmic reticulum (left). They also regulate trafficking of protein toxins, viruses and viral proteins, and the secretion of enzymes and proteins to the extracellular matrix (right).

promote de-phosphorylation of endosome-associated internalised EGF receptors by an ER-localised phosphatase [29]. Disruption of contacts, by interfering with NAADP signalling either molecularly (through knock down of TPC1 but not TPC2) or chemically with Ned-19, can delay de-phosphorylation resulting in EGF receptor hyperactivation [31\*\*]. This was again associated with profound changes in endo-lysosomal morphology. As contact sites are formed without membrane fusion, this work links TPCs to non-vesicular traffic and provides scope for investigating Ca<sup>2+</sup> fluxes in the control of lipid transfer that is heavily reliant on inter-organelle contact sites [32].

### The TPC interactome - towards mechanism

How do TPCs regulate the various aspects of membrane traffic discussed above? The TPC interactome is providing some key insights. To date, three studies have taken unbiased approaches to define proteins that associate with TPC1, TPC2 or both [14,19,22] (reviewed in Ref. [33]). One common feature that emerges is the interaction of TPCs with SNARE proteins that regulate membrane fusion. These include both Q-SNARES and R-SNARES,

such as syntaxins and VAMP isoforms. Such associations are consistent with a role for TPCs in vesicular fusion, which could potentially account for TPC2-dependent lysosomal expansion [14,17] and TPC2-dependent lysosomal exocytosis [34]. A notable omission from these data sets is the synaptotagmins that serve as Ca<sup>2+</sup> sensors for fusion elsewhere.

A fusogenic role for TPCs could underlie the halted transit of internalised receptors, protein toxins and viruses upon TPC blockade. The identity of the compartment(s) in which cargoes stall is unclear at present. For EBOV, there is evidence both for [35] and against [23\*\*] the virus accumulating in TPC2-positive compartments that are also positive for NPC1, an essential host factor for EBOV fusion. Additionally, we know little of the relationship between this compartment and compartments that accumulate receptors and protein toxins in the absence of TPC activity.

TPC2 also associates with a number of Rab proteins, including the endo-lysosomal Rab - Rab7, through an identified binding site in its N-terminus [14]. Rab

proteins are key regulators of both vesicular and non-vesicular traffic [36]. Consequently, TPC2 might act as a Rab effector. Consistent with this, mutant channels unable to bind Rabs are less able to support NAADP-evoked  $\text{Ca}^{2+}$  signals [14]. Furthermore, morphology defects associated with TPC2 overexpression and mutant LRRK2 are reversed by Rab7 inhibition [14,17]. Rab7 and several isoforms of the annexin family of  $\text{Ca}^{2+}$  binding proteins associate with TPCs and regulate contact site formation between endocytic compartments and the ER [37,38]. These interactors could thus act upstream and downstream of TPCs, respectively to direct non-vesicular traffic.

Interestingly, Gunaratne *et al.* provided several lines of evidence indicating that interfering with TPC action impairs the activity of furin, a  $\text{Ca}^{2+}$ -dependent endoprotease, which may mediate the proteolytic priming of MERS virus spike protein [24]. This could underpin the decrease in MERS pseudovirus entry upon TPC blockade. However, furin is reportedly dispensable for the activation of MERS-CoV envelope protein [39]. Moreover, this mechanism is unlikely to apply to EBOV fusion/entry since EBOV glycoprotein (GP) is processed by endogenous cathepsins rather than furin [40]. Additionally, tetrandrine and Ned-19 inhibited the entry of EBOV virus-like particles (VLPs) bearing cleaved GP, suggesting that TPCs regulate a step distal to proteolysis [23\*\*].

In summary, TPCs likely regulate both vesicular and non-vesicular trafficking through physical association with regulatory proteins but further work is required to fully define these interactions and relate them to  $\text{Ca}^{2+}$  fluxes.

### Chemical targeting of TPCs

Despite the increasingly recognised pathophysiological significance of TPCs [41], their pharmacology is poorly defined. The fact that TPCs are essential host factors required by EBOV and MERS-CoV to enter target cells is significant. This is because the available literature on viral entry blockers offers a rich source of potential TPC modulators since some of the compounds with unknown mechanisms of action may elicit their antiviral effects through inhibition of TPCs.

In this context, Penny *et al.* screened a library of FDA-approved drugs against an *in silico* model of TPC2 to identify lead compounds targeting the pore region of the channel [42\*]. To further refine the hits from the virtual screen, the highest-ranking compounds were compared with the hits from two independent high-throughput screens for EBOV antivirals [43,44]. This cross-referencing returned drugs targeting oestrogen and dopamine receptors as common hits [42\*]. The selected hits inhibited endogenous NAADP-evoked  $\text{Ca}^{2+}$  release and recombinant TPC2 channel activity. Mechanistically,

the drugs behaved as TPC2 pore blockers, reducing the mean open time of the channel. Notably, the relative potency of the drugs in electrophysiological studies directly correlated with their potencies for inhibition of EBOV VLP entry [42\*]. Gunaratne *et al.* undertook an independent screening campaign using sea urchin egg homogenates and identified several novel inhibitors of NAADP-dependent  $\text{Ca}^{2+}$  signalling that were active in mammalian cells [45]. Importantly, the compounds blocked NAADP-evoked  $\text{Ca}^{2+}$  signals and MERS-CoV pseudovirus translocation with similar potencies [24]. Whether these compounds act directly on TPCs remains to be established.

Collectively, these findings underscore the importance of NAADP signalling through TPCs in viral entry and uncover novel blocking drugs for manipulating TPC-dependent trafficking events.

### Outlook

To date most of the effects of TPCs have been ascribed to local  $\text{Ca}^{2+}$  fluxes. However, the downstream effectors that translate these fluxes to function are largely unknown. Further mining of TPC interactomes and functional validation is thus warranted not least due to their potential in combatting disease. Moreover, the permeability of the channels to  $\text{Na}^+$ , associated with activation by  $\text{PI}(3,5)\text{P}_2$  in a functional context is still largely unexplored. Early work suggested that TPCs are tonically inhibited by the nutrient sensing protein kinase mTOR under nutrient replete conditions but become activated during starvation to regulate the lysosome membrane potential through  $\text{Na}^+$  permeability [46]. This in turn was proposed to maintain lysosomal  $\text{H}^+$ , amino acid flux and exercise endurance *in vivo*. Recent work suggests an additional role for TPC-associated  $\text{Na}^+$  flux in regulating the volume of macropinosomes [47\*\*]. This osmoregulatory role may apply more generally to the endocytic system and associated trafficking functions. Recent work also suggests that  $\text{Na}^+$  flux through TPCs may selectively couple to lysosomal exocytosis [11\*\*], even though this process is thought to be  $\text{Ca}^{2+}$ -dependent [12]. Understanding and maintaining endocytic well-being necessitate teasing apart fluxes of  $\text{Ca}^{2+}$  and  $\text{Na}^+$  in endocytic trafficking, defining molecularly the sequelae of events that these fluxes initiate, and developing strategies for their selective manipulation.

### Conflict of interest statement

Nothing declared.

### Acknowledgements

This work was funded by a BBSRC PhD studentship, BBSRC grant BB/N01524X/1 and Medical Research Council funding to the MRC-UCL LMCB University Unit (MC\_UU\_12018/1).

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as

- of special interest
- of outstanding interest

1. Rahman T, Cai X, Brailoiu GC, Abood ME, Brailoiu E, Patel S: **Two-pore channels provide insight into the evolution of voltage-gated Ca<sup>2+</sup> and Na<sup>+</sup> channels.** *Sci Signal* 2014, **7**:ra109.
  2. Patel S: **Function and dysfunction of two-pore channels.** *Sci Signal* 2015, **8**:re7.
  3. Calcra PJ, Ruas M, Pan Z, Cheng X, Arredouani A, Hao X, Tang J, Rietdorf K, Teboul L, Chuang KT *et al.*: **NAADP mobilizes calcium from acidic organelles through two-pore channels.** *Nature* 2009, **459**:596-600.
  4. Brailoiu E, Churamani D, Cai X, Schrlau MG, Brailoiu GC, Gao X, Hooper R, Boulware MJ, Dun NJ, Marchant JS, Patel S: **Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling.** *J Cell Biol* 2009, **186**:201-209.
  5. Cai X, Patel S: **Degeneration of an intracellular ion channel in the primate lineage by relaxation of selective constraints.** *Mol Biol Evol* 2010, **27**:2352-2359.
  6. Luzio JP, Pryor PR, Bright NA: **Lysosomes: fusion and function.** *Nat Rev Mol Cell Biol* 2007, **8**:622-632.
  7. Wu H, Carvalho P, Voeltz GK: **Here, there, and everywhere: the importance of ER membrane contact sites.** *Science* 2018:361.
  8. Churchill GC, Okada Y, Thomas JM, Genazzani AA, Patel S, Galione A: **NAADP mobilizes Ca<sup>2+</sup> from reserve granules, lysosome-related organelles, in sea urchin eggs.** *Cell* 2002, **111**:703-708.
  9. Wang X, Zhang X, Dong XP, Samie M, Li X, Cheng X, Goschka A, Shen D, Zhou Y, Harlow J *et al.*: **TPC proteins are phosphoinositide-activated sodium-selective ion channels in endosomes and lysosomes.** *Cell* 2012, **151**:372-383.
  10. She J, Guo J, Chen Q, Zeng W, Jiang Y, Bai XC: **Structural insights into the voltage and phospholipid activation of the mammalian TPC1 channel.** *Nature* 2018, **556**:130-134.
  11. Gerndt S, Chen C-C, Chao Y-K, Yuan Y, Burgstaller S, Scotto
  - Rosato A, Krogsaeter E, Urban N, Jacob K, Nguyen ONP *et al.*: **Agonist-mediated switching of ion selectivity in TPC2 differentially promotes lysosomal function.** *eLife* 2020, **9**:e54712.
- This paper identified an unusual property of the TPCs whereby TPC2 exhibits an agonist-dependent switch in its ion selectivity. The authors demonstrated that TPC2 is a non-selective Ca<sup>2+</sup>-permeable channel in the presence of NAADP but Na<sup>+</sup>-selective in the presence of PI(3,5)P<sub>2</sub>. Notably, a high throughput screen yielded two cell permeable small molecules that effectively mirrored each agonist in switching ion selectivity. These findings resolve opposing views on the activation and permeability properties of these abundant cation channels, and provide new molecular tools to strategically manipulate them.
12. Xu H, Ren D: **Lysosomal physiology.** *Annu Rev Physiol* 2015, **77**:57-80.
  13. te Vrugte D, Speak AO, Wallom KL, Al Eisa N, Smith DA, Hendriksz CJ, Simmons L, Lachmann RH, Cousins A, Hartung R *et al.*: **Relative acidic compartment volume as a lysosomal storage disorder-associated biomarker.** *J Clin Invest* 2014, **124**:1320-1328.
  14. Lin-Moshier Y, Keebler MV, Hooper R, Boulware MJ, Liu X, Churamani D, Abood ME, Walseth TF, Brailoiu E, Patel S, Marchant JS: **The two-pore channel (TPC) interactome unmasks isoform-specific roles for TPCs in endolysosomal morphology and cell pigmentation.** *Proc Natl Acad Sci U S A* 2014, **111**:13087-13092.
  15. Brailoiu E, Rahman T, Churamani D, Prole DL, Brailoiu GC, Hooper R, Taylor CW, Patel S: **An NAADP-gated two-pore channel targeted to the plasma membrane uncouples triggering from amplifying Ca<sup>2+</sup> signals.** *J Biol Chem* 2010, **285**:38511-38516.
  16. Ruas M, Rietdorf K, Arredouani A, Davis LC, Lloyd-Evans E, Koegel H, Funnell TM, Morgan AJ, Ward JA, Watanabe K *et al.*: **Purified TPC isoforms form NAADP receptors with distinct roles for Ca<sup>2+</sup> signaling and endolysosomal trafficking.** *Curr Biol* 2010, **20**:703-709.
  17. Hockey LN, Kilpatrick BS, Eden ER, Lin-Moshier Y, Brailoiu GC, Brailoiu E, Futter C, Schapira AH, Marchant JS, Patel S: **Dysregulation of lysosomal morphology by pathogenic LRRK2 is corrected by TPC2 inhibition.** *J Cell Sci* 2015, **128**:232-238.
  18. Klein AD, Mazzulli JR: **Is Parkinson's disease a lysosomal disorder?** *Brain* 2018, **141**:2255-2262.
  19. Grimm C, Holdt LM, Chen CC, Hassan S, Muller C, Jors S, Cuny H, Kissing S, Schroder B, Butz E *et al.*: **High susceptibility to fatty liver disease in two-pore channel 2-deficient mice.** *Nat Commun* 2014, **5**:4699.
  20. Ruas M, Chuang KT, Davis LC, Al-Douri A, Tynan PW, Tunn R, Teboul L, Galione A, Parrington J: **TPC1 has two variant isoforms and their removal has different effects on endo-lysosomal functions compared to loss of TPC2.** *Mol Cell Biol* 2014, **34**:3981-3992.
  21. Nguyen ON, Grimm C, Schneider LS, Chao YK, Atzberger C, Bartel K, Watermann A, Ulrich M, Mayr D, Wahl-Schott C *et al.*: **Two-pore channel function is crucial for the migration of invasive cancer cells.** *Cancer Res* 2017, **77**:1427-1438.
  22. Castonguay J, Orth JHC, Muller T, Sleman F, Grimm C, Wahl-Schott C, Biel M, Mallmann RT, Bildl W, Schulte U, Klugbauer N: **The two-pore channel TPC1 is required for efficient protein processing through early and recycling endosomes.** *Sci Rep* 2017, **7**:10038.
  23. Sakurai Y, Kolokolstov AA, Chen CC, Tidwell MW, Bauta WE,
  - Klugbauer N, Grimm C, Wahl-Schott C, Biel M, Davey RA: **Two-pore channels control Ebola virus host cell entry and are drug targets for disease treatment.** *Science* 2015, **347**:995-998.
- This paper first identified a link between TPCs and Ebola virus (EBOV) infection. Following macropinocytosis and proteolytic cleavage of the EBOV envelope glycoprotein, EBOV requires the Niemann-Pick disease type C protein 1 for fusion and entry into cells. Using gene knock outs, siRNA and small molecule inhibitors, this work shows that TPCs are also required for EBOV entry. One particular inhibitor of TPCs, tetrandrine, efficiently inhibited EBOV infection in tissue culture and in a mouse model, indicating the potential that drugs against TPCs might be used to inhibit EBOV infection.
24. Gunaratne GS, Yang Y, Li F, Walseth TF, Marchant JS: **NAADP-dependent Ca(2+) signaling regulates Middle East respiratory syndrome-coronavirus pseudovirus translocation through the endolysosomal system.** *Cell Calcium* 2018, **75**:30-41.
  25. Ou X, Liu Y, Lei X, Li P, Mi D, Ren L, Guo L, Guo R, Chen T, Hu J *et al.*: **Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV.** *Nat Commun* 2020, **11**:1620.
  26. Khan N, Halcrow PW, Lakpa KL, Afghah Z, Miller NM, Dowdy SF, Geiger JD, Chen X: **Two-pore channels regulate Tat endolysosome escape and Tat-mediated HIV-1 LTR transactivation.** *FASEB J* 2020, **34**:4147-4162.
  27. Galione A, Receptors NAADP: *Cold Spring Harb Perspect Biol* 2019, **11**.
  28. Patel S, Brailoiu E: **Triggering of Ca<sup>2+</sup> signals by NAADP-gated two-pore channels: a role for membrane contact sites?** *Biochem Soc Trans* 2012, **40**:153-157.
  29. Eden ER, White IJ, Tsapara A, Futter CE: **Membrane contacts between endosomes and ER provide sites for PTP1B-epidermal growth factor receptor interaction.** *Nat Cell Biol* 2010, **12**:267-272.
  30. Kilpatrick BS, Eden ER, Schapira AH, Futter CE, Patel S: **Direct mobilisation of lysosomal Ca<sup>2+</sup> triggers complex Ca<sup>2+</sup> signals.** *J Cell Sci* 2013, **126**:60-66.
  31. Kilpatrick BS, Eden ER, Hockey LN, Yates E, Futter CE, Patel S: **An endosomal NAADP-sensitive two-pore Ca<sup>2+</sup> channel regulates ER-endosome membrane contact sites to control growth factor signaling.** *Cell Rep* 2017, **18**:1636-1645.

This paper revealed an essential role for TPC1 in strengthening membrane contact sites between endosomes and the ER. The authors demonstrated that molecular or chemical interference with TPC1 or NAADP signalling reduced contact site formation and delayed de-phosphorylation of endocytosed EGF receptors leading to their hyperactivation. The study highlights the importance of endo-lysosomal Ca<sup>2+</sup> in the structural and functional wellbeing of the endocytic system through regulating non-vesicular traffic.

32. Wong LH, Gatta AT, Levine TP: **Lipid transfer proteins: the lipid commute via shuttles, bridges and tubes.** *Nat Rev Mol Cell Biol* 2019, **20**:85-101.
33. Krogsaeter EK, Biel M, Wahl-Schott C, Grimm C: **The protein interaction networks of mucolipins and two-pore channels.** *Biochim Biophys Acta Mol Cell Res* 2019, **1866**:1111-1123.
34. Davis LC, Morgan AJ, Chen JL, Snead CM, Bloor-Young D, Shenderov E, Stanton-Humphreys MN, Conway SJ, Churchill GC, Parrington J *et al.*: **NAADP activates two-pore channels on T cell cytolytic granules to stimulate exocytosis and killing.** *Curr Biol* 2012, **22**:2331-2337.
35. Simmons JA, D'Souza RS, Ruas M, Galione A, Casanova JE, White JM: **Ebolavirus glycoprotein directs fusion through NPC1+ endolysosomes.** *J Virol* 2016, **90**:605-610.
36. Zerial M, McBride H: **Rab proteins as membrane organizers.** *Nat Rev Mol Cell Biol* 2001, **2**:107-117.
37. Rocha N, Kuijl C, van der Kant R, Janssen L, Houben D, Janssen H, Zwart W, Neeffjes J: **Cholesterol sensor ORP1L contacts the ER protein VAP to control Rab7-RILP-p150 Glued and late endosome positioning.** *J Cell Biol* 2009, **185**:1209-1225.
38. Eden ER, Sanchez-Heras E, Tsapara A, Sobota A, Levine TP, Futter CE: **Annexin A1 tethers membrane contact sites that mediate ER to endosome cholesterol transport.** *Dev Cell* 2016, **37**:473-483.
39. Matsuyama S, Shirato K, Kawase M, Terada Y, Kawachi K, Fukushi S, Kamitani W: **Middle east respiratory syndrome coronavirus spike protein is not activated directly by cellular Furin during viral entry into target cells.** *J Virol* 2018:92.
40. Chandran K, Sullivan NJ, Felbor U, Whelan SP, Cunningham JM: **Endosomal proteolysis of the Ebola virus glycoprotein is necessary for infection.** *Science* 2005, **308**:1643-1645.
41. Patel S, Kilpatrick BS: **Two-pore channels and disease.** *Biochim Biophys Acta Mol Cell Res* 2018, **1865**:1678-1686.
42. Penny CJ, Vassileva K, Jha A, Yuan Y, Chee X, Yates E, Mazzon M, Kilpatrick BS, Muallem S, Marsh M, Rahman T, Patel S: **Mining of Ebola virus entry inhibitors identifies approved drugs as two-pore channel pore blockers.** *Biochimica et biophysica acta. Mol Cell Res* 2019, **1866**:1151-1161.

This paper builds on the work of Sakurai *et al.* [23]. Using *in silico* screening of a library of FDA approved drugs against a model of the human TPC2 pore, and cross-referencing the 'hits' against two high through put screens for inhibitors of Ebola virus entry, the authors identified a number of compounds targeting dopamine and oestrogen receptors as common leads. These compounds inhibited channel activity of TPC2 as well as the entry of Ebola virus-like particles, providing further support for a key role for TPCs in virus entry.

43. Kouznetsova J, Sun W, Martinez-Romero C, Tawa G, Shinn P, Chen CZ, Schimmer A, Sanderson P, McKew JC, Zheng W, Garcia-Sastre A: **Identification of 53 compounds that block Ebola virus-like particle entry via a repurposing screen of approved drugs.** *Emerg Microbes Infect* 2014, **3**:e84.
44. Johansen LM, DeWald LE, Shoemaker CJ, Hoffstrom BG, Lear-Rooney CM, Stossel A, Nelson E, Delos SE, Simmons JA, Grenier JM *et al.*: **A screen of approved drugs and molecular probes identifies therapeutics with anti-Ebola virus activity.** *Sci Transl Med* 2015, **7**:290ra289.
45. Gunaratne GS, Johns ME, Hintz HM, Walseth TF, Marchant JS: **A screening campaign in sea urchin egg homogenate as a platform for discovering modulators of NAADP-dependent Ca (2+) signaling in human cells.** *Cell Calcium* 2018, **75**:42-52.
46. Cang C, Zhou Y, Navarro B, Seo YJ, Aranda K, Shi L, Battaglia-Hsu S, Nissim I, Clapham DE, Ren D: **mTOR regulates lysosomal ATP-sensitive two-pore Na(+) channels to adapt to metabolic state.** *Cell* 2013, **152**:778-790.
47. Freeman SA, Uderhardt S, Saric A, Collins RF, Buckley CM, Mylvaganam S, Boroumand P, Plumb J, Germain RN, Ren D, Grinstein S: **Lipid-gated monovalent ion fluxes regulate endocytic traffic and support immune surveillance.** *Science* 2020, **367**:301-305.

This very interesting paper shows that TPCs contribute to maintaining the volume of endocytic organelles. The authors find that in macrophages, removal of extracellular Na<sup>+</sup> or chemical inhibition, or knockout, of TPCs prevents the shrinkage of macropinosomes and associated tubules formed during macropinocytosis. The authors conclude that Na<sup>+</sup> flux through TPCs and associated osmotic movement of water across the organellar membrane is important for volume 'resolution'.