



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Original Article

A familial cluster of COVID-19 infection in a northern Chinese region

Cai-xia Liu^{a,1}, Zhi-hui Liu^{b,1}, Lu-yao Sun^{a,1}, Kai-yu Zhang^{a,*}, Yi-zhe Sun^{b,**}^a Department of Infectious Diseases, the First Hospital of Jilin University, Changchun, 130021, China^b Department of Infectious Diseases, Gongzhuling Central Hospital, Jilin Province, 136100, China

ARTICLE INFO

Article history:

Received 28 April 2021

Received in revised form 7 July 2021

Accepted 10 July 2021

Keywords:

Coronavirus disease 2019

Family cluster

Incubation period

Quarantine

ABSTRACT

Objective: Currently, coronavirus disease 2019 (COVID-19) has spread worldwide and become a global health concern. Here, we report a familial cluster of six patients infected with severe acute respiratory coronavirus 2 (SARS-CoV-2) in a northern Chinese region and share our local experience with regard the control of COVID-19.

Methods: The demographic data, clinical features, laboratory examinations, and epidemiological characteristics of enrolled cases were collected and analyzed. Two family members (Cases 1 and 2) had Hubei exposure history and were admitted to the hospital with a confirmed diagnosis of COVID-19; eight familial members who had contact with them during the incubation period underwent quarantine in a hospital.

We closely followed up all the family members and analyzed their clinical outcome.

Results: Case 3 had negative SARS-CoV-2 reverse transcription-polymerase chain reaction (RT-PCR) results but was suspected to have COVID-19 because of radiographic abnormalities. Cases 4 and 5 developed symptomatic COVID-19. Case 6 was considered an asymptomatic carrier as his SARS-CoV-2 RT-PCR result was positive. The other four family members with close contacts to COVID-19 patients had no evidence of SARS-CoV-2 infection.

Conclusions: Our findings suggest that COVID-19 has infectivity during the incubation period and preventive quarantine is effective for controlling an outbreak of COVID-19 infection.

© 2021 The Author(s). Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

In December 2019, a number of cases with viral pneumonia of unknown cause were reported in Wuhan, Hubei Province, China. On February 8, 2020, the disease was named as “Novel Coronavirus Pneumonia (NCP)” by the health administration of China [1]. Subsequently, the disease was recognized as coronavirus disease 2019 (COVID-19) and declared an epidemic on March 11, 2020 by the World Health Organization (WHO) [2]. As of June 30, 2021,

about 180 million cases of confirmed COVID-19 infection have been reported by the WHO and more than 3.9 million individuals have died from the disease. The causative agent of this viral pneumonia of unknown cause, which was provisionally named 2019 novel coronavirus (2019-nCoV), was ultimately designated severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) based on phylogeny, taxonomy, and established practice.

Currently, accumulating evidence has indicated person-to-person transmission [3]. In most cases, respiratory and direct contact with contaminated surfaces is the two main infection routes of COVID-19 [4]. Recently, COVID-19 nucleic acid was detected in fecal samples, and in an epidemiological and clinical investigation of childhood COVID-19, which raised the possibility of fecal-oral transmission [5,6]. Jilin Province is located at the north-east of China and is about 2000 miles from Hubei Province. In the early stage of the COVID-19 epidemic, imported infection was the most common source in Jilin Province. However, in the later stage, sporadic clusters of SARS-CoV-2 infection have become a major source of the epidemic. According to the Jilin Centers for Disease Control and Prevention report, a total of 93 confirmed COVID-19 cases were reported during the epidemic period. Among them, 54 cases (60%) can be interpreted as clustering SARS-CoV-2 infection.

Abbreviations: COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory coronavirus 2; NCP, Novel Coronavirus Pneumonia; RT-PCR, reverse transcription-polymerase chain reaction; WHO, World Health Organization; 2019-nCoV, 2019 novel coronavirus; WBCs, white blood cells; CRP, C-reactive protein; ESR, erythrocyte sedimentation rate; CT, Computed tomography; ACE2, angiotensin-converting enzyme 2.

* Corresponding author at: Department of Infectious Diseases, the First Hospital of Jilin University, No. 1 Xinmin Street, Changchun, 130021, China.

** Corresponding author.

E-mail addresses: kaiyu@jlu.edu.cn (K.-y. Zhang), jlsyz4420@163.com

(Y.-z. Sun).

¹ These authors contributed equally to this work.

<https://doi.org/10.1016/j.jiph.2021.07.011>

1876-0341/© 2021 The Author(s). Published by Elsevier Ltd on behalf of King Saud Bin Abdulaziz University for Health Sciences. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

To better understand epidemiological characteristics of COVID-19 and the evolution of clustering infection will be helpful in controlling this pandemic.

Hence, we report a familial cluster of COVID-19 infection in Jilin, a northern Chinese region, and share our experience in controlling COVID-19. Our findings help to highlight the importance of quarantine measures for the control of COVID-19.

Materials and methods

Ethics

The study protocol adhered to the declaration of Helsinki and was approved by the ethic committee of the Gongzhuling Central Hospital of Jilin Province (Jilin, China). Written informed consent was obtained for this prospective study.

Patients

Between January 20, 2020 and February 23, 2020, a familial cluster of six patients infected with SARS-CoV-2 and four close contacts were included for analysis. The demographic data, clinical features, laboratory examinations, and epidemiological characteristics of enrolled cases were collected and analyzed.

Diagnostic criteria

COVID-19 pneumonia was diagnosed according to the Diagnosis and Treatment Guideline for New Coronavirus Pneumonia (7th edition), issued by the National Health Commission of China [7]. All patients with COVID-19 pneumonia were confirmed by performing reverse transcription-polymerase chain reaction (RT-PCR) on their samples (e.g., pharyngeal swab and fecal sample) for SARS-CoV-2. A cluster of SARS-CoV-2 infection was defined as two or more cases reported from the same address (e.g., family, school, and office) and diagnosed within a 2-week interval.

Results

Table 1 shows the clinical and epidemiological characteristics of confirmed and suspected COVID-19 cases. Timeline of exposure and chronology of symptom onset in the family cluster of SARS-CoV-2 infection are depicted in Fig. 1. The relationship of relatives to cases is shown in Fig. 2.

Index cases

Case 1 was a 50-year-old woman, who lived in Hubei Province. On January 16, 2020, she took the train, returned from Xiantao (Hubei Province), and stopped in Wuhan (Hubei Province) before arriving in Gongzhuling (Jilin Province). On January 25, she had a fever (37.8 °C). Two days later, she was admitted to our hospital for fever, cough, and sore throat. RT-PCR for the detection of SARS-CoV-2 was performed and COVID-19 was confirmed. Laboratory examination revealed decreased levels of white blood cells (WBCs) ($2.15 \times 10^9/L$, reference: $3.5\text{--}9.5 \times 10^9/L$), lymphocytes ($0.46 \times 10^9/L$, reference: $1.1\text{--}3.2 \times 10^9/L$), and neutrophils ($1.64 \times 10^9/L$, reference: $1.8\text{--}6.3 \times 10^9/L$); and increased levels of C-reactive protein (CRP) (>10 mg/L, reference: 2–10 mg/L; Table 1) and erythrocyte sedimentation rate (ESR) (55 mm/h, reference: 0–20 mm/h). Computed tomography (CT) images showed patchy ground-glass opacities in the lungs (Fig. 3A). Case 2 was the husband of Case 1; he arrived in Gongzhuling together with Case 1. On January 30, 2020, he had a temporary fever. Chest CT showed lung infection; the radiological findings are shown in Fig. 3B. One

day later, he was diagnosed with COVID-19 because of the positive SARS-CoV-2 RT-PCR results. He was given an anti-viral and also treated for his symptoms. On March 2, his clinical condition significantly improved, as revealed by CT findings.

A familial cluster of secondary cases

After Case 1 was confirmed to have COVID-19 on January 27, 2020, close contacts were identified, immediately isolated, and screened for SARS-CoV-2 in the fever clinic of our hospital. Respiratory samples were collected and tested using RT-PCR for detection of SARS-CoV-2. Case 3 had dinner with Cases 1, 2, and 4 on January 20. On January 31, she became ill with fatigue and cough. Chest CT scan was performed and multiple patchy shadows were observed (Fig. 3C). Although RT-PCR for detection of SARS-CoV-2 was performed several times, the results were all negative. Due to the COVID-19 contact history, clinical symptoms, and abnormal radiographic findings, the patient was diagnosed with suspected COVID-19. Case 4 presented with mild symptoms of fatigue. On February 2, COVID-19 was confirmed by RT-PCR. Chest CT revealed multiple patchy, flocculent, fuzzy, and high-density shadows in the lungs (Fig. 3D). Case 5 had contact with Cases 1 and 2 on January 18 and 23, respectively. She became ill on February 4, presenting with fever, transcutaneous oxygen saturation 86% at rest, and patchy consolidation on CT images (Fig. 3E). Subsequently, COVID-19 was confirmed by RT-PCR. Case 6 had contact with Cases 1 and 2 on January 19, and no symptoms were observed. On February 6, pharyngeal swabs, including fecal samples, were tested for SARS-CoV-2 and positive results were confirmed. In addition, a chest CT scan was done and appeared normal. Hence, the patient was considered to have asymptomatic infection. The remaining four cases, who had close contact with the two index cases, were not found to have COVID-19. The detailed investigation was follows. C1D (daughter of Cases 1 and 2) lived with Case 1 between January 17 and 19, reporting no symptoms, and had persistently negative RT-PCR for SARS-CoV-2, and normal radiographic appearance. C5H (husband of Case 5) invited Cases 1 and 2 for a dinner at the home and was isolated on January 28. He was also considered to not have infection, based on the combination of clinical characteristics, SARS-CoV-2 RT-PCR results, and chest CT scan. Similarly, C5S (son of Case 3, husband of C1D) and C6H (husband of Case 6) were also found to not have SARS-CoV-2 infection. The four cases were all followed until March 15. They were tested for antibodies (IgG and IgM) against SARS-CoV-2 and negative results were observed in all cases. In addition, the familial cluster of SARS-CoV-2 infection was all reported at home or at our hospital and no other COVID-19 patients in the district were reported. Therefore, the early quarantine of close contacts of index patients, especially within a family, plays an important role in preventing the transmission of COVID-19.

Discussion and conclusions

Due to the epidemiological features of the infection, the case series described in our study were considered a familial cluster of COVID-19 infection. The cluster of COVID-19 infection, including two imported cases and four secondary cases, occurred within 12 days. In this cluster, our findings suggested that: (1) during the incubation period, COVID-19 patients could infect close contacts; (2) the early quarantine for those with close contact with COVID-19 patients remains a useful tool for controlling the epidemic; and (3) asymptomatic infection may be more common than previously thought. In fact, in the later stage of COVID-19 epidemic in Jilin, sporadic clusters of SARS-CoV-2 infection became a major source. Therefore, knowing the characteristics of COVID-19 clustering is helpful for guiding the controlling measures.

Table 1
Clinical and epidemiological characteristics of confirmed and suspected COVID-19 cases.

	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6
Sex	Female	Male	Female	Male	Female	Female
Age (years)	50	57	54	28	50	51
Native place	Hubei Province	Jilin Province	Jilin Province	Jilin Province	Jilin Province	Jilin Province
Occupation	Building worker	Building worker	Housewife	Worker	Housewife	Housewife
Types of COVID-19	Common	Common	Suspected	Common	Severe	Asymptomatic infection
Medical history	Autoimmune hemolytic anemia	No	No	No	Encephalorrhagia	No
Incubation period (estimated, days)	9	14	11	13	17	17
Clinical characteristics						
Abnormal chest CT findings	Yes	Yes	Yes	Yes	Yes	No
Symptoms						
Fever	Yes	Yes	No	No	Yes	No
Cough	Yes	No	Yes	No	No	No
Fatigue	No	No	Yes	Yes	No	No
Rhinobyon	No	No	No	No	No	No
Sneeze	No	No	No	No	No	No
Sore throat	Yes	No	No	No	No	No
Chest pain	No	No	No	No	No	No
Diarrhea	No	No	No	No	No	No
Laboratory examinations						
White blood cell ($\times 10^9/L$)	2.15	5.16	4.48	3.82	6.69	4.73
Neutrophil ($\times 10^9/L$)	1.64	3.03	2.11	1.76	5.61	3.05
Lymphocyte ($\times 10^9/L$)	0.46	3.18	1.94	1.82	0.78	1.47
Platelet ($\times 10^9/L$)	141	171	257	245	215	377
C-reactive protein (mg/L)	>10	<10	ND	ND	70.8	ND
Erythrocyte sedimentation rate (mm/h)	55	ND	ND	ND	10	ND
COVID-19 RT-PCR						
Throat swab (conversion)	+, Jan. 27	+, Jan. 31	-	+, Feb. 2	+, Feb. 4	+Feb. 6
Throat swab (reversion)	-, Feb. 3	-, Feb. 26	-	-, Feb. 3	-, Feb. 9	-, Feb. 14
Viral shedding period (Throat swab) (days)	7	26	-	1	5	8
Fecal samples	-	-	-	-	ND	+, Feb. 11
Antibody assays against COVID-19						
Date	Mar. 15	Mar. 26	Mar. 23	ND	March 15	ND
IgG	+	-	-	-	+	-
IgM	+	-	-	-	+	-




Fig. 1. Timeline of exposure and chronology of symptom onset in the family cluster of SARS-CoV-2 infection.




Fig. 2. The relationship of relatives to the index cases (Cases 1 and 2).




Fig. 3. Chest scan of five cases. Case 1: patchy ground-glass opacities in the lungs (A); Case 2: abnormal findings with suspected lung infection (B); Case 3: multiple patchy shadows (C); Case 4: multiple patchy, flocculent, fuzzy, and high-density shadows in the lungs (D); Case 5: patchy consolidation (E).

Respiratory and contact routes are proven transmission mechanisms for COVID-19 infection. However, in a recent study, SARS-CoV-2 RNA was detected in the fecal samples of COVID-19 patients, indicating the possibility of transmission via the fecal-oral route [6,8]. In our study, respiratory and contact should be considered as the two main transmission mechanisms responsible for this cluster of infections. Moreover, an asymptomatic carrier was also suspected of being a source for transmission of COVID-19 [9], and this key point was proved by Zhang et al. [10]. Although these new findings make the prevention and control of COVID-19 more complicated, they will lead to improved and more effective strategies. To avoid acquiring COVID-19, specific attention should be paid to factors such as environmental, direct contact, and social distance [7,11].

Person-to-person transmission has been confirmed. It is unclear if during the incubation period, the virus is infective. A previous study by Yu et al. [12] showed possible transmission of COVID-19 during the incubation period. Similarly, our data also supported that the virus remains infective during the incubation period. In this study, before the onset of symptoms of Case 1 (January 25), Cases 3 and 4 only had dinner with Case 1, as well as a dinner party, and Case 5 had another social contact with Case 1. Three cases were all infected with COVID-19. This special characteristic mentioned above is different from the transmission of SARS infection. In our study, the incubation period was relatively long and ranged from 9 to 17 days. Remarkably, although the incubation period of Case 1 was estimated at 9 days, the possibility remained of it being extended to longer than 9 days because of her residential location (Xiantao, Hubei). The incubation periods of Cases 5 and 6 were estimated from the first contact with Case 1, with the longest period being 17 days. Likewise, the incubation period of a secondary case has been reported over 14 days [13], even reaching up to 20 days [14]. Hence, a longer quarantine period, such as 2 or 3 weeks, may be required.

Usually, patients with COVID-19 present with fever, cough, or lung infiltrates [15]. In some cases, they may also have mild respiratory symptoms. More recently, asymptomatic infection has been reported. In Japan, the prevalence of asymptomatic infection is estimated at 41.6% (95% confidence interval: 16.7%, 66.7%) among COVID-19-infected individuals [16]. However, in China, a study of 72,314 COVID-19 patients showed that asymptomatic carriers comprised 1.2% of all COVID-19 cases [17]. The difference in the prevalence of asymptomatic infection may be caused by sample size estimated and different subjects included between the two studies (high-risk vs. confirmed). In our study, due to positive SARS-CoV-2 RT-PCR results from fecal and pharyngeal samples, Case 6 was considered asymptomatic infection, the time of viral shedding was 8 days, longer than the time for Cases 4 and 5. We believe that asymptomatic infection is common. A similar result was reported in another familiar cluster of COVID-19 infection. Due to SARS-CoV-2 RT-PCR (+) and IgM assays (+), four cases without clinical symptoms were considered asymptomatic infection [18]. Hence, on one hand, good personal hygiene is emphasized; on the other hand, due to no clinical symptoms and normal radiography, asymptomatic infection could be easily neglected. Hence, more attention should be given to improving the diagnostics and management of asymptomatic carriers, this would improve the prevention and control of the COVID-19 epidemic [19].

For the diagnosis of COVID-19 patients, the sensitivities of IgM and IgG assays were 77.3%, and 83.3%, respectively [20], and RT-PCR for detection of COVID-19 has a lower sensitivity of 35% [21]. It was then concluded that a significant proportion of COVID-19 would be diagnosed as suspected cases. Case 3, as a suspected COVID-19 case, presented with mild respiratory symptoms and abnormal radiography. Although RT-PCR for SARS-CoV-2 detection was performed several times, the case was not confirmed. However,

due to contact history with the index case, Case 3 was identified, isolated, and treated as a suspected case. Fortunately, until now, several new assays, such as an automated chemiluminescent immunoassay, a reverse transcription loop-mediated isothermal amplification assay, and antibody detection assay for the diagnosis of COVID-19, have been evaluated and good performance has been confirmed [20,22–25]. These new assays may improve the diagnostic dilemma of current assays for COVID-19.

Accumulated evidence suggests that SARS-CoV-2 is more infectious than SARS-CoV and MERS-CoV [18]. However, this study of a familiar cluster of COVID-19 infection demonstrated a differential susceptibility. Although C1D and C5S lived with Cases 1 and 2, it remains to be found without COVID-19 infection. The result suggests a difference in the susceptibility to COVID-19 among individuals. The SARS-CoV-2 has preferential tropism to human airway epithelial cells through the same cellular receptor as that for SARS, angiotensin-converting enzyme 2 (ACE2), which is a central body receptor for the surface glycoprotein S of the virus [26]. Therefore, the down regulation of ACE2 expression is thought to explain the lower susceptibility to COVID-19. In addition, the differential susceptibility suggested by the study implicated that interventions, such as vaccine trials and preventive measures, for the COVID-19 control should be evaluated their feasibility.

China is one of the countries hard-hit by COVID-19 at the beginning of its outbreak. After the impose of strict quarantine measures, we successfully stopped the exacerbation of the condition [27]. After that, COVID-19 only occurred as sporadic cases across China, in response to which, quarantine and close follow up of individuals with close contact with COVID-19 patients were implemented. All these measures play an critical role in stifle the wide spread of COVID-19. Currently, vaccination against COVID-19 is widely carried out all over the world, which would decrease the incidence of COVID-19 [28,29]. Sporadic cases may be more common during the vaccination era. Effective quarantine measures can help to accelerate the eradication of the COVID-19.

In terms of the control of sporadic clusters of COVID-19 infection, several recommendations can be made as a result of the study. (1) Cases who are exposed to COVID-19 patients during the incubation period should also be included as close contacts. (2) The diagnosis of COVID-19 infection could not reasonably rely on RT-PCR and antibody detection and all suspected cases should be treated as confirmed patients. Moreover, further research should be performed to look for a rapid, accurate, economic diagnostic assay for COVID-19. (3) A differential susceptibility to COVID-19 between individuals exists. Further investigation for the mechanism of the difference may be helpful to improve the management of the susceptible population. (4) Finally, it is better that all close contacts are isolated and observed in a centralized setting [30]. Otherwise, home isolation and observation are required and a community supervision mechanism is needed.

Our study enhance our knowledge with regard the epidemiology characteristics of COVID-19 especially the characteristics of clustering infection, and highlighted the importance of quarantine for the COVID-19. It is true that transmission of COVID-19 had been well characterized by previous studies, and the information in our study is a little bit dated. However, few case reports emphasize the importance of preventive quarantine for COVID-19 control, especially when sporadic cases remain prevalent. We believe our findings will help to better guide strategy of COVID-19 controlling and accelerate the termination of pandemic.

Consent for publication

All data published here are under the consent for publication.

Availability of data and material

The datasets generated and analyzed during the present study are available from the corresponding author on reasonable request.

Competing interests

None declared.

Ethics approval and consent to participate

The study protocol adhered to the declaration of Helsinki and was approved by the ethic committee of the Gongzhuling Central Hospital of Jilin Province (Jilin, China). Written informed consent was obtained for this prospective study.

Funding

This work was supported by the Emergency Scientific Research Fund for the Prevention and Control of Novel Coronavirus Pneumonia of the Science and Technology Department of Jilin Province [grant number 20200901008SF].

Authors contributions

Kai-yu Zhang and Cai-xia Liu designed/performed most of the investigation, data analysis and wrote the manuscript; Yi-zhe Sun and Zhi-hui Liu provided the detail information of the cases; Lu-yao Sun contributed to interpretation of the data and analyses. All of the authors have read and approved the manuscript.

Acknowledgements

None.

References

- China NHCoPsRo. Notification of novel coronavirus pneumonia temporarily named by the national health and Health Commission. <http://www.nhc.gov.cn/xcs/zhengcwj/202002/18c1bb43965a4492907957875de02ae7.shtml>. 8 February 2020.
- Organization WH. Novel Coronavirus(2019-nCoV) Situation Report-22. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200211-sitrep-22-ncov.pdf?sfvrsn=fb6d49b1_2. 11 February 2020.
- Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern. *Lancet* 2020;395:470–3, [http://dx.doi.org/10.1016/S0140-6736\(20\)30185-9](http://dx.doi.org/10.1016/S0140-6736(20)30185-9), 2020/01/28.
- Rothan HA, Byrareddy SN. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J Autoimmun* 2020;109:102433, <http://dx.doi.org/10.1016/j.jaut.2020.102433>, 2020/03/03.
- Holshue ML, DeBolt C, Lindquist S, Lofy KH, Wiesman J, Bruce H, et al. Washington State 2019-nCoV Case Investigation Team. First case of 2019 novel coronavirus in the United States. *N Engl J Med* 2020;382:929–36, <http://dx.doi.org/10.1056/NEJMoa2001191>.
- Xu Y, Li X, Zhu B, Liang H, Fang C, Gong Y, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nat Med* 2020;26:502–5, <http://dx.doi.org/10.1038/s41591-020-0817-4>, 2020/04/15.
- China NHCo. New coronavirus pneumonia prevention and control program (7th edn). 2020 [cited 2020 7th, May]; Available from: <http://www.nhc.gov.cn/yzygj/s7653p/202003/46c9294a7dfe4cef80dc7f5912eb1989.shtml>.
- Jiang X, Luo M, Zou Z, Wang X, Chen X, Qiu J. Asymptomatic SARS-CoV-2 infected case with viral detection positive in stool but negative in nasopharyngeal samples lasts for 42 days. *J Med Virol* 2020;92:1807–9, <http://dx.doi.org/10.1002/jmv.25941>, 2020/04/25.
- Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, et al. Presumed asymptomatic carrier transmission of COVID-19. *JAMA* 2020;323:1406–7, <http://dx.doi.org/10.1001/jama.2020.2565>, 2020/02/23.
- Zhang J, Tian S, Lou J, Chen Y. Familial cluster of COVID-19 infection from an asymptomatic. *Crit Care* 2020;24:119, <http://dx.doi.org/10.1186/s13054-020-2817-7>, 2020/03/30.
- Special Expert Group for Control of the Epidemic of Novel Coronavirus Pneumonia of the Chinese Preventive Medicine Association. An update on the epidemiological characteristics of novel coronavirus pneumonia (COVID-19). *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:139–44, <http://dx.doi.org/10.3760/cma.j.issn.0254-6450.2020.02.002>, 2020/02/15.
- Yu P, Zhu J, Zhang Z, Han Y. A familial cluster of infection associated with the 2019 novel coronavirus indicating potential person-to-person transmission during the incubation period. *J Infect Dis* 2020;221:1757–61, <http://dx.doi.org/10.1093/infdis/jiaa077>.
- Guan Q, Liu M, Zhuang YJ, Yuan Y, Wang SS, Li J, et al. Epidemiological investigation of a family clustering of COVID-19. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:629–33, <http://dx.doi.org/10.3760/cma.j.cn112338-20200223-00152>, 2020/03/10.
- Yang HY, Xu J, Li Y, Liang X, Jin YF, Chen SY, et al. The preliminary analysis on the characteristics of the cluster for the Corona Virus Disease. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:623–8, <http://dx.doi.org/10.3760/cma.j.cn112338-20200223-00153>, 2020/03/09.
- Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506, [http://dx.doi.org/10.1016/S0140-6736\(20\)30183-5](http://dx.doi.org/10.1016/S0140-6736(20)30183-5), 2020/01/28.
- Nishiura H, Kobayashi T, Miyama T, Suzuki A, Jung SM, Hayashi K, et al. Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-19). *Int J Infect Dis* 2020;94:154–5, <http://dx.doi.org/10.1016/j.ijid.2020.03.020>, 2020/03/18.
- Epidemiology Working Group for NCIP Epidemic Response, Chinese Center for Disease Control and Prevention. [The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:145–51, <http://dx.doi.org/10.3760/cma.j.issn.0254-6450.2020.02.003>, 2020/02/18.
- Bai SL, Wang JY, Zhou YQ, Yu DS, Gao XM, Li LL, et al. Analysis of the first cluster of cases in a family of novel coronavirus pneumonia in Gansu Province. *Zhonghua Yu Fang Yi Xue Za Zhi* 2020;54:E005, <http://dx.doi.org/10.3760/cma.j.issn.0253-9624.2020.0005>, 2020/02/18.
- Epidemiology Working Group, Strategy and Policy Working Group for NCIP Epidemic Response, Chinese Center for Disease Control and Prevention. Cluster investigation technical guideline for the 2019 Novel Coronavirus Pneumonia (COVID-19), China (1st Trial Version). *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:293–5, <http://dx.doi.org/10.3760/cma.j.issn.0254-6450.2020.03.001>, 2020/02/18.
- Xiang F, Wang X, He X, Peng Z, Yang B, Zhang J, et al. Antibody detection and dynamic characteristics in patients with COVID-19. *Clin Infect Dis* 2020;71:1930–4, <http://dx.doi.org/10.1093/cid/ciaa461>, 2020/04/20.
- Mardani R, Ahmadi Vasmehjani A, Zali F, Gholami A, Mousavi Nasab SD, Kaghazian H, et al. Laboratory parameters in detection of COVID-19 patients with positive RT-PCR; a diagnostic accuracy study. *Arch Acad Emerg Med* 2020;8:e43, 2020/04/08.
- Yan C, Cui J, Huang L, Du B, Chen L, Xue G, et al. Rapid and visual detection of 2019 novel coronavirus (SARS-CoV-2) by a reverse transcription loop-mediated isothermal amplification assay. *Clin Microbiol Infect* 2020;26:773–9, <http://dx.doi.org/10.1016/j.cmi.2020.04.001>, 2020/04/11.
- Yang AP, Liu JP, Tao WQ, Li HM. The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients. *Int Immunopharmacol* 2020;84:106504, <http://dx.doi.org/10.1016/j.intimp.2020.106504>, 2020/04/19.
- Baek YH, Um J, Antigua KJC, Park JH, Kim Y, Oh S, et al. Development of a reverse transcription-loop-mediated isothermal amplification as a rapid early-detection method for novel SARS-CoV-2. *Emerg Microbes Infect* 2020;9:998–1007, <http://dx.doi.org/10.1080/22221751.2020.1756698>, 2020/04/21.
- Infantino M, Grossi V, Lari B, Bambi R, Perri A, Manneschi M, et al. Diagnostic accuracy of an automated chemiluminescent immunoassay for anti-SARS-CoV-2 IgM and IgG antibodies: an Italian experience. *J Med Virol* 2020;92:1671–5, <http://dx.doi.org/10.1002/jmv.25932>, 2020/04/25.
- Munster VJ, Koopmans M, van Doremalen N, van Riel D, de Wit E. A novel coronavirus emerging in China – key questions for impact assessment. *N Engl J Med* 2020;382:692–4, <http://dx.doi.org/10.1056/NEJMp2000929>, 2020/01/25.
- Peng F, Tu L, Yang Y, Hu P, Wang R, Hu Q, et al. Management and treatment of COVID-19: the Chinese experience. *Can J Cardiol* 2020;36:915–30, <http://dx.doi.org/10.1016/j.cjca.2020.04.010>, 2020/05/23.
- Chung JY, Thone MN, Kwon YJ. COVID-19 vaccines: the status and perspectives in delivery points of view. *Adv Drug Deliv Rev* 2021;170:1–25, <http://dx.doi.org/10.1016/j.addr.2020.12.011>, 2020/12/29.
- Frederiksen LSF, Zhang Y, Foged C, Thakur A. The long road toward COVID-19 herd immunity: vaccine platform technologies and mass immunization strategies. *Front Immunol* 2020;11:1817, <http://dx.doi.org/10.3389/fimmu.2020.01817>, 2020/08/15.
- Qiu YY, Wang SQ, Wang XL, Lu WX, Qiao D, Li JB, et al. Epidemiological analysis on a family cluster of COVID-19. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41:494–7, <http://dx.doi.org/10.3760/cma.j.cn112338-20200221-00147>, 2020/03/07.