

Intravitreal homocysteine-thiolactone injection leads to the degeneration of multiple retinal cells, including photoreceptors

Han-Hsin Chang,^{1,2} David Pei-Cheng Lin,^{2,3,4} Ying-Shan Chen,⁵ Hsiang-Jui Liu,⁶ Wei Lin,⁷ Zih-Jay Tsao,^{1,3} Mei-Ching Teng,⁸ Bo-Yie Chen^{2,3}

¹School of Nutrition, Chung Shan Medical University, Taichung, Taiwan, ROC; ²Department of Ophthalmology, Chung Shan Medical University Hospital, Taichung, Taiwan, ROC; ³School of Optometry, Chung Shan Medical University, Taichung, Taiwan, ROC; ⁴School of Medical Laboratory and Biotechnology, Chung Shan Medical University, Taichung, Taiwan, ROC; ⁵Department of Ophthalmology, Hsinchu Cathay General Hospital, Hsinchu, Taiwan, ROC; ⁶Department of Optometry, Jen-Teh Junior College of Medicine, Nursing and Management, Miaoli, Taiwan, ROC; ⁷Department of Optometry, Central Taiwan University of Science and Technology, Taichung, Taiwan, ROC; ⁸Department of Ophthalmology, Kaohsiung Chang Gung Memorial Hospital and Chang Gung University College of Medicine, Kaohsiung, Taiwan, ROC

Purpose: Hyperhomocysteinemia is known to cause degeneration of retinal ganglion cells, but its influence on photoreceptors remains largely unknown. In particular, the role of homocysteine-thiolactone (Hcy-T)—the physiologic metabolite of homocysteine that has been proven to be more cytotoxic than homocysteine itself—as a factor that causes retinopathy, has not been defined. This study aimed to investigate the toxic effects of excessive Hcy-T in a mouse model.

Methods: A total of 60 six-week-old female ICR mice were used in this study. The mice were divided into 3 experimental groups and 2 control groups. The mice in the experimental groups were subjected to intravitreal injections of Hcy-T to reach final estimated intravitreal concentrations at 5, 25, and 200 μ M, respectively. Mice without injection (blank) and with 0.9 NaCl injections (sham injection) were used as controls. The mice with 200 μ M Hcy-T were sacrificed at days 7, 15, 45, and 90 after injection and the mice with 5 or 25 μ M Hcy-T were sacrificed at day 90, with the controls sacrificed at day 15 or 90 for comparison. Semi-quantitative dot-blot analysis was performed for confirmation of retinal homocysteinylation. The mouse retinas were evaluated microscopically, with the thickness of total and specific retinal layers determined. Immunohistochemical analysis was performed and the labeled cells were quantified to determine the effects of excessive Hcy-T on specific retinal cells.

Results: Dose-dependent retinal homocysteinylation after Hcy-T injection was confirmed. The homocysteinylation was localized in the outer and inner segments of photoreceptors and the ganglion cell layer (GCL). Retinal cell degenerations were found in the GCL, inner nuclear layer, and outer nuclear layer at day 90 after 200 μ M Hcy-T injection. Significant thickness reduction was found in the total retina, outer nuclear layer, and the outer and inner segment layers. A trend of thickness reduction was also found in the GCL and inner nuclear layer, although this was not statistically significant. The rhodopsin⁺ photoreceptors and the calbindin⁺ horizontal cells were significantly reduced at day 15, and were nearly ablated at day 90 after 200 μ M Hcy-T injection ($p < 0.001$ for both day 15 and day 90), which was not seen in the sham injection controls. The Chx-10⁺ or the Islet-1⁺ bipolar cells and the Pax-6⁺ amacrine cells were severely misarranged at day 90, but no significant reduction was found for both cell types. The GFAP⁺ Müller cells were activated at day 15, but were not significantly increased at day 90 after the injection.

Conclusions: Excessive retinal homocysteinylation by Hcy-T, a condition of hyperhomocysteinemia, could lead to degeneration of photoreceptors, which might lead to retinopathies associated with severe hyperhomocysteinemia or diabetes mellitus. Received: April 1, 2011 Accepted: July 14, 2011

Homocysteine (Hcy) is a sulfur-containing amino acid derived from methionine metabolism as an intermediate metabolite. Clinically, the normal range of total plasma Hcy falls between 5 to 15 μ mol/l. Patients with Hcy concentrations greater than 15 μ mol/l are considered to have hyperhomocysteinemia (hHcy) and the condition is commonly associated with diabetes mellitus [1]. In recent

years, hHcy has been implicated in systematic hypertension, stroke, and other cardiovascular diseases [2,3]. In the ocular system, many lines of evidence indicated hHcy as a risk factor in a variety of diseases, including retinal arteriosclerosis [4], glaucoma [5,6], exudative age-related macular degeneration [7], and macular and optic atrophy due to retinal vascular occlusion or non-arteritic ischemic optic neuropathy [8–10].

Notably, most hHcy-induced retinopathies are indirectly due to retinal arteriosclerosis and thrombosis. At the cellular level, previous studies have indicated involvement of hHcy in vascular endothelial damage and proliferation of vascular smooth muscle cells [11,12]. At the molecular level, hHcy was

Correspondence to: Bo-Yie Chen, School of Optometry, Chung Shan Medical University, No. 110, Chien-Kuo N. Rd., Taichung 402, Taiwan, ROC Phone: +886-4-24730022 ext. 17230; FAX: +886-4-2324-8131; email: boychen@csmu.edu.tw

reported to be associated with elevated lipid peroxidation, endoplasmic reticulum stress, DNA methylation, and collagen accumulation [13,14]. Direct Hcy attack on retina has also been reported. Ganapathy et al. [15] used the cystathionine beta-synthase (cbs) mutant mouse to study the effects of endogenous homocysteine elevation on the retina. They found an approximate sevenfold elevation of Hcy in the cbs^{-/-} retina. As a result of Hcy elevation, distinct alterations were observed in the ganglion, inner plexiform, inner nuclear, and epithelial layers in retinas of cbs^{-/-} and 1-year-old cbs^{+/-} mice. High levels of Hcy were concomitantly located in these retinal layers, particularly the ganglion cell layer (GCL), suggesting direct attack by Hcy [15,16]. Another study by Lee et al. [17] showed that short-term hHcy-induced oxidative stress can activate retinal glial cells and increase VEGF expression in the retina.

The accumulation of Hcy in human cells depends on the activities of methionine synthase, cystathionine β -synthase, and methionyl-tRNA synthetase. Hcy can be metabolized by methionine synthase through its activity to convert Hcy into methionine. Hcy can also be converted into cysteine by cystathionine β -synthase. These two pathways deplete intracellular Hcy in a healthy manner. A third metabolic pathway for Hcy is its conversion into homocysteine -thiolactone (Hcy-T) by methionyl-tRNA synthetase, which occurs under normal physiologic reactions in all human cell types investigated [18]. Both Hcy and Hcy-T can attack proteins through either S-homocysteinylation or N-homocysteinylation [2]. S-homocysteinylation is undertaken by Hcy through attack on the Cys-SH group of proteins, while N-homocysteinylation involves acylation of Lys ϵ -amino group by the activated carboxyl group of Hcy-T [1,19]. Both types of homocysteinylation will cause protein damage and will lead to impairment of protein function [2]. As such, retinal proteins are liable to be attacked by homocysteinylation under the influence of hHcy. Besides, Hcy-T has been reported to possess stronger cytotoxicity and pro-inflammatory properties than Hcy [20]. Therefore, the third metabolic pathway for Hcy will potentially lead to more severe retinopathies. Yet, there has been no report regarding the direct role of Hcy-T in causing retinopathies.

Another interesting, important, and unanswered question is how hHcy can attack retina under in vivo conditions. There has so far been no report on hHcy-induced photoreceptor damage, although an earlier report indicated a non-detectable electroretinogram in a case of combined methylmalonic aciduria and homocystinuria [21]. Another report also showed some abnormalities on behavior tests assessing visual function in six cases of combined methylmalonic aciduria and homocystinuria [22]. Potentially, these clinical symptoms can be a result of damage to the photoreceptors and to the ganglion cells, under the influence of hHcy. Yet, there has been no data reported on this area.

In this study, we conducted intravitreal Hcy-T injections in a mouse model to investigate the direct role of Hcy-T in causing retinopathies. Following the injections, Hcy-T remained detected at day 90 in the retina in a dose-dependent manner, suggesting occurrence of homocysteinylation. The toxic level of intravitreal Hcy-T started at 25 μ M in this mouse model. The ganglion cell layer (GCL), inner nuclear layer (INL), outer nuclear layer (ONL), and the outer segment and inner segment (OS/IS) of photoreceptors were severely affected. With intravitreal Hcy-T level at 200 μ M, the ONL were all eradicated at day 90 after the injection. Total retinal thickness was reduced as compared to that of the blank or the sham injection controls. This reduction was also observed in the GCL, INL, ONL, and OS/IS layers, respectively. Immunohistochemical staining localized Hcy-T, mainly in the GCL and OS/IS layers, which persisted at day 90 after the injection. Photoreceptor and horizontal cells were the most affected as indicated by the disappearance of their specific markers. The bipolar, amacrine, and Müller glial cells were all disorganized. Our results indicate that excessive Hcy-T leads to more extensive retinal homocysteinylation than hHcy and most retinal cells, including photoreceptors, are damaged under the influence of excessive Hcy-T.

METHODS

Animals: A total of 60 six-week-old female ICR mice were used in this study. The mice were purchased from National Laboratory Animal Center (Taipei, Taiwan, ROC). The mice weighed about 25 g on arrival and were fed ad libitum and kept under standard conditions with a 12 h:12 h light-dark cycle. The mice were acclimatized and habituated to the laboratory for at least one week before the experiments. All animal maintenance and experiment protocols were reviewed and approved by the Animal Care and Use Committee of Chung Shan Medical University and were performed in agreement with the Association for Research in Vision and Ophthalmology (ARVO) Resolution on the Use of Animals in Research.

Study groups and intravitreal homocysteine-thiolactone injections: Prior to intravitreal injection, the mice were anesthetized with intraperitoneal injections of sodium pentobarbital (45 mg/kg bodyweight). Intravitreal injection procedures and criteria were applied following the details published by Moore [23]. Briefly, both eyes were injected using a 5 μ l Hamilton glass syringe (Hamilton Bonaduz AG, Bonaduz, Switzerland). The injections were made posterior to the iris and into the vitreous under a Nikon SMZ-645 dissection microscope (Nikon, Tokyo, Japan; Figure 1A). Caution was taken to avoid direct contact of the needle with the lens. The mice were injected intravitreally with either 0.9% NaCl (sham injection control) or Hcy-T (Sigma-Aldrich, St. Louis, MO) and some mice were left without injection to serve as blank controls (Table 1). The Hcy-T was prepared in 0.9% NaCl, pH 7.4. The final intravitreal




Figure 1. The preparation of intravitreal homocysteine-thiolactone (Hcy-T) injection is illustrated in **A** and retinal homocystinylation as confirmed by a semi-quantitative dot-blot analysis is shown in **B** and **C**. **A**: In each retinal section, the thickness measurement was made at two sites, located approximately 200 to 300 μm apart, on either side of the optic nerve. **B**: Dose-dependent retinal homocystinylation was observed on day 90 after injection. **C**: Quantification of relative intensity (pixels) on dot-blot membranes using an arbitrary unit of retinal homocystinylation as compared to that of the blank control. The sample size was 5 in all groups. The error bars indicate standard error of the means (SEMs).

TABLE 1. SUMMARY OF EXPERIMENTS

Numbers	Days after injection when sacrificed	Number of mice	Final intravitreal concentrations
1	90	8	Non-injection control (Blank control)
2	15	5	0.9% NaCl
3	90	8	0.9% NaCl
4	90	8	Homocysteine thiolactone (5 μM)
5	90	8	Homocysteine thiolactone (25 μM)
6	7	5	Homocysteine thiolactone (200 μM)
7	15	5	Homocysteine thiolactone (200 μM)
8	45	5	Homocysteine thiolactone (200 μM)
9	90	8	Homocysteine thiolactone (200 μM)

concentration of Hcy-T was calculated based on an assumed vitreal volume at 25 μl per eye. Hcy-T was prepared in three stock solutions of 0.25, 1.25, and 10 mM. The injection of 0.5 μl into the vitreous gave a final estimated intravitreal concentration of 5, 25, and 200 μM , respectively. Since the purpose of this study was to assess the consequences of excessive levels of Hcy-T on the retinal cells, the intravitreal Hcy-T concentrations tested in this study were set above a physiologic range or were equivalent to those of the retinopathogenic conditions [24,25].

Semi-quantitative dot-blot analysis for retinal homocystinylation: Dot-blot analysis was used to test the levels of retinal homocystinylation after intravitreal Hcy-T injections. After the mice were euthanized, their eyes were extracted, followed by removal of the cornea and the lens tissues and separation of the retinas from the eye cups. The retinas were homogenized in radioimmunoprecipitation (RIPA) buffer (Upstate Biotechnology, Lake Placid, NY) containing a protease inhibitor cocktail and were then

centrifuged at 10,000 \times g for 15 min at 4 $^{\circ}\text{C}$. The supernatants were collected and assayed for their protein levels with a Bio-Rad Protein Assay kit (Life Science Research, Hercules, CA) using BSA as standard. Equal aliquots (30 μg) of the retinal proteins were applied to a 96-well dot blot apparatus (Life Science Research) and were then transferred to a nitrocellulose membrane by vacuum filtration. The preparation was immersed in 10% non-fat milk for 30 min at room temperature and was then incubated with the rabbit anti-homocysteine antibody (1/1500, cat. ab15154; Abcam, Cambridge, MA) overnight at 4 $^{\circ}\text{C}$, followed by incubation with the peroxidase-linked antirabbit IgG antibody (1/1000; Jackson ImmunoResearch Laboratories, Inc., West Grove, PA) for 1 h at room temperature. After thorough washing, chemiluminescence was developed (Immobilon Western chemiluminescent horseradish peroxidase substrate; Millipore, Temecula, CA) and detected with a digital imaging system (IS4000R; Kodak, New Haven, CT). Intensities of dots stained with anti-homocysteine were determined by

calibration of the total amounts of pixels using the ImageJ image processing program (National Institute of Health, Bethesda, MD). The retinal homocysteinylation levels in the injection groups were represented as the relative amounts of pixels compared with those of the retinas without injection (blank controls). Six independent samples from each injection group were determined. To ensure independent observation, only one eye from each mouse was randomly selected for assessment of retinal homocysteinylation.

Microscopic evaluation and quantification of thickness for total and specific retinal layers: For microscopic evaluations, the mice were sacrificed by cervical dislocation. One eye from each mouse was randomly selected and extracted. The extracted eyes were fixed in 4% formalin for at least 24 h, washed with 0.9% NaCl, and processed through ethanol and xylene solutions. The preparations were then embedded in paraffin, cut at 5- μ m thickness, and mounted on glass slides following conventional procedures [26–28]. Hematoxylin-Eosin stain was performed for histopathological examinations [28,29]. The retinal tissue sections were scanned for evidence of gross defects and were then used to measure the thickness of the total retina, GCL, INL, ONL, and the OS/IS of the photoreceptors. In each retinal section, the measurement of thickness was made on two sites, approximately 200 to 300 μ m apart, on either side of the optic nerve. The results of these measurements were then averaged for that particular section. For each retina analyzed, three separate eye sections were measured by two observers without prior knowledge of the study groups. All measurements were performed with a Nikon E100 microscope (Nikon,) equipped with a digital camera and the ImageJ image processing program (National Institute of Health).

Immunohistochemical analysis and quantification of labeled cells: For immunohistochemical analysis, the retinal tissue sections were boiled in a citrate buffer (pH 6.0) for 20 min for antigen retrieval and were then incubated, respectively, with one of the following antibodies: rabbit anti-homocysteine (1/50, cat. ab15154; Abcam); mouse anti-rhodopsin (1/400, cat. Ab81702; Abcam); mouse anti-calbindin (1/50, cat. sc-74462; Santa Cruz Biotechnology, Santa Cruz, CA); mouse anti-Pax-6 antibody (1/50, cat. sc-32766; Santa Cruz Biotechnology); rabbit anti-Islet-1 (1/150, cat. Ab20670; Abcam); sheep anti-Chx-10 (1/150, cat. ab16141; Abcam); or rabbit anti-GFAP (1/250, cat. 2301-1; Epitomics, Burlingame, CA). The preparations were then incubated with a horseradish peroxidase-conjugated secondary antibody (1/200), either antimouse, antirabbit, or antisheep IgG (Jackson ImmunoResearch Laboratories, Inc.). After incubation, the preparations were washed thoroughly, incubated in diaminobenzidine tetrahydrochloride solution for color detection, and were counterstained with hematoxylin. Positively labeled cells were counted on two sites, approximately 200 to 300 μ m, apart on either side of the optic nerve. The areas for counting were fixed at 100 μ m in

width for each site. The Rhodopsin positive cells were counted in the ONL. The calbindin positive cells were counted in the OPL. The Islet-1 or Chx-10 or Pax-6 positive cells were counted in the INL. The glial fibrillary acidic protein (GFAP) positive cells were counted in the whole thickness of the retina. The results of counting were then averaged for that particular section. For each retina analyzed, three separate sections were measured by two observers without prior knowledge of the study groups. All quantifications of labeled cells were performed with a Nikon E100 microscope (Nikon) equipped with a digital camera and the ImageJ image processing program (National Institute of Health.).

Statistical analysis: All data were obtained from triple repeats and are presented as the means \pm standard error of the means (SEMs) and were compared among groups. The retinal thickness and the means of labeled retinal cells were compared and analyzed by the Mann-Whitney test. All statistical analyses were performed using the Prism program (GraphPad Software, San Diego, CA).

RESULTS

Retinal homocysteinylation after intravitreal homocysteine thiolactone injection: We first tried to determine whether intravitreal Hcy-T injection can lead to homocysteinylation in the retina. Dot-blot analysis was performed using the supernatant of the whole retina at day 90 after the injection. Retinal homocysteinylation was found in a dose-dependent manner, with the highest extent of homocysteinylation found in the retinas injected with Hcy-T to reach a final concentration of 200 μ M (Figure 1B). Quantitative analysis on the relative density of dot blots showed that Hcy-T injections for final 200 μ M resulted in approximately a fivefold increase in retinal homocysteinylation compared to the sham injection controls ($p < 0.001$; Figure 1C). The results indicated that intravitreal Hcy-T injections could lead to retinal homocysteinylation and the doses of Hcy in the vitreous fluid will determine the extent of homocysteinylation.

Retinal degeneration and reduced thickness of total retina and specific retinal layers after retinal homocysteinylation: With retinal homocysteinylation confirmed after Hcy-T injections for final concentrations of 5, 25, and 200 μ M, we then examined the effects after this manipulation. There was no evidence of the degeneration of specific retinal layers following Hcy-T injection for 5 μ M. Hcy-T injection for 25 μ M, however, started to result in retinal cell degeneration, particularly in the GCL, INL, and ONL, where most of the retinal cell nuclei are located (Figure 2D). Furthermore, the inner plexiform layer (IPL) generally became loosely deployed, suggesting occurrence of edema. Notably, all of the cells in the GCL, INL, and ONL exhibited nucleus condensation and lost their orderly organization as a layer. Quantitative analysis showed a trend of thickness reduction in the total retina, with significant difference found in the




Figure 2. The effects of an intravitreal homocysteine-thiolactone (Hcy-T) injection on mouse retinas at day 90 after preparation. All retinas were prepared with hematoxylin-eosin staining for histological assessment and for measurement of the thickness of specific layers. **A:** Blank control without any injection. **B:** 0.9% NaCl sham injection control. **C:** Final intravitreal Hcy-T at 5 μM . **D:** Final intravitreal Hcy-T at 25 μM . **E:** Final intravitreal Hcy-T at 200 μM . **F:** Total retinal thickness. **G:** Ganglion cell layer (GCL) thickness. **H:** Inner nuclear layer (INL) thickness. **I:** Outer nuclear layer (ONL) thickness. **J:** Outer and inner segments of photoreceptors (OS/IS) thickness. The sample size for total retinal thickness was 5 in all groups. The error bars indicate standard error of the means (SEMs). Scale bar: 25 μm .

group injected with 200 μM of Hcy-T, as compared to that of the sham injection control group (Figure 2F). With Hcy-T injection, the GCL and INL layers were also reduced in thickness, although the results were not statistically significant (Figure 2G,H). In contrast, the thickness of the ONL and OS/IS layers were already significantly reduced after Hcy-T injection for 25 μM , and Hcy-T injection for 200 μM lead to additional reduction (both $p < 0.001$; Figure 2I,L). These results indicated that Hcy-T injections caused reduction of total retinal thickness, which was likely due to the effects on multiple retinal layers.

Localization of retinal homocysteinylation after Hcy-T injection: To further understand the effects of excessive Hcy-T on the retina, we compared the localization of retinal homocysteinylation by immunohistochemistry at days 7, 15, 45, and 90 after injections for a final intravitreal Hcy-T concentration at 200 μM . Generally, the sites of retinal homocysteinylation were most evident in the OS/IS and the GCL layers (Figure 3C-F), as compared to the absence of evident localization in the blank and the sham injection

controls (Figure 3A,B). Evidence of accumulation of the wreckage of degenerated photoreceptors was seen on day 45 and day 90 after Hcy-T injection (Figure 3E-F).

The effect of intravitreal Hcy-T injections on specific retinal cell types: Since the thickness reduction was observed in multiple layers of the retina, it is important to identify the specific retinal cell types affected after intravitreal Hcy-T injection for 200 μM . We therefore conducted immunohistochemistry using specific markers for photoreceptors (rhodopsin), horizontal (calbindin), bipolar (Chx-10 or Islet-1), amacrine (Pax-6), and Müller cells (GFAP) at day 15 and day 90 following the injections. At day 15, the rhodopsin⁺ photoreceptor cells were significantly reduced (Figure 4C) as compared to those of the sham injection control (Figure 4A). This reduction of photoreceptors was aggravated to the extent that almost all photoreceptors were ablated at day 90 after the injection (Figure 4D), which was not seen in the sham injection controls (Figure 4B). The calbindin⁺ horizontal cells were also affected at day 15 after the injection (Figure 4G) and were also totally




Figure 3. The immunohistochemical localization of retinal homocystinylation sites at days 7, 15, 45, 90 after a homocysteine-thiolactone (Hcy-T) injection to reach a final intravitreal Hcy-T concentration at 200 μ M. **A:** Blank control without any injection. **B:** 0.9% NaCl sham injection control at day 90. **C:** Day 7 after injection. **D:** Day 15 after injection. **E:** Day 45 after injection. **F:** Day 90 after injection. Scale bar: 40 μ m.

ablated in most retinas at day 90 after the injection (Figure 4H), in contrast to the normal presence and deployment of horizontal cells in the sham injection controls at day 15 (Figure 4E) and day 90 (Figure 4F), respectively. We used two markers, Chx-10 and Islet-1, for the identification of bipolar cells (Figure 4I,P). At day 15 after the Hcy-T injection, misarrangement of bipolar cells was readily found (Figure 4K,O), which was not found in the sham injection controls (Figure 4I,M). The same misarrangement of bipolar cells was also observed at day 90 after the Hcy-T injection (Figure 4L,P), in contrast to the normal deployment of bipolar cells seen in the sham injection controls at the same stage (Figure 4J,N). The Pax-6⁺ amacrine cells appeared normal at day 15 after the injection (Figure 4S), but they were generally misarranged at day 90 (Figure 4T), as compared to the sham injection controls at day 15 (Figure 4Q) and day 90 (Figure 4R). The GFAP⁺ Müller cells were activated at day 15 after the injection (Figure 4W), which was not found in the sham injection controls (Figure 4U). However, the activated Müller cells were generally not seen at day 90 after the injection (Figure 4X), as compared to the results in the sham injection control group (Figure 4V).

To further confirm our observations on the histological level, we quantified specific retinal cell types according to the expression of their markers. At day 15 after intravitreal Hcy-T injections, the cell populations of rhodopsin⁺ photoreceptor cells and calbindin⁺ horizontal cells were both significantly reduced (Figure 5A,B; $p < 0.001$ for both), as compared to those of the sham injection controls. Significant reduction of the rhodopsin⁺ photoreceptor cells and the calbindin⁺ horizontal cells was also seen at day 90 ($p < 0.001$ for both). Reduction of cell populations was also observed in the Chx-10⁺ (Figure 5C) or the Islet-1⁺ (Figure 5D) bipolar cells, but not in the Pax-6⁺ amacrine cells (Figure 5E). Interestingly, the GFAP⁺ Müller cells were significantly activated at day 15 after intravitreal Hcy-T injections, but the activation was not seen at day 90 as there was no significant increase in the

numbers of Müller cells detected (Figure 5F). The results of the quantification of specific cell types in the retinas after intravitreal Hcy-T injection confirmed our observations on the tissue level.

DISCUSSION

Hyperhomocysteinemia (hHcy) has been implicated in human cardiovascular and neurodegenerative diseases, particularly under diabetic conditions [2,3,16]. In recent years, many lines of evidence have indicated high levels of Hcy in the plasma or vitreous fluid [24,25,30] or deficiency of methionine synthase [22] as risk factors of clinical retinopathies and disturbed visual functions [24,31–33]. Hcy, like the neurotransmitter amino acid glutamate, is excitotoxic when its plasma concentration is elevated [13,14]. Hcy-T, the metabolically active form of Hcy, is known to be more cytotoxic than Hcy itself [1,2,18,19] and increased Hcy-T levels have been associated with severe diabetic retinopathy [30]. In previous literature, although the neurotoxic effects of Hcy-T and Hcy on retinal ganglion cells have been reported [15,16,23,34,35], little is known about their toxic effect on photoreceptors and other retinal cells. Clinically, however, a methionine synthase deficiency child with severe hyperhomocysteinemia was reported to show photopic ERG responses related to impaired photoreceptor function [36]. Earlier reports had also associated the combined conditions of methylmalonic aciduria and homocystinuria with abnormal electroretinograms and visual function [21,22]. Thus, in this study, we established a mouse hyperhomocysteinemia model to determine the effects of Hcy-T. The retinas administered with an intravitreal Hcy-T injection were analyzed and compared with their age-matched blank controls and sham injection controls.

Our present study leads to several important findings in support of photoreceptor degeneration induced by excessive Hcy-T. First, the thickness of the ONL and OS/IS of photoreceptors was reduced in the retinas following Hcy-T




Figure 4. The cytotoxic effects of homocysteine-thiolactone (Hcy-T) on specific retinal cells as reflected by alterations of **A-D**: rhodopsin positive photoreceptors, **E-H**: calbindin positive horizontal cells, **I-L**: Chx- 10 positive bipolar cells, **M-P**: Islet-1 positive bipolar cells, **Q-T**: Pax-6 positive amacrine cells, and **U-X**: glial fibrillary acidic protein (GFAP) positive Müller cells at day 15 and day 90 following the intravitreal injections. Abbreviations: GCL represents ganglion cell layer; INL represents inner nuclear layer; IPL represents inner plexiform layer; ONL represents outer nuclear layer; OPL represents outer plexiform layer; OS/IS represents outer and inner segments of photoreceptors. Scale bar: 45 μ m.




Figure 5. Quantification of specific retinal cells per 100 μm length according to their specific markers. The markers were used to detect at day 90 following the intravitreal injections. **A:** Rhodopsin positive cells per 100 μm length in outer nuclear layer (ONL). **B:** Calbindin positive cells per 100 μm length in the retina. **C:** Chx-10 positive cells per 100 μm length in inner nuclear layer (INL). **D:** Islet-1 positive cells per 100 μm length in INL. **E:** Pax-6 positive cells per 100 μm length in INL. **F:** Glial fibrillary acidic protein (GFAP) positive cells per 100 μm length in the retina. The sample size for all markers was 5 in all groups. The error bars indicate standard error of the means (SEMs).

injections for final intravitreal concentrations at 25 μM and 200 μM, as compared to the age-matched 0.9%NaCl sham injection controls or the blank controls, suggesting

photoreceptor cell loss in those layers. Second, the loss of photoreceptors was further supported by specific localization of retinal homocysteinylation on the photoreceptor OS/IS

junction at day 7 after Hcy-T injection for 200 μ M (Figure 3C), which likely started to damage the photoreceptors, resulting in the accumulation of the wreckage of degenerated photoreceptors as seen at day 45 and day 90 (Figure 3E-F). Third, another line of evidence also showed a significant loss of rhodopsin⁺ photoreceptors in the ONL after Hcy-T injection for 200 μ M, as compared to the age-matched 0.9% NaCl sham injection controls (Figure 4A-D and Figure 5A).

As photoreceptors were found to be degenerated after Hcy-T injection, several important implications can be derived from this study. First, the integrity of the photoreceptor OS/IS layer in the retina is highly critical as vision stimuli are initially triggered when the photosensitive proteins in the outer segment absorb photons and generate a signal that will eventually be transmitted to the visual cortex. Notably, the junction region of OS/IS serves as the site for generation of new outer segment discs, where newly synthesized rhodopsins are incorporated. Clinically, the integrity of the photoreceptor OS/IS layer is known to correlate with visual function and visual acuity in patients with retinitis pigmentosa [37,38]. In this study, we observed reduced thickness of the OS/IS layers (Figure 2D,E,J) and loss of rhodopsin protein incorporation into the OS/IS layers (Figure 4C,D) at day 90 after Hcy-T injection. These effects are likely to be directly caused by homocysteinylation, since homocysteinylation was localized in the OS/IS layers (Figure 3C,D,F). When the OS/IS structures are targeted directly by homocysteinylation, the physiologic renewal process for installment of photosensitive proteins will be disrupted. Thus, the status of retinal homocysteinylation by Hcy-T might be regarded as an important risk factor for poor visual acuity or vision loss in clinical retinopathies associated with hyperhomocysteinemia. Second, the horizontal cells are second-order neurons of the vertebrate retina, being responsible for the lateral integration of signal transmission between photoreceptor and bipolar cells. Retinal cone photoreceptors have been reported to drive the horizontal cells to express N-methyl-D-aspartate (NMDA) receptors [39,40]. It has also been reported that overstimulation of NMDA receptors by excessive glutamate or homocysteine on the retinal ganglion cells in vitro leads the cells into a tract of rapid death [23,34]. Thus, the loss of horizontal cells after Hcy-T injection in vivo (Figure 4E-H and Figure 5B) might mediate the same pathway to cell death. This hypothesis, however, remains to be confirmed. Third, Hcy-T might induce endoplasmic reticulum (ER) stress and cause damage to photoreceptors, resulting in a reduction in ONL thickness (Figure 2D-E,I). This induced ER stress and cell death has been well documented in the vascular endothelial cells [41, 42] and lens epithelial cells [43] under the influence of excessive Hcy-T or Hcy. Previous reports have implicated ER stress as a critical factor in retinal degeneration [44,45], but without referring to hyperhomocysteinemia. It is likely that

the same mode of action occurs in the retina after Hcy-T injection.

Apart from photoreceptor degeneration, significant loss of horizontal cells (calbindin⁺ cells) in the OPL was also found after Hcy-T injection for 200 μ M as compared to the age-matched 0.9% NaCl sham injection controls (Figure 4E-H and Figure 5B). Reduction of cell populations was also observed in the Chx-10⁺ (Figure 5C) or the Islet-1⁺ (Figure 5D) bipolar cells. The disruption process of the horizontal cells (calbindin⁺ cells) and bipolar cells after Hcy-T injection seemed to differ from that of the OS/IS photoreceptor layers. Hcy-T might directly target the OS/IS layers since homocysteinylation was localized there. In contrast, the disruption of the horizontal cells and bipolar cells appeared to be due to indirect effects, as no apparent homocysteinylation was localized within these cells. Functionally, bipolar and horizontal cells act as interneurons to transfer information from photoreceptors to ganglion cells [46]. Reduction or disruption of these interneurons will lead to lower sensitivity to contrast and differences in intensity, resulting in lower visual acuity [46–48]. Therefore, it is likely that hyperhomocysteinemia, commonly found in patients with diabetes mellitus [1,24,25], is a factor causing the reduced visual acuity found in diabetic retinopathy.

Interestingly, the GFAP⁺ Müller cells were significantly activated at day 15 after intravitreal Hcy-T injections, but the activation was not seen at day 90, as no significant increase in GFAP⁺ Müller cells was detected (Figure 5F). Müller cells contain multiple functions vital to the health of other retinal neuron cells [49,50]. They may be involved in both the phagocytosis of neuronal debris and the maintenance of other retinal cells. The early increase in GFAP⁺ Müller cells at day 15 may represent a functional response to deplete the damaged retinal cells after Hcy-T injection. This early effort obviously failed as the GFAP⁺ Müller cell population was reduced to nearly the same level as that in the sham-injection control at day 90. The result from this mouse model indicates the risk of chronic hyperhomocysteinemia in a period of just 90 days.

We conclude that excessive Hcy-T could cause degeneration of photoreceptors in the mouse retina. Our results support the necessity of evaluating photoreceptors and the OS/IS junction in patients with severe hyperhomocysteinemia or diabetes mellitus.

ACKNOWLEDGMENTS

This work was supported by a grant (NSC 98–2314-B-040–017-) to B.Y. Chen and partly by a grant (NSC 96–2321-B-040–007-MY3) to H.H. Chang, from National Science Council, Taiwan. A substantial part of this work was performed in the Instrument Center of Chung Shan Medical University, which is supported by National Science Council, Ministry of Education and Chung Shan Medical University.

REFERENCES

- Perla-Kaján J, Twardowski T, Jakubowski H. Mechanisms of homocysteine toxicity in humans. *Amino Acids* 2007; 32:561-72. [PMID: 17285228]
- Karolczak K, Olas B. Mechanism of action of homocysteine and its thiolactone in hemostasis system. *Physiol Res* 2009; 58:623-33. [PMID: 19093716]
- Dionisio N, Jardin I, Salido GM, Rosado JA. Homocysteine, intracellular signaling and thrombotic disorders. *Curr Med Chem* 2010; 17:3109-19. [PMID: 20629621]
- Ghorbanihaghjo A, Javadzadeh A, Argani H, Nezami N, Rashtchizadeh N, Rafeey M, Rohbaninoubar M, Rahimi-Ardabili B. Lipoprotein(a), homocysteine, and retinal arteriosclerosis. *Mol Vis* 2008; 14:1692-7. [PMID: 18806883]
- Micheal S, Qamar R, Akhtar F, Khan MI, Khan WA, Ahmed A. MTHFR gene C677T and A1298C polymorphisms and homocysteine levels in primary open angle and primary closed angle glaucoma. *Mol Vis* 2009; 15:2268-78. [PMID: 19936026]
- Clement CI, Goldberg I, Healey PR, Graham SL. Plasma homocysteine, MTHFR gene mutation, and open-angle glaucoma. *J Glaucoma* 2009; 18:73-8. [PMID: 19142139]
- Javadzadeh A, Ghorbanihaghjo A, Bahreini E, Rashtchizadeh N, Argani H, Alizadeh S. Plasma oxidized LDL and thiol-containing molecules in patients with exudative age-related macular degeneration. *Mol Vis* 2010; 16:2578-84. [PMID: 21151596]
- Gerth C, Morel CF, Feigenbaum A, Levin AV. Ocular phenotype in patients with methylmalonic aciduria and homocystinuria, cobalamin C type. *J AAPOS* 2008; 12:591-6. [PMID: 18848477]
- Stanger O, Weger M, Obeid R, Temmel W, Meinitzer A, Steinbrugger I, Schmut O, Herrmann W. Impairment of homocysteine metabolism in patients with retinal vascular occlusion and non-arteritic ischemic optic neuropathy. *Clin Chem Lab Med* 2005; 43:1020-5. [PMID: 16197292]
- Kuo JZ, Lai CC, Ong FS, Shih CP, Yeung L, Chen TL, Chen KJ, Wu WC. Central retinal vein occlusion in a young Chinese population: risk factors and associated morbidity and mortality. *Retina* 2010; 30:479-84. [PMID: 20010454]
- Bellamy MF, McDowell IF, Ramsey MW, Brownlee M, Bones C, Newcombe RG, Lewis MJ. Hyperhomocysteinemia after an oral methionine load acutely impairs endothelial function in healthy adults. *Circulation* 1998; 98:1848-52. [PMID: 9799203]
- Mujumdar VS, Aru GM, Tyagi SC. Induction of oxidative stress by homocyst(e)ine impairs endothelial function. *J Cell Biochem* 2001; 82:491-500. [PMID: 11500925]
- Tyagi SC. Homocyst(e)ine and heart disease: pathophysiology of extracellular matrix. *Clin Exp Hypertens* 1999; 21:181-98. [PMID: 10225475]
- Torres L, Garcia-Trevijano ER, Rodriguez JA, Carretero MV, Bustos M, Fernandez E, Eguinoa E, Mato JM, Avila MA. Induction of TIMP-1 expression in rat hepatic stellate cells and hepatocytes: a new role for homocysteine in liver fibrosis. *Biochim Biophys Acta* 1999; 1455:12-22. [PMID: 10524225]
- Ganapathy PS, Moister B, Roon P, Mysona BA, Duplantier J, Dun Y, Moister TK, Farley MJ, Prasad PD, Liu K, Smith SB. Endogenous elevation of homocysteine induces retinal neuron death in the cystathionine-beta-synthase mutant mouse. *Invest Ophthalmol Vis Sci* 2009; 50:4460-70. [PMID: 19357353]
- Ganapathy PS, Roon P, Moister TK, Mysona B, Smith SB. Diabetes accelerates retinal neuronal cell death in a mouse model of endogenous hyperhomocysteinemia. *Ophthalmol Eye Dis* 2009; 1:3-11. [PMID: 20407615]
- Lee I, Lee H, Kim JM, Chae EH, Kim SJ, Chang N. Short-term hyperhomocysteinemia-induced oxidative stress activates retinal glial cells and increases vascular endothelial growth factor expression in rat retina. *Biosci Biotechnol Biochem* 2007; 71:1203-10. [PMID: 17485853]
- Jakubowski H. Homocysteine thiolactone: metabolic origin and protein homocysteinylation in humans. *J Nutr* 2000; 130:377S-81S. [PMID: 10721911]
- Jakubowski H. Pathophysiological consequences of homocysteine excess. *J Nutr* 2006; 136:1741S-9S. [PMID: 16702349]
- Kerkeni M, Tnani M, Chuniaud L, Miled A, Maaroufi K, Trivin F. Comparative study on in vitro effects of homocysteine thiolactone and homocysteine on HUVEC cells: evidence for a stronger proapoptotic and proinflammatory homocysteine thiolactone. *Mol Cell Biochem* 2006; 291:119-26. [PMID: 16715183]
- Ticho BH, Feist RM, Fishman GA. Nondetectable electroretinogram in combined methylmalonic aciduria and homocystinuria. *Ann Ophthalmol* 1992; 24:180-1. [PMID: 1637127]
- Ricci D, Pane M, Deodato F, Vasco G, Rando T, Caviglia S, Dionisi-Vici C, Mercuri E. Assessment of visual function in children with methylmalonic aciduria and homocystinuria. *Neuropediatrics* 2005; 36:181-5. [PMID: 15944903]
- Moore P, El-sherbeny A, Roon P, Schoenlein PV, Ganapathy V, Smith SB. Apoptotic cell death in the mouse retinal ganglion cell layer is induced in vivo by the excitatory amino acid homocysteine. *Exp Eye Res* 2001; 73:45-57. [PMID: 11428862]
- Aydemir O, Turkcuoglu P, Guler M, Celiker U, Ustundag B, Yilmaz T, Metin K. Plasma and vitreous homocysteine concentrations in patients with proliferative diabetic retinopathy. *Retina* 2008; 28:741-3. [PMID: 18463519]
- Coral K, Angayarkanni N, Gomathy N, Bharathselvi M, Pukhraj R, Rupak R. Homocysteine levels in the vitreous of proliferative diabetic retinopathy and rhegmatogenous retinal detachment: its modulating role on lysyl oxidase. *Invest Ophthalmol Vis Sci* 2009; 50:3607-12. [PMID: 19369240]
- Chen BY, Lin DP, Liu JY, Chang H, Huang PH, Chen YL, Chang HH. A mouse prostate cancer model induced by Hedgehog overexpression. *J Biomed Sci* 2006; 13:373-84. [PMID: 16374524]
- Chen BY, Liu JY, Chang HH, Chang CP, Lo WY, Kuo WH, Yang CR, Lin DP. Hedgehog is involved in prostate basal cell hyperplasia formation and its progressing towards tumorigenesis. *Biochem Biophys Res Commun* 2007; 357:1084-9. [PMID: 17466949]
- Chang HH, Chen BY, Wu CY, Tsao ZJ, Chen YY, Chang CP, Yang CR, Lin DP. Hedgehog overexpression leads to the formation of prostate cancer stem cells with metastatic

- property irrespective of androgen receptor expression in the mouse model. *J Biomed Sci* 2011; 18:6. [PMID: 21241512]
29. Chen BY, Chang HH, Chen ST, Tsao ZJ, Yeh SM, Wu CY, Lin DP. Congenital eye malformations associated with extensive periocular neural crest apoptosis after influenza B virus infection during early embryogenesis. *Mol Vis* 2009; 15:2821-8. [PMID: 20029647]
 30. Barathi S, Angayarkanni N, Pasupathi A, Natarajan SK, Pukraj R, Dhupper M, Velpandian T, Muralidharan C, Sivashanmugham M. Homocysteinethiolactone and paraoxonase: novel markers of diabetic retinopathy. *Diabetes Care* 2010; 33:2031-7. [PMID: 20551012]
 31. Vine AK. Hyperhomocysteinemia: a risk factor for central retinal vein occlusion. *Am J Ophthalmol* 2000; 129:640-4. [PMID: 10844057]
 32. Pianka P, Almog Y, Man O, Goldstein M, Sela BA, Loewenstein A. Hyperhomocysteinemia in patients with nonarteritic anterior ischemic optic neuropathy, central retinal artery occlusion, and central retinal vein occlusion. *Ophthalmology* 2000; 107:1588-92. [PMID: 10919914]
 33. Lattanzio R, Sampietro F, Ramoni A, Fattorini A, Brancato R, D'Angelo A. Moderate hyperhomocysteinemia and early-onset central retinal vein occlusion. *Retina* 2006; 26:65-70. [PMID: 16395141]
 34. Martin PM, Ola MS, Agarwal N, Ganapathy V, Smith SB. The sigma receptor ligand (+)-pentazocine prevents apoptotic retinal ganglion cell death induced in vitro by homocysteine and glutamate. *Brain Res Mol Brain Res* 2004; 123:66-75. [PMID: 15046867]
 35. Dun Y, Thangaraju M, Prasad P, Ganapathy V, Smith SB. Prevention of excitotoxicity in primary retinal ganglion cells by (+)-pentazocine, a sigma receptor-1 specific ligand. *Invest Ophthalmol Vis Sci* 2007; 48:4785-94. [PMID: 17898305]
 36. Poloschek CM, Fowler B, Unsold R, Lorenz B. Disturbed visual system function in methionine synthase deficiency. *Graefes Arch Clin Exp Ophthalmol* 2005; 243:497-500. [PMID: 15931548]
 37. Mitamura Y, Aizawa S, Baba T, Hagiwara A, Yamamoto S. Correlation between retinal sensitivity and photoreceptor inner/outer segment junction in patients with retinitis pigmentosa. *Br J Ophthalmol* 2009; 93:126-7. [PMID: 19098050]
 38. Aizawa S, Mitamura Y, Baba T, Hagiwara A, Ogata K, Yamamoto S. Correlation between visual function and photoreceptor inner/outer segment junction in patients with retinitis pigmentosa. *Eye (Lond)* 2009; 23:304-8. [PMID: 18188175]
 39. Shen Y, Zhang M, Jin Y, Yang XL. Functional N-methyl-D-aspartate receptors are expressed in cone-driven horizontal cells in carp retina. *Neurosignals* 2006–2007; 15:174-9. [PMID: 17047395]
 40. Wang XL, Jiang XD, Liang PJ. Intracellular calcium concentration changes initiated by N-methyl-D-aspartic acid receptors in retinal horizontal cells. *Neuroreport* 2008; 19:675-8. [PMID: 18382285]
 41. Roybal CN, Yang S, Sun CW, Hurtado D, Vander Jagt DL, Townes TM, Abcouwer SF. Homocysteine increases the expression of vascular endothelial growth factor by a mechanism involving endoplasmic reticulum stress and transcription factor ATF4. *J Biol Chem* 2004; 279:14844-52. [PMID: 14747470]
 42. Raposo B, Rodriguez C, Martinez-Gonzalez J, Badimon L. High levels of homocysteine inhibit lysyl oxidase (LOX) and downregulate LOX expression in vascular endothelial cells. *Atherosclerosis* 2004; 177:1-8. [PMID: 15488859]
 43. Ikesugi K, Yamamoto R, Mulhern ML, Shinohara T. Role of the unfolded protein response (UPR) in cataract formation. *Exp Eye Res* 2006; 83:508-16. [PMID: 16643900]
 44. Yang LP, Wu LM, Guo XJ, Tso MO. Activation of endoplasmic reticulum stress in degenerating photoreceptors of the rd1 mouse. *Invest Ophthalmol Vis Sci* 2007; 48:5191-8. [PMID: 17962473]
 45. Yang LP, Wu LM, Guo XJ, Li Y, Tso MO. Endoplasmic reticulum stress is activated in light-induced retinal degeneration. *J Neurosci Res* 2008; 86:910-9. [PMID: 17929311]
 46. Masland RH. The fundamental plan of the retina. *Nat Neurosci* 2001; 4:877-86. [PMID: 11528418]
 47. Zhang AJ, Wu SM. Receptive fields of retinal bipolar cells are mediated by heterogeneous synaptic circuitry. *J Neurosci* 2009; 29:789-97. [PMID: 19158304]
 48. Burkhardt DA. Contrast processing by ON and OFF bipolar cells. *Vis Neurosci* 2011; 28:69-75. [PMID: 21092350]
 49. García M, Vecino E. Role of Müller glia in neuroprotection and regeneration in the retina. *Histol Histopathol* 2003; 18:1205-18. [PMID: 12973689]
 50. Bringmann A, Iandiev I, Pannicke T, Wurm A, Hollborn M, Wiedemann P, Osborne NN, Reichenbach A. Cellular signaling and factors involved in Müller cell gliosis: neuroprotective and detrimental effects. *Prog Retin Eye Res* 2009; 28:423-51. [PMID: 19660572]

Articles are provided courtesy of Emory University and the Zhongshan Ophthalmic Center, Sun Yat-sen University, P.R. China. The print version of this article was created on 14 July 2011. This reflects all typographical corrections and errata to the article through that date. Details of any changes may be found in the online version of the article.