

Kirenol production in hairy root culture of *Siegesbeckea orientalis* and its antimicrobial activity

Jian-Ping Wang^{1,2}, Ya-Ming Zhou², Yong-Hui Zhang¹

¹Key Laboratory of Natural Medicinal Chemistry and Resources Evaluation of Hubei Province, Department of Traditional Chinese Medicine, College of Pharmacy, Huazhong University of Science and Technology, Wuhan, People's Republic of China, ²Institut für Pharmazeutische Biologie und Biotechnologie, Heinrich-Heine-Universität Düsseldorf, Germany

Submitted: 02-09-2011

Revised: 10-09-2011

Published: 23-05-2012

ABSTRACT

Background: Despite the excellent anti-inflammatory and anti-rheumatic efficacy associated with kirenol generation, the content of kirenol in *Siegesbeckea orientalis* is quite low. **Objective:** This study was designed to establish a reliable kirenol production protocol by transformed root cultures of *S. orientalis* and to investigate the antimicrobial activities of kirenol, hairy root, and *S. orientalis*. **Materials and Methods:** Transformed root cultures of *S. orientalis* were established by the transformation of *Agrobacterium rhizogenes* A4. Transgenic status of the roots was confirmed by *polymerase chain reaction* (PCR) using *rolB* specific primers. The biomass and kirenol accumulation of hairy root clones were assessed using four different culture media: MS, MS/2, B5, and white. The antimicrobial activities of kirenol, hairy root, and *S. orientalis* were evaluated by the disc diffusion method. **Results:** The optimum media for kirenol synthesis was MS. The content of kirenol in transformed hairy roots made up about 80% of that observed in natural leaves of *S. orientalis* (1.6 mg/g dry weight). All tested samples displayed antimicrobial activity against Gram-positive pathogens including *Staphylococcus epidermidis*, *Staphylococcus aureus*, and *Acinetobacter baumannii*, with MIC ranging from 78 to 625 $\mu\text{g/mL}$. **Discussion and Conclusion:** The high level of kirenol contents was obtained from hairy roots of *S. orientalis*. Kirenol was effective against gram-positive bacteria. Interestingly, the extract from hairy roots showed a diverse antimicrobial effect from that of kirenol and *S. orientalis*.

Key words: Antimicrobial activity, hairy root culture, kirenol, *Siegesbeckia orientalis*

INTRODUCTION

The plants of the genus *Siegesbeckia* (Compositae) are annual herbs widely distributed throughout tropical, subtropical, and temperate regions of the world. *Siegesbeckia orientalis* is a common weed in China, and its aerial part has been used traditionally to treat arthritis, hypertension, malaria, neurasthenia, and snakebite.^[1] Phytochemical studies carried out with this plant revealed that *ent*-kaurane and *ent*-pimarane type diterpenoids and sesquiterpenoids were the primary components of the aerial parts.^[2-4] Kirenol (8(14)-pimarane-2, 15, 16, 19-tetrol) [Figure 1] was a main and common *ent*-pimarane type diterpenoid in *S. orientalis* and had been suggested as the main anti-inflammatory and




Figure 1: Chemical structure of kirenol

anti-rheumatic active compound in *S. orientalis*.^[5] Moreover, kirenol was selected as the reference substance to quantify and qualify *S. orientalis* for Chinese Pharmacopoeia.^[1]

Access this article online

Website:

www.phcog.com

DOI:

10.4103/0973-1296.96569

Quick Response Code:



Address for correspondence:

Prof. Yong-Hui Zhang, Key Laboratory of Natural Medicinal Chemistry and Resources Evaluation of Hubei Province, College of Pharmacy, Huazhong University of Science and Technology, Wuhan, People's Republic of China.
E-mail: jpwang1001@163.com

Despite the excellent efficacy associated with kirenol generation, the content of kirenol in *S. orientalis* is quite low (about 0.5 mg/g, dry wt); the increased production of this compound is therefore to be demonstrated.

Hairy root cultures have been proposed as an alternative method of producing plant secondary metabolites because of their genetic and biochemical stability, rapid growth rate, and their capability to synthesize secondary products at levels comparable to those of the original plants.^[6] The primary objective of this study is to establish a reliable protocol of transformed root cultures of *S. orientalis* and to investigate their utilities in kirenol production.

In addition, our previous research confirmed the topical anti-inflammatory and wound healing activities of kirenol isolated from *S. orientalis*, which correlates well with their antimicrobial properties.^[7,8] However, the antimicrobial activity of kirenol and *S. orientalis* has not been reported before. Herein, we evaluated the antimicrobial activities of kirenol, hairy root culture extract, and conventional extract of *S. orientalis* by the disc diffusion method.

MATERIALS AND METHODS

Plant material, chemicals and microorganisms

S. orientalis (Compositae) was obtained in autumn 2009 from the Herbal Plant Garden of School of Pharmacy, Huazhong University of Science and Technology, China, and authenticated by Professor Jia-Chun Chen (School of Pharmacy, Huazhong University of Science and Technology, China). A voucher specimen (No. 09012-S) was preserved at the herbarium of School of Pharmacy, Huazhong University of Science and Technology. The reference substance of kirenol (batch 0774-200901, purity > 99%) was obtained from the National Institute for the Control of Pharmaceutical and Biological Products (NICBP), Beijing, China. The ¹H and ¹³C NMR data are in agreement with the literature [Figure 2].

The microbial strains used as test organisms were: Gram-positive bacteria, *Bacillus subtilis* (ATCC 6051), *Staphylococcus aureus* (ATCC 29213), *Staphylococcus epidermidis* (ATCC 14990), and *Streptococcus oralis* (ATCC 3507); Gram-negative bacteria, *Acinetobacter baumannii* (ATCC BAA 747), *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853); and fungi, *Candida albicans* (ATCC 90028), and *Candida glabrata* (ATCC MYA 2950). All microorganism cultures were obtained from China General Microbiological Culture Collection Center (CGMCC), Beijing, China. The strains were sub-cultured on an appropriate agar plate 24 h prior to antimicrobial experiments.

Agrobacterium rhizogene

In our previous experiments, four different wild strains of *Agrobacterium rhizogenes* (A4, R1601, LBA9402, and ATCC15834) were used for infection, with A4 being the most efficient one.

Therefore, we chose *A. rhizogenes strain A4*, which was grown in a 40 mL Yeast Extract Broth (YEB) medium at 28°C while shaking at 100 rpm. When OD₆₀₀ reached 1, the culture was centrifuged and the cells were resuspended in a 40 mL hormone-free Murashige and Skoog (MS) liquid medium, supplemented with 20 μM acetosyringone.^[9]

Establishment of hairy root clones

Young leaves of *S. orientalis* were sterilized with 70% (v/v) ethanol for 30 s and 1% (w/v) NaOCl for 5–10 min, and then rinsed four times with sterile distilled water. The leaves carefully wounded with a scalpel were used as explants. The explants were submerged in the bacterial suspension for 40 min, washed with 50 mL sterilized water, and blot-dried on the sterile filter paper. The explants were then transferred to a hormone-free MS medium containing 300 mg/L of cefotaxime. Hairy roots emerging from the explants were excised. Aseptically weighed hairy roots (100 mg) were transferred to flasks containing 50 mL of four different liquid culture media: MS, half strength MS (MS/2), B5, and white.^[10] The cultures were maintained in darkness at 25°C on orbital shakers (80 rpm). During a period of 49 days, following establishment of the hairy root cultures in the liquid medium, a regular subculturing routine was used to maintain the culture. Once a week, a portion of the root clump was aseptically removed and transferred to fresh culture medium.

Identification of transformed hairy roots using polymerase chain reaction

Genomic DNA was extracted and purified from both fresh transformed and native roots according to the method outlined by Thomson and Henry.^[11] Approximately 200 ng genomic DNA was used for PCR. The following primers were used to amplify the TL-DNA *rolB* sequences: 5'-ATGTCGCAAGGACGTAAGCCGA-3' (forward primer) and 5'-GGAGTCTTTCAGCATGGAGCAA-3' (reverse primer). For amplification, initial denaturation was performed at 94°C for 3 min, followed by 35 cycles of amplification (each comprising 1 min at 94°C, 1 min at 55°C, and 1 min at 72°C) and a final extension at 72°C for 5 min. The amplified products were separated on a 1.5% (w/v) agarose gel with 0.5 μg ethidium bromide/mL and detected under UV light.

Culture experiments for hairy roots

The biomass and metabolite accumulation of hairy root clones were assessed using four different culture media as




Figure 2: The spectra of kirenol: H NMR (a), C NMR (b), MS (c)

described above. Four independent growth experiments were separately run, and all experiments were performed in triplicate.

For culture and time-course experiments, roots were harvested and both growth (fresh and dry weight) and kirenol contents were measured at different intervals.

To measure fresh weight, harvested hairy roots were washed twice with doubled-distilled water, then gently pressed onto filter papers to remove excess water and weighed. Dry weight was measured by drying the fresh hairy roots in an oven at 50°C until constant weight was achieved.

To measure kirenol content, slightly modified methods of PCCh^[1] were used. In brief, 100 mg powdered dry hairy roots were extracted with 50 mL of methanol using a reflux condenser. A HPLC system consisting of a Hypersil

ODS-2 column (ThermoElectron Corporation, USA; 250 mm × 4.6 mm, particle size 5 μm) and a SPD-10A variable wavelength programmable UV/Vis detector was used. The mobile phase was acetonitrile–water (25:75 v/v). The temperature of the column was 30°C. The detective wavelength was set at 215 nm. The flow rate was 1.0 mL/min. Calibration tests were performed, and a suitable calibration curve was determined. A correlation coefficient value higher than 0.9997 indicated a linearity for the tested concentrations. Samples from hairy roots, native roots, and leaves (10 μL) were injected and their peak areas were compared against the calibration curves.

Antimicrobial activity

The hairy roots were cultured in MS medium for 49 days. Samples of *S. orientalis* (hairy roots, leaves, roots, and stem) were dried, powdered, and then extracted with methanol (50 mL) by using a reflux condenser. The resulting extracts

were dried, dissolved in DMSO, and used for estimation of minimum inhibitory concentration (MIC). Assays were carried out as described previously.^[12] The MIC of an extract was defined as the lowest concentration where microbial growth was inhibited.^[13] All experiments were performed in triplicate, and MIC values were expressed in $\mu\text{g}/\text{mL}$.

Statistical analysis

Analysis of variance (one-way ANOVA) was performed to test the significance of differences between means obtained among the treatments in each experiment at the 5% level of significance ($P < 0.05$).

RESULTS AND DISCUSSION

Hairy root cultures of *S. orientalis* were initiated by inoculation of leaf discs with *A. rhizogenes* A4 and grown in the dark in liquid MS, B5, white, and MS/2 culture media. These roots exhibited typical characteristics of transformed roots: rapid growth, extensive lateral branch, and a lack of geotropism^[14] [Figure 3]. The transformed roots were confirmed by PCR to determine the presence of a T-DNA sequence in their genomes [Figure 4]. The PCR products from the hairy roots for *rolB* regions but not from untransformed roots gave the expected 540 bp PCR product. This finding indicated that the *rolB* genes from the Ri plasmid of *A. rhizogenes* were integrated into the genome of *S. orientalis* hairy roots.

The hairy roots grown in MS medium showed a higher biomass increase, expressed either in fresh weight or dry weight when compared with those grown in White and MS/2 media. Fresh weight growth curves of the hairy roots grown in the B5 medium were similar to those of




Figure 3: Hairy root cultures of *S. orientalis*. Hairy roots emerging from the leaves after *A. rhizogenes* A4 infection (a) and exhibiting a normal phenotype with a large percentage of branched fine roots and an intermediate growth rate when cultured in liquid media (b)

the hairy roots grown in the MS medium [Figure 5a and b].

Following 49 days of cultivation, time course of kirenol production in different hairy root culture media was achieved. The different hairy root clones showed variation in kirenol contents, with the white medium culture having the lowest and MS medium culture having the highest kirenol content [Figure 5c].

With the objective to achieve high biomass and kirenol production, kirenol production increased with biomass on four different media like MS, MS/2, B5, and white. Hairy roots grown on MS medium accumulated the maximum amount of kirenol content followed by MS/2, B5 and White. It suggested that MS medium contains higher ionic concentrations of inorganic salts compared to MS/2, B5, and White, favoring growth and kirenol production. Altering inorganic nutrients of the cultured medium are known to influence the production of bioactive molecules from cultured tissues.^[15,16]

The kirenol content was detected in methanol extracts of nontransformed roots, stem, and leaves as well as hairy roots using HPLC analysis. Retention time of the peak obtained with the kirenol standard was used to identify the corresponding peaks in extracts [Figure 6].

The presence of kirenol was detected in all tested samples. The kirenol content of hairy roots cultured in MS medium for 49 days was compared with those of various origins of *S. orientalis* [Table 1]. The medium was free of kirenol throughout the cultivation process. The kirenol content of 1.6 mg/g dry wt was obtained using *S. orientalis* hairy root cultures, which was comparable with amounts present in the leaves and roots of the parent plant (2.1 and 0.3 mg/g




Figure 4: Electrophoretogram of *rol* genes PCR identification of hairy roots. Lane 1: molecular weight marker (100 bp ladder), lane 2 and 4: DNA from non-transformed roots (negative control), lane 3: *Agrobacterium rhizogenes* DNA (positive control), lane 5–10: DNA from different lines of hairy roots




Figure 5: Fresh weight (a), dry weight (b) and kirenol content of hairy root clones (c) growth curves of hairy roots grown in MS (■), B5 (○), White (▲) and MS/2 (×) liquid media

Table 1: Kirenol contents from various origins of *S. orientalis* and hairy roots

Material	Kirenol content ^a (mg/g dry wt)
Leaves of parent plant	2.1 ± 0.1
Roots of parent plant	0.3 ± 0.1
Stem of parent plant	0.8 ± 0.4
Hairy roots	1.6 ± 0.5

^aResults are mean ± SEM of three parallel measurements (n = 3)

dry wt, respectively). Much higher level of the kirenol content was obtained from hairy roots than the native roots of *S. orientalis*.

Although the hairy roots under these conditions produced approximately 20% less kirenol than the leaves of *S. orientalis*,




Figure 6: HPLC chromatograms of kirenol standard (a), methanol extract from hairy roots (b) and methanol extract from leaves of *S. orientalis* (c)

the growing time was much shorter when compared to field-grown plants. With hairy root cultures, product quality and quantity are easy to control because natural variances in seasonal climates and geographical environments are excluded and culture conditions and process variables are easily optimized.^[17] Several strategies for the enhancement of biomass and kirenol content have been adopted in which the effects of medium compositions, culture conditions, and elicitations on root growth and kirenol accumulation are currently under further investigation.

For the isolation and purification of kirenol from the cultured hairy roots, the methods of Wang *et al.*^[18] with small modifications were adopted. Briefly, the hairy roots were separated from culture media and soaked in MeOH overnight. Cells were destroyed using Ultra Turrax for 10 min, followed by filtration and exhaustive extraction. The MeOH extract was mixed with H₂O to form a suspension and then partitioned successively with petroleum ether, EtOAc, and *n*-BuOH. The EtOAc-soluble part was subjected to silica gel column chromatography (CC) eluted with petroleum ether–acetone (30:1, 15:1, 8:1, 4:1, 2:1, 1:1, 0:100, and MeOH) to give eight fractions (A–H). Fraction E was separated by silica gel CC eluted with CHCl₃–MeOH (10:1) to give subfraction E', which was recrystallized with acetone to yield kirenol.

The results of the antimicrobial assays have been reported [Table 2]. All tested samples were found to be active on *S. epidermidis*, *S. aureus*, and *A. baumannii*, and the MIC ranged from 78 to 625 µg/mL. Kirenol showed the greatest antibacterial activity against *B. subtilis* and *S. oralis* with a

Table 2: MICs ($\mu\text{g/ml}$) of kirenol, hairy root culture extract, and conventional extract of *Siegesbeckia orientalis* against microorganism

	Microorganisms	Ampicillin ^a	Kirenol	MHR ^b	MES ^c	MER ^d	MEL ^e
Gram-positive	<i>B. subtilis</i>	3.0	39.0	312.5	625.0	–	312.5
	<i>S. epidermidis</i>	6.0	156.0	625.0	39.0	625.0	78.0
	<i>S. aureus</i>	3.0	78.0	78.0	78.0	625.0	39.0
	<i>S. oralis</i>	1.5	39.0	78.0	78.0	–	312.5
Gram-negative	<i>A. baumannii</i>	3.0	625.0	312.5	312.5	312.5	78.0
	<i>E. coli</i>	2.5	312.5	–	–	–	312.5
	<i>P. aeruginosa</i>	1.5	625.0	312.5	–	–	625.0
Fungi	<i>C. albicans</i>	7.5	–	312.5	312.5	–	78.0
	<i>C. glabrata</i>	6.0	–	312.5	–	–	625.0

(–) >625 $\mu\text{g/ml}$. ^aAntibiotic used as a positive control in antibacterial tests. The same MIC values were obtained in three separate experiments, ^bMHR: methanolic extract of hairy roots, ^cMEL: methanolic extract of leaf, ^dMER: methanolic extract of root, ^eMES: methanolic extract of stem

MIC of 39 $\mu\text{g/ml}$. MIC of kirenol on bacteria ranged from 39 $\mu\text{g/ml}$ to 625 $\mu\text{g/ml}$, though it showed almost no activity on the two studied fungi species. Antimicrobial activity seemed to be higher for methanolic extracts from leaves of *S. orientalis* compared to extracts from roots and stem of *S. orientalis*. Previous results from our laboratory showed that kirenol production was tissue-specific with highest amounts present in leaves.

The extract from hairy roots showed a very different antimicrobial effect from that of kirenol and *S. orientalis*. The transformed roots showed distinguished HPLC chromatograms, implying their possible contribution to synthesize and store significant quantities of secondary metabolites. This suggests either an interaction between compounds from hairy roots or the synthesis of new constituents in hairy roots, which might be responsible for this activity. Interestingly, kirenol, hairy root extract, and *S. orientalis* extract were more effective against gram-positive bacteria (such as *Stypholococcus* and *Streptococcus*) than Gram-negative bacteria (*Escherichia*, *Pseudomonas*). This phenomenon could be explained by the fact that the structures of cell envelope differ substantially between gram-positive and gram-negative bacteria. Gram-negative bacteria possess an outer membrane surrounding the cell wall, which restricts diffusion of hydrophobic compounds through its lipopolysaccharide covering. The cell wall of gram-positive bacteria without outer membrane can be permeated more easily, and the constituents of hairy root extract and *S. orientalis* extract, including kirenol, can disturb the various molecular targets such as the cytoplasmic membrane, proton motive force (PMF), electron flow, active transport, and coagulation of cell contents.^[19]

ACKNOWLEDGMENTS

This work was supported by the Fundamental Research Funds for the Central Universities (2011QN243). Jian-Ping Wang is grateful

to Daniela Ackermann (Institut für Pharmazeutische Biologie und Biotechnologie, Heinrich-Heine-Universität Düsseldorf, Germany) for linguistic revision.

REFERENCES

1. Pharmacopoeia Commission of Peoples Republic of China (PCCh). Pharmacopoeia of People's Republic of China (PCh). Vol. 1. Beijing: Chemical Industry; 2010. p. 255-6.
2. Zdero C, Bohlmann F, King RM, Robinson H. Sesquiterpene lactones and other constituents from *Siegesbeckia orientalis* and *Guizotia scabra*. *Phytochemistry* 1991;30:1579-84.
3. Wang LL, Hu LH. Chemical constituents of *Siegesbeckia orientalis* L. *J Integr Plant Biol* 2006;48:991-5.
4. Wang JP, Cai YL, Wu YX. Antiinflammatory and analgesic activity of topical administration of *Siegesbeckia pubescens*. *Pak J Pharm Sci* 2008;21:89-91.
5. Hu HH, Tang LX, Li XM. Experimental research of effect of crude and processed *Herba Siegesbeckiae* on anti-inflammation and anti-rheumatism. *Chin J Chin Mat Med* 2004;29:542-5.
6. Georgiev M, Pavlov A, Bley T. Hairy root type plant *in vitro* systems as sources of bioactive substances. *Appl Microbiol Biotechnol* 2007;74:1175-85.
7. Devib P, Meeraa R. Study of antioxidant, antiinflammatory and wound healing activity of extracts of *Litsea glutinosa*. *J Pharm Sci Res* 2010;2:55-163.
8. Dickson RA, Fleischer TC, Ekuadzi E, Mensah AY, Annan K, Woode E. Antibacterial, antioxidant and anti-inflammatory properties of *Margaritaria discoidea*, a wound healing remedy from Ghana. *Pharmacognosy J* 2010;17: 29-32.
9. Rahnama H, Hasanloo T, Shams MR, Sepehrifar R. Silymarin production by hairy root culture of *Silybum marianum* (L.) Gaertn. *Iran J Biotechnol* 2008;6:113-9.
10. Owen HR, Miller RA. An examination and correction of plant tissue culture basal medium formulations. *Plant Cell Tissue Organ Cult* 1992;28:147-50.
11. Thomson D, Henry R. Single-step protocol for preparation of plant tissue for analysis by PCR. *Biotechniques* 1995;19: 394-400.
12. Teeyapant R, Woerdenbag HJ, Kreis P, Hacker J, Wray V, Witte L, et al. Antibiotic and cytotoxic activity of brominated compounds from the marine sponge *Verongia aerophoba*. *Z Naturforsch C* 1993;48:939-45.
13. Andrews JM. Determination of minimum inhibitory concentrations. *J Antimicrob Chemother* 2001;48:5-16.

14. Alpizar E, Dechamp E, Lapeyre-Montes F, Guilhaumon C, Bertrand B, Jourdan C, *et al.* Agrobacterium rhizogenes-transformed roots of coffee (*Coffea arabica*): Conditions for long-term proliferation, and morphological and molecular characterization. *Ann Bot* 2008;101:929-40.
15. Pan XW, Xu HH, Liu X, Gao X, Lu YT. Improvement of growth and camptothecin yield by altering nitrogen source supply in cell suspension cultures of *Camptotheca acuminata*. *Biotechnol Lett* 2004;26:1745-8.
16. Sivakumar G, Yu KW, Hahn EJ, Paek KY. Optimisation of organic nutrients for ginseng hairy roots production in large scale bioreactors. *Curr Sci* 2005;89:641-9.
17. Payne GF, Bringi V, Prince, C, Shuler ML. *Plant Cell and Tissue Culture in Liquid Systems*. New York: Hanser; 1992. p. 225-79.
18. Wang JP, Zhou YM, Ye YJ, Shang XM, Cai YL, Xiong CM, *et al.* Topical anti-inflammatory and analgesic activity of kirenol isolated from *Siegesbeckia orientalis*. *J Ethnopharmacol* 2011;137:1089-94.
19. Tian F, Li B, Ji BP, Yang JH, Zhang GZ, Chen Y, *et al.* Antioxidant and antimicrobial activities of consecutive extracts from *Galla chinensis*: The polarity affects the bioactivities. *Food Chem* 2009;113:173-9.

Cite this article as: Wang J, Zhou Y, Zhang Y. Kirenol production in hairy root culture of *Siegesbeckia orientalis* and its antimicrobial activity. *Phcog Mag* 2012;8:149-55.

Source of Support: Fundamental Research Funds for the Central Universities (2011QN243), **Conflict of Interest:** None declared.

New features on the journal's website

Optimized content for mobile and hand-held devices

HTML pages have been optimized of mobile and other hand-held devices (such as iPad, Kindle, iPod) for faster browsing speed.

Click on **[Mobile Full text]** from Table of Contents page.

This is simple HTML version for faster download on mobiles (if viewed on desktop, it will be automatically redirected to full HTML version)

E-Pub for hand-held devices

EPUB is an open e-book standard recommended by The International Digital Publishing Forum which is designed for reflowable content i.e. the text display can be optimized for a particular display device.

Click on **[EPub]** from Table of Contents page.

There are various e-Pub readers such as for Windows: Digital Editions, OS X: Calibre/Bookworm, iPhone/iPod Touch/iPad: Stanza, and Linux: Calibre/Bookworm.

E-Book for desktop

One can also see the entire issue as printed here in a 'flip book' version on desktops.

Links are available from Current Issue as well as Archives pages.

Click on  View as eBook