



# Unraveling the hidden paleobiodiversity of the Middle Devonian (Emsian) crinoids (Crinoidea, Echinodermata) from Poland

William I. Ausich<sup>1</sup>, Mariusz A. Salamon<sup>2</sup>, Bartosz J. Płachno<sup>3</sup>, Tomasz Brachaniec<sup>2</sup>, Wojciech Krawczyński<sup>2</sup>, Andrzej Boczarowski<sup>2</sup>, Karolina Paszcza<sup>2</sup>, Magdalena Łukowiak<sup>4</sup> and Przemysław Gorzelak<sup>5</sup>

<sup>1</sup> University of Columbus, Columbus, OH, USA

<sup>2</sup> University of Silesia in Katowice, Sosnowiec, Poland

<sup>3</sup> Jagiellonian University in Kraków, Kraków, Poland

<sup>4</sup> Insitute of Paleobiology, Warszawa, Poland

<sup>5</sup> Insitute of Paleobiology Polish Academy of Science, Warszawa, Poland

## ABSTRACT

Most previous publications on Devonian crinoids from the Holy Cross Mountains in Poland have concentrated on crinoid columns, and until now, little has been published about crinoid cups and calyxes. Herein, five crinoid taxa are described from an abundant occurrence of aboral cups and partial crowns from the Bukowa Góra Member (Emsian) in the Holy Cross Mountains of southern Poland. The following taxa are described: *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* (Bohatý, 2005), *Halocrinites schlotheimii* Steininger, 1831, and a single brachial plate from a flexible crinoid placed in *Flexibilia incertae sedis*. Simple discoid holdfasts are also present encrusted to cylindrical stromatoporoids. These taxa are the first crinoids described from the remains of partial crowns from Emsian strata of Poland.

**Subjects** Marine Biology, Paleontology

**Keywords** Devonian, Emsian, Poland, Crinoids, Echinodermata, Diversity

## INTRODUCTION

Crinoid remains are abundant in Devonian (Emsian-Famennian) strata of Poland (Holy Cross Mountains, southern Poland; Cracow-Silesian area, southern Poland; Sudetes, southwestern Poland; Pomerania, northern Poland). Polish Devonian crinoids were mentioned initially by *Dames* (1868), *Zeuschner* (1867), *Zeuschner* (1869), *Gürich* (1896), and *Sobolev* (1909). Much later, *Kongiel* (1958) and *Piotrowski* (1977) described the occurrence of the genus *Ammonocrinus* in the Holy Cross Mountains (see also *Gorzelak, Gluchowski & Salamon, 2014; Bohatý, 2011*). In a series of subsequent papers, *Gluchowski* (*Gluchowski, 1980, Gluchowski, 1981a, Gluchowski, 1981b, Gluchowski, 1981c, Gluchowski, 1993a, Gluchowski, 1993b, Gluchowski, 2002, Gluchowski, 2003*; see also *Hauser, 2002, Bohatý, 2005, Bohatý, 2009a, Bohatý, 2011*) listed ~50 crinoid taxa from Devonian (Lochkovian, Emsian-Famennian) strata of Poland (for summary see [Fig. 1](#)). With the exception of taxa listed below, most of them were based on isolated skeletal remains,

Submitted 13 December 2021

Accepted 6 January 2022

Published 9 February 2022

Corresponding author

Mariusz A. Salamon,  
paleo.crinoids@poczta.fm

Academic editor

Haijun Song

Additional Information and  
Declarations can be found on  
page 21

DOI 10.7717/peerj.12842

© Copyright  
2022 Ausich et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

Lochkovian	Pragian	Emsian	Eifelian	Givetian	Frasnian	Famenn.	Crinoid taxa
							<i>Kotanocrinus</i> cf. <i>balaensis</i> (col.)
							<i>Poteriocrinites?</i> <i>morlieresensis</i>
							<i>Formosocrinus</i> cf. <i>formosus</i> (col.)
							<i>Acanthocrinus</i> sp. (col.)
							<i>Marettocrinus subbiconcavus</i> (col.)
							<i>Laudonomphalus humilicarinatus</i> (col.)
							<i>Procupressocrinites gracilis</i>
							<i>Mediocrinus microgrumosus</i> (col.)
							<i>Kazachstanocrinus acutilobus</i> (col.)
							<i>Asperocrinus brevispinosus</i> (col.)
							<i>Ricebocrinus kulagaiensis</i> (col.)
							<i>Ammonocrinus sulcatus</i> (col.)
							<i>Eocampocrinus fragilis</i> (col.)
							<i>Gilbertocrinus vetulus</i> (col.)
							<i>Myelodactylus canaliculatus</i> (col.)
							<i>Pentagonostipes petaloides</i> (col.)
							<i>Platycrinites minimalis</i> (col.)
							<i>Noctocrinus?</i> <i>varius</i> (col.)
							<i>Marettocrinus angustannulus</i> (col.)
							<i>Halocrinites schlotheimi</i>
							<i>Praerocrinus polonicus</i> (col.)
							<i>Stenocrinus</i> cf. <i>bifurcatus</i> (col.)
							<i>Stenocrinus degratus</i> (col.)
							<i>Tantalocrinus scutellus</i> (col.)
							<i>Tjeecrinus crassijugatus</i> (col.)
							<i>Schyschcatocrinus creber</i> (col.)
							<i>Kazachstanocrinus tenuis</i> (col.)
							<i>Stenocrinus raricostatus</i> (col.)
							<i>Marettocrinus kartzevae</i> (col.)
							<i>Ammonocrinus kongieli</i> (col.)
							<i>Amurocrinus imatschensis</i> (col.)
							<i>Haplocrinites minor</i>
							<i>Laudonomphalus pinguicostatus</i> (col.)
							<i>Haplocrinites aremoricanus</i>
							<i>Anthinocrinus brevicostatus</i> (col.)
							<i>Anthinocrinus primaevus</i> (col.)
							<i>Halocrinites inflatus</i>
							<i>Onocrinus gracilis</i> (col.)
							<i>Urushicrinus perbellus</i> (col.)
							<i>Ricebocrinus parvus</i> (col.)
							<i>Tjeecrinus simplex</i> (col.)
							<i>Schyschcatocrinus multiformis</i> (col.)
							<i>Haplocrinites</i> sp.
							<i>Schyschcatocrinus delicatus</i> (col.)
							<i>Wenjukowicrinus wenjukowi</i> (col.)
							<i>Caleocrinus bicostatus</i> (col.)
							<i>Glyphidocrinus infimus</i> (col.)
							<i>Caleocrinus kielcensis</i> (col.)
							<i>Melocrinites</i> cf. <i>hieroglyphicus</i>
							<i>Melocrinites</i> cf. <i>gibbosus</i>
							<i>Exaesisdiscus compositus</i> (col.)
							<i>Haplocrinites gluchowskii</i>
							<i>Tjeecrinus insectus</i> (col.)
							<i>Cosmocrinus polonicus</i> (col.)

**Figure 1** Stratigraphic ranges of crinoid taxa recorded in Devonian of Poland. Data compiled after: Dames (1868), Zeuschner (1867), Zeuschner (1869), Gürich (1896), Sobolev (1909), Kongiel (1958), Piotrowski (1977), Gluchowski (1980), Gluchowski (1981a)–Gluchowski (1981c), Gluchowski (1993a), Gluchowski (1993b), Gluchowski (2002), Gluchowski (2003) and Hauser (2002).

Full-size  DOI: 10.7717/peerj.12842/fig-1

mainly columnals; and they were described using the principles of artificial classification of crinoid remains proposed by Moore & Jeffords (1968). Gluchowski (2003) added that the applicability of crinoid stems may be useful for stratigraphic and correlation purposes.

To date, only nine crinoid taxa have been identified on the basis of complete (or nearly complete) crowns and aboral cups with column from the Devonian of Poland. Among these is *Haplocrinites* sp. from Givetian-Frasnian of Holy Cross Mountains recorded by Gluchowski (1993a) and Gluchowski (2003). Specimens of this taxon from late Frasnian Detrital Beds of the Holy Cross Mountains were later designated by Hauser (2002) as *Haplocrinites gluchowskii* (Hauser, 2002). According to Gluchowski, Casier & Olempska (2006) Givetian and early Frasnian *Haplocrinites* sp. specimens differ from *H. gluchowskii* in having distinctly less prominent radial facets. Another haplocrinitid (Haplocrinitidae) species is *Haplocrinites aremoricensis* Le Menn, 1985 from the uppermost

Givetian of the Holy Cross Mountains ([Głuchowski, 1993a](#)). *Platyhexacrinus?* was identified by [Głuchowski \(1993a\)](#) (see [Bohatý, 2009a](#)). Also four cupressocrinitid (Cupressocrinitidae) taxa have been described from the Holy Cross Mountains by [Głuchowski \(1993a\)](#). These are *Cupressocrinites* cf. *abbreviatus* Goldfuss (late Eifelian-late Givetian) (now *Halocrinites schlotheimii* [Steininger, 1831](#)), *H. geminatus* ([Bohatý, 2005](#)) (these specimens were originally described as *Cupressocrinites* cf. *abbreviatus* in ([Głuchowski, 1993a](#), fig. 6g–h), *C. inflatus* ([Schultze, 1866](#)) (late Givetian) (now *Halocrinites inflatus*), and *C. sampelayoi* ([Almela & Revilla, 1950](#)) (now *Halocrinites minor* ([Schultze, 1866](#)) known from late Givetian ([Głuchowski, 1993a](#); [Głuchowski, 2002](#)). The remaining two crinoid species belong to the Melocrinitidae. These are *Melocrinites* cf. *gibbosus* [Goldfuss, 1831](#) and *M. cf. hieroglyphicus* [Goldfuss, 1831](#), which were found in the sediments of the uppermost Frasnian.

Only four crinoid taxa are known from the Emsian of the Holy Cross Mountains and all were documented on the basis of isolated columnals or their impressions (casts). In particular, [Głuchowski, \(1981b\)](#), [Głuchowski, \(2002\)](#) listed the following taxa from the Bukowa Góra shales: *Acanthocrinus* sp. (col.) and *Formosocrinus* cf. *formosus* (col.) ([Yeltyschewa & Sisova, 1973](#)), *Laudonomphalus humilicarinatus* (col.) (Yeltyschewa in [Yeltyschewa & Dubalotova, 1961](#)) [now *Hexacrinites? humilicarinatus* (col.) (note that an affiliation with the crown-based genus, *Hexacrinites*, cannot be verified.)], and *Marettocrinus subbiconcavus* (col.) ([Stukalina, 1965](#)).


Here we report complete or almost complete Emsian cups associated with numerous isolated calyx and column remains from the Bukowa Góra Member in the Holy Cross Mountains of southern Poland. These include *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* ([Bohatý, 2005](#)), *Halocrinites schlotheimii* ([Steininger, 1831](#)), and *Flexibilia* incertae sedis. Simple discoid holdfasts are also present. Remains of unidentifiable specimens indicate that several other crinoids also existed in the Bukowa Góra Member fauna.

## GEOLOGIC FRAMEWORK

The Holy Cross Mountains are located in the southern part of Poland. Their main element is the Paleozoic core, divided into two parts: the Łysogóry region (northern, connected with the Łysogóry Block) and the Kielce region (southern, connected with the Małopolska Block; see [Fig. 2A](#)). These regions differ from each other by facies development of contemporaneous deposits. Devonian sediments of the Łysogóry region were formed in the deeper basin in contrast to the shallower facies exposed in the Kielce region ([Szulczewski, 1995](#)).

Outcrops of Lower Devonian rocks in the Łysogóry Region are connected with the southern limb of the Bodzentyn Syncline. The sedimentary rocks of the upper Emsian are best exposed in the active quarry “Bukowa Góra”, located about 16 km northeast of Kielce (see [Fig. 2A](#)). The section includes sediments belonging to *patulus* Conodont Zone and *douglastownense-euryptero*ta Miospore Zone [Malec, 2005](#); [Filipiak, 2011](#); [Fijałkowska-Mader & Malec, 2011](#)).

In the lower part of the section, the 110 m thick Zagórze “formation” comprised of siliciclastic deposits is present (see [Fig. 2B](#)). They are mostly represented by quartzitic



**Figure 2** (A) The map of Poland with the Holy Cross Mountains area marked as grey rectangle. (B) The lithostratigraphical scheme of Middle and Upper Devonian in the Bukowa Góra Quarry. Compiled after *Marynowski, Salamon & Narkiewicz (2002)*, *Malec (2005)*, *Szulcowski & Porębski (2008)*, *Wójcik (2015)*, and *Salamon et al. (2018)*.

Full-size DOI: 10.7717/peerj.12842/fig-2

sandstones with abundant trace fossils and by claystones. Within sandstones, there are storm originated brachiopod coquinas with gastropods, bivalves, tentaculitids, crinoids, rarely trilobites, rugose corals, nautiloids, and ostracodes. At the top of the Zagórze “formation”, conglomerates and sandstones of estuary facies crop out. Deposits of the Zagórze “formation” were formed in the shallow sea environment from the lagoon to the shoreface (*Szulczewski & Porębski, 2008*; see also *Lobanowski, 1971, Lobanowski, 1981*).

As a result of the progressive deepening of the marine basin, deposits of the Grzegorzowice Formation were formed, which is also present in the Kielce region (*Malec, 2005; Wójcik, 2015*). In the Bukowa Góra quarry section the two members are present: Bukowa Góra Member and the Kapkazy Member. The Bukowa Góra Member occurs only within the *patulus* Conodont Zone in the western part of Bodzentyn Syncline (see *Fig. 2B*). In the eastern part of the Bodzentyn Syncline, the Bukowa Góra Member appears earlier, *i.e.*, in the *serotinus* Conodont Zone (*Malec, 2005*). *Malec (2005)* marked the lower boundary of the Bukowa Góra Member in the bottom of the complex of dark claystones, whereas *Szulczewski & Porębski (2008)* put this boundary on the pebble conglomerate that begins in the lower shoreface to offshore transitional series.

In the lower part of interbedded sandstones, the Bukowa Góra Member is comprised of sandstones, siltstones, and claystones, which are about 7 m thick (see *Szulczewski & Porębski, 2008*). Above them appears the offshore facies represented by black to dark-gray claystones and silty claystones with a thickness of ~13 m and containing discontinuous beds of dolomitic limestones up to 10 cm thick (see *Fig. 2B*). Both claystones and limestones contain a rich faunal assemblage related to the colonization of the soft sea bottom. There are massive colonies of both stromatoporoids and tabulate corals accompanied by solitary rugoses, brachiopods, crinoids, ostracods, gastropods, and trilobites (*Malec, 2005*; see also *Gluchowski, 1993b; Fijałkowska-Mader et al., 1997*).

Claystones of the Bukowa Góra Member are overlain by sandstones of the Kapkazy Member, which is ~34 m thick. The lower part of the Kapkazy Member is comprised of coarse-grained and conglomeratic sandstones, containing rare crinoids, brachiopods, and gastropods. Above this is fine-grained sandstone, which is indicative of a clear shallowing of the sea basin (*Malec, 2005*).

## MATERIALS AND METHODS

The studied material from Bukowa Góra Quarry was collected in 2019 and 2021. The first step consisted of examination of slab surfaces in the field. At this stage, numerous crinoid remains (isolated columnals and complete or nearly complete crowns) were collected. The next step consisted of soaking the respective samples (11 shales samples weighing each ca. 10 kg) only with hot water. Limy samples (4 samples weighing each ca. 5 kg) were soaked with Glauber’s salt. These samples were then boiled and frozen (2–3 times). The residues were finally washed with running tap water and sieved on a sieve column (Ø1.0, 0.315 and 0.1 mm mesh). The final step consisted of drying the shaly and limy residues at 160 °C. Residue was hand-picked from each macerated sample for microscopic study.

All crinoids were photographed by a SONY DSC-RX10M3 digital camera. Specimens discussed here are deposited in the University of Silesia in Katowice, Faculty of Natural

Sciences, Institute of Earth Sciences, Poland (GIUS 4-3696) and in the Senckenberg Forschungsinstitute und Naturmuseum, Frankfurt am Main, Germany (SMF).

“The electronic version of this article in Portable Document Format (PDF) will represent a published work according to the International Commission on Zoological Nomenclature (ICZN), and hence the new names contained in the electronic version are effectively published under that Code from the electronic edition alone. This published work and the nomenclatural acts it contains have been registered in ZooBank, the online registration system for the ICZN. The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix <http://zoobank.org/>. The LSID for this publication is: [zoobank.org:pub:B89FD16E-2084-431A-ACE9-4E4362C6C3CD](http://zoobank.org/pub:B89FD16E-2084-431A-ACE9-4E4362C6C3CD), and for taxonomic registration it is: [urn:lsid:zoobank.org:act:66DBF909-CF1C-479C-BDA0-F324F4FFC15F](http://zoobank.org/act:66DBF909-CF1C-479C-BDA0-F324F4FFC15F). The online version of this work is archived and available from the following digital repositories: PeerJ, PubMed Central SCIE and CLOCKSS”.

## RESULTS

More than 1,000 columnals and pluricolumnals, dozens of disarticulated ossicles from cups and arms, and 26 complete (or nearly complete) cups/calyces were collected. As a result of our investigations, the following taxa were identified: *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* (Bohatý, 2005), *Halocrinites schlotheimii* Steininger, 1831, and *Flexibilia incertae sedis*. Simple discoid holdfasts are also described.

## SYSTEMATIC PALEONTOLOGY

Abbreviations used for specimen measurements include ACH, aboral cup height; ACdistW, distal width of aboral cup; ACmaxW, maximum width of aboral cup; ACproxW, proximal width of aboral cup; BConW, basal concavity width; BH, basal plate height; BW, basal plate maximum width; CrW, crown width; 1stPBH, first primibrachial height; 1stBrW, first primibrachial width; 2ndPBH, second primibrachial height; 2ndPBdistW, second primibrachial distal width; 2ndPBproxW, second primibrachial distal width, 5thPBH, fourth primibrachial height; 3rdPBW, third primibrachial width; 3rdPBH, third primibrachial height; 5thSBW, fifth primibrachial width. All measurements are in mm. Terminology for encrusting organisms follows the recommendations of Taylor & Wilson (2002).

Class Crinoidea *Miller, 1821*  
Subclass Pentacrinoidea *Jaekel, 1918*  
Infraclass Inadunata *Wachsmuth and Springer, 1885*  
Parvclass Cladida *Moore & Laudon, 1943*  
Magnorder Euclidida *Wright, 2017*  
Superorder Cyathiformes *Wright et al., 2017*  
Superfamily Codiacrinoidea *Bather, 1890*  
Family Codiocrinidae *Bather, 1890*  
Subfamily Codiocrininae *Bather, 1890*  
Genus *Codiacrinus* *Schultze, 1866*

### Type species

*Codiacrinus granulatus* *Schultze, 1866*

### Included species

*C. granulatus* (*Schultze, 1866*); *C. nicolli* *Jell & Jell, 1999*; *C. ornatus* (*Prokop, 1973*);  
*C. piriformis* *Le Menn, 1997*; *C. procerus* (*Prokop, 1973*); *P. rarus* (*Jell & Holloway, 1983*);  
*C. robustus* *Le Menn, 1997*; *C. schultzei* *Follman, 1887*; *C. secundus* *Jell, 1999*.

*Codiacrinus sevastopuloi* sp. nov.

[Figs. 3A1–3A4, 3B1–3B3, 3C](#)

### Diagnosis

Aboral cup medium bowl shape; three or more radiating ridges from center of basal plates that project onto radial and infrabasal plates, also very fine nodose sculpturing across calyx plates; basal plates largest plates of aboral cup; radial facets ~50% of distal width of radial plates (arms and column characters not known).

### Types


Holotype: GIUS4-3693/Codiacrinus1; paratypes: GIUS4-3693/Codiacrinus2, GIUS4-3693/Codiacrinus3.

### Occurrence

Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland.

### Description

Aboral cup medium globe shaped ([Figs. 3A4, 3B1](#)), height to width ratio ~1.0, maximum width at middle aboral cup height; three or more radiating ridges from center of basal plates that project onto radial and infrabasal plates ([Fig. 3A1](#)), also very fine nodose sculpturing across calyx plates. Infrabasal circlet ~9% of aboral cup height, extends proximally in a short neck that is truncate proximally with a shallow, circular basal concavity that occupies



**Figure 3** (A1–A4) *Codiocrinus sevastopuloi* sp. nov. GIUS 4-3696/Codiocrinus1, holotype; (A1) lateral view of aboral cup; (A2) basal view of aboral cup, note basal concavity bordered by ridge; (A3), oral view of aboral cup; (A4), lateral lateral view of aboral cup. (B1–B3) *Codiocrinus sevastopuloi* sp. nov. GIUS 4-3696/Codiocrinus2, paratype; (B1) lateral view of aboral cup; (B2) basal view of aboral cup; (B3) oral view of aboral cup. (C) lateral view of an incomplete and compressed specimen of *Codiocrinus sevastopuloi* sp. nov. GIUS 4-3696/Codiocrinus3. (D1–D2) compressed specimen of *Halocrinites* sp. GIUS 4-3696/Hsp; (D1) lateral view of aboral cup with plate boundaries visible; (D2) basal view of cup. (E) *Bactrocrinites* sp. GIUS 4-3696/Bactrocrinites1; lateral view of incomplete aboral cup, note small radial plates and large basal plates. (continued on next page...)

Full-size  DOI: 10.7717/peerj.12842/fig-3



**Figure 3 (...continued)**

(F1–F3) compressed aboral cup of *Halocrinites schlotheimii*. GIUS 4- 3696/Hschloth1 (F1) oblique basal view; (F2, F3) lateral views. All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

~75% of proximal aboral cup width (Figs. 4A2, 4B2). Five pentagonal infrabasal plates, ~3.8 times wider than high, outer surface concave, sculpturing irregular nodose. Basal circlet ~55% of aboral cup height; basal plates largest plates in aboral cup, hexagonal, ~1.2 times higher than wide; sculpturing with radiating ridges and nodes, ridges from near the center of the plates to ridges on adjoining proximal and distal plates. Radial circlet ~36% of aboral cup height; radial plates ~1.2 times wider than high, pentagonal; plate sculpturing with ridges and nodes, ridges diagonal from base of radial facet to like ridges on adjoining basal plates. Radial facets angustary (~52% of radial plate distal width), horseshoe shaped (Figs. 4A3, 4B3). Radial facets, arms, and column unknown.

**Etymology**

The species name is in recognition of the substantial contributions that George D. Sevastopulo made to crinoid paleobiology, as well as paleontology and stratigraphy in general.

**Measurements**


GIUS4-3693/*Codiocrinus*1 (holotype): ACH, 9.8; ACmaxW, 10.4; IH, 1.5; IW, 2.5; BH, 5.4; BW, 7.6; RH, 5.2; RmaxW, 6.0, RdistW, 5.3, RFW, 3.1. GIUS4-3693/*Codiocrinus*2 (paratype): ACH, 11.25; ACmaxW, 11.25\*; IH, 1.4; IW, 5.3; BH, 8.4; BW, 6.8; RH, 5.4; RmaxW, 6.6, RdistW, 4.8, RFW, 2.5.

**Remarks**

One well-preserved and two poorly preserved aboral cups are assigned to *Codiocrinus sevastopuloi* sp. nov. Both poorly preserved specimens have their shapes distorted through compaction.

Ten species of *Codiocrinus*, including *C. sevastopuloi*, are recognized herein. *C. weyeri* is excluded, and it is regarded either as an aberrant individual or a member of another genus. Of these ten species, only three have arms and proximal columnals preserved (*C. robustus*, *C. schultzei*, and *C. secundus*). Thus, species diagnoses are largely based on characters of the aboral cup, which vary widely.

The most noticeable character used to differentiate species of *Codiocrinus* is the aboral cup shape, which may be low bowl, high cone, medium globe, high globe, or medium vase in shape. *Codiocrinus granulatus*, *C. nicolli*, *C. robustus*, and *Codiocrinus sevastopuloi* sp. nov. all have a medium globe-shaped aboral cup. *Codiocrinus granulatus* has a medium globe-shaped aboral cup, two poorly defined radiating ridges from the base of the radial facet onto each subjacent basal plate and perhaps some poorly developed concentric ridges, radial plates are the largest plates in the aboral cup, and the radial facets occupy ~50% of the distal radial plate width. *Codiocrinus nicolli* has a medium globe-shaped aboral cup, very fine nodose sculpturing, radial plates are the largest plates in the aboral cup, and



**Figure 4** Plate drawing of the posterior of *Bactrocrinites* sp. Black pattern, radial plates; cross-hatched pattern, radial plate; dotted pattern and X, anal X plate; dotted line, outer margin of radial facets.

Full-size [DOI: 10.7717/peerj.12842/fig-4](https://doi.org/10.7717/peerj.12842/fig-4)

the radial facets occupy ~60% of the distal radial plate width. *Codiocrinus robustus* has a medium globe-shaped aboral cup, three radiating ridges from the basal plate center and otherwise smooth sculpturing, basal plates are the largest plates in the aboral cup, and the radial facets occupy ~50% of the distal radial plate width.

Alternatively, *Codiocrinus sevastopuloi* sp. nov. has a medium globe-shaped aboral cup, three or more radiating ridges from center of basal plates that project onto radial and infrabasal plates, also very fine nodose sculpturing across calyx plates, basal plates are the largest plates in the aboral cup, and the radial facets occupy ~50% of the distal radial plate width.

Superfamily Dendrocrinacea *Wachsmuth and Springer, 1886*

Family Dendrocrinidae *Wachsmuth and Springer, 1886*

Genus *Bactrocrinites* *Schnur in Steininger, 1849*

### Type species

*Poteriocrinus fusiformis* *Roemer, 1844*.

### Included species

*Bactrocrinites birmanicus* *Reed, 1908*; *B. cyathus* (*Schmidt, 1942*); *B. depressus* (*Schultze, 1866*); *B. fieldi* (*Springer & Slocum, 1906*); *B. fusiformis* (*Roemer, 1844*); *B. jaekeli* (*Schmidt, 1934*); *B. muelleri* (*Jaekel, 1895*); *B. oklahomaensis* *Strimple, 1952*; *B. onondagensis* *Goldring,*

1954; *B. penaneachensis* Le Menn, 1985; *B. porectus* (Bohatý, 2005); *B. tenuis* (Jaekel, 1895); *B.?* *trabicus* (Schmidt, 1934); *B. zeileri* (Mueller in Zeiler & Wirtgen, 1855).

*Bactrocrinites* sp.

Figs. 3E, 4

## Description

Relatively large aboral cup, aboral cup plates with pustulose plate sculpturing (Fig. 3E). Infrabasal plates not known. Basal plates partially preserved, inferred to be the dominant plate circlet in aboral cup (Fig. 3E). C radial plate supported beneath by radianal plate and BC basal plate; D radial plate larger than C radial plate, supported beneath by CD and DE basal plates. Radial facets large, semicircular, angustary, declivate. Two anal plates in aboral cup. Radianal presumably tetragonal, below and to the left of the C radial plate and supports the anal X plate on the upper left (Fig. 4). Anal X plate hexagonal, supported beneath by the CD basal plate and the radianal, separates and articulates with lateral sides of the C and D radial plates.

Other aspects of the aboral cup, anal sac, arms, and column are not known.

## Occurrence

Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland.

## Remarks

Species diagnostic characters for Devonian *Bactrocrinites* include shape of the aboral cup, plate sculpturing, relative heights of aboral cup plates, and the dimensions of the infrabasal and basal plates. Unfortunately, the single specimen of *Bactrocrinites* from the Emsian of Poland is not complete (GIUS4-3693/*Bactrocrinites*), so aboral cup shape, relative proportions of aboral cup plates and the dimensions of the basal plates cannot be determined. The pustulose aboral cup plate sculpturing and what are inferred to be prominent (high) basal plates most closely ally this specimen with *B. fusiformis*. However, a more complete accounting of the morphology of this Emsian specimen is required before a confident species assignment can be made.

Superfamily Gasterocomoidea Roemer, 1854

Family Cupressocrinitidae Roemer, 1854

Subfamily Cupressocrinitinae Bohatý, 2006

Genus *Halocrinites* Steininger, 1831

## Type species

*Halocrinites schlotheimi schlotheimi* Steininger, 1831.

## Included species

*Halocrinites altus* (Schultze, 1866); *H. assimilis* (Dubatolova, 1964); *H. geminatus* (Bohatý, 2005); *H. gibber* (Bather, 1919); *H. heinorum* Bohatý & Ausich (2021); *H. inflatus inflatus* (Schultze, 1866); *H. inflatus convexus* (Hauser, 2001); *H. inflatus cuneatus* (Bohatý, 2006); *H. inflatus depressus* (Hauser, 2001); *H. minor* (Schultze, 1866); *H. nodosus* (Sandberger and Sandberger, 1856); *H. rectangularis* (Schmidt, 1941); *H. schlotheimii schlotheimii* (Steininger, 1831); *H. schlotheimii granulatus* Schultze, 1866; *H. schreueri* Bohatý, 2006; *H. tesserula* (Goldfuss, 1831); *H. townsendi* (König, 1825); and *H. urogali* Roemer, 1850.


## Remarks

As discussed in Bohatý & Ausich (2021), generic and specific assignments of the Cupressocrinitidae have been varied, commonly changed, and confused until recently (e.g., Bohatý, 2005; Bohatý, 2006; Bohatý, 2009b; Bohatý & Herbig, 2010; Bohatý & Ausich, 2021). Two species of *Halocrinites* are recognized from the Bukowa Góra Member in Poland, including *H. geminatus* (Bohatý, 2005) and *H. schlotheimii* (Steininger, 1831). Most *Halocrinites* specimens from Poland are lacking the exoplacoid layer or are sufficiently worn that the character of the exoplacoid layer cannot be determined. One exception is specimen GIUS 4-3696Hscholth6 (Fig. 5E), although even this specimen is worn. They are differentiated on the basis of aboral cup shape, basal plate morphology, size of the infrabasal cirlet relative to the size of the proximalmost columnal, and the size of the basal concavity, as described below. *Halocrinites schlotheimii* has a bowl-shaped aboral cup with a ratio of aboral cup diameter versus crown height  $\sim 1:1.15\text{--}2.0$ ; aboral cup typically  $\sim 2.0$  times wider than high; infrabasal plates fused into a single pentagonal plate that is confined to the basal concavity; brachials wider than high (height to width ratio  $\sim 1:2.0\text{--}2.5$ ); proximal columnal circular not filling entire basal concavity. In contrast, *H. geminatus* aboral cup bowl to moderately conical in shape; ratio of aboral cup diameter versus crown height  $\sim 1:1.15\text{--}2.0$ ; typically 2.0 times wider than high; infrabasal plates fused into a single pentalobate plate that is confined to the basal concavity; brachials wider than high (height to width ratio  $\sim 1:2.0\text{--}2.5$ ); proximal columnal circular not filling entire basal concavity.

Similar to the Cupressocrinitidae described by Bohatý (2009b) and Bohatý & Ausich (2021), *Halocrinites* from Poland have a variety of epizoans encrusting the outer surface of crown plates. These include trepostome bryozoans encrusted on aboral cup and brachial plates. A presumable microconchid that is attached to a radial plate, and a juvenile pelmatozoan holdfast is attached to a different radial plate. These encrustations did not induce a recognizable response from the crinoid host, so it is probable that these encrustations occurred after the death of the crinoid and, thus, are episkeletozoans (see Taylor & Wilson, 2002).

*Halocrinites schlotheimii* Steininger, 1831

Figs. 3F1–F3, 5A1–A3, 5B1–B2, 5C1–C2, 53, 6A1, A2



**Figure 5** (A1–A3, B1–B2, C1–C2) *Halocrinites schlotheimii* GIUS 4-3696/Hschloth2, 5 and 4 respectively; (A1, B1, C1) lateral views of aboral cup; (A2, B2, C2) basal views of aboral cup; (A3) oral view of aboral cup. (D, H) *Halocrinites* gem GIUS 4-3696/Hgem2 and 1 respectively; (D1) lateral view of aboral cup; (D2) basal view of aboral cup. (E) *Halocrinites* with good exoplacoid sculpturing preserved; GIUS 4-3696/Hschloth6. (F, G) Crinoidea indeterminate (presumably remains of a camerate crinoid.) (continued on next page...)

Full-size  DOI: 10.7717/peerj.12842/fig-5

**Figure 5 (...continued)**

GIUS 4-3696/indet1 and 2 respectively. All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

**Type**

The type specimens for this taxon are not known.

**Diagnosis**

*Halocrinites* with bowl-shaped aboral cup with a ratio of aboral cup diameter versus aboral cup height  $\sim 1:1.15\text{--}2.0$ ; aboral cup typically  $\sim 2.0$  times wider than high; infrabasal plates fused into a single pentagonal plate that is confined to the basal concavity; brachials wider than high (height to width ratio  $\sim 1:2.0\text{--}2.5$ ); proximal columnal circular not filling entire basal concavity.

**Occurrence**

In Poland, *H. schlotheimii* is from the Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, Poland. Previously, this species has been described from the Eifelian and Givetian of Australia, China, Germany, Poland, and Spain ([Webster & Webster, 2019](#)).

**Description**

Calyx medium sized. Aboral cup low to very low bowl shape in adults with height to maximum width ratio  $0.44\text{--}0.66$  ([Fig. 3F](#)); perfect pentameral symmetry; deep, subpentalobate basal concavity occupies  $62\text{--}66\%$  of proximal aboral cup width ([Fig. 5E](#)). Plates gently convex; coarse multilaminar exoplacoid sculpturing preserved on only a few specimens.


Infrabasal circling completely in basal concavity, pentagonal. Infrabasal plates presumably five. Basal circling  $\sim 58\text{--}60\%$  of aboral cup height, present on base and on vertical sides of aboral cup ([Figs. 3D, 3F, 5A–5C](#)). Five basal plates, equal in size, wider than high, smaller than radial plates. Radial circling  $40\text{--}42\%$  of calyx height ([Fig. 6A](#)). Radial plates five, pentagonal, largest plates in aboral cup, height to width ratio  $0.48\text{--}0.58$ . Radial facets plenary, planate; radial facet topography not known. Posterior interray plates absent from aboral cup; anal sac, if present, unknown.

Arms five, atomous, brachials uniserial; V-shaped in cross section across width of brachial plate; incompletely known (preserved only through sixth primibrachial) ([Fig. 6A](#)). First primibrachial (articular plate, see [Bohatý & Ausich, 2021](#)), very low, full width of radial facet; subsequent brachials, flat sided, equal in width to distal edge of second primibrachials; height to width ratio  $\sim 0.65$ .

Proximal column narrow, attachment to base of aboral cup circular, occupies slightly more than one half of infrabasal circling; remainder of column unknown.

**Measurements**

GIUS4-3693/Hschloth3: CrH,  $33.0^*$ ; ACH, 7.5; ACmaxW, 11.3; 4thPBH, 6.0; 4thPBW, 4.0. GIUS4-3693/Hschloth4: ACH, 6.3; ACdistW, 14.0; BH, 6.3; BW, 6.5; RH, 4.3; RW, 8.9.



**Figure 6** (A1–A2) *Halocrinites schlotheimii*; (A1) lateral view of partial crown, aboral cup plate boundaries distinct; (A2) basal view of a moderately compressed aboral cup. GIUS 4-3696/Hschloth3. (B–D) Crinoidea indeterminate; GIUS 4-3696/Hschloth3. (B–D) Crinoidea indeterminate; GIUS 4-3696/indet3, 4 and 5 respectively; (B, C) partial arms of an unknown cladid crinoids; (D) in upper left of specimen a pentalobate column presumably infrabasals plates attached; probably an unknown cladid). All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

Full-size DOI: 10.7717/peerj.12842/fig-6

## Remarks

Six specimens of *H. schlotheimii* are known from the Emsian of Poland (GIUS4-3693/Hschloth1 to GIUS4-3693/Hschloth6). In the collection of Polish specimens, small individuals tend to have more pronounced convex basal plates that nearly produce a central node.

*Halocrinites geminatus* Bohatý, 2005  
Figs. 5D, 5H

## Type

Holotype is SMF 75308 (see Bohatý, 2005).

## Diagnosis

*Halocrinites* with aboral cup bowl to moderately conical in shape; ratio of aboral cup diameter versus aboral cup height  $\sim 1:1.15\text{--}2.0$ ; typically 2.0 times wider than high; infrabasal plates fused into a single pentalobate plate that is confined to the basal concavity; brachials wider than high (height to width ratio  $\sim 1:2.0\text{--}2.5$ ; proximal columnal circular not filling entire basal concavity.

## Occurrence

In Poland, *H. geminatus* is from the Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains. Previously, it was known from the Eifelian to early Givetian of Germany (Webster & Webster, 2019).

## Description

Calyx medium sized. Aboral cup very low bowl shape (Fig. 5D1), height to maximum width ratio 0.43; perfect pentameral symmetry; shallow, subpentagonal basal concavity occupies  $\sim 73\%$  of proximal aboral cup width (Fig. 5D2). Plates gently convex with multilaminar exoplacoid sculpturing (see Gluchowski, 1993a).

Infrabasal circlet completely concealed in basal concavity; outer margin of basal concavity subtetragonal, entirely covered by proximal columnal. Infrabasal plates presumably five. Basal circlet  $\sim 56\%$  of aboral cup height, present on base and on vertical sides of aboral cup. Five basal plates, equal in size, wider than high, much smaller than radial plates. Radial circlet  $\sim 44\%$  of calyx height. Radial plates five, pentagonal, largest plates in aboral cup, height to width ratio 0.60. Radial facets plenary, planate; radial facet topography not known. Posterior interray plates absent from aboral cup; anal sac, if present, unknown.

Arms five, atomous, brachials uniserial (Fig. 5H); V-shaped in cross section across width of brachial plate; incompletely known (preserved only through sixth primibrachial). First primibrachial (articular plate, see Bohatý & Ausich, 2021), very low (height to width ratio 0.16), full width of radial facet; subsequent brachials, flat sided, equal in width to distal edge of second primibrachials; height to width ratio  $\sim 1.2$ .

Proximal column attachment to base of aboral cup wide, tetralobate, fills entire basal concavity covering infrabasal plates; remainder of column unknown.



### Measurements

GIUS4-3969/Hgem1: CrH, 41.0\*; ACH, 9.0; ACmaxW, 21.0; ACproxW, 9.0; BConW, 6.5; BH, 7.6; BW, 7.25; RH, 5.9; RW, 12.4; 1stPbH, 1.5; 1stPbW, 13.0; 2ndPbH, 6.0; 2ndPbproxW, 13.0; 2ndPbdistW, 10.0; 3rdPBH, 5.9.0; 3rdPBW, 5.0.

### Remarks

[Bohatý \(2005\)](#) illustrated individuals of *H. geminatus* with a wide variety of shapes. The description above is for the Poland specimens that all have a very low bowl shape.

Crinoidea Incertae Sedis

[Figs. 5F, 5G, 6B–6D, 7A–7I, 8C, 8D,](#)

### Occurrence

Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland.

### Remarks

A single brachial plate is identified as a flexible crinoid (GIUS4-3693/flexible; [Fig. 7I](#)). The brachial plate is ~4.0 times deeper than high with, only the distal facet is visible and part of the sides of the plate are visible that includes the aboral indentation where a patelloid process from the distal adjoining brachial would reside. The brachial plate is as wide as deep. A crenulated articular ridge is present along the abaxial portion of the facet, and the lateral sides of the facet are crenulated. A narrow, shallow aboral groove is present along the adaxial margin of the facet.

Poorly preserved and unidentifiable remains of several additional taxa also occur in the Bukowa Góra Member. In addition to the flexible crinoid and holdfasts mentioned below, others include camerate crinoids ([Figs. 5F, 5G](#)) and various distinctive cladid crinoids ([Figs. 6B–6D](#)). Distinctive columnals and pluricolumnals are also present ([Figs. 7A, 8C, 8D](#)). The pluricolumnals illustrated in [Fig. 8C](#) undoubtedly belong to the Platycrinidae and may be *Platycrinites minimalis* (col.) ([Głuchowski, 1993a](#)).

Simple discoid holdfasts


[Figs. 8A, 8B](#)

### Occurrence

Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, Poland.

### Remarks

Solitary rugose coral and a presumable stromatoporoid specimens associated with the described crinoids have small, discoid holdfasts cemented to their outer surface ([Figs. 8A, 8B](#)). These holdfasts are subcircular in outline and some have a slightly digitate outer



**Figure 7** (A–H) crinoid pluricolumnals. GIUS 4-3696/indet6-13. (A, B) pluricolumnal with numerous nodes around the periphery of each columnal; (C) set of pluricolumnals; (D) pluricolumnal with nodes and perhaps some spines around the periphery of each columnal; (E) numerous pluricolumnals of a column that lacks nodes; (F) lateral view of three-columnal pluricolumnal with a few nodes around the periphery of columnals that are offset in position from one columnal to the next; (continued on next page...)

Full-size  DOI: [10.7717/peerj.12842/fig-7](https://doi.org/10.7717/peerj.12842/fig-7)

**Figure 7 (...continued)**

(G) columnal facet with a narrow peripheral lumen and a narrow, raised perilumen; (H) columnal with a wide lumen and a narrow, raised perilumen; (I) Flexible crinoid brachial. GIUS 4-3696/flexible; note crenulate sides and on the upper margin a notch to hold a patelloid process from the next highest columnal. All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. All scale bar equals 10 mm but in case of flexible crinoid brachial it is 1 mm.


margin. In one example, the holdfast articulation to the column was canted toward the long axis of a rugose coral, suggesting the crinoid was encrusted to the coral when both were alive. Therefore, these holdfasts should be considered epizoozoans ([Taylor & Wilson, 2002](#)). It is not possible to speculate on the identity of the crown that was attached to these holdfasts, and the smaller specimens may have been juveniles or from multiple radices of a single adult.

## CONCLUDING REMARKS

The first Emsian crinoids described on the basis of aboral cups and crowns are reported here from the Bukowa Góra Member in the Holy Cross Mountains of southern Poland. Named taxa include *Bactrocrinites* sp., *Codiocrinus sevastopuloi* sp. nov., *Halocrinites geminatus* ([Bohatý, 2005](#)), *Halocrinites schlotheimii* ([Steininger, 1831](#)). Taxa that can only be recognized as incertae sedis include one flexible crinoid, as many as three camerate crinoids, as many as four additional cladid crinoids, and a number of distinctive holdfasts, columnals, and pluricolumnals that cannot be matched with the crown to which they were attached. Additional collecting in the Bukowa Góra Member should yield remains of many crinoids.

Previously, *Halocrinites* (including *H. schlotheimii*) have been described from younger Devonian strata in Poland (see [Fig. 1](#)). Further, species of *Bactrocrinites*, *Codiocrinus*, and *Halocrinites* occur in other Devonian crinoid faunas from Germany and Spain ([Webster & Webster, 2019](#)).

The new crinoids reported here are from the Bukowa Góra Member of Poland (Emsian) and are an extension of the Lower to Middle Devonian crinoid faunas from across Europe, which are best represented by Emsian to Givetian crinoids from Germany and Spain (e.g., [Bohatý, 2005](#); [Bohatý, 2006](#); [Bohatý, 2009b](#); [Bohatý & Herbig, 2010](#); [Hauser, 2001](#); [Hauser, 2002](#); [Hauser, 2007](#)). *Halocrinites* has been reported from Germany, Spain, Belgium, and Russia (Eifelian–Frasnian); whereas both *Codiocrinus* and *Bactrocrinites* have longer ranges and are cosmopolitan in distribution. In addition, to Western Europe, *Codiocrinus* is known from Gondwana terrane (northern Africa and Australia). The oldest recognized species of *Bactrocrinites* is from the middle Silurian of North America, and this genus is only known from North America and Europe. As known, *Bactrocrinites* became extinct at the Givetian–Frasnian extinction and *Bactrocrinites* and *Halocrinites* became extinct at the Frasnian–Famennian extinction.



**Figure 8** (A, B) Crinoid holdfast on stromatoporoids. GIUS 4-3696/holdfast1 and 2, respectively. (C) Crinoid pluricolumnal presumably from the dististele with broken radices; (D) pluricolumnal with elliptical columnals, presumably from a member of the Platycrinitidae; GIUS 4-3696/indet14 and 15 respectively. All specimens are from Bukowa Góra Member (Emsian), Bukowa Góra quarry, Holy Cross Mountains, southern Poland. Scale bar equals 10 mm.

Full-size  DOI: [10.7717/peerj.12842/fig-8](https://doi.org/10.7717/peerj.12842/fig-8)

## ACKNOWLEDGEMENTS

We thank Michał Stachacz, Kamil Moskowczenko, Damian Kuźma, Janusz Kucharski and Mateusz Syncerz for help during the field work. Jan Bohatý generously helped with identifications of taxa. Paweł Bącal is acknowledged for SEM imaging. We also thank Jan Bohatý and David L. Meyer, who provided critical reviews that significantly improved this paper.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the NCN (National Science Centre, Poland) Grant no. 2018/31/B/ST10/00387. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
NCN (National Science Centre, Poland): 2018/31/B/ST10/00387.

### Competing Interests

Bartosz Płachno is an Academic Editor for PeerJ.

### Author Contributions

- William I. Ausich, Bartosz J. Płachno and Przemysław Gorzelak conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Mariusz A. Salamon conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Tomasz Brachaniec performed the experiments, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Wojciech Krawczyński conceived and designed the experiments, prepared figures and/or tables, and approved the final draft.
- Andrzej Boczarowski and Magdalena Łukowiak performed the experiments, prepared figures and/or tables, and approved the final draft.
- Karolina Paszcza performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

*Codiocrinus sevastopuloi* sp. nov.

Holotype: GIUS4-3693/Codiocrinus1; paratypes: GIUS4-3693/Codiocrinus2, GIUS4-3693/Codiocrinus3.

The specimen is deposited in the University of Silesia in Katowice, Faculty of Natural Sciences, Institute of Earth Sciences, Poland.

## New Species Registration

The following information was supplied regarding the registration of a newly described species:

Publication LSID: urn:lsid:zoobank.org:pub:B89FD16E-2084-431A-ACE9-4E4362C6C3CD

Codiacrinus sevastopuloi LSID: urn:lsid:zoobank.org:act:66DBF909-CF1C-479C-BDA0-F324F4FFC15F

## REFERENCES

- Almela A, Revilla J. 1950.** Especies fósiles nuevas del Devoniano de León. *Notas y Comunicaciones del Instituto Geológico y Minero de España* **20**:45–60.
- Bather FA. 1890.** British fossil crinoids. II, The classification of the Inadunata Fistulata (cont'd). *Annals and Magazine of Natural History, Ser. 6* **5**:373–388  
[DOI 10.1080/00222939009460849](https://doi.org/10.1080/00222939009460849).
- Bather FA. 1919.** *Cupressocrinus gibber* n. sp. du Dévonien supérieur de Belgique. *Bulletin de la Société Belge de Géologie, de Paléontologie et l'Hydrologie* **28**:129–136.
- Bohatý J. 2005.** Doppellagige Kronenplatten: Ein neues anatomisches Merkmal paläozoischer Crinoiden und Revision der Familie Cupressocrinitidae (Devon). *Paläontologische Zeitschrift* **79**(2):201–225 [DOI 10.1007/BF02990185](https://doi.org/10.1007/BF02990185).
- Bohatý J. 2006.** Neue Cupressocrinitidae (Crinoidea) aus den mitteldevonischen Kalkmulden der Eifel (linksrheinisches Schiefergebirge, Deutschland). *Senckenbergiana Lethaea* **86**(2):151–189 [DOI 10.1007/BF03043488](https://doi.org/10.1007/BF03043488).
- Bohatý J. 2009a.** First report of aboral cups of the crinoid family Parahexacrinidae from the Middle and Upper Devonian of the Eifel (Germany) and the Holy Cross Mountains (Poland). *Paleontological Journal* **43**:1569–1577  
[DOI 10.1134/S0031030109120028](https://doi.org/10.1134/S0031030109120028).
- Bohatý J. 2009b.** Pre- and postmortem skeletal modifications of the Cupressocrinitidae (Crinoidea, Cladida). *Journal of Paleontology* **83**:45–62 [DOI 10.1666/07-125R.1](https://doi.org/10.1666/07-125R.1).
- Bohatý J. 2011.** Revision of the flexible crinoid genus *Ammonicrinus* and a new hypothesis on its life mode (Crinoidea, Flexibilia). *Acta Palaeontologica Polonica* **56**(3):615–639 [DOI 10.4202/app.2010.0020](https://doi.org/10.4202/app.2010.0020).
- Bohatý J, Ausich WI. 2021.** Revision of two Devonian cupressocrinitids from the Schultze Collection (Museum of Comparative Zoology, Harvard University) and description of a new *Halocrinites* (Crinoidea, Euclidida). *Journal of Paleontology* 1–17 [DOI 10.1017/jpa.2021.65](https://doi.org/10.1017/jpa.2021.65).
- Bohatý J, Herbig H-G. 2010.** Middle Givetian echinoderms from the Schlade Valley (Rhenish Massif, Germany): habitats, taxonomy and ecostratigraphy. *Paläontologische Zeitschrift* **84**:365–386 [DOI 10.1007/s12542-010-0072-y](https://doi.org/10.1007/s12542-010-0072-y).
- Dames W. 1868.** Über die in der Umgebung Freiburgs in Nieder-Schlesien auftretenden devonischen Ablagerungen. *Zeitschrift der Deutschen Geologischen Gesellschaft* **21**:469–508.
- Dubatolova YuA. 1964.** *Morskije lilii devona Kuzbassa*. Akademiya Nauk SSSR, Sibirskoe Otdeleniye Trudy Instituta Geologii i Geofiziki, 153 pp.

- Fijałkowska-Mader A, Malec J. 2011.** Biostratigraphy of the Emsian to Eifelian in the Holy Cross Mountains (Poland). *Geological Quarterly* **55**:109–138.
- Fijałkowska-Mader A, Malec J, Tarnowska M, Turnau E. 1997.** Stratygrafia dolnego dewonu w rejonie Bodzentyna –region łysogórski Gór Świętokrzyskich. *Posiedzenia Naukowe Państwowego Instytutu Geologicznego* **53**:122–125.
- Filipiak P. 2011.** Palynology of the Lower and Middle Devonian deposits in southern and central Poland. *Review of Palaeobotany and Palynology* **166**:213–252  
DOI [10.1016/j.revpalbo.2011.05.010](https://doi.org/10.1016/j.revpalbo.2011.05.010).
- Follman O. 1887.** Unterdevonische Crinoidea, Verhandlungen des Naturhistorischen Vereins der Preussischen Rheinlande, Westfalens, and des Regierungs-bezirks Osnabruck, 26 pp.
- Głuchowski E. 1980.** New taxa of Devonian and Carboniferous crinoid stem parts from Poland. *Bulletin of the Polish Academy of Science* **28**:43–49.
- Głuchowski E. 1981a.** Stratigraphic significance of Paleozoic crinoid columnals from Poland. *Zeszyty Naukowe AGH* **7**:89–110.
- Głuchowski E. 1981b.** Paleozoic crinoid columnals and pluricolumnals from Poland. *Zeszyty Naukowe AGH* **7**:29–57.
- Głuchowski E. 1981c.** Człony łodyg liliowców z serii skalskiej (żywet) Gór Świętokrzyskich, Sprawozdanie z posiedzeń Komisji Nauk Geologicznych. *PAN* **22**:428–429.
- Głuchowski E. 1993a.** Crinoid assemblages in the Polish Givetian and Frasnian. *Acta Palaeontologica Polonica* **39**:35–92.
- Głuchowski E. 1993b.** Upper Emsian crinoids from the Bukowa Góra quarry in the Klonów Range, Holy Cross Mts. *Prace Naukowe Uniwersytetu Śląskiego*, 1331. *Geologia* **12/13**:159–174.
- Głuchowski E. 2002.** Crinoids from the Famennian of the Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* **47**:319–328.
- Głuchowski E. 2003.** Gromada Crinoidea (Miller, 1821). In: Malinowska L, ed. *Budowa Geologiczna Polski. Atlas skamieniałości przewodnich i charakterystycznych. T. III. Dewon*. Warszawa: Państwowy Instytut Wydawniczy, 458–475.
- Głuchowski E, Casier J-G, Olempska E. 2006.** Crinoid and ostracod succession within the Early–Middle Frasnian interval in the Wietrznia quarry, Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* **51**:695–706.
- Goldfuss GA. 1826-1844.** Petrefacta Germaniae, tam ea, Quae in Museo Universitatis Regiae Borussicae Fridericiae Wilhelmae Rhenanea, serventur, quam alia quaecunque in Museis Hoeninghusiano Muensteriano aliisque, extant, iconibus et descriptions illustrata. –Abbildungen und Beschreibungen der Petrefacten Deutschlands und der Angränzende Länder, unter Mitwirkung des Hern Grafen Georg zu Münster, herausgegeben von August Goldfuss. v. 1 (1826–1833), Divisio prima. Zoophytorum reliquiae, p. 1–114; Divisio secunda. Radiariorum reliquiae, p. 115–221 [Echinodermata]; Divisio tertia. Annulatorium reliquiae, p. 222–242; v. 2 (1834–40), Divisio quarta. Molluscorum acephalicorum reliquiae. I. Bivalvia, p. 65–286; II. Brachiopoda, p. 287–303; III. (1841–44), Divisio quinta. Molluscorum gasteropodum reliquiae, p. 1–121; Düsseldorf, Arnz & Co.

- Goldring W. 1954.** Devonian crinoids, new and old, II. *New York State Museum Circular* 37:51 pp.
- Gorzalak P, Gluchowski E, Salamon MA. 2014.** Reassessing the improbability of a muscular crinoid stem. *Scientific Reports* 4:4069.
- Gürich G. 1896.** Das Paläozoicum im polnische Mittelgebirge. In: *Russischkaiserlichen mineralogischen gesellschaft zu. St. Petersburg* 2, Egger und cie, 168 pp.
- Hauser J. 2001.** Neubeschreibung mitteldevonischer Eifelcrinoiden aus der Sammlung Schultze (Museum of Comparative Zoology, The Agassiz Museum, Harvard University, Massachusetts, USA) nebst einer Zusammenstellung der Eifelcrinoiden (Holotypen) der Goldfuss-Sammlung. Published by author, Bonn, 86 pp.
- Hauser J. 2002.** *Die Crinoiden der Frasnies-Stufe (Oberdevon) von Wallerheim/Loch (Prümer Mulde, Eifel) nebst einer Zusammenstellung sämtlicher Melocriniten weltweit.* Bonn: Published by author, 69 pp.
- Hauser J. 2007.** *Die Crinoidenwelt der Eifel vor 350.000.000 Jahren-Neue Crinoiden aus dem Mitteldevon der Eifel, Teil 2.* Bonn: Published by author, 83 pp.
- Jaekel O. 1895.** Beiträge zur Kenntniss der palaeozoischen Crinoiden Deutschlands. *Paläontologische Abhandlungen* 7:1–116.
- Jaekel O. 1918.** Phylogenie und System der Pelmatozoen. *Paläontologische Zeitschrift* 3:1–128.
- Jell PA. 1999.** Silurian and Devonian crinoids from central Victoria. *Memoirs of the Queensland Museum* 43:114.
- Jell PA, Holloway DJ. 1983.** Devonian and ?Late Silurian palaeontology of the Winneke Reservoir site, Christman Hills, Victoria. *Proceedings of the Royal Society of Victoria* 95:1–21.
- Jell PA, Jell JS. 1999.** Crinoids, a blastoid and a cyclocystoid from the Upper Devonian reef complex of the Canning Basin, Western Australia. *Memoirs of the Queensland Museum* 43:201–236.
- Kongiel R. 1958.** Nowy gatunek *Ammonicrinus* i jego występowanie w Polsce. *Prace Muzeum Ziemi* 2:31–40.
- König CDE. 1825.** *Icones fossilium sectiles.* London, Londini in vico Regent Street, 4.
- Le Menn J. 1985.** Les crinoides du Dévonien inférieur et moyen du massif Armoricaïn. *Mémoires de la Société Géologique et Minéralogique de Bretagne* 30:268.
- Le Menn J. 1997.** Crinoïdes devoniens d’Afrique du nord, revision systematique et affinites paleobiogeographiques. *Annales de la Société Géologique du Nord, 2nd ser* 5:129–139.
- Lobanowski H. 1971.** The Lower Devonian in the western part of the Klonów belt (Holy Cross Mts), Part 1. Upper Emsian. *Acta Geologica Polonica* 21:629–686.
- Lobanowski H. 1981.** Bukowa Góra quarry: lower Devonian sandstones. In: Żakowa H, ed. *Przewodnik 53 Zjazdu Polskiego Towarzystwa Geologicznego w Kielcach.* Warszawa: Wydawnictwa Geologiczne, 249–255.
- Malec J. 2005.** Lithostratigraphy of the Lower and Middle Devonian boundary interval in the Łysogóry Region of the Holy Cross Mts (Poland). *Biuletyn Państwowego Instytutu Geologicznego* 415:5–28.



- Marynowski L, Salamon MA, Narkiewicz M. 2002.** Thermal maturity and depositional environments of organic matter in the post-Variscan succession on the Holy Cross Mountains. *Geological Quarterly* **46**:25–36.
- Miller JS. 1821.** A natural history of the Crinoidea, or lily-shaped animals; with observations on the genera, Asteria, Euryale, Comatula and Marsupites. Bristol, C. Frost, 150 pp.
- Moore RC, Jeffords RM. 1968.** Classification and nomenclature of fossil crinoids based on studies of dissociated parts of their columns. *The University of Kansas Paleontological Contributions* **46**:1–86.
- Moore RC, Laudon LR. 1943.** Evolution and classification of Paleozoic crinoids. *Geological Society of America Special Paper* **46**:151 pp.
- Piotrowski A. 1977.** Genus *Ammonicrinus* (Crinoidea) from the Middle Devonian of the Holy Cross Mountains, Poland. *Acta Palaeontologica Polonica* **22**:205–218.
- Prokop RJ. 1973.** *Elicrinus* n. gen. from the Lower Devonian of Bohemia (Crinoidea). *Věstník Ustředního Ústavu Geologického* **48**(4):221–224.
- Reed FRC. 1908.** The Devonian faunas of the Northern Shan states. *India Geological Survey, Memoirs* **2**(5):157 pp.
- Roemer CF. 1844.** *Das Rheinische Übergangsgebirge*. Hannover, Hahn: Eine palaeontologisch-geognostische Darstellung, 96 pp.
- Roemer CF. 1850.** Über *Stephanocrinus*, eine fossile Crinoiden-Gattung aus der Familie der Cystideen. *Wiegmanns Archiv für Naturgeschichte, Jahrgang* **16**:365–375.
- Roemer CF. 1852–1854.** Erste Periode, Kohlen-Gebirge. In: HG Bronn, ed. *Lethaea Geognostica*. 1851–1856, 3rd edition. 2. Stuttgart, E. Schweizerbart: 788 pp.
- Salamon MA, Gerrienne P, Steemans P, Gorzelak P, Filipiak P, Hérissé AL, Paris F, Cascales-Miñana B, Brachaniec T, Misz-Kennan M, Niedzwiedzki R, Trela W. 2018.** Putative Late Ordovician land plants. *New Phytologist* **4**:1305–1309 DOI [10.1111/nph.15091](https://doi.org/10.1111/nph.15091).
- Sandberger G, Sandberger F. 1850-1856.** Die Versteinerungen des rheinischen Schichtensystems in Nassau. In: *Wiesbaden, Kreidel and Niedner*. 564 pp.
- Schmidt WE. 1934.** Die Crinoideen des Rheinischen Devons, I. Teil; Die Crinoideen des Hunsrückschiefers. *Abhandlung der Preussischen Geologischen Landesanstalt* **163**:149.
- Schmidt WE. 1941.** Die Crinoideen des Rheinischen Devons II. Teil. A. Nachtrag zu: Die Crinoideen des Hunsrückschiefers; B. Die Crinoideen des Unterdevons bis zur *cultrijugatus*-Zone (mit Ausschluss des Hunsrückschiefers). *Abhandlungen der Preussischen Geologischen Landesanstalt* **182**:253.
- Schmidt WE. 1942.** Die Crinoideen des Rheinischen Devons. II. Teil; A. Nachtrag zu Die Crinoideen des Hunsrückschiefers; B. Die Crinoideen des Unterdevon bis zur *Cultrijugatus* -Zone (mit Ausschluss des Hunsrückschiefers). *Abhandlungen der Reichsstelle für Bodenforschung, n. s.* **182**:253.
- Schultze LJT. 1866.** *Monographie der Echinodermen des Eifler Kalkes*. Vienna: Carl Gerold's Sohn, 118 pp.
- Sobolev D. 1909.** Sredniy devon kelecko-sandomirskoho krjaza. *Materialy dlia Geologii Rossiji* **24**:41–536.

- Springer F, Slocum AW. 1906.** *Hypsocrinus*, a new genus of crinoids from the Devonian. *Field Columbia Museum Publication*, 114, *Geology Series* **2(9)**:267–271.
- Steininger J. 1831.** *Bemerkungen über die Versteinerungen, welche in dem Uebergangs-Kalkgebirge der Eifel gefunden werden.* Trier: Eine Beilage zum Gymnasial-Programm zu Trier, 44 pp.
- Steininger J. 1849.** Die Versteinerungen des Uebergangs-Gebirges der Eifel. *Jahresbericht über den Schul-Cursus 1848/49 an dem Gymnasium zu Trier* **2**:1–22.
- Strimple HL. 1952.** Some new species of crinoids from the Henryhouse Formation of Oklahoma. *Journal of the Washington Academy of Science* **42**:75–79.
- Stukalina GA. 1965.** Novye vidji *Hexacrinites* tsentralnogo Kazakhstana [New species of *Hexacrinites* from central Kazakhstan]. *Ezhogodnik Vsesoyuznogo Paleontologičeskogo Obščestva* **17**:188–195 (In Russian).
- Szulcowski M. 1995.** Depositional evolution of the Holy Cross Mts. (Poland) in the Devonian and Carboniferous – a review. *Geological Quarterly* **39**:159–174.
- Szulcowski M, Porębski S. 2008.** Stop 1 –Bukowa Góra. Lower Devonian. In: Pieńkowski G, Uchman A, eds. *Ichnological Sites of Poland –the Holy Cross Mountains and the Carpathian Flysch: The Second International Congress on Ichnology, The Pre-Congress and Post-Congress Field Trip Guide Book.* Cracow, 18–37.
- Taylor PD, Wilson MA. 2002.** A new terminology for marine organisms inhabiting hard substrates. *Palaios* **17**:522–525  
DOI [10.1669/0883-1351\(2002\)017<0522:ANTFMO>2.0.CO;2](https://doi.org/10.1669/0883-1351(2002)017<0522:ANTFMO>2.0.CO;2).
- Wachsmuth C, Springer F. 1880–1886.** Revision of the Palaeocrinoidea. Proceedings of the Academy of Natural Sciences of Philadelphia Pt. I. The families Ichthyocrinidae and Cyathocrinidae (1880), p. 226–378, (separate repaged p. 1–153). Pt. II. Family Sphaeroidocrinidae, with the sub-families Platycrinidae, Rhodocrinidae, and Actinocrinidae (1881), p. 177–411 (separate repaged, p. 1–237). Pt. III, Sec. 1. Discussion of the classification and relations of the brachiote crinoids, and conclusion of the generic descriptions (1885), p. 225–364 (separate repaged, 1–138). Pt. III, Sec. 2. Discussion of the classification and relations of the brachiote crinoids, and conclusion of the generic descriptions (1886), p. 64–226 (separate repaged to continue with section 1, 139–302).
- Webster GD, Webster DW. 2019.** Bibliography and index of Paleozoic crinoids, corinoids, and hemistreptocrinids. 1758–2012. Available at <http://crinoids.azurewebsites.net/> (accessed on September 2021).
- Wójcik K. 2015.** The uppermost Emsian and lower Eifelian in the Kielce Region of the Holy Cross Mts. Part I: Lithostratigraphy. *Acta Geologica Polonica* **65**:141–179  
DOI [10.1515/agp-2015-0006](https://doi.org/10.1515/agp-2015-0006).
- Wright DF. 2017.** Bayesian estimation of fossil phylogenies and the evolution of early to middle Paleozoic crinoids (Echinodermata). *Journal of Paleontology* **91**:799–814  
DOI [10.1017/jpa.2016.141](https://doi.org/10.1017/jpa.2016.141).
- Wright DF, Ausich WI, Cole SR, Peter ME, Rhenberg EC. 2017.** Phylogenetic taxonomy and classification of the Crinoidea (Echinodermata). *Journal of Paleontology* **91**:829–846  
DOI [10.1017/jpa.2016.142](https://doi.org/10.1017/jpa.2016.142).

- Yeltschewa RS, Dubalotova YA. 1961.** Morskie lilii V kn. Biostratigrafiia paleozoiia Sayano-Altaiiskoi gornoj oblasti, Srednii paleozoi, t. II [Crinoids. In Biostratigraphy of the Paleozoic of the Sayano Altai mountainous country, middle Paleozoic, v. II]. Moscow, Izdatel'stvo Nauka, Trudy, pt. 20:294–296, 552–560. (In Russian).
- Yeltschewa RS, Sisova EN. 1973.** Anthinocrinidae-Novoe Semeistvo Srednepaleozoiskikh Morskikh Lilii V kn. Voprosy Stratigrafii i Tektoniki Vostochnogo Kazakhstana [Anthinocrinidae New Family of Middle Paleozoic Marine Lily. In Aleksandrova MI, ed, Questions on the Stratigraphy and Tectonics of East Kazakhstan]. Trudy Vsesoyuznogo Ordena Lenina Nauchno-Issledovatel'skogo Geologicheskogo Instituta, n. s. 160:86–99. (In Russian).
- Zeiler F, Wirtgen PW. 1855.** Bemerkungen über die Petreefacten der älteren devonischen Gebirge am Rheine, insbesondere über die Echinodermen in der Umgegend von Coblenz und in dem Eifeler Kalke vorkommenden Arten. *Verhandlungen des Naturhistorisch Vereins der Preussischen Rheinlands und Westfalen* **12**:79–85.
- Zeuschner L. 1867.** Devonische Formation im Sandomirer-Chenciner Gebirge. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie* **67**:478–508.
- Zeuschner L. 1869.** Geognostische Beschreibung der mittleren devonischen Schichten zwischen Grzegorzowice und Skały-Zagaje, bei Nowa Słupia. *Zeitschrift der Deutschen Geologischen Gesellschaft* **21**:263–274.