



Draft Genome Sequence of *Bacillus cereus* ET31, Isolated from an Extreme Environment

Panagiota Stathopoulou,^a Sofia Nikolaki,^a Maria Lanara,^a George Tsiamis^a

^aDepartment of Environmental Engineering, University of Patras, Agrinio, Greece

ABSTRACT *Bacillus cereus* is a Gram-positive, widely distributed bacterium that has a high level of metabolite production. Here, we report the draft genome sequence of a *B. cereus* strain exhibiting high and diverse hydrolytic potential that was isolated from glacial water samples from Svalbard, Norway.

Bacterial communities associated with glaciers found all over the world are being increasingly studied for their potential biotechnological applications (1). The Svalbard archipelago (74° to 81° N, 10° to 35° E) comprises an area of 62,248 km² with more than 2,000 glaciers, and ice covers 60% of its surface (2). As a result, the draft genome sequence of *Bacillus cereus* ET31, which was isolated from glacier sediment from Svalbard Islands and has a high and diverse hydrolytic potential, is reported here.

Bacillus cereus is a Gram-positive, rod-shaped, aerobic or facultatively anaerobic, motile, beta-hemolytic, spore-forming bacterium that is widely distributed environmentally due to its ability to form thermoresistant endospores and biofilms and to grow over a broad temperature range (3, 4). Moreover, *B. cereus* has a high level of production of diverse metabolites, including bacteriocins, autolysins, and enzymes, particularly proteases (5–9).

This article reports the draft genome sequences of *B. cereus* strain ET31, isolated from glacial water in Svalbard, Norway (10). Bacteria were isolated on Antarctic bacterial medium (ABM) agar (0.5% [wt/vol] peptone, 0.2% [wt/vol] yeast extract, and 1.5% [wt/vol] agar, pH 7.0) at 12°C under aerobic conditions. *Bacillus cereus* ET31 was identified among 300 bacterial isolates obtained from glacier water from geologically and hydrologically diverse regions of southern Svalbard. Genomic DNA extraction from one culture, grown as described previously, was performed according to Haught et al. (11). The DNA was sent to MicrobesNG, a next-generation sequencing (NGS) service provider in the United Kingdom, for 16S rRNA and whole-genome sequencing. DNA was quantified in triplicates with the Quant-iT double-stranded DNA (dsDNA) high-sensitivity (HS) assay (Invitrogen) in an Eppendorf AF2200 plate reader. The genomic DNA library was prepared using the Nextera XT library prep kit (Illumina, San Diego, CA), following the manufacturer's protocol with the following modifications: 2 ng of DNA was used as input instead of 1 ng, and PCR elongation time was increased to 1 min from 30 s. DNA quantification and library preparation were carried out on a Hamilton Microlab Star automated liquid-handling system. The library was quantified using the Kapa Biosystems library quantification kit for Illumina on a Roche LightCycler 96 quantitative PCR (qPCR) machine. The library was sequenced with 52-fold coverage on the Illumina HiSeq platform using a 250-bp paired-end protocol, producing 714,745 reads. Reads were adapter trimmed using Trimmomatic v. 0.30 with a sliding window quality cutoff of Q15 (12). FastQC v. 0.11.8 with default parameters was used to check the quality of the validated reads (13). Reads were *de novo* assembled into 34 contigs using SPAdes v. 3.7 (14). The quality of the assembly was evaluated with Quast (15). Taxonomic assignment of the reads was calculated using Kraken (16), and Prokka v. 1.11

Citation Stathopoulou P, Nikolaki S, Lanara M, Tsiamis G. 2019. Draft genome sequence of *Bacillus cereus* ET31, isolated from an extreme environment. Microbiol Resour Announc 8:e01061-19. <https://doi.org/10.1128/MRA.01061-19>.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2019 Stathopoulou et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to George Tsiamis, gtsiamis@upatras.gr.

Received 14 September 2019

Accepted 16 October 2019

Published 7 November 2019

(17) was used for genome annotation. Protein function was predicted by performing sequence similarity queries against the UniProt Knowledgebase (UniProtKB) (18).

The assembled draft genome had a length of 5,875,600 bp, with a GC content of 34.9%. The largest contig was 2,021,752 bp long, the L_{50} was 2, and the N_{50} value equaled 1,600,101 bp. The majority of reads (94.07%) were classified into the genus *Bacillus*, and 59.55% were placed in the *B. cereus* group. The draft genome contained 5,859 protein coding sequences, with 2,535 of them identified as encoding hypothetical proteins, 99 tRNA sequences, 13 sequences identified as 5S rRNAs, and 1 as a 16S rRNA.

Data availability. The whole-genome shotgun project of *B. cereus* ET31 has been deposited at DDBJ/ENA/GenBank under accession number [VSSM00000000](https://www.ncbi.nlm.nih.gov/nuccore/VSSM00000000), BioProject number [PRJNA560934](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA560934), and BioSample number [SAMN12603168](https://www.ncbi.nlm.nih.gov/biosample/SAMN12603168). The raw sequencing reads were deposited to the Sequence Read Archive (SRA) under accession number [SRR9997721](https://www.ncbi.nlm.nih.gov/sra/SRR9997721). The version described in this paper is the first version, VSSM01000000.

ACKNOWLEDGMENT

This work was cofunded by Greece and the European Union in the framework of the action titled “Supporting researchers with an emphasis on young researchers” of the operational program “Human resources development, education and lifelong learning” (code CCI2014GR059OP001) of the National Strategic Reference Framework (NSRF) 2014–2020 under contract MIS5005675.

REFERENCES

- García-Lopez E, Rodríguez-Lorente I, Alcazar P, Cid C. 2019. Microbial communities in coastal glaciers and tidewater tongues of Svalbard Archipelago, Norway. *Front Mar Sci* 5:512. <https://doi.org/10.3389/fmars.2018.00512>.
- Hagen JO, Kohler J, Melvold K, Winther J-G. 2003. Glaciers in Svalbard: mass balance, runoff and freshwater flux. *Polar Res* 22:145–159. <https://doi.org/10.3402/polar.v22i2.6452>.
- Bottone EJ. 2010. *Bacillus cereus*, a volatile human pathogen. *Clin Microbiol Rev* 23:382–398. <https://doi.org/10.1128/CMR.00073-09>.
- Setlow B, Setlow P. 1994. Heat inactivation of *Bacillus subtilis* spores lacking small, acid-soluble spore proteins is accompanied by generation of abasic sites in spore DNA. *J Bacteriol* 176:2111–2113. <https://doi.org/10.1128/jb.176.7.2111-2113.1994>.
- Ouertani A, Chaabouni I, Mosbah A, Long J, Barakat M, Mansuelle P, Mghirbi O, Najjari A, Ouzari H-I, Masmoudi AS, Maresca M, Ortet P, Gigmes D, Mabrouk K, Cherif A. 2018. Two new secreted proteases generate a casein-derived antimicrobial peptide in *Bacillus cereus* food born isolate leading to bacterial competition in milk. *Front Microbiol* 9:1148. <https://doi.org/10.3389/fmicb.2018.01148>.
- Kotlar C, Ponce A, Roura S. 2015. Characterization of a novel protease from *Bacillus cereus* and evaluation of an eco-friendly hydrolysis of a brewery byproduct. *J Inst Brew* 121:558–565. <https://doi.org/10.1002/jib.257>.
- Chaabouni I, Barkallah I, Hamdi C, Jouini A, Saidi M, Mahillon J, Cherif A. 2015. Metabolic capacities and toxigenic potential as key drivers of *Bacillus cereus* ubiquity and adaptation. *Ann Microbiol* 65:975–983. <https://doi.org/10.1007/s13213-014-0941-9>.
- Raddadi N, Cherif A, Mora D, Brusetti L, Borin S, Boudabous A, Daffonchio D. 2005. The autolytic phenotype of the *Bacillus cereus* group. *J Appl Microbiol* 99:1070–1081. <https://doi.org/10.1111/j.1365-2672.2005.02713.x>.
- Bizani D, Dominguez APM, Brandelli A. 2005. Purification and partial chemical characterization of the antimicrobial peptide cerein 8A. *Lett Appl Microbiol* 41:269–273. <https://doi.org/10.1111/j.1472-765X.2005.01748.x>.
- Ntougias S, Polkowska Z, Nikolaki S, Dionyssopoulou E, Stathopoulou P, Doudoumis V, Ruman M, Kozak K, Namieśnik J, Tsiamis G. 2016. Bacterial community structures in freshwater polar environments of Svalbard. *Microbes Environ* 31:401–409. <https://doi.org/10.1264/jsme2.ME16074>.
- Haught C, Wilkinson DL, Zgafas K, Harrison RG. 1994. A method to insert a DNA fragment into a double-stranded plasmid. *Biotechniques* 16:46–48.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Wood DE, Salzberg SL. 2014. Kraken: ultrafast metagenomic sequence classification using exact alignments. *Genome Biol* 15:R46. <https://doi.org/10.1186/gb-2014-15-3-r46>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
2019. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 47:D506–D515. <https://doi.org/10.1093/nar/gky1049>.