

Letter to the Editor

In Response to *Clinical Features of Parosmia Associated with COVID-19 Infection*

**Key Words:** anosmia, coronavirus, COVID-19, olfactory dysfunction, olfactory function, parosmia, SARS-CoV-2, smell.  
*Laryngoscope*, 132:E28-E29, 2022

Dear Editor:

We have read with interest the recent article by Lerner et al.,<sup>1</sup> which analyzed the clinical features of parosmia following coronavirus disease 2019 (COVID-19). Congratulating the authors for the quality of their study, we would like to discuss some important points regarding pathogenetic hypotheses and the epidemiological implications emerging from their results. COVID-19 qualitative olfactory dysfunctions (ODs) represent a relatively recent topic that emerged along with persistent olfactory dysfunctions (POD). To date, research studies were mainly limited to the study of OD in the course of infection, which are mostly quantitative. The most recent theories emerging from histopathological studies on animal models<sup>2-4</sup> and humans<sup>5,6</sup> correlate these disorders to damage to the olfactory neuroepithelium (ONE) without involvement of the olfactory bulb.<sup>2</sup> In the same way, the recovery of the olfactory function may occur through the regeneration of the ONE, which takes place starting from the basal cells<sup>5</sup> and takes about 30 days.<sup>7</sup> The regeneration of the olfactory neurons is fundamental for the quantitative recovery of the olfactory function, but it is not sufficient for this perception to be qualitatively normal. In fact, part of the axonal regeneration starting from the new olfactory receptors can be disordered, leading to a misguided signal conduction to the higher olfactory centers and therefore to parosmia.<sup>8,9</sup>

Such evidences fit with the results detected by Lerner et al.<sup>1</sup> and may explain the fact that the onset of parosmia is not generally immediate and may be associated with a quantitatively normal olfactory function. Precisely, it is important to underline the importance of this latest finding. Persistent quantitative ODs are proving to be one of the most frequent symptoms of long-COVID-19.<sup>10,11</sup> The study by Lerner et al. stresses the possibility that all these studies published so far, although based on solid psychophysical tests, may underestimate the prevalence of residual OD through the lack of detection of the purely qualitative disorders that can occur in patients with normal

psychophysical scores. Unfortunately, a specific psychophysical test for parosmia has only recently been proposed<sup>12</sup> and it is not yet available in clinical practice. In the future, it will be appropriate to deepen the threshold-discriminative-identification assessment with the evaluation of qualitative perception so as to identify patients with parosmia, which is a fundamental determinant in the reduction of the quality of life of patients with POD, even more than purely quantitative disorders.<sup>1,13</sup>

LUIGI ANGELO VAIRA, MD 

Maxillofacial Surgery Operative Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy  
Biomedical Science Department, PhD School of Biomedical Science, University of Sassari, Sassari, Italy

GIACOMO DE RIU, MD, FEBOMFS

Maxillofacial Surgery Operative Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy

GIOVANNI SALZANO, MD 

ENT and Maxillofacial Surgery Unit, Istituto Tumori G. Pascale of Naples, Naples, Italy  
Maxillofacial Surgery Unit, University Hospital of Naples "Federico II", Naples, Italy

FABIO MAGLITTO, MD

Maxillofacial Surgery Unit, University Hospital of Naples "Federico II", Naples, Italy

PAOLO BOSCOLO-RIZZO, MD

Department of Medical, Surgical and Health Sciences, Section of Otolaryngology, University of Trieste, Trieste, Italy

JEROME R. LECHIEN, MD, PhD 

Department of Anatomy and Experimental Oncology, Mons School of Medicine, UMONS Research Institute for Health Sciences and Technology, University of Mons (UMons), Mons, Belgium  
Department of Otolaryngology-Head Neck Surgery, Elsan Hospital, Paris, France

The authors have no funding, financial relationships, or conflicts of interest to disclose.

REFERENCES

1. Lerner DK, Garvey KL, Arrighi-Allisan AE, et al. Clinical features of parosmia associated with COVID-19 infection. *Laryngoscope*. 2022;132:633-639.

Send correspondence to: Luigi Angelo Vaira, MD, Viale San Pietro 43/B, Sassari, Italy.  
E-mail: [lavaira@uniss.it](mailto:lavaira@uniss.it)

DOI: 10.1002/lary.30134

2. Khan M, Yoo SJ, Clijsters M, et al. Visualizing in deceased COVID-19 patients how SARS-CoV-2 attacks the respiratory and olfactory mucosae but spares the olfactory bulb. *Cell*. 2021;184:5932-5949.e15.
3. Vaira LA, Hopkins C, Sandison A, et al. Olfactory epithelium histopathological findings in long-term coronavirus disease 2019 related anosmia. *J Laryngol Otol*. 2020;134:1123-1127.
4. Kirschenbaum D, Imbach LL, Ulrich S, et al. Inflammatory olfactory neuropathy in two patients with COVID-19. *Lancet*. 2020;396:166.
5. Reyna RA, Kishimoto M, Urata S, Makishima T, Paessler S, Maruyama J. Recovery of anosmia in hamsters infected with SARS-CoV-2 is correlated with repair of the olfactory epithelium. *Sci Rep*. 2022;12:628.
6. Yu P, Deng W, Bao L, et al. Comparative pathology of the nasal epithelium in K18-hACE2 Tg mice, hACE2 Tg mice, and hamsters infected with SARS-CoV-2. *Vet Pathol*. 2022. <https://doi.org/10.1177/03009858211071016>.
7. Graziadei PPC, Monti Graziadei GA. Neurogenesis and neuron regeneration in the olfactory system of mammals. I. Morphological aspects of differentiation and structural organization of the olfactory sensory neurons. *J Neurocytol*. 1979;8:1-18.
8. Saniasiaya J, Narayanan P. Parosmia post COVID-19: an unpleasant manifestation of long COVID syndrome. *Postgrad Med J*. 2022;98:e96.
9. Parker JK, Kelly CE, Gane SB. Molecular mechanism of parosmia. medRxiv. 2021. doi:<https://doi.org/10.1101/2021.02.21251085>.
10. Boscolo-Rizzo P, Hummel T, Hopkins C, et al. High prevalence of long-term olfactory, gustatory, and chemesthesis dysfunction in post-COVID-19 patients: a matched case-control study with one-year follow-up using a comprehensive psychophysical evaluation. *Rhinology*. 2021;59:517-527.
11. Vaira LA, Salzano G, Le Bon S, et al. Prevalence of persistent olfactory disorders in patients with COVID-19: a psychophysical case-control study with 1-year follow-up. *Otolaryngol Head Neck Surg*. 2021;1945998211061511. <https://doi.org/10.1177/01945998211061511>.
12. Liu DT, Welge-Lussen A, Besser G, Mueller CA, Renner B. Assessment of odor hedonic perception: the Sniffin' sticks parosmia test (SSParoT). *Sci Rep*. 2020;22:18019.
13. Vaira LA, Gessa C, Deiana G, et al. The effects of persistent olfactory and gustatory dysfunctions on quality of life in long-COVID-19 patients. *Life*. 2022;12:141.