Contents lists available at ScienceDirect



Molecular Genetics and Metabolism Reports

journal homepage: www.elsevier.com/locate/ymgmr



Correspondence

Investigating the neurobehavioral symptoms of neuronopathic Hunter syndrome

ARTICLE INFO

Keywords: Mucopolysaccharidosis II Hunter syndrome Neuronopathic phenotype Neurobehavioral symptoms Pain Siblings

Managing the neurobehavioral symptoms of Hunter syndrome (mucopolysaccharidosis II) can be particularly challenging. Most individuals with the neuronopathic phenotype have significant language impairment and are unable to verbally communicate the reasons for their behavior [1]. Accordingly, it is uncertain whether the neurobehavioral symptoms are indicative of a psychiatric disorder associated with Hunter syndrome or occur in response to underlying somatic complications such as pain [2]. Eisengart et al. provide an important description of the neurobehavioral symptoms of neuronopathic Hunter syndrome; however, some additional questions and implications must be considered [3].

Eisengart et al. report that neurobehavioral symptoms include hitting and biting. While these behaviors may be intentionally aggressive, they may also occur in non-aggressive contexts such as when attempting to communicate with others [3]. However, the authors do not describe the factors that helped them distinguish between a behavior that was intentionally aggressive over one that was an attempt to engage socially. This information is needed and may help elucidate the sources that stimulate these challenging behaviors.

If challenging behaviors occur in attempts to communicate with others, what are affected individuals trying to communicate? The authors report that caregivers worried pain was a source for behavioral outbursts [3]. This implies that pain management should be considered when treating neurobehavioral symptoms. More studies are needed to assess the association of pain and neurobehavioral symptoms in Hunter syndrome.

Furthermore, siblings often provide care for their brothers and sisters with mucopolysaccharidosis [4]. Eisengart et al. describe the impact of neurobehavioral symptoms on parents but do not comment on the experiences of siblings. Future studies must assess the perspectives of siblings. While Eisengart et al. provide much-needed information, these recommendations may guide future studies to dig even deeper to understand the contributing pathophysiological process and overall impact of the neurobehavioral symptoms of Hunter syndrome.

Funding

No funding was received.

Declaration of Competing Interest

There are no conflicts of interest.

References

- J.B. Holt, M.D. Poe, M.L. Escolar, Natural progression of neurological disease in mucopolysaccharidosis type II, Pediatrics 127 (2011) e1258–e1265, https://doi.org/ 10.1542/peds.2010-1274.
- [2] J. Roberts, C. Stewart, S. Kearney, Management of the behavioural manifestations of Hunter syndrome, Br. J. Nurs. 25 (2016) 22, 24, 26–30, https://doi.org/10.12968/ bjon.2016.25.1.22.
- [3] J.B. Eisengart, K.E. King, E.G. Shapiro, C.B. Whitley, J. Muenzer, The nature and impact of neurobehavioral symptoms in neuronopathic Hunter syndrome, Mol. Genet. Metab. Rep. 22 (2019) 100549, https://doi.org/10.1016/j.ymgmr.2019. 100549.
- [4] N. Grant, Sibling and family caregivers, BMJ 362 (2018) k3158, https://doi.org/10. 1136/bmj.k3158.

Nathan Grant*

Department of Molecular and Cellular Biology, Department of Anthropology, Harvard University, Cambridge, MA, USA E-mail address: grantn@college.harvard.edu.

https://doi.org/10.1016/j.ymgmr.2020.100566

^{*} Corresponding author at: Department of Molecular and Cellular Biology, Department of Anthropology, Harvard University, 97 Dunster Mail Center, 945 Memorial Drive, Cambridge, MA 02138, USA.

Received 13 January 2020; Accepted 13 January 2020

^{2214-4269/ © 2020} The Author. Published by Elsevier Inc. This is an open access article under the CC BY license (http://creativecommons.org/licenses/BY/4.0/).