










BMJ Open Risk assessment models for venous thromboembolism in pregnancy and in the puerperium: a systematic review

Abdullah Pandor ¹, Jahnavi Daru ², Beverley J Hunt ³, Gill Rooney ¹,
Jean Hamilton ¹, Mark Clowes ¹, Steve Goodacre ¹,
Catherine Nelson-Piercy ³, Sarah Davis ¹

To cite: Pandor A, Daru J, Hunt BJ, *et al.* Risk assessment models for venous thromboembolism in pregnancy and in the puerperium: a systematic review. *BMJ Open* 2022;**12**:e065892. doi:10.1136/bmjopen-2022-065892

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-065892>).

Received 23 June 2022

Accepted 15 September 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

¹SCHARR, The University of Sheffield, Sheffield, UK

²Institute of Population Health Sciences, Queen Mary University of London, London, UK

³Guy's and St Thomas' NHS Foundation Trust, London, UK

Correspondence to

Abdullah Pandor;
a.pandor@sheffield.ac.uk

ABSTRACT

Objectives To assess the comparative accuracy of risk assessment models (RAMs) to identify women during pregnancy and the early postnatal period who are at increased risk of venous thromboembolism (VTE).

Design Systematic review following Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.

Data sources MEDLINE, Embase, Cochrane Library and two research registers were searched until February 2021.

Eligibility criteria All validation studies that examined the accuracy of a multivariable RAM (or scoring system) for predicting the risk of developing VTE in women who are pregnant or in the puerperium (within 6 weeks post-delivery).

Data extraction and synthesis Two authors independently selected and extracted data. Risk of bias was appraised using PROBAST (Prediction model Risk Of Bias ASsessment Tool). Data were synthesised without meta-analysis.

Results Seventeen studies, comprising 19 externally validated RAMs and 1 internally validated model, met the inclusion criteria. The most widely evaluated RAMs were the Royal College of Obstetricians and Gynaecologists guidelines (six studies), American College of Obstetricians and Gynecologists guidelines (two studies), Swedish Society of Obstetrics and Gynecology guidelines (two studies) and the Lyon score (two studies). In general, estimates of sensitivity and specificity were highly variable with sensitivity estimates ranging from 0% to 100% for RAMs that were applied to antepartum women to predict antepartum or postpartum VTE and 0% to 100% for RAMs applied postpartum to predict postpartum VTE. Specificity estimates were similarly diverse ranging from 28% to 98% and 5% to 100%, respectively.

Conclusions Available data suggest that external validation studies have weak designs and limited generalisability, so estimates of prognostic accuracy are very uncertain.

PROSPERO registration number CRD42020221094.

INTRODUCTION

Venous thromboembolism (VTE) remains an important cause of maternal morbidity and mortality in the developed world.¹ While uncommon, VTE complications can occur

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ A number of risk assessment models for venous thromboembolism (VTE) in pregnancy and puerperium have been developed using a variety of methods and based on a variety of predictor variables.
- ⇒ This systematic review provides a comprehensive review of risk assessment models for predicting the risk of developing VTE in women who are pregnant or in the puerperium (within 6 weeks post-delivery).
- ⇒ The newly developed PROBAST (Prediction model Risk Of Bias ASsessment Tool) was used to evaluate the risk of bias and applicability of the available evidence.
- ⇒ Heterogeneity in the included studies (participants, inclusion criteria, clinical condition, outcome definition and measurement) and variable reporting of items precluded meta-analysis.
- ⇒ Limitations of the existing evidence and areas of future research are highlighted.

at a rate of 1–2 per 1000 deliveries and can develop at any time during pregnancy.^{2–4} The risks substantially increase during the postpartum period (6 weeks post-delivery)⁵ and can be as high as 60-fold in some individuals compared with age-matched non-pregnant women.⁶ Preventative treatment with low-dose anticoagulation (thromboprophylaxis) has the potential to reduce the risk of symptomatic and asymptomatic VTE in pregnancy and the postpartum period.⁵ Consequently, various prominent international guidelines recommend targeted thromboprophylaxis for pregnant and puerperal women deemed to be at high risk of VTE.^{5 7–13} However, these expert-based consensus guidelines vary substantially with regards to the threshold of risk (based on certain risk factors) and the timing, dose and duration of pharmacological thromboprophylaxis.

Risk assessment models (RAMs) have been developed to help stratify the risk of VTE during pregnancy and the early

postnatal period. These models use clinical information from the patient's history and examination to identify those with an increased risk of developing VTE who are most likely to benefit from pharmacological thromboprophylaxis. Inappropriate use of VTE prophylaxis may not reduce VTE rates and may cause unnecessary harm especially through bleeding and bruising.¹⁴ While RAMs could improve the ratio of benefit to risk and benefit to cost, it is unclear which VTE RAM are best applied to guide decision-making for thromboprophylaxis in clinical practice and thereby optimise patient care.

The aim of this systematic review was to identify primary validation studies and determine the accuracy of individual RAMs that identify pregnant and postpartum women at increased risk of developing VTE who could be selected for thromboprophylaxis.

METHODS

A systematic review was undertaken in accordance with the general principles recommended in the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.¹⁵ This review was part of a larger project on Thromboprophylaxis in pregnancy and after delivery¹⁶ and was registered on the International Prospective Register of Systematic Reviews (PROSPERO) database.

Eligibility criteria

All studies evaluating the accuracy (eg, sensitivity, specificity, C-statistic) of a multivariable RAM (or scoring system) for predicting the risk of developing VTE were eligible for inclusion. We primarily sought and selected studies that included validation of the model in a group of patients that were not involved in the development of the prediction model. Although the included studies could have reported derivation of the model (for internal validation), we only used the external validation data to estimate accuracy, where appropriate. The study population of interest in our review consisted of pregnant and postpartum (within 6 weeks post-delivery) women who are at increased risk of developing a VTE and receiving care in both hospital, community and primary care settings. Studies that focused on non-pregnant women were excluded as these patient groups have VTE risk profiles that differ markedly from the obstetric population.

Data sources and searches

Potentially relevant studies were identified through searches of several electronic databases and research registers. This included MEDLINE (OvidSP from 1946), Embase (OvidSP from 1974), the Cochrane Library (<https://www.cochranelibrary.com> from inception), ClinicalTrials.gov (US National Institutes of Health from 2000) and the International Clinical Trials Registry Platform (WHO from 1990). All searches were conducted

from inception to February 2021. The search strategy used free text and thesaurus terms and combined synonyms relating to the condition (eg, VTE in pregnant and postpartum women) with risk prediction modelling terms.¹⁷ No language or date restrictions were used. Searches were supplemented by hand-searching the reference lists of all relevant studies (including existing systematic reviews); forward citation searching of included studies; contacting key experts in the field; and undertaking targeted searches of the World Wide Web using the Google search engine. Further details on the search strategy can be found in the online supplemental appendix S1.

Study selection

All titles were examined for inclusion by one reviewer (GR) and any citations that clearly did not meet the inclusion criteria (eg, non-human, unrelated to VTE in pregnancy and the puerperium) were excluded (for quality assurance a random subset of 20% was checked by a second reviewer (AP)). All abstracts and full-text articles were then examined independently by two reviewers (GR and AP). Any disagreements in the selection process were resolved through discussion or if necessary, arbitration by a third reviewer (JD) or the wider group (BJH, CN-P, SG) and included by consensus.

Data extraction and quality assessment

For eligible studies, data relating to study design, methodological quality and outcomes were extracted by one reviewer (GR) into a standardised data extraction form and independently checked for accuracy by a second reviewer (AP). Any discrepancies were resolved through discussion, or if this was unsuccessful, a third reviewer's opinion was sought (JD). Where multiple publications of the same study were identified, data were extracted and reported as a single study.

The methodological quality of each included study was assessed using PROBAST (Prediction model Risk Of Bias ASessment Tool).^{18 19} This instrument includes four key domains: participants (eg, study design and patient selection), predictors (eg, differences in definition and measurement of the predictors), outcome (eg, differences related to the definition and outcome assessment) and statistical analysis (eg, sample size, choice of analysis method and handling of missing data). Each domain is assessed in terms of risk of bias and the concern regarding applicability to the review (first three domains only). To guide the overall domain-level judgement about whether a study is at high, low or an unclear (in the event of insufficient data in the publication to answer the corresponding question) risk of bias, subdomains within each domain include several signalling questions to help judge with bias and applicability concerns. An overall risk of bias for each individual study was defined as low risk when all domains were judged as low; and high risk of bias when one or more domains were considered as high. Studies were assigned an unclear risk of bias if one or more domains were unclear, and all other domains were low.

Data synthesis and analysis

Due to significant levels of heterogeneity between studies (study design, participants, inclusion criteria) and variable reporting of items, a meta-analysis was not considered possible. As a result, a prespecified narrative synthesis approach^{20 21} was undertaken, with data being summarised in tables with accompanying narrative summaries that included a description of the included variables, statistical methods and performance measures (eg, sensitivity, specificity and C-statistic (a value between 0.7 to 0.8 and >0.8 indicated good and excellent discrimination, respectively; and values <0.7 were considered weak)),²² where applicable. All analyses were conducted using Microsoft Excel 2010 (Microsoft Corporation, Redmond, Washington, USA).

Patient and public involvement

Patients and the public were not involved in the design or conduct of this systematic review.

RESULTS

Study flow

Figure 1 summarises the process of identifying and selecting relevant literature. Of the 2268 citations identified, 16 studies^{23–38} investigating 19 unique externally validated RAMs met the inclusion criteria. Only one of these studies³⁵ presented data on model development and external validation (this study used UK Clinical Practice Research Data linked to Hospital Episode Statistics to develop a risk prediction model and externally validated using Swedish medical birth registry data). The remaining studies focused on external validation with no description of the initial derivation methodology.^{23–34 36–38} Due to the lack of model derivation studies with external validation, we also identified and included one internal validation study for completeness (ie, prediction model development without external validation).³⁹ This study used a bootstrap validation approach to capture optimism in model performance^{40 41} when applied to similar future patients. Most of the full-text articles (n=97)




Figure 1 Study flow chart (adapted). RAM, risk assessment model; VTE, venous thromboembolism.

were excluded primarily based on not using an RAM for predicting the risk of developing VTE during pregnancy or the puerperium, having no useable or relevant outcome data or an inappropriate study design (eg, reviews, commentaries or study protocols). A full list of excluded studies with reasons for exclusion is provided in online supplemental appendix S2.

Study and patient characteristics

The design and participant characteristics of the 17 included studies are summarised in [table 1](#). All studies were published between 2000 and 2020 and were undertaken in North America (n=4),^{24 37–39} Southeast Asia (n=1),³³ Europe (n=10),^{23 25–30 32 34 36} South America (n=1)³¹ and one study was multicountry.³⁵ Sample sizes ranged from 52³¹ to 662 387³⁵ patients in 14 observational cohort studies (6 prospective^{25 27 28 31 33 36} (all single centre) and 8 retrospective^{24 26 29 30 34 35 37 39} [2 of which were multicentre] in design). Sample sizes in two, single centre case–control studies^{32 38} ranged from 76³⁸ to 2421³² patients and one study used a non-randomised multicentre study design.²³ The mean age ranged from 27.8years³⁹ to 34years^{25 29} (not reported in 7 studies).^{24 27 32 34 36–38}

The majority of studies were conducted across antenatal and postnatal periods,^{23 27–29 31 34 36 38} or postpartum period only^{24–26 30 32 33 35 37 39} and generally included women at increased risk of VTE.^{23–25 28 29 31–33 38 39} One study excluded women with a history of VTE³⁵ and six studies^{26 27 30 34 36 37} included all pregnant women who delivered. Thromboprophylaxis was employed in about half (n=9)^{23 25 28–31 33 35 36} of the studies, with the proportion receiving thromboprophylaxis ranging from 3%³⁵ to 100%.^{23 28} The remaining studies did not report data on thromboprophylaxis use.

VTE definition and case ascertainment

Only a few studies^{23 27 32 36} defined the VTE endpoint (deep vein thrombosis and/or pulmonary embolism) as being confirmed by objective testing. Of the remainder, 3 studies^{35 37 39} had no objective confirmation of VTE and 10 studies^{24–26 28–31 33 34 38} did not report the methods for diagnosis confirmation. Although 9 studies^{23 24 27 29 32–34 36 39} did not report the VTE risk period, the majority of the remaining studies used the RAMs to predict the occurrence of VTE up to 3 months after delivery.^{25 28 30 31} Despite differences in study design, study participants, definitions, different criteria for the use of thromboprophylaxis and differences between doses of low molecular weight heparin (LMWH), the reported overall incidence of VTE in pregnancy and the puerperium was <1.3%.

RAMs

The studies included in this review evaluated 19 externally validated RAMs^{23–38} and 1 internally validated risk model.³⁹ While most RAMs focused solely on the estimate of thromboembolic risk, RAMs varied in design,

structure, threshold, dosage and duration for pharmacological prophylaxis. In addition, the individual predictors and their weighting varied markedly between RAMs. The most commonly used tools were the Royal College of Obstetricians and Gynaecologists guidelines (six studies),^{24 30 33–35 37} American College of Obstetricians and Gynecologists (ACOG) guidelines (two studies),^{30 33} Swedish Society of Obstetrics and Gynecology guidelines (two studies)^{32 35} and the Lyon score (two studies).^{28 29} A simplified summary of their associated characteristics and composite clinical variables is provided in online supplemental appendix S3.

Risk of bias and applicability assessment

The overall methodological quality of the 17 included studies is summarised in [table 2](#) and [figure 2](#). The methodological quality of the included studies was variable, with most studies having high or unclear risk of bias in at least one item of the PROBAST. The main risk of bias limitations was related to patient selection factors (arising from retrospective data collection,^{24 26 29 30 32 34 37–39} unclear exclusions/incomplete patient enrolment^{24 26 27 31–34 36 38 39} or unclear criteria for patients receiving VTE prophylaxis)^{23 30 35}; predictor and outcome bias (due to a general lack of details on the definition^{24–26 28–31 33 34 38} and methods of outcome determination^{24 26 28–31 33 34 37–39} and whether all predictors were available at the models intended time of use^{23 24 29 31 32 34 36–39} or influenced by the outcome measurement)^{23–28 30–39} and analysis factors (low event rates,^{23–31 33–37 39} unclear handling of missing data^{23–29 31–34 36–39} and failure in reporting relevant performance measures such as calibration and discrimination).^{23–34 36–38}

Assessment of applicability to the review question led to the majority of studies being classed either as unclear (n=13)^{23 26–30 32 34–39} or high (n=4)^{24 25 31 33} risk of inapplicability. These assessments were generally related to patient selection (highly selected study populations, for example, selected women at increased risk of VTE, caesarean delivery only, single disease pathologies, single site settings), predictors (inconsistency in definition, assessment or timing of predictors) and outcome determination.

Predictive performance of VTE RAMs (summary of results)

[Table 3](#) and [table 4](#) shows the sensitivity and specificity of RAMs that were applied to antepartum women to predict antepartum or postpartum VTE or applied postpartum to predict postpartum VTE, respectively, with the results grouped by RAM. However, any meaningful comparisons between these alone is difficult, without considering the models' corresponding discrimination and calibration metrics, which were not universally reported. Only one external validation study considered model discrimination and calibration. In this study by Sultan *et al*,³⁵ their recalibrated novel risk prediction model (also known as the Maternity Clot Risk) provided good discrimination and was able to discriminate postpartum women with

Table 1 Study and population characteristics

Author, year	Country	Design	Single/ multicentre	Sample size	Population	Period	Mean age (years)	VTE prophylaxis	RAMs evaluated	Target condition, definition (risk period)	Incidence
Antepartum and postpartum following vaginal and caesarean delivery											
Bauersachs <i>et al</i> , 2007 ²³	Germany	P, NRS	Multi	810	Women at increased risk of VTE (due to thromboembolic status and prior VTE)	March 1999 to December 2002	30.8	100%	► ETHiG	Antepartum and postpartum VTE, symptomatic (NR)	0.62% (antepartum); 0.25%; postpartum: 0.37%)
Chaleur <i>et al</i> , 2008 ²⁷	France	P, CS	Single	2685	All women who delivered	July 2002 to June 2003	NR (median, 29)	NR	► STRATHEGE	Antepartum and postpartum VTE (NR)	0.34% (antepartum); 0.19%; postpartum: 0.15%)
Dargaud <i>et al</i> , 2017 ²⁶	France	P, CS	Single	445	Women at increased risk of VTE (due to thromboembolic status and prior VTE)	January 2005 to January 2015	33	100%	► Lyon	Antepartum and postpartum VTE, not defined (pregnancy and 3 months postpartum)	1.35%
Dargaud <i>et al</i> , 2005 ²⁹	France	R, CS	Single	116	Women at increased risk of VTE (due to thromboembolic status and prior VTE)	2001 to 2003	34	53%	► Lyon	Antepartum and postpartum VTE, not defined (NR)	0.86% (antepartum only)
Hase <i>et al</i> , 2018 ³¹	Brazil	P, CS	Single	52	Hospitalised pregnant women with cancer	1 December 2014 to 31 July 2016	31	57.7%	► RCOG (modified)	Antepartum and postpartum VTE, not defined (pregnancy and 3 months postpartum)	Unable to estimate—no VTE
Shcaluga and Rayment, 2019 (correspondence) ³⁴	Wales	R, CS	Single	42 000	All managed pregnancies	2009 to 2015	NR	NR	► All Wales RCOG	Antepartum and postpartum VTE, not defined (NR)	0.08% (antepartum); 0.04%; postpartum: 0.04%)
Testa <i>et al</i> , 2015 ³⁶	Italy	P, CS	Single	1719	All pregnant women enrolled in Pregnancy Healthcare Program	January 2008 to December 2010	NR (median 33)	4.6%	► Novel (Testa)	Antepartum and postpartum VTE (NR)	Unable to estimate—no VTE
Weiss and Bernstein, 2000 ³⁸	USA	CC	Single	19 cases: 57 control*	Women with (confirmed cases) and without (unmatched control) VTE	1987 to 1998	NR	NR	► Novel (Weiss)	Antepartum and postpartum VTE, not defined (pregnancy and 6 weeks postpartum)	–
Postpartum only following vaginal and caesarean delivery											
Chau <i>et al</i> , 2019 ²⁸	France	R, CS	Single	1069 (time period 2012: 557; 2015: 512)	All women who delivered	February to April 2012 and February to April 2015	2012: 29 2015: 29	NR	► Novel (Chau)	Postpartum VTE, not defined (8 weeks)	2012: 0.18% 2015: 0.20%
Ellis-Kahana <i>et al</i> , 2020 ³⁹ †	USA	R, CS	Multi	83 500	All obese women (BMI >30 kg/m ²) who delivered	2002 to 2008	27.8	NR	► Novel (Ellis-Kahana)	Postpartum VTE (NR)	0.13%
Gassmann <i>et al</i> , 2021 ⁴⁰	Switzerland	R, CS‡	Single	344	All women who delivered	1–31 January 2019	32.2	24%	► RCOG ► ACOG ► ACCP ► ASH	Postpartum VTE, not defined (3 months)	Unable to estimate—no VTE

Continued

Table 1 Continued

Author, year	Country	Design	Single/ multicentre	Sample size	Population	Period	Mean age (years)	VTE prophylaxis	RAMs evaluated	Target condition, definition (risk period)	Incidence
Lindqvist <i>et al</i> , 2008 ³²	Sweden	CC	Single	37 cases; 2384 control	All women with (confirmed cases) and without (unselected population-based control) VTE	1990 to 2005	NR	NR	▲ SFOG (Swedish guidelines)	Postpartum VTE (NR)	–
Sultan <i>et al</i> , 2016 ³⁵	England (derivation) [§] and, Sweden (validation)	R, CS	Multi	662387 (validation cohort) [§]	All women (with no history of VTE) who delivered	1 July 2005 to 31 December 2011	30.32	3%	▲ Novel (Sultan) ▲ RCOG [§] ▲ SFOG (Swedish Guidelines)	Postpartum VTE (6 weeks)	0.08% (validation cohort)
Tran <i>et al</i> , 2019 ³⁷	USA	R, CS	Single	6094	All women who delivered after 14 weeks	01 January 2015 to 31 December 2016	NR	NR	▲ RCOG ▲ Padua ▲ Caprini	Postpartum VTE (6 months)	0.05%
Postpartum following caesarean delivery											
Binstock and Larkin, 2019 (abstract) ³⁴	USA	R, CS	Single	2875	Postpartum women following CD	2011	NR	NR	▲ Novel (Binstock) ▲ RCOG	Postpartum VTE, not defined (NR)	0.38%
Cavazza <i>et al</i> , 2012 ²⁵	Italy	P, CS	Single	501	Postpartum women following CD	2007 to 2009	34	53.5%	▲ Novel (Cavazza)	Postpartum VTE, symptomatic, not defined (90 days)	0.20%
Lok <i>et al</i> , 2019 ³³	Hong Kong	P, CS	Single	859	Postpartum women following CD	May 2017 to April 2018	32.9	3.3%	▲ Novel (Lok) ▲ RCOG ▲ ACOG	Postpartum VTE, symptomatic, not defined (NR)	Unable to estimate—no VTE

*Retrospective case-control study of pregnant and postpartum women, but data reported for antepartum period only due to low number of postpartum VTE events (n=2).

†Internal validation study (ie, prediction model development without external validation).

‡Prospective cohort study with retrospective analysis, thus classified as retrospective cohort study.

§RCOG was applied to an English derivation cohort, n=433 353; incidence, 0.07% (312 events).

ACCP, American College of Chest Physicians; ACOG, American College of Obstetricians and Gynecologists; ASH, American Society of Hematology; BMI, body mass index; CC, case-control; CD, caesarean delivery; CS, cohort study; EThIG, Efficacy of Thromboprophylaxis as an Intervention during Gravidity Investigators; NR, not reported; NRS, non-randomised study; P, prospective; R, retrospective; RAM, risk assessment model; RCOG, Royal College of Obstetricians and Gynaecologists; SFOG, Swedish Society of Obstetrics and Gynecology; VTE, venous thromboembolism.

Table 2 Summary of each study's risk of bias and applicability concern using the PROBAST (Prediction model Risk Of Bias Assessment Tool) — review authors' judgements

Author, year	Risk of bias					Applicability					Overall	
	Participant selection	Predictors	Outcome	Analysis	Participant selection	Predictors	Outcome	Risk of bias	Applicability	Risk of bias	Applicability	
Bauersachs <i>et al</i> , 2007 ²³	?	?	+	-	?	?	+	-	?	-	?	
Binstock and Larkin, 2019 ²⁴	?	?	?	-	-	?	?	-	-	-	-	
Cavazza <i>et al</i> , 2012 ²⁵	-	?	?	-	-	+	?	-	-	-	-	
Chau <i>et al</i> , 2019 ²⁶	?	?	?	-	?	?	?	-	?	-	?	
Chauleur <i>et al</i> , 2008 ²⁷	?	?	?	-	?	?	?	-	?	-	?	
Dargaud <i>et al</i> , 2017 ²⁸	?	?	?	-	?	?	?	-	?	-	?	
Dargaud <i>et al</i> , 2005 ²⁹	-	?	?	-	?	+	?	-	?	-	?	
Ellis-Kahana <i>et al</i> , 2020 ³⁹	-	?	?	-	?	?	?	-	?	-	?	
Gassmann <i>et al</i> , 2021 ³⁰	?	?	?	-	?	?	?	-	?	-	?	
Hase <i>et al</i> , 2018 ³¹	?	?	?	-	-	?	?	-	-	-	-	
Lindqvist <i>et al</i> , 2008 ³²	-	?	?	-	-	?	?	-	-	-	?	
Lok <i>et al</i> , 2019 ³³	?	?	-	-	-	+	?	-	-	-	-	
Shacaluga and Rayment, 2019 ³⁴	-	?	?	-	?	?	?	-	?	-	?	
Sultan <i>et al</i> , 2016 ³⁵	-	?	+	+	-	?	+	-	-	-	?	
Testa <i>et al</i> , 2015 ³⁶	?	?	?	-	?	?	?	-	?	-	?	
Tran <i>et al</i> , 2019 ³⁷	-	?	?	-	?	?	?	-	?	-	?	
Weiss and Bernstein, 2000 ³⁸	-	?	?	-	?	?	?	-	?	-	?	

+ indicates low risk of bias/low concern regarding applicability; -, indicates high risk of bias/high concern regarding applicability; and ? indicates unclear risk of bias/unclear concern regarding applicability




Figure 2 PROBABST (Prediction model Risk Of Bias ASsessment Tool) assessment summary graph—review authors' judgements.

and without VTE in the external Swedish cohort with a C-statistic of 0.73 (95% CI: 0.71 to 0.75), and calibration, of observed and predicted VTE risk, close to ideal (calibration slope of 1.11 (95% CI: 1.01 to 1.20)). In the remaining studies, interpretation was further limited by marked heterogeneity, which was exacerbated when different thresholds were reported by different studies evaluating the same model. In general, model accuracy was generally poor, with high sensitivity usually reflecting a threshold effect, as indicated by corresponding low specificity values (and vice versa).

DISCUSSION

Summary of results

This systematic review identified 19 externally validated RAMs (and 1 internally validated risk model) that aimed to predict the risk of VTE in pregnant and postpartum women and who could be selected for thromboprophylaxis. Although various risk models (based on a variety of predictor variables) are being used, most of these lacked rigorous development and evaluation. The predictive accuracy of the RAMs was highly variable, and the substantial risk of bias concerns and the general lack of methodological clarity and unclear applicability make meaningful comparisons of the evidence difficult.

Interpretation of results

Despite the development and use of various RAMs to predict the risk of developing VTE in women who are

pregnant or in the puerperium (within 6 weeks post-delivery), VTE remains the leading cause of direct maternity mortality in the UK (MBRRACE-UK report 2021). Several explanations for this are possible: the risk assessment tools are inadequate; the application of these tools is incomplete or inaccurate; the underlying VTE risks of the pregnant population (increasing age, body mass index and comorbidities) are changing from when the RAMs were developed; or all three problems are operating.

The use of thromboprophylaxis was reported in nine studies^{23 25 28–31 33 35 36} (ranging from 3%³⁵ to 100%^{23 28}). This may lead to underestimation of predictive accuracy if a given RAM was to predict VTE events that were subsequently prevented by thromboprophylaxis. In the remaining studies (n=8) where thromboprophylaxis use was not reported (n=8), further analysis of its impact on the performance of the RAMs was not possible. This also suggests that the degree to which thromboprophylaxis reduces the risk of VTE in those who received it cannot be accurately estimated. Moreover, the lack of data on the predictive performance of weight-based LMWH dosing, dosage change throughout pregnancy and D-dimer testing in the included studies also precluded further analysis of its association with VTE.

Comparison to the existing literature

To our knowledge, there are no previous systematic reviews on this topic. However, recently several large registries have been interrogated in an attempt to derive robust prediction rules for this population, although with some methodological concerns. Sultan *et al.*³⁵ developed (using a large English-based registry database covering 6% of the population) and validated (using a Swedish national database registry) a risk prediction tool to estimate the absolute risk of VTE in postpartum women according to their individual risk factor combinations. Despite the low incidence of VTE in both cohorts (<0.08%), their model showed good discrimination in the external cohort and poor sensitivity at predicting those at risk of experiencing VTE. In addition, their model lacked some important VTE risk factors (eg, thrombophilia, antepartum immobilisation), and possibly underestimated the risks due to diagnosis limited to diagnostic coding (eg, varicose veins, severity of comorbidities) and the use of thromboprophylaxis in both cohorts.⁴² Ellis-Kahana *et al.*³⁹ also derived (using a large national database from the USA) a risk prediction model for VTE in obese pregnant women and indicated strong discrimination. However, this model still requires external validation.

Strengths and limitations

This systematic review has several strengths. It is the first systematic review to evaluate RAMs for predicting the risk of developing VTE in women during pregnant and the puerperium periods, and was conducted with robust methodology in accordance with the PRISMA statement¹⁵ and the protocol was registered with the PROSPERO register. Clinical experts, in addition to the core review

Table 3 Performance of RAMs applied antepartum to predict VTE

Risk assessment models	Threshold or cut-off	Endpoint	Data source	Performance measures						
				TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)	
Predicting either antepartum or postpartum VTE										
All Wales (one study)	NR	VTE	Shacaluga and Rayment ³⁴	25	NR	9	NR	NR	0.74 (0.57 to 0.85)	NR
ETHIG (one study)	High/very high risk	VTE	Bauersachs et al ²³	5	580	0	225	225	1.00 (0.57 to 1.00)	0.28 (0.25 to 0.31)
Lyon (two studies)	Risk score ≥ 3	VTE	Dargaud et al ²⁸	5	282	1	157	157	0.83 (0.44 to 0.97)	0.36 (0.31 to 0.4)
Lyon	Risk score ≥ 3	VTE	Dargaud et al ²⁹	1	56	0	59	59	1.00 (0.21 to 1.00)	0.51 (0.42 to 0.6)
RCOG (modified) (one study)	Risk score ≥ 3	VTE	Hase et al ³¹	0	34	0	18	18	unable to estimate – no VTE	0.35 (0.23 to 0.48)
STRATHEGE (one study)	Risk score ≥ 3	VTE	Chauleur et al ²⁷	0	54	9	2622	2622	0.00 (0.00 to 0.3)	0.98 (0.97 to 0.99)
Testa 2015 (one study)	Risk score ≥ 2.5	VTE	Testa et al ³⁶	0	85	0	1634	1634	unable to estimate – no VTE	0.95 (0.94 to 0.96)
Predicting antepartum VTE										
ETHIG (one study)	High/very high risk	VTE	Bauersachs et al ²³	2	583	0	225	225	1.00 (0.34 to 1.00)	0.28 (0.25 to 0.31)
Lyon (one study)	Risk score ≥ 3	VTE	Dargaud et al ²⁸	1	286	1	157	157	0.50 (0.09 to 0.91)	0.35 (0.31 to 0.4)
STRATHEGE (one study)	Risk score ≥ 1	VTE	Chauleur et al ²⁷	0	54	4	2627	2627	0.00 (0.00 to 0.49)	0.98 (0.97 to 0.99)
Weiss 2000 (one study)	Risk score ≥ 2	VTE	Weiss and Bernstein ³⁸	4	3	15	54	54	0.21 (0.09 to 0.43)	0.95 (0.86 to 0.98)
Predicting postpartum VTE										
ETHIG (one study)	High/very high risk	VTE	Bauersachs et al ²³	3	582	0	225	225	1.00 (0.44 to 1.00)	0.28 (0.25 to 0.31)
Lyon (one study)	Risk score ≥ 3	VTE	Dargaud et al ²⁸	4	283	0	158	158	1.00 (0.51 to 1.00)	0.36 (0.31 to 0.4)
STRATHEGE (one study)	Risk score ≥ 1	VTE	Chauleur et al ²⁷	0	54	5	2626	2626	0.00 (0.00 to 0.43)	0.98 (0.97 to 0.98)

ETHIG, Efficacy of Thromboprophylaxis as an Intervention during Gravidity Investigators; FN, false negative; FP, false positive; NR, not reported; RAMs, risk assessment models; RCOG, Royal College of Obstetricians and Gynaecologists; TN, true negative; TP, true positive; VTE, venous thromboembolism.



Table 4 Performance of RAMs applied postpartum to predict VTE

Risk assessment models	Threshold or cut-off	Endpoint	Data source	Performance measures						Specificity (95% CI)
				TP	FP	FN	TN	Sensitivity (95% CI)		
Predicting postpartum VTE following vaginal and caesarean delivery										
ACCP (one study)	NR	VTE	Gassmann <i>et al</i> ³⁰	0	34	0	310	unable to estimate – no VTE	0.90 (0.86 to 0.93)	
ACOG (one study)	NR	VTE	Gassmann <i>et al</i> ³⁰	0	30	0	314	unable to estimate – no VTE	0.91 (0.88 to 0.94)	
ASH (one study)	NR	VTE	Gassmann <i>et al</i> ³⁰	0	0	0	344	unable to estimate – no VTE	1.00 (0.99 to 1.00)	
Caprini (one study)	Risk score ≥2	VTE	Tran <i>et al</i> ³⁷	3	5780	0	311	1.00 (0.44 to 1.00)	0.05 (0.05 to 0.06)	
Caprini	Risk score ≥3	VTE	Tran <i>et al</i> ³⁷	1	3066	2	3025	0.33 (0.06 to 0.79)	0.50 (0.48 to 0.51)	
Caprini	Risk score ≥4	VTE	Tran <i>et al</i> ³⁷	0	1257	3	4834	0.00 (0.00 to 0.56)	0.79 (0.78 to 0.80)	
Padua (one study)	Risk score ≥4	VTE	Tran <i>et al</i> ³⁷	0	50	3	6041	0.00 (0.00 to 0.56)	0.99 (0.99 to 0.99)	
RCOG (three studies)	NR	VTE	Gassmann <i>et al</i> ³⁰	0	138	0	206	unable to estimate – no VTE	0.60 (0.55 to 0.65)	
RCOG	Risk score ≥2	VTE	Tran <i>et al</i> ³⁷	1	3837	2	2254	0.33 (0.06 to 0.79)	0.37 (0.36 to 0.38)	
RCOG	≥2 low risk factors or 1 high risk factor	VTE	Sultan <i>et al</i> ³⁵	197	149205	115	283836	0.63 (0.58 to 0.68)	0.66 (0.65 to 0.66)	
SFOG (two studies)	Risk score ≥2	VTE	Lindqvist <i>et al</i> ³²	18	111	19	2273	0.49 (0.33 to 0.64)	0.95 (0.94 to 0.96)	
SFOG	≥2 risk factors	VTE	Sultan <i>et al</i> ³⁵	109	41145	412	620721	0.21 (0.18 to 0.25)	0.94 (0.94 to 0.94)	
Chau, 2019 (one study*)	Risk score ≥3 (2012 data set)	VTE	Chau <i>et al</i> ²⁶	0	101	1	456	0.00 (0.00 to 0.79)	0.82 (0.78 to 0.85)	
Chau, 2019	Risk score ≥3 (2015 data set)	VTE	Chau <i>et al</i> ²⁶	0	113	1	393	0.00 (0.00 to 0.79)	0.78 (0.74 to 0.81)	
Ellis-Kahana, 2020 (full model) (one study†)	Risk score >3 (high risk)	VTE	Ellis-Kahana <i>et al</i> ²⁹	68	7942	41	75449	0.62 (0.53 to 0.71)	0.90 (0.90 to 0.91)	
Ellis-Kahana, 2020 (without antepartum thromboembolic disorder)	Risk score >3 (high risk)	VTE	Ellis-Kahana <i>et al</i> ²⁹	63	9926	46	73465	0.58 (0.48 to 0.67)	0.88 (0.88 to 0.88)	
Sultan, 2016 (one study‡)	≥2 risk factors: top 35% (threshold: 7.2 per 10 000 deliveries)	VTE	Sultan <i>et al</i> ³⁵	355	231480	166	430386	0.68 (0.64 to 0.72)	0.65 (0.65 to 0.65)	
Sultan, 2016	≥2 risk factors: top 25% (threshold: 8.7 per 10 000 deliveries)	VTE	Sultan <i>et al</i> ³⁵	310	164976	211	496890	0.60 (0.55 to 0.64)	0.75 (0.75 to 0.75)	
Sultan, 2016	≥2 risk factors: top 20% (threshold: 9.8 per 10 000 deliveries)	VTE	Sultan <i>et al</i> ³⁵	278	131921	243	529945	0.53 (0.49 to 0.58)	0.80 (0.80 to 0.80)	
Sultan, 2016	≥2 risk factors: top 10% (threshold: 14 per 10 000 deliveries)	VTE	Sultan <i>et al</i> ³⁵	185	66053	336	595813	0.36 (0.32 to 0.40)	0.90 (0.90 to 0.90)	
Sultan, 2016	≥2 risk factors: top 6% (threshold: 18 per 10 000 deliveries)	VTE	Sultan <i>et al</i> ³⁵	158	41096	363	620770	0.30 (0.27 to 0.34)	0.94 (0.94 to 0.94)	
Sultan, 2016	≥2 risk factors: top 5% (threshold: 19.7 per 10 000 deliveries)	VTE	Sultan <i>et al</i> ³⁵	139	32980	382	628886	0.27 (0.23 to 0.31)	0.95 (0.95 to 0.95)	
Sultan, 2016	≥2 risk factors: top 1% (threshold: 41.2 per 10 000 deliveries)	VTE	Sultan <i>et al</i> ³⁵	47	6576	474	655290	0.09 (0.07 to 0.12)	0.99 (0.99 to 0.99)	
Predicting postpartum VTE following caesarean delivery only										
ACOG (one study)	Risk score ≥3	VTE	Lok <i>et al</i> ³³	0	0	0	859	unable to estimate – no VTE	1.00 (1.00 to 1.00)	
RCOG (two studies)	NR	VTE	Binstock and Larkin (abstract) ²⁴	11	2692	0	172	1.00 (0.74 to 1.00)	0.06 (0.05 to 0.07)	
RCOG	Risk score ≥3	VTE	Lok <i>et al</i> ³³	0	649	0	210	unable to estimate – no VTE	0.24 (0.22 to 0.27)	

Continued

Table 4 Continued

Risk assessment models	Threshold or cut-off	Endpoint	Data source	Performance measures					
				TP	FP	FN	TN	Sensitivity (95% CI)	Specificity (95% CI)
Binstock, 2019 (one study)	NR	VTE	Binstock and Larkin (abstract) ⁴⁴	11	2635	0	229	1.00 (0.74 to 1.00)	0.08 (0.07 to 0.09)
Cavazza, 2012 (one study)	Moderate/high/very high	VTE	Cavazza <i>et al</i> ²⁵	0	268	1	232	0.00 (0.00 to 0.79)	0.46 (0.42 to 0.51)
Lok, 2019 (one study)	Risk score ≥ 3	VTE	Lok <i>et al</i> ⁶³	0	28	0	831	unable to estimate – no VTE	0.97 (0.95 to 0.98)

*Data discrepancy in paper—text states analysis included 1069 women: 557 in the 2012 time frame and 512 in the 2015 time frame; however, data in tables suggest 558 women included in the 2012 time frame and 507 in the 2015 time frame.
 †Internal validation study, Full risk prediction model: C-statistic, 0.817 (95% CI: 0.768 to 0.865) with Hosmer-Lemeshow p value=0.297; model without antepartum thromboembolic disorder: C-statistic, 0.778 (95% CI: 0.729 to 0.826) with Hosmer-Lemeshow p value=0.114.
 ‡Sultan *et al*,³⁵ final risk prediction model in external Swedish cohort: C-statistic, 0.73 (95% CI: 0.71 to 0.75) and calibration slope, 1.11 (95% CI: 1.01 to 1.20).
 ACCP, American College of Chest Physicians; ACOG, American College of Obstetricians and Gynecologists; ASH, American Society of Hematology; FN, false negative; FP, false positive; NR, not reported; RAMs, risk assessment models; RCOG, Royal College of Obstetricians and Gynaecologists; SFOG, Swedish Society of Obstetrics and Gynecology; TN, true negative; TP, true positive; VTE, venous thromboembolism.

team, were involved and consulted throughout as advisors and to assess the validity and applicability of research findings during the review processes.

The main limitations of this study related to the observational nature of the studies reviewed and their own limitations. Most of the included risk prediction studies were retrospective cohorts. Retrospective cohort studies of large health database registries are limited by poor data quality and failure to accurately ascertain outcomes and case-control designs are prone to bias including uncontrolled confounding, temporal and selection bias.⁴³ Conversely, better quality data may be obtained with prospective cohorts, but smaller sample sizes will lack statistical power. In addition, most of the external validation studies evaluated predictive performance of risk models that were not statistically derived (ie, without model development and internal validation). This process is vital, as risk models with only external validation may be subject to overfitting and optimism.⁴⁰ Similarly, the absence of model performance measures such as calibration or discrimination hinders the full appraisal of models.⁴¹

Due to the high levels of heterogeneity between studies, we were unable to undertake any meta-analysis or statistical examination of the causes of heterogeneity due to the small number of external validation studies per risk model. Potential sources of heterogeneity include variation in study design, the study population, risk model implementation, outcome definition and measurement and the use of thromboprophylaxis. As a result, we reported descriptive statistics to provide a better understanding of the evidence base applicable to the subject matter, and shortcomings regarding reliability and validity of the data. Finally, assessments on study relevance, information gathering and validity of articles were unblinded and could potentially have been influenced by preformed opinions. However, masking is resource intensive with uncertain benefits in protecting against bias decisions.⁴⁴

Implications for policy, practice and future research

VTE risk assessment is challenging for numerous reasons. Many risk factors for VTE are pre-existing and non-modifiable (such as parity and inherited thrombophilia). These are then often combined with evolving risk factors which can change over the course of a pregnancy or post-natal period. Despite wide scale awareness of VTE being a major contributor to maternal mortality, numerous challenges with VTE risk stratification have been highlighted. In the UK, the MBRRACE-UK report (Saving Lives, Improving Mothers' Care 2018)⁴⁵ shows that doctors and midwives find existing risk scoring systems difficult to apply consistently in clinical practice. There is a need for development of an RAM that is simpler and more reproducible. National Institute for Health and Care Excellence guidelines on the use of thromboprophylaxis (NG89)⁴⁶ concluded that the tool described by Sultan *et al*³⁵ showed poor sensitivity compared with their prespecified target of 90% sensitivity. However, this high level of sensitivity

may not be realistic because there is evidence that only 70% of women having antenatal pulmonary embolism had any identifiable classic risk factors suggesting that sensitivity rates above 70% may not be achievable.⁴⁷ In addition, a high sensitivity rate is usually associated with a lower specificity rate and the overall balance of benefits and harms may be undesirable if that means exposing a high proportion of women to thromboprophylaxis.

Despite lack of evidence, many guidelines and clinical care bundles include the use of RAMs to guide VTE prophylaxis. Recently published ACOG guidelines state that most RAMs have not been validated prospectively in the obstetrical population and that current usage of such models is based on extrapolations from non-pregnant women, who differ biologically from pregnant women. The practice bulletin emphasises the need for more research to identify optimal models.³⁷ Although further research is clearly needed the routine use of thromboprophylaxis may present a barrier to generating accurate and precise estimates of the prognostic accuracy of RAMs. Further work to improve RAMs to help stratify the risk of VTE in women who are pregnant or in the puerperium could focus on using decision-analytical modelling to compare the effects, harms and costs of giving thromboprophylaxis to patients with varying risks of VTE. This would allow determination of the risk threshold at which thromboprophylaxis provides optimal overall benefit. Subsequent work to validate these findings would require primary research. Despite the limitations of undertaking accuracy studies in populations where thromboprophylaxis is routinely used, future research could focus on selected higher risk groups who are more likely to benefit from prophylaxis and, with a higher prevalence of VTE, are more amenable to an appropriately powered prospective study. However, given the uncertain benefits and harms of VTE thromboprophylaxis during pregnancy and the postpartum period,^{14 48} risk prediction studies should be undertaken alongside (or as a part of) randomised trials of prophylaxis in targeted groups deemed to be at higher risk of VTE.

CONCLUSIONS

Currently, there are a number of risk assessment models for assessing risk of VTE in pregnancy and the puerperium. Our review has shown that none of these models has been adequately validated and they have limited abilities to detect those at risk of VTE.

Acknowledgements The authors would like to thank all additional members of the core project group for NIHR131021 for input and commentary throughout the work. We are also indebted to Donna Davis for assistance with logistics and administration.

Contributors AP coordinated the review. SD, AP, JD, JH, MC, BJH, CN-P and SG were responsible for the conception, design and obtaining funding for the study. MC developed the search strategy, undertook searches and organised the retrieval of papers. AP, GR, JH, BJH, JD, CN-P and SG were responsible for the acquisition, analysis and interpretation of data. BJH, JD, CN-P and SG helped interpret and provided a methodological, policy and clinical perspective on the data. AP, BJH, JD and SD were responsible for the drafting of this paper, although all authors provided

comments on the drafts, read and approved the final version. AP is the guarantor for the paper.

Funding This study was funded by the UK National Institute for Health Research Health Technology Assessment (NIHR HTA) Programme (project number 131021). The views expressed in this paper are those of the authors and not necessarily those of the NIHR HTA Programme. Any errors are the responsibility of the authors. The funders had no role in the study design, in the collection, analysis and interpretation of data; in the writing of the manuscript; and in the decision to submit the manuscript for publication.

Competing interests CN-P reports personal fees from Sanofi, and is the lead developer of the RCOG Green Top Guideline on thromboprophylaxis in pregnancy (37a). All other authors declare no competing interests.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Abdullah Pandor <http://orcid.org/0000-0003-2552-5260>
 Jahnvi Daru <http://orcid.org/0000-0001-5816-2609>
 Beverley J Hunt <http://orcid.org/0000-0002-4709-0774>
 Gill Rooney <http://orcid.org/0000-0002-8388-9444>
 Jean Hamilton <http://orcid.org/0000-0003-3326-9842>
 Mark Clowes <http://orcid.org/0000-0002-5582-9946>
 Steve Goodacre <http://orcid.org/0000-0003-0803-8444>
 Catherine Nelson-Piercy <http://orcid.org/0000-0001-9311-1196>
 Sarah Davis <http://orcid.org/0000-0002-6609-4287>

REFERENCES

- Say L, Chou D, Gemmill A, *et al*. Global causes of maternal death: a who systematic analysis. *Lancet Glob Health* 2014;2:e323–3.
- Heit JA, Kobbervig CE, James AH, *et al*. Trends in the incidence of venous thromboembolism during pregnancy or postpartum: a 30-year population-based study. *Ann Intern Med* 2005;143:697–706.
- James AH, Jamison MG, Brancaccio LR, *et al*. Venous thromboembolism during pregnancy and the postpartum period: incidence, risk factors, and mortality. *Am J Obstet Gynecol* 2006;194:1311–5.
- Lindqvist P, Dahlbäck B, Marsál K. Thrombotic risk during pregnancy: a population study. *Obstet Gynecol* 1999;94:595–9.
- Royal College of Obstetricians & Gynaecologists. Thrombosis and embolism during pregnancy and the puerperium, reducing the risk (Green-top guideline No. 37A), 2015. Available: <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg37a/> [Accessed 04 Apr 2022].
- Pomp ER, Lenselink AM, Rosendaal FR, *et al*. Pregnancy, the postpartum period and prothrombotic defects: risk of venous thrombosis in the MEGA study. *J Thromb Haemost* 2008;6:632–7.
- Bates SM, Rajasekhar A, Middeldorp S, *et al*. American Society of hematology 2018 guidelines for management of venous

- thromboembolism: venous thromboembolism in the context of pregnancy. *Blood Adv* 2018;2:3317–59.
- 8 Bates SM, Greer IA, Middeldorp S, *et al.* VTE, thrombophilia, antithrombotic therapy, and pregnancy: antithrombotic therapy and prevention of thrombosis, 9th ED: American College of chest physicians evidence-based clinical practice guidelines. *Chest* 2012;141:e691S–736.
 - 9 Chan W-S, Rey E, Kent NE, *et al.* Venous thromboembolism and antithrombotic therapy in pregnancy. *J Obstet Gynaecol Can* 2014;36:527–53.
 - 10 D'Alton ME, Friedman AM, Smiley RM, *et al.* National partnership for maternal safety: consensus bundle on venous thromboembolism. *Obstet Gynecol* 2016;128:688–98.
 - 11 James A, Birsner M, Kaimal A. ACOG practice bulletin No. 196 summary: thromboembolism in pregnancy. *Obstet Gynecol* 2018;132:243–8.
 - 12 Lindqvist PG, Hellgren M. Obstetric thromboprophylaxis: the swedish guidelines. *Adv Hematol* 2011;2011:157483.
 - 13 McLintock C, Brighton T, Chunilal S, *et al.* Recommendations for the prevention of pregnancy-associated venous thromboembolism. *Aust N Z J Obstet Gynaecol* 2012;52:3–13.
 - 14 Middleton P, Shepherd E, Gomersall JC. Venous thromboembolism prophylaxis for women at risk during pregnancy and the early postnatal period. *Cochrane Database Syst Rev* 2021;3:CD001689.
 - 15 Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–9.
 - 16 Davis S, Goodacre S, Pandor A. Estimating the value of future research to improve guidelines on thromboprophylaxis for women during pregnancy and after delivery, 2021. Available: <https://fundingawards.nihr.ac.uk/award/NIHR131021> [Accessed 04 Apr 2022].
 - 17 Wilczynski N. HiRU's Approach to Search Filter Development McMaster University; 2022. https://hiru.mcmaster.ca/hiru/HIRU_Hedges_home.aspx [Accessed 04 Apr 2022].
 - 18 Moons KGM, Wolff RF, Riley RD, *et al.* PROBAST: a tool to assess risk of bias and applicability of prediction model studies: explanation and elaboration. *Ann Intern Med* 2019;170:W1–33.
 - 19 Wolff RF, Moons KGM, Riley RD, *et al.* PROBAST: a tool to assess the risk of bias and applicability of prediction model studies. *Ann Intern Med* 2019;170:51–8.
 - 20 Centre for Reviews and Dissemination. *Systematic Reviews: CRD's Guidance for Undertaking Reviews in Health Care*. York, 2009.
 - 21 McKenzie JE, Brennan SE, Ryan RE. Chapter 9: Summarizing study characteristics and preparing for synthesis. In: Higgins JPT, Thomas J, Chandler J, *et al.*, eds. *Cochrane Handbook for systematic reviews of interventions version 6.3*. 2022. Cochrane, 2022.
 - 22 Hosmer DW, Lemeshow S. *Applied logistic regression*. 2nd Edition. New York: John Wiley & Sons, 2000.
 - 23 Bauersachs RM, Dudenhausen J, Faridi A, *et al.* Risk stratification and heparin prophylaxis to prevent venous thromboembolism in pregnant women. *Thromb Haemost* 2007;98:1237–45.
 - 24 Binstock AB, Larkin JC. Development of a venous thromboembolism prevention tool in the postpartum period [5E]. *Obstet Gynecol* 2019;133:52S.
 - 25 Cavazza S, Rainaldi MP, Adduci A, *et al.* Thromboprophylaxis following cesarean delivery: one site prospective pilot study to evaluate the application of a risk score model. *Thromb Res* 2012;129:28–31.
 - 26 Chau C, Campagna J, Vial M, *et al.* Use of a personalized iterative score to evaluate risk of venous thromboembolism during pregnancy and puerperium. *Int J Gynaecol Obstet* 2019;144:277–82.
 - 27 Chaleur C, Quenet S, Varlet M-N, *et al.* Feasibility of an easy-to-use risk score in the prevention of venous thromboembolism and placental vascular complications in pregnant women: a prospective cohort of 2736 women. *Thromb Res* 2008;122:478–84.
 - 28 Dargaud Y, Rugeri L, Fleury C, *et al.* Personalized thromboprophylaxis using a risk score for the management of pregnancies with high risk of thrombosis: a prospective clinical study. *J Thromb Haemost* 2017;15:897–906.
 - 29 Dargaud Y, Rugeri L, Ninet J, *et al.* Management of pregnant women with increased risk of venous thrombosis. *Int J Gynaecol Obstet* 2005;90:203–7.
 - 30 Gassmann N, Viviano M, Righini M, *et al.* Estimating the risk thresholds used by guidelines to recommend postpartum thromboprophylaxis. *J Thromb Haemost* 2021;19:452–9.
 - 31 Hase EA, Barros VIPVde, Igai AMK, *et al.* Risk assessment of venous thromboembolism and thromboprophylaxis in pregnant women hospitalized with cancer: preliminary results from a risk score. *Clinics* 2018;73:e368.
 - 32 Lindqvist PG, Torsson J, Almqvist A, *et al.* Postpartum thromboembolism: severe events might be preventable using a new risk score model. *Vasc Health Risk Manag* 2008;4:1081–7.
 - 33 Lok WY, Kong CW, To WWK, . A local risk score model for venous thromboembolism prophylaxis for caesarean section in Chinese women and comparison with international guidelines. *Taiwan J Obstet Gynecol* 2019;58:520–5.
 - 34 Shacaluga A, Rayment R. Venous thromboembolic risk assessment in pregnancy: comparison of the All-Wales maternity risk assessment tool with guidance from the Royal College of obstetrics and gynaecology. *Br J Haematol* 2019;185:162–5.
 - 35 Sultan AA, West J, Grainge MJ, *et al.* Development and validation of risk prediction model for venous thromboembolism in postpartum women: multinational cohort study. *BMJ* 2016;355:i6253.
 - 36 Testa S, Passamonti SM, Paoletti O, *et al.* The "pregnancy health-care program" for the prevention of venous thromboembolism in pregnancy. *Intern Emerg Med* 2015;10:129–34.
 - 37 Tran JP, Stribling SS, Ibezim UC, *et al.* Performance of risk assessment models for peripartum thromboprophylaxis. *Reprod Sci* 2019;26:1243–8.
 - 38 Weiss N, Bernstein PS. Risk factor scoring for predicting venous thromboembolism in obstetric patients. *Am J Obstet Gynecol* 2000;182:1073–5.
 - 39 Ellis-Kahana J, Sparks AD, Gimovsky AC, *et al.* Developing a model for predicting venous thromboembolism in obese pregnant women in a national study. *Thromb Res* 2020;191:42–9.
 - 40 Collins GS, Reitsma JB, Altman DG, *et al.* Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD statement. *BMC Med* 2015;13:1.
 - 41 Moons KGM, de Groot JAH, Bouwmeester W, *et al.* Critical appraisal and data extraction for systematic reviews of prediction modelling studies: the charms checklist. *PLoS Med* 2014;11:e1001744.
 - 42 Andrew L, Ní Áinle F, Blondon M, *et al.* Preventing postpartum venous thromboembolism: a call to action to reduce undue maternal morbidity and mortality. *Thromb Res* 2020;193:190–7.
 - 43 Reps JM, Ryan PB, Rijnbeek PR, *et al.* Design matters in patient-level prediction: evaluation of a cohort vs. case-control design when developing predictive models in observational healthcare datasets. *J Big Data* 2021;8.
 - 44 Higgins JPT, Thomas J, Chandler J. *Cochrane Handbook for systematic reviews of interventions version 6.3*. Cochrane, 2022.
 - 45 Knight M, Bunch K, Tuffnell D. *Saving Lives, Improving Mothers' Care - Lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2014-16*. Oxford: National Perinatal Epidemiology Unit, University of Oxford, 2018.
 - 46 National Institute for Health and Care Excellence. Venous thromboembolism in over 16s: reducing the risk of hospital-acquired deep vein thrombosis or pulmonary embolism - NICE Guideline 89. London UK NICE; 2019.
 - 47 Knight M, UKOSS. Antenatal pulmonary embolism: risk factors, management and outcomes. *BJOG* 2008;115:453–61.
 - 48 Lu MY, Blanchard CT, Ausbeck EB, *et al.* Evaluation of a risk-stratified, heparin-based, obstetric thromboprophylaxis protocol. *Obstet Gynecol* 2021;138:530–8.