



Published in final edited form as:

Obesity (Silver Spring). 2014 November ; 22(11): 2434–2440. doi:10.1002/oby.20865.

Association of the Degree of Adiposity and Duration of Obesity with Measures of Cardiac Structure and Function: The CARDIA Study

Jared P. Reis, PhD¹, Norrina Allen, PhD², Bethany B. Gibbs, PhD³, Samuel S. Gidding, MD⁴, Joyce M. Lee, MD, MPH⁵, Cora E. Lewis, MD, MSPH⁶, Joao Lima, MD⁷, Donald Lloyd-Jones, MD, ScM², Catherine M. Loria, PhD¹, Tiffany M. Powell-Wiley, MD, MPH⁸, Shishir Sharma, MD², Gina Wei, MD, MPH¹, and Kiang Liu, PhD²

¹Division of Cardiovascular Sciences, National Heart, Lung, and Blood Institute, Bethesda, MD

²Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL

³Department of Health and Physical Activity, University of Pittsburgh, Pittsburgh, PA

⁴A.I. DuPont Hospital for Children, Wilmington, DE

⁵Division of General Pediatrics, University of Michigan, Ann Arbor, MI

⁶Division of Preventive Medicine, Department of Medicine, University of Alabama at Birmingham, Birmingham, AL

⁷Departments of Cardiology and Radiology, Johns Hopkins University School of Medicine, Baltimore, MD

⁸Cardiovascular and Pulmonary Branch, National Heart, Lung, and Blood Institute, Bethesda, MD

Abstract

OBJECTIVE—Examine whether there are independent influences of a greater degree of adiposity and longer duration of obesity on cardiac structure and function.

METHODS—Participants of CARDIA were 18-30 years when they underwent a baseline examination in 1985-86. Seven follow-up examinations were conducted every 2-5 years.

RESULTS—Among 2,547 participants who underwent an echocardiogram at the year 25 examination and were not obese at baseline, 34.4% and 35.5% were overall (BMI ≥ 30 kg/m²) and abdominally obese (waist circumference: men: >102 cm; women: >88 cm) at year 25, respectively. A greater degree of overall and abdominal adiposity at year 25 were each associated with a greater left ventricular (LV) mass ($p<0.001$), LV volume ($p<0.001$), LV mass-to-volume

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Name and address for correspondence: Jared P. Reis, PhD, Division of Cardiovascular Sciences, National Heart, Lung, and Blood Institute, 6701 Rockledge Dr, Suite 10186, Bethesda, MD, 20892, (301) 435-1291, FAX: (301) 435-1291, reisjp@mail.nih.gov.

Disclosures

The authors declare no conflict of interest.

ratio ($p<0.001$), left atrial dimension ($p<0.001$), and ejection fraction ($p<0.05$) after adjustment for duration of obesity and other risk factors. In contrast, a longer duration of overall obesity was associated with a greater LV mass ($p=0.003$) and a trend for a lower ejection fraction ($p=0.07$).

CONCLUSIONS—A greater degree of adiposity is strongly associated with concentric LV remodeling in midlife, while the cumulative effects of a longer duration of overall obesity during young adulthood contribute to concentric remodeling predominantly by increasing LV mass.

Keywords

cardiovascular risk; obesity; heart; echocardiography

Introduction

Obesity has been linked to a spectrum of cardiovascular changes ranging from a hyperdynamic circulation, through subclinical cardiac structural changes to overt heart failure (1-4). Existing studies have shown that left ventricular (LV) hypertrophy, LV systolic function, and LV diastolic function worsen with a greater degree of overall and abdominal adiposity (5). Recent evidence also suggests obesity is associated with a greater LV mass to a greater extent than chamber size (6-8). While the pattern of remodeling associated with a greater degree of adiposity has been documented, little information exists on the consequences of a longer duration of obesity.

Understanding the consequences of a longer duration of obesity is critical, given the obesity epidemic. With a doubling of obesity rates for adults and a tripling of rates for children and adolescents over the last three decades, younger individuals are experiencing a greater degree and longer duration of exposure to excess adiposity over their lifetime (9, 10). Preliminary data suggest that a longer duration of obesity contributes to a greater LV mass and worse LV function in most (11-13), but not all studies (14). Important limitations of these studies include small clinic-based samples, lack of simultaneous adjustment for the degree of adiposity, and the dependence upon the recall of body weight at previous ages to determine the duration of obesity, a methodology that may be susceptible to a substantial amount of measurement error.

The present study was conducted to examine whether there are independent influences of a greater degree of adiposity and longer duration of obesity beginning in early adulthood as a result of the obesity epidemic on cardiac structure and function measured via echocardiography in middle-age. Participants were black and white adults recruited in 1985-1986 as part of the Coronary Artery Risk Development in Young Adults (CARDIA) Study. The CARDIA Study provided an ideal setting in which to address this question because participants were recruited and followed largely during the obesity epidemic of the last three decades in the United States as well as during the period of greatest gain in adiposity during the life course (9, 15).

Methods

Study sample

CARDIA is a multicenter community-based longitudinal cohort study of the development and determinants of cardiovascular disease over time in 5,115 young adults initially aged 18–30 years in 1985–1986. Black and white adults were recruited from 4 cities in the United States with population-based samples approximately balanced within center by sex, age, race, and education. To date, participants have been re-examined 2, 5, 7, 10, 15, 20, and 25 years after baseline and retention rates across examinations were 91%, 86%, 81%, 79%, 74%, 72%, and 72%, respectively.

Of the 3,498 participants who completed the year 25 follow-up examination, we excluded those who did not complete an echocardiogram (n=24) and those without available measures of cardiac structure and function (n=353); those who were overall [body mass index (BMI) $\geq 30 \text{ kg/m}^2$] or abdominally obese ($>102 \text{ cm}$ in men and $>88 \text{ cm}$ in women (16)) at baseline (n=341); were pregnant during any examination (n=160); those who experienced angina, a myocardial infarction, or heart failure during follow-up (n=54); and those who had bariatric surgery (n=19). The remaining 2,547 participants formed the sample population for analysis.

Clinical measurements

Standardized protocols for data collection were used across study centers and examinations. Participants were asked to fast for at least 12 hours and to avoid smoking or engaging in heavy physical activity for at least 2 hours before each examination. Anthropometry Weight and height were measured with participants wearing light examination clothes and no shoes. Body weight was measured to the nearest 0.2 kg with a calibrated balance-beam scale. Height was measured with a vertical ruler to the nearest 0.5 cm. BMI was calculated as weight in kilograms divided by height in meters squared. Waist circumference was measured with a tape in duplicate to the nearest 0.5 cm around the minimal abdominal girth identified laterally midway between the iliac crest and the lowest portion of the rib cage and anteriorly midway between the xiphoid process and the umbilicus.

Duration of overall and abdominal obesity

For each participant, the number of years of overall and abdominal obesity was calculated separately based upon the presence (or absence) of overall or abdominal obesity at each follow-up examination and the preceding examination. For example, a participant who was not overall obese at year 2, but was overall obese at year 5 and all subsequent follow-up examinations was assigned 0 years of overall obesity at year 2, 3 years at year 5, 2 years at year 7, 3 years at year 10, and so on. This scoring methodology was applied at each follow-up examination and the cumulative duration of obesity across examination years for each participant was then calculated (17).

Echocardiography

Echocardiography was performed at year 25 using an Artida cardiac ultrasound machine (Toshiba Medical Systems, Tokyo, Japan) and a previously described protocol (18, 19). Briefly, sonographers at each field center underwent initial, centralized training followed by

quality assurance and control procedures to assess intra- and inter-sonographer reproducibility. Measurements of LV and left atrium (LA) dimensions were acquired from 2-dimensional (2D)-guided M-mode echocardiograms obtained from optimized parasternal short-axis views. Studies were electronically transmitted to a core reading laboratory at Johns Hopkins University. LV mass was calculated using the Devereux formula (20). LV end-diastolic volume was acquired from a 2D 4-chamber view. Mass-to-volume ratio was calculated as the ratio of LV mass to end-diastolic volume. Ejection fraction was calculated as the ratio of stroke volume to end-diastolic volume. Heart rate was recorded from the electrocardiogram during the echocardiographic examination. Quality control procedures included participant re-examination and blind re-reading to allow assessment of intra- and inter-sonographer variability and intra- and inter-reader variability, respectively. Technical errors for intra- and inter-sonographer variability ranged from 5-11% and 6-12%, respectively. Intra- and inter-reader variability ranged from 3-9% and 6-11%, respectively.

The assessment of other measurements can be found in the Supplemental file.

Statistical analysis

Participant characteristics overall, and according to the degree of adiposity and duration of obesity were described using means, medians, and proportions as appropriate. Differences and trends were tested using linear regression models and χ^2 analyses for continuous and categorical characteristics, respectively. The Kruskal-Wallis test was used for characteristics with skewed distributions. Multivariable linear regression models were used to estimate the association between the degree of adiposity at year 25 and duration of obesity with each of the measures of cardiac structure and function. The degree of adiposity and duration of obesity were modeled in one of two exposure forms: as continuous variables assuming a linear dose-response association, and as five-level categorical variables for overall and abdominal (i.e., sex-specific quintiles) adiposity and a six-level categorical variable for duration of obesity. Models were simultaneously adjusted for degree of adiposity and duration of obesity. Analyses were further adjusted for age, sex, race, educational attainment, CARDIA field center, smoking, and average physical activity and alcohol intake during follow-up. A second model adjusted additionally for use of antihypertensive medication, diabetes, heart rate, and average systolic blood pressure during follow-up. The additional adjustment variables in the second model could be in the causal pathway; we regarded these models as possibly explanatory. Models using BMI as an exposure did not additionally adjust for height. Tests for a linear trend were performed by entering the degree of adiposity or duration of obesity into the multivariable models as a continuous variable. Potential effect modification by race-sex was explored in stratified analyses and evaluated by testing the statistical significance of multiplicative interaction terms in models that also included lower-order terms.

In a sensitivity analysis, we determined the influence of missing BMI and waist circumference values (7.8% of all measurements for each) on the associations between the degree of adiposity and duration of obesity with measures of cardiac structure and function. Multiple imputation was used to impute missing BMI and waist circumference values using the sequential regression imputation approach that is implemented in the software package

IVEware (21). Five datasets were generated using all available BMI and waist circumference data. Each data set was analyzed separately and results from the 5 analyses were combined using the rules of Little and Rubin (22).

Tests of statistical significance were two-tailed, with an alpha level of 0.05. Due to the increased number of formal tests of interaction, statistical significance for effect modification was defined at an alpha level of 0.01. SAS version 9.3 (SAS Institute, Cary, NC) was used to perform all analyses.

Results

Of the 2,547 eligible participants, 44.4% were black and 53.0% were women. At year 25, the prevalence of a BMI <18.5, 18.5-24.9, 25.0-29.9, 30.0-34.9, and ≥ 35.0 kg/m² was 0.9%, 28.8%, 35.9%, 21.9%, and 12.5%, respectively. The prevalence of abdominal obesity at year 25 was 35.5%. The average duration of overall and abdominal obesity during follow-up was 14.1 (SD=6.7) and 12.7 (SD=6.3) years, respectively. Mean age at onset of obesity was 36.1 (SD=8.1) and 38.3 (SD=7.6) years, respectively. The prevalence of overall obesity duration of 1-5, 6-10, 11-15, 16-20, and >20 years was 8.4%, 7.4%, 8.0%, 8.2%, and 7.1%, respectively. For abdominal obesity duration, these estimates were 11.3%, 8.7%, 8.0%, 7.9%, and 4.4%, respectively.

Those with a longer exposure to overall obesity were more likely to be younger at baseline, black, achieve less education, have a higher average waist circumference during follow-up, and were less physically active and consumed less alcohol during follow-up (Table 1). In addition, a longer duration of obesity was associated with a higher heart rate, average blood pressures, glucose and insulin levels, and were more likely to use antihypertensive medication and have diabetes. Sex and smoking were unrelated to obesity duration. Similar results were observed when the duration of overall obesity was replaced with the degree of overall obesity, degree of abdominal obesity, and the duration of abdominal obesity except there was no association of age at baseline with the degree of abdominal adiposity or duration of abdominal obesity, and women were more likely to have a longer duration of abdominal obesity (data not shown).

Mean (SD) LV mass, LV end-diastolic volume, LV mass-to-volume ratio, LA dimension, and ejection fraction were 164.0 (48.1) g, 126.0 (30.3) ml, 1.32 (0.32) g/ml, 3.66 (0.47) cm, and 69.6% (7.8), respectively. A greater degree of overall adiposity at year 25 was strongly associated with a greater LV mass, LV end-diastolic volume, LV mass-to-volume ratio, and LA dimension (p-trend <0.001, for all) (Table 2). With each five-unit higher BMI, an 11.4 g greater LV mass, a 5.9 ml greater LV end-diastolic volume, a 0.04 g/ml higher LV mass-to-volume ratio, and a 0.2 cm larger LA was observed. It was also associated with a higher ejection fraction (0.5% higher with each five-unit higher BMI; p-trend 0.03). In contrast, the duration of overall obesity was associated with a greater LV mass (p-trend 0.003) and a lower ejection fraction that approached statistical significance (p-trend 0.05). With each additional 5-years of overall obesity, a 2.5 g greater LV mass and a 0.3% lower ejection fraction was observed. Adjustment for antihypertensive medication use, heart rate, average systolic blood pressure, and diabetes during follow-up had minimal influence on these

associations (Table 2; model 2). Results were also consistent when LV mass was indexed to height^{2.7} (data not shown).

A greater degree of abdominal adiposity at year 25 was strongly associated with a greater left ventricular mass, LV end-diastolic volume, LV mass-to-volume ratio, and LA dimension (p-trend <0.001, for all) (Table 3). With each 10 cm higher waist circumference, an 11.0 g greater LV mass, a 4.0 ml greater LV end-diastolic volume, a 0.05 g/ml higher LV mass-to-volume ratio, and a 0.2 cm larger LA was observed. Similarly, a greater degree of abdominal adiposity was associated with a higher ejection fraction that approached statistical significance (0.4% higher with each 10 cm higher waist circumference; p-trend 0.06). In contrast, the duration of abdominal obesity was associated with a greater LV end-diastolic volume only (p-trend 0.03). Adjustment for antihypertensive medication use, heart rate, average systolic blood pressure, and diabetes during follow-up had minimal influence on these associations (Table 3; model 2).

Associations did not vary by race-sex except for the association of the duration of overall obesity with LV mass-to-volume ratio (p-interaction 0.007) and the degree of overall adiposity with LA dimension (p-interaction <0.001). However, similar to findings in the total sample, no association of the duration of overall obesity with LV mass-to-volume ratio was observed within each race-sex group [slope: 0.01, -0.02, 0.01, and 0.01 for black men, black women, white men, and white women, respectively (p >0.2, for all), per 5-year longer duration of overall obesity]. Measures of association were in a similar direction for the association of the degree of overall adiposity with LA dimension [slope: 0.17, 0.09, 0.01, 0.19 for black men, black women, white men, and white women, respectively (p <0.001, for all), per 5-unit higher BMI].

We observed similar results when we further adjusted for weight change over the follow-up period (quintiles) and when we imputed missing values using all available measures of BMI and waist circumference during follow-up on the sample of 2,547 participants (data not shown).

Discussion

In this community-based sample of adults recruited and followed largely during the obesity epidemic in the United States, a greater degree of overall and abdominal adiposity were each strongly associated with numerous measures of cardiac structure, including a greater left ventricular mass, left ventricular end diastolic volume, and left atrial dimension. A greater LV mass-to-volume ratio suggested an adverse pattern of concentric LV remodeling with a greater degree of adiposity. A longer duration of overall obesity, on the other hand, was strongly associated with a greater LV mass. These findings were independent of sociodemographic characteristics, lifestyle behaviors, and other cardiovascular risk factors. In addition, results did not vary consistently between white and black men and women. These results suggest that a greater degree of adiposity is strongly associated with concentric LV remodeling in midlife, while the cumulative effects of a longer duration of overall obesity during young adulthood may contribute to concentric remodeling predominately by increasing LV mass.

Findings of the current study suggest that the degree of adiposity may be a more important determinant of cardiac structure than a longer duration of obesity. This was evidenced by strong dose-response associations of the degree of overall and abdominal adiposity with nearly all measures of LV structure and LA size. Obesity has long been considered as a state of chronic volume overload because the heart is required to circulate blood through the large depot of adipose tissue (23). A greater degree of adiposity contributes to a higher cardiac output and workload, and a greater LV mass (24, 25). Early studies suggested that a greater degree of adiposity was associated with eccentric LV remodeling (26). However, the results of the current study as well as a number of other recent studies suggest a pattern of concentric LV remodeling whereby the LV mass may increase to a greater extent than cavity size (higher LV mass-to-volume ratio) (8, 27).

Studies estimating the association of a longer duration of obesity on cardiac structure have been mixed. These studies have relied exclusively on the self-report of body weight at previous ages to determine the duration of obesity as opposed to the objective assessment in the current study, and did not account for the degree of adiposity. In a small, clinic-based sample of morbidly obese patients prior to weight loss surgery, Alpert et al. (11) showed the duration of morbid obesity correlated significantly with a higher LV mass and poorer LV systolic function. In a sample of 35 obese patients, a longer duration of obesity was associated with LV enlargement and wall thickening independent of body weight (12). However, among 109 overweight and obese adults attending a weight management center, the duration of obesity showed no association with numerous measures of LV structure and function (14). In the current study, we found a longer duration of overall obesity was associated with a greater LV mass independent of the degree of adiposity, potential confounders, and intermediate factors. Our findings suggest that while the degree of adiposity may be a more important determinant of cardiac structure in midlife, the long-term cumulative effects of sustained obesity and metabolic dysregulation during young adulthood may also contribute to concentric remodeling primarily by increasing LV mass.

In the current study, we found significant, positive associations of the degree of overall and abdominal adiposity with ejection fraction that were independent of the duration of obesity, potential confounders, and risk factors. Conversely, we observed an inverse association of a longer duration of overall obesity with a lower ejection fraction that approached statistical significance. Findings from available studies of the degree of adiposity with LV systolic function have varied considerably suggesting that ejection fraction is lower (28), similar (29), or higher (30) among obese subjects as compared to those who are not obese. Existing studies of the duration of obesity have generally shown consistent inverse associations of prolonged obesity with reduced fractional shortening (11), stroke volume (12), and ejection fraction (13). Findings from the current study suggest that there may be a compensatory increase in systolic function in middle-age with a greater degree of adiposity that may subsequently become impaired with a longer duration of obesity. Future studies are needed to confirm these findings particularly with more sensitive indicators of LV contractile function.

In addition to the increased cardiac work load and resultant increased LV mass that occurs to meet the enhanced metabolic demands of both lean and fat mass with obesity, a number of

other potential mechanisms may help to explain the association of a greater degree of adiposity with structural and functional changes in the heart. Intra-abdominal fat located in close proximity to the portal circulation may mediate concentric LV remodeling by secreting a number of pro-inflammatory cytokines that may play a role in the myocardial remodeling process by directly influencing hypertrophy, fibrosis, and ultimately contractility (31, 32). Hyperinsulinemia and insulin resistance, which may promote myocardial hypertrophy via the growth-stimulating effects of insulin, may also mediate the increased LV mass and LA size in obesity (14, 33). Obstructive sleep apnea may also contribute to LV structural changes by exacerbating daytime and nighttime hypertension, increased sympathetic tone, and chronic hypoxemia (34). Additional mechanisms that may explain, at least in part, the association of a greater degree of adiposity with LV concentric remodeling include increases in leptin and angiotensin II, and reductions in adiponectin (35-37). The cumulative effects of these hemodynamic and metabolic changes as a result of prolonged obesity may also help to at least partially explain the association of a longer duration of overall obesity with increases in LV mass and impaired systolic function over time.

Strengths of the current study include a community-based sampling method; cohort recruitment and follow-up largely during the obesity epidemic of the last three decades; a biracial cohort; extensive data on potential confounders; a large sample size well balanced with respect to age, sex, race, and education that increased precision and permitted simultaneous adjustment and stratification by multiple variables; repeated assessments of BMI, waist circumference, and potential confounding factors; a high retention rate; and the standardized data collection protocols and rigorous quality control of the CARDIA study. Nevertheless, three limitations deserve mention. First, our study relied on relatively simple estimates of anthropometry that are unable to distinguish between fat and fat-free mass. However, these measures are reliable, efficient, cost-effective, and widely applicable for research in large studies and clinical practice. Second, our estimation of the duration of overall and abdominal obesity during follow-up was based on the measurement of BMI and waist circumference, respectively, every two to five years. It is likely that a more frequent number of assessments would have led to a more accurate estimation of the duration of obesity during follow-up. Third, since our study collected data over a 25-year follow-up period, some participants were missing at least one eligible measurement of BMI and waist circumference. However, we noted similar results between our multiple imputed datasets and our primary dataset.

In conclusion, in this community-based longitudinal cohort study of adults recruited and followed during the obesity epidemic over the last three decades in the United States, we found the degree of overall and abdominal adiposity to be important factors associated with a greater LA dimension and a pattern of LV concentric remodeling. These findings were independent of the duration of obesity and numerous potential confounding factors and cardiovascular risk factors. A longer duration of overall obesity, on the other hand, was associated with a greater left ventricular mass and a trend for a lower ejection fraction. Thus, these findings suggest that cardiac remodeling throughout young adulthood results largely as a consequence of a greater degree of adiposity, while the cumulative effects of a longer duration of overall obesity may contribute to concentric remodeling predominately by increasing LV mass. Future public health interventions focused on cardiovascular disease

prevention should focus on younger populations in order to prevent or at least delay the onset of obesity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Sources of funding

The Coronary Artery Risk Development in Young Adults Study (CARDIA) is supported by contracts HHSN268201300025C, HHSN268201300026C, HHSN268201300027C, HHSN268201300028C, HHSN268201300029C, and HHSN268200900041C from the National Heart, Lung, and Blood Institute (NHLBI), the Intramural Research Program of the National Institute on Aging (NIA), and an intra-agency agreement between NIA and NHLBI (AG0005).

References

- Alexander JK. Obesity and the heart. *Heart Dis Stroke*. Jul+aug;1993 2(4):317–21. [PubMed: 8156183]
- Alpert MA. Obesity cardiomyopathy: pathophysiology and evolution of the clinical syndrome. *Am J Med Sci*. Apr; 2001 321(4):225–36. [PubMed: 11307864]
- Kenchiah S, Evans JC, Levy D, Wilson PW, Benjamin EJ, Larson MG, et al. Obesity and the risk of heart failure. *The New England journal of medicine*. Aug 1; 2002 347(5):305–13. [PubMed: 12151467]
- Lauer MS, Anderson KM, Levy D. Separate and joint influences of obesity and mild hypertension on left ventricular mass and geometry: the Framingham Heart Study. *Journal of the American College of Cardiology*. Jan; 1992 19(1):130–4. [PubMed: 1729324]
- Vasan RS. Cardiac function and obesity. *Heart (British Cardiac Society)*. Oct; 2003 89(10):1127–9. [PubMed: 12975393]
- Lauer MS, Anderson KM, Kannel WB, Levy D. The impact of obesity on left ventricular mass and geometry. The Framingham Heart Study. *Jama*. Jul 10; 1991 266(2):231–6. [PubMed: 1829117]
- Peterson LR, Waggoner AD, Schechtman KB, Meyer T, Gropler RJ, Barzilai B, et al. Alterations in left ventricular structure and function in young healthy obese women: assessment by echocardiography and tissue Doppler imaging. *Journal of the American College of Cardiology*. Apr 21; 2004 43(8):1399–404. [PubMed: 15093874]
- Turkbey EB, McClelland RL, Kronmal RA, Burke GL, Bild DE, Tracy RP, et al. The impact of obesity on the left ventricle: the Multi-Ethnic Study of Atherosclerosis (MESA). *JACC Cardiovasc Imaging*. Mar; 2010 3(3):266–74. [PubMed: 20223423]
- Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *Jama*. Feb 1; 2012 307(5):491–7. [PubMed: 22253363]
- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *Jama*. Feb 1; 2012 307(5):483–90. [PubMed: 22253364]
- Alpert MA, Lambert CR, Panayiotou H, Terry BE, Cohen MV, Massey CV, et al. Relation of duration of morbid obesity to left ventricular mass, systolic function, and diastolic filling, and effect of weight loss. *Am J Cardiol*. Dec 1; 1995 76(16):1194–7. [PubMed: 7484912]
- Nakajima T, Fujioka S, Tokunaga K, Hirobe K, Matsuzawa Y, Tarui S. Noninvasive study of left ventricular performance in obese patients: influence of duration of obesity. *Circulation*. Mar; 1985 71(3):481–6. [PubMed: 3971522]

13. Scaglione R, Diciara MA, Indovina A, Lipari R, Ganguzza A, Parrinello G, et al. Left ventricular diastolic and systolic function in normotensive obese subjects: influence of degree and duration of obesity. *Eur Heart J*. Jun; 1992 13(6):738–42. [PubMed: 1623860]
14. Wong CY, O'Moore-Sullivan T, Leano R, Byrne N, Beller E, Marwick TH. Alterations of left ventricular myocardial characteristics associated with obesity. *Circulation*. Nov 9; 2004 110(19): 3081–7. [PubMed: 15520317]
15. Sheehan TJ, DuBrava S, DeChello LM, Fang Z. Rates of weight change for black and white Americans over a twenty year period. *Int J Obes Relat Metab Disord*. Apr; 2003 27(4):498–504. [PubMed: 12664083]
16. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation*. Oct 25; 2005 112(17):2735–52. [PubMed: 16157765]
17. Reis JP, Loria CM, Lewis CE, Powell-Wiley TM, Wei GS, Carr JJ, et al. Association between duration of overall and abdominal obesity beginning in young adulthood and coronary artery calcification in middle age. *Jama*. Jul 17; 2013 310(3):280–8. [PubMed: 23860986]
18. Gardin JM, Wagenknecht LE, Anton-Culver H, Flack J, Gidding S, Kurosaki T, et al. Relationship of cardiovascular risk factors to echocardiographic left ventricular mass in healthy young black and white adult men and women. The CARDIA study. *Coronary Artery Risk Development in Young Adults*. *Circulation*. Aug 1; 1995 92(3):380–7. [PubMed: 7634452]
19. Gidding SS, Liu K, Colangelo LA, Cook NL, Goff DC, Glasser SP, et al. Longitudinal determinants of left ventricular mass and geometry: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Circ Cardiovasc Imaging*. Sep; 2013 6(5):769–75. [PubMed: 23922005]
20. Devereux RB, Alonso DR, Lutas EM, Gottlieb GJ, Campo E, Sachs I, et al. Echocardiographic assessment of left ventricular hypertrophy: comparison to necropsy findings. *Am J Cardiol*. Feb 15; 1986 57(6):450–8. [PubMed: 2936235]
21. Raghunathan TE, Lepkowski JM, VanHoewyk J, Solenberger P. A multivariate technique for multiply imputing missing values using a sequence of regression models. *Survey Methodology*. 2001; 27:85–95.
22. Little, RJA.; Rubin, DB. *Statistical Analysis with Missing Data*. 2nd edition ed.. John Wiley; New York: 2002.
23. Aurigemma GP, de Simone G, Fitzgibbons TP. Cardiac remodeling in obesity. *Circ Cardiovasc Imaging*. Jan 1; 2013 6(1):142–52. [PubMed: 23322729]
24. Bella JN, Devereux RB, Roman MJ, O'Grady MJ, Welty TK, Lee ET, et al. Relations of left ventricular mass to fat-free and adipose body mass: the strong heart study. The Strong Heart Study Investigators. *Circulation*. Dec 8; 1998 98(23):2538–44. [PubMed: 9843460]
25. de Simone G, Devereux RB, Kizer JR, Chinali M, Bella JN, Oberman A, et al. Body composition and fat distribution influence systemic hemodynamics in the absence of obesity: the HyperGEN Study. *The American journal of clinical nutrition*. Apr; 2005 81(4):757–61. [PubMed: 15817849]
26. Messerli FH, Sundgaard-Riise K, Reisin ED, Dreslinski GR, Ventura HO, Oigman W, et al. Dimorphic cardiac adaptation to obesity and arterial hypertension. *Annals of internal medicine*. Dec; 1983 99(6):757–61. [PubMed: 6651022]
27. Powell BD, Redfield MM, Bybee KA, Freeman WK, Rihal CS. Association of obesity with left ventricular remodeling and diastolic dysfunction in patients without coronary artery disease. *Am J Cardiol*. Jul 1; 2006 98(1):116–20. [PubMed: 16784933]
28. Ferraro S, Perrone-Filardi P, Desiderio A, Betocchi S, D'Alto M, Liguori L, et al. Left ventricular systolic and diastolic function in severe obesity: a radionuclide study. *Cardiology*. Jul-Aug;1996 87(4):347–53. [PubMed: 8793172]
29. Avelar E, Cloward TV, Walker JM, Farney RJ, Strong M, Pendleton RC, et al. Left ventricular hypertrophy in severe obesity: interactions among blood pressure, nocturnal hypoxemia, and body mass. *Hypertension*. Jan; 2007 49(1):34–9. [PubMed: 17130310]

30. Garavaglia GE, Messerli FH, Nunez BD, Schmieder RE, Grossman E. Myocardial contractility and left ventricular function in obese patients with essential hypertension. *Am J Cardiol.* Sep 15; 1988 62(9):594–7. [PubMed: 3414551]
31. Aukrust P, Yndestad A, Damas JK, Gullestad L. Inflammation and chronic heart failure–potential therapeutic role of intravenous immunoglobulin. *Autoimmun Rev.* Mar; 2004 3(3):221–7. [PubMed: 15110235]
32. Hilfiker-Kleiner D, Landmesser U HD. Molecular mechanisms in heart failure: focus on cardiac hypertrophy, inflammation, angiogenesis, and apoptosis. *Journal of the American College of Cardiology.* 2006; 48:A56–A66.
33. Karason K, Sjostrom L, Wallentin I, Peltonen M. Impact of blood pressure and insulin on the relationship between body fat and left ventricular structure. *Eur Heart J.* Aug; 2003 24(16):1500–5. [PubMed: 12919774]
34. Abel ED, Litwin SE, Sweeney G. Cardiac remodeling in obesity. *Physiol Rev.* Apr; 2008 88(2): 389–419. [PubMed: 18391168]
35. Barouch LA, Berkowitz DE, Harrison RW, O'Donnell CP, Hare JM. Disruption of leptin signaling contributes to cardiac hypertrophy independently of body weight in mice. *Circulation.* Aug 12; 2003 108(6):754–9. [PubMed: 12885755]
36. Engeli S, Negrel R, Sharma AM. Physiology and pathophysiology of the adipose tissue renin-angiotensin system. *Hypertension.* Jun; 2000 35(6):1270–7. [PubMed: 10856276]
37. Liao Y, Takashima S, Maeda N, Ouchi N, Komamura K, Shimomura I, et al. Exacerbation of heart failure in adiponectin-deficient mice due to impaired regulation of AMPK and glucose metabolism. *Cardiovasc Res.* Sep 1; 2005 67(4):705–13. [PubMed: 15907819]

What is already known on this topic?

- The degree of adiposity (i.e., body mass index or waist circumference) is associated with adverse cardiac remodeling
- Less information is available on whether a longer duration of obesity also contributes independently to cardiac remodeling
- The few studies available that have investigated the duration of obesity have been limited by small clinic-based samples, lack of simultaneous control for the degree of adiposity, and the self-report of the duration of obesity

What this study adds?

- Few, if any, long-term studies are able to objectively estimate the duration of obesity from repeat assessments of adiposity
- This study compared the relative importance of both the degree of adiposity and duration of obesity in a large community-based sample
- The degree of adiposity was strongly associated with adverse cardiac remodeling, while the cumulative effects of a longer duration of obesity contribute to concentric left ventricular remodeling by increasing left ventricular mass

Table 1

Participant characteristics* according to duration of overall obesity† in 5-year increments, The CARDIA Study (n = 2,547).

Variable*	Duration of overall obesity (years)						p-trend
	0 (n = 1,552)	1-5 (n = 215)	6-10 (n = 188)	11-15 (n = 203)	16-20 (n = 208)	>20 (n = 181)	
Selected characteristics							
Baseline age (years)	25.1 (3.5)	24.5 (3.5)	25.1 (3.6)	24.9 (3.8)	24.2 (3.8)	24.8 (3.6)	0.003
Women, % (n)	51.7 (802)	54.4 (117)	54.3 (102)	54.7 (111)	54.8 (114)	58.0 (105)	0.07
Black race, %	35.4 (550)	49.3 (106)	53.2 (100)	58.6 (119)	61.5 (128)	70.7 (128)	< 0.001
Educational attainment (years)	16.0 (2.6)	15.7 (2.5)	15.5 (2.6)	15.6 (2.6)	15.5 (2.5)	15.1 (2.4)	< 0.001
Ever smoker, % (n)	39.0 (605)	41.4 (89)	39.4 (74)	34.5 (70)	36.1 (75)	33.7 (61)	0.09
Physical activity (exercise units)‡	366 (243-524)	324 (200-511)	316 (205-474)	293 (176-477)	289 (186-454)	273 (152-428)	< 0.001
Alcohol (ml/day)‡	7.3 (1.6-17.2)	5.2 (0.7-16.4)	7.1 (1.0-17.0)	4.2 (0.6-11.3)	3.7 (0.4-12.8)	2.9 (0.3-10.6)	< 0.001
Clinical and physical characteristics							
Body mass index (kg/m ²)‡	23.3 (2.2)	26.5 (1.5)	27.4 (1.4)	28.7 (1.5)	30.6 (2.2)	33.4 (3.1)	< 0.001
Waist circumference (cm)‡	77.7 (7.9)	84.2 (7.1)	86.1 (7.3)	88.2 (7.3)	92.3 (7.6)	96.8 (8.1)	< 0.001
Heart rate (bpm)	63.6 (10.4)	66.4 (10.2)	66.9 (9.6)	67.5 (10.3)	67.7 (10.4)	68.1 (10.6)	< 0.001
Systolic blood pressure (mmHg)‡	109.3 (9.2)	110.9 (9.4)	110.8 (8.5)	113.5 (9.2)	112.9 (8.5)	114.9 (8.7)	< 0.001
Diastolic blood pressure (mmHg)‡	69.1 (7.1)	70.3 (6.6)	70.5 (6.3)	72.6 (7.2)	71.9 (6.6)	74.1 (6.5)	< 0.001
Ever antihypertensive medication, % (n)	15.7 (243)	28.4 (61)	23.9 (45)	32.5 (66)	31.7 (66)	44.8 (81)	< 0.001
Ever diabetes, % (n)‡	5.4 (83)	12.6 (27)	10.0 (18)	18.2 (37)	21.6 (45)	31.5 (57)	< 0.001
Fasting blood concentration‡							
Glucose (mg/dl)	89.2 (8.4)	91.4 (10.6)	91.7 (10.4)	93.7 (12.1)	93.7 (11.7)	97.7 (15.9)	< 0.001
Insulin (μU/ml)	9.6 (3.8)	12.8 (5.8)	13.2 (4.6)	14.4 (5.6)	16.0 (6.9)	17.9 (7.9)	< 0.001

* Presented as the mean (SD), median (IQR), or % (n) as appropriate.

† Overall obesity defined as a body mass index ≥ 30 kg/m².

‡ Based upon average values during follow-up.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Diabetes defined as a self-report of oral hypoglycemic medications or insulin at examination years 2, 5, 7, 10, 15, 20, and 25; fasting glucose levels >7.0 mmol/l (126 mg/dL) at examination years 7, 10, 15, 20, or 25; a 2-h postload glucose ≥ 11.1 mmol/l (200 mg/dL) at examination years 10, 20, and 25; or a glyated hemoglobin A_{1c} $\geq 6.5\%$ at years 20 and 25.

Table 2

Mean (standard error) left ventricular mass, left ventricular end-diastolic volume, left atrial dimension, and ejection fraction according to body mass index (BMI) at year 25 and duration of overall obesity* during follow-up. The CARDIA Study (n = 2,547).

	Left ventricular mass (g)		Left ventricular end-diastolic volume (ml)		Left ventricular mass-to-volume ratio (g/ml)		Left atrial dimension (cm)		Ejection Fraction (%)	
	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2
BMI (kg/m ²)										
<18.5	146.9 (8.7)	149.5 (8.6)	115.4 (6.0)	115.2 (6.0)	1.28 (0.07)	1.30 (0.07)	3.3 (0.09)	3.3 (0.09)	69.5 (1.7)	69.7 (1.7)
18.5-24.9	164.3 (3.7)	163.6 (3.7)	127.5 (2.5)	126.7 (2.6)	1.30 (0.03)	1.31 (0.03)	3.6 (0.04)	3.6 (0.04)	68.4 (0.7)	68.3 (0.7)
25.0-29.9	179.8 (3.3)	178.3 (3.3)	136.4 (2.3)	135.5 (2.3)	1.35 (0.03)	1.35 (0.03)	3.8 (0.03)	3.8 (0.04)	69.1 (0.7)	69.0 (0.7)
30.0-34.9	177.9 (2.0)	177.6 (2.2)	129.3 (1.4)	128.7 (1.5)	1.40 (0.02)	1.40 (0.02)	3.8 (0.02)	3.8 (0.02)	69.8 (0.4)	69.8 (0.4)
35.0	186.8 (2.8)	186.2 (3.0)	136.2 (2.0)	136.0 (2.1)	1.40 (0.02)	1.40 (0.02)	3.9 (0.03)	3.9 (0.03)	70.1 (0.6)	70.0 (0.6)
p-trend	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.03	0.04
Per 5 kg/m ²	11.4 (1.2)	11.2 (1.2)	5.9 (0.8)	6.2 (0.8)	0.04 (0.01)	0.03 (0.01)	0.2 (0.01)	0.2 (0.01)	0.5 (0.2)	0.5 (0.2)
Duration of overall obesity (years)*										
0	155.3 (2.6)	156.1 (2.9)	118.5 (1.8)	117.4 (2.0)	1.33 (0.02)	1.35 (0.02)	3.5 (0.03)	3.5 (0.03)	70.1 (0.5)	70.1 (0.6)
1-5	161.9 (3.6)	162.2 (3.6)	126.6 (2.5)	126.2 (2.5)	1.30 (0.03)	1.31 (0.03)	3.6 (0.04)	3.6 (0.04)	70.0 (0.7)	69.9 (0.7)
6-10	169.9 (3.8)	170.7 (3.9)	127.6 (2.6)	127.2 (2.7)	1.34 (0.03)	1.36 (0.03)	3.7 (0.04)	3.7 (0.04)	69.7 (0.7)	69.8 (0.8)
11-15	173.7 (3.8)	172.9 (3.9)	130.8 (2.7)	130.3 (2.7)	1.34 (0.03)	1.34 (0.03)	3.7 (0.04)	3.7 (0.04)	69.4 (0.8)	69.3 (0.8)
16-20	183.9 (4.0)	183.6 (4.1)	133.9 (2.8)	133.5 (2.8)	1.40 (0.03)	1.40 (0.03)	3.7 (0.04)	3.7 (0.04)	68.2 (0.8)	68.3 (0.8)
>20	182.1 (4.4)	180.7 (4.4)	136.4 (3.0)	136.0 (3.0)	1.35 (0.03)	1.35 (0.03)	3.8 (0.05)	3.8 (0.05)	68.8 (0.9)	68.8 (0.9)
p-trend	0.003	0.003	0.17	0.11	0.32	0.46	0.23	0.23	0.05	0.07
Per 5 years	2.5 (0.8)	2.5 (0.8)	0.8 (0.6)	0.9 (0.6)	0.01 (0.01)	0.004 (0.007)	0.01 (0.01)	0.01 (0.01)	-0.3 (0.2)	-0.3 (0.2)

Model 1 simultaneously adjusts for BMI at year 25 (kg/m²) and duration of overall obesity (years) as well as age, sex, race, educational attainment, study center, current smoking, and average physical activity and alcohol intake during follow-up.

Model 2 adjusts additionally for use of antihypertensive medication, diabetes, heart rate, and average systolic blood pressure during follow-up.

* Overall obesity defined as a body mass index ≥ 30 kg/m².

Table 3

Mean (standard error) left ventricular mass, left ventricular end-diastolic volume, left atrial dimension, and ejection fraction according to waist circumference at year 25 and duration of abdominal obesity* during follow-up, The CARDIA Study (n = 2,547).

	Left ventricular mass (g)		Left ventricular end-diastolic volume (ml)		Left ventricular mass-to-volume ratio (g/ml)		Left atrial dimension (cm)		Ejection Fraction (%)	
	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2	Model 1	Model 2
Waist circumference quintiles [†]										
I	151.6 (3.3)	150.7 (3.4)	124.8 (2.3)	123.2 (2.4)	1.23 (0.03)	1.24 (0.03)	3.5 (0.03)	3.4 (0.04)	68.5 (0.7)	68.2 (0.7)
II	166.1 (3.2)	164.6 (3.3)	133.7 (2.2)	132.1 (2.3)	1.26 (0.03)	1.27 (0.03)	3.6 (0.03)	3.6 (0.04)	69.6 (0.7)	69.3 (0.7)
III	170.3 (2.9)	168.8 (3.0)	133.9 (2.0)	132.7 (2.1)	1.30 (0.02)	1.30 (0.02)	3.8 (0.03)	3.7 (0.03)	69.7 (0.6)	69.5 (0.6)
IV	175.6 (2.1)	174.8 (2.3)	131.7 (1.5)	131.2 (1.6)	1.36 (0.02)	1.36 (0.02)	3.8 (0.02)	3.8 (0.02)	69.5 (0.4)	69.3 (0.5)
V	189.0 (2.1)	189.1 (2.2)	132.8 (1.4)	133.2 (1.6)	1.44 (0.02)	1.44 (0.02)	3.9 (0.02)	3.9 (0.02)	70.0 (0.4)	70.0 (0.5)
p-trend	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	0.06	0.04
Per 10 cm	11.0 (1.0)	11.1 (1.0)	4.0 (0.7)	4.6 (0.1)	0.05 (0.008)	0.04 (0.008)	0.2 (0.01)	0.2 (0.01)	0.4 (0.2)	0.5 (0.2)
Duration of abdominal obesity (years) [*]										
0	163.5 (1.7)	163.8 (2.1)	122.2 (1.2)	121.6 (1.4)	1.36 (0.01)	1.37 (0.02)	3.6 (0.02)	3.6 (0.02)	69.7 (0.3)	69.5 (0.4)
1-5	169.9 (2.6)	169.2 (2.8)	129.4 (1.8)	128.5 (1.9)	1.33 (0.02)	1.34 (0.02)	3.7 (0.03)	3.7 (0.03)	69.7 (0.5)	69.5 (0.6)
6-10	164.9 (3.0)	164.4 (3.2)	126.9 (2.1)	126.1 (2.2)	1.32 (0.02)	1.32 (0.03)	3.7 (0.03)	3.7 (0.03)	69.7 (0.6)	69.6 (0.7)
11-15	168.3 (3.4)	167.4 (3.5)	131.7 (2.4)	130.9 (2.4)	1.29 (0.03)	1.30 (0.03)	3.7 (0.04)	3.7 (0.03)	69.3 (0.7)	69.1 (0.7)
16-20	177.7 (3.7)	175.8 (3.7)	137.4 (2.5)	136.0 (2.6)	1.33 (0.03)	1.33 (0.03)	3.8 (0.04)	3.7 (0.04)	68.3 (0.7)	68.1 (0.8)
>20	178.9 (4.5)	176.8 (4.5)	140.7 (3.1)	139.8 (3.1)	1.29 (0.04)	1.29 (0.04)	3.8 (0.05)	3.7 (0.05)	69.9 (0.9)	69.8 (0.9)
p-trend	0.13	0.19	0.03	0.04	0.38	0.32	0.90	0.94	0.15	0.17
Per 5 years	1.4 (0.9)	1.2 (0.9)	1.3 (0.6)	1.3 (0.6)	-0.006 (0.01)	-0.007 (0.01)	0.001 (0.01)	-0.001 (0.01)	-0.3 (0.1)	-0.2 (0.2)

Model 1 simultaneously adjusts for waist circumference at year 25 (cm) and duration of abdominal obesity (years) as well as age, sex, race, educational attainment, height, study center, current smoking, and average physical activity and alcohol intake during follow-up.

* Abdominal obesity defined as a waist circumference >102 cm in men and >88 cm in women.

[†] Cut-points were 86.0, 92.5, 98.3, and 105.5 cm for men and 74.3, 81.5, 89.5, and 97.5 cm for women.