

Meeting abstract

Open Access

Compartmentalization of TNF-receptor I signalling: acid sphingomyelinase is activated by Caspase-8 in internalized TNF-R1 receptosomes

B Edelman*, U Bertsch, C Hallas, V Tchikov, S Winoto-Morbach, M Jakob, S Adam and S Schütze

Address: Institut für Immunologie, UKSH Campus Kiel, Kiel, Germany

* Corresponding author

from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

Cell Communication and Signaling 2009, **7**(Suppl 1):A13 doi:10.1186/1478-811X-7-S1-A13

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A13>

© 2009 Edelman et al; licensee BioMed Central Ltd.

In a recently identified novel TNF-induced apoptotic signaling pathway the generation of ceramide by acid sphingomyelinase (A-SMase) and the activation of the aspartic protease cathepsin D (CTSD) occur in the same endosomal compartment [1]. CTSD subsequently mediates cleavage of Bid leading to activation of caspase-9 and -3. Since activation of A-SMase is linked to the death domain of TNF-receptor 1 (TNF-R1) and since the death domain adapter proteins FADD and caspase-8 are recruited during internalization of TNF-R1 in endosomes (TNF-receptosomes) [2], we addressed the question, whether A-SMase can be activated directly by caspase-8 within this compartment. We here show by confocal laser scan microscopy and in immunomagnetically isolated TNF-receptosome preparations that the active form of caspase-8 colocalizes with A-SMase within TNF-receptosomes. Activation of caspase-8 correlates with cleavage of a 70/72 kDa pro-A-SMase molecule, paralleled by enhanced A-SMase activity, activation of CTSD as well as Bid-cleavage in isolated TNF-receptosomes. The functional link between caspase-8 activity and A-SMase stimulation is revealed by the lack of TNF-induced A-SMase activation in caspase-8 deficient Jurkat cells, which can be restored in vivo after retransfection of caspase-8 as well as in vitro by the addition of exogenous caspase-8 to lysates from caspase-8 deficient cells.

References

1. Heinrich M, Neumeyer J, Jakob M, Hallas C, Tchikov V, Winoto-Morbach S, Wickel M, Schneider-Brachert W, Trauzold A, Hethke A, Schütze S: **Cathepsin D links TNF-induced acid sphingomyelinase to Bid-mediated caspase-9 and -3 activation.** *Cell Death Differ* 2004, **11**(5):550-563.
2. Schneider-Brachert W, Tchikov V, Neumeyer J, Jakob M, Winoto-Morbach S, Held-Feindt J, Heinrich M, Merkel O, Ehrenschwender M, Adam D, Mentlein R, Kabelitz D, Schütze S: **Compartmentalization of TNF receptor I signaling: internalized TNF receptosomes as death signaling vesicles.** *Immunity* 2004, **21**(3):415-428.