

# The effect of air pollution on hospitalization of individuals with respiratory and cardiovascular diseases in Jinan, China

Cai Chen, BD<sup>a</sup>, Xianfeng Wang, MS<sup>b</sup>, Chenguang Lv, MS<sup>a</sup>, Wei Li, PHD<sup>a,\*</sup>, Dedong Ma, MD<sup>c,d,\*</sup>, Qi Zhang, PHD<sup>e</sup>, Leilei Dong, BD<sup>a</sup>

## Abstract

To analyze the short-term effects of air pollution on the hospitalization rates of individuals with acute exacerbation of chronic obstructive pulmonary disease (AECOPD), stroke, and myocardial infarction (MI) after adjusting for confounding factors including weather, day of the week, holidays, and long-term trends in Jinan, China.

Hospitalization information was extracted based on data from the primary class 3-A hospitals in Jinan from 2013 to 2015. The concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, and O<sub>3</sub> were obtained from Jinan Environment Monitoring Center. The relative risk and 95% confidence intervals of AECOPD, stroke, and MI were estimated using generalized additive models with quasi-Poisson distribution in the mgcv package, using R software, version 1.0.136.

The incremental increased concentrations of particulate pollutants including PM<sub>2.5</sub> and PM<sub>10</sub> were significantly associated with increased risk of hospitalization of AECOPD, stroke, and MI, and the adverse influences of PM<sub>2.5</sub> on these diseases were generally stronger than that of PM<sub>10</sub>. The incremental increased concentrations of gaseous pollutants including SO<sub>2</sub>, NO<sub>2</sub>, and O<sub>3</sub> were significantly associated with increased risk of hospitalization of stroke and MI in this population.

Air pollution has significant adverse effects on hospitalization rates of individuals with AECOPD, stroke, and MI in Jinan, China.

**Abbreviations:** AECOPD = acute exacerbation of chronic obstructive pulmonary disease, CI = confidence interval, MI = myocardial infarction, NO<sub>2</sub> = nitrogen dioxide, O<sub>3</sub> = Ozone, RR = relative risk, SO<sub>2</sub> = sulfur dioxide.

**Keywords:** acute exacerbation of chronic obstructive pulmonary disease, ambient pollutants, generalized additive model, myocardial infarction, stroke

## 1. Introduction

Accumulating evidence suggests that air pollution is a primary risk factor for hospitalization of individuals with respiratory and cardiovascular diseases in the general population.<sup>[1–4]</sup> Acute respiratory and cardiovascular diseases, such as acute exacerbation of chronic obstructive pulmonary disease (AECOPD), stroke, and myocardial infarction (MI), have become the leading

causes of mortality worldwide. Substantial epidemiological evidence has demonstrated that ground-level ambient pollutants are associated with AECOPD, stroke, and MI in developed countries such as in Europe and North America.<sup>[5–7]</sup> However, for developing countries like China, research has been inconsistent, probably due to the limited sample sizes of the studies. Moreover, to the best of our knowledge, none of the previous

Editor: Miao Liu.

This study was approved by the ethics committee of Biomedical Engineering Institute, School of Control Science and Engineering, Shandong University. All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

This study was supported in part by grants from the National Natural Science Foundation (#21728701), the Ministry of Education Postdoctoral Fund (#2015M572044), Shandong Province Science and Technology Development Project (#GG201709260070).

The authors declare that they have no conflict of interest.

Supplemental Digital Content is available for this article. <http://links.lww.com/MD/C977>

<sup>a</sup> Biomedical Engineering Institute, School of Control Science and Engineering, Shandong University, Jinan, <sup>b</sup> Department of Ecology and Environment of the People's Republic of Shandong, <sup>c</sup> Department of Pulmonary and Critical Care Medicine Qilu Hospital Shandong University, <sup>d</sup> NHC Key Laboratory of Otorhinolaryngology (Shandong University), Jinan, PR China, <sup>e</sup> Department of Environmental Toxicology, University of California, Davis, CA.

\* Correspondence: Wei Li, Biomedical Engineering Institute, School of Control Science and Engineering, Shandong University, Jinan 250061, China (e-mail: cindy@sdu.edu.cn); Dedong Ma, Department of Pulmonary and Critical Care Medicine Qilu Hospital Shandong University, NHC Key Laboratory of Otorhinolaryngology (Shandong University), China (e-mail: mdd@sdu.edu.cn).

Copyright © 2019 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Medicine (2019) 98:22(e15634)

Received: 15 October 2018 / Received in final form: 29 March 2019 / Accepted: 18 April 2019

<http://dx.doi.org/10.1097/MD.00000000000015634>

studies simultaneously evaluated the influence of ground-level ambient pollutants on the hospitalization rate of AECOPD, stroke, and MI. Therefore, large epidemiologic studies are needed to determine the effects of ground-level ambient pollutants on hospitalization rates of individuals with AECOPD, stroke, and MI in China.

Jinan, the capital city of Shandong Province in China, has been reported to be one of the most polluted cities in China in recent years. Overall, Jinan is a northern inland city with complex natural terrain and poor airflow and ventilation, which ranks it among the top 10 cities with the worst air quality in China. The characteristic terrain in Jinan includes higher elevations in the southern area and lower elevations in the northern area, surrounded by mountains that hinder the management and diffusion of ambient pollutants. Accordingly, the objective of this study is to explore the association between the air pollution and hospitalization rates of individuals with AECOPD, stroke, and MI in Jinan from 2013 to 2015. Results of our study are expected to clarify the influence of ground-level ambient pollutants on hospitalizations attributed to acute respiratory and cardiovascular diseases in China, and to provide pertinent advice and guidance for the prevention of acute respiratory and cardiovascular diseases.

## 2. Materials

### 2.1. Study population

Information on 6981 AECOPD, 56,922 stroke, and 11,583 MI patients, including age, gender, date of admission, date of discharge, clinical symptoms, disease classification, workplace, and current residence, were obtained from Jinan Qilu Hospital, the Provincial Hospital of Shandong Province, and the Central Hospital of Shandong Province database, respectively. Patients who did not reside or work in Jinan were excluded from the present study. According to epidemiological studies about AECOPD, stroke, and MI in China,<sup>[8–10]</sup> patients who met the following criteria were included:

- (1)  $\geq 18$  years of age;
- (2) resided and worked in the study area (Jinan City) during study period.

The exclusion criteria were as follows:

- (1)  $< 18$  years old;
- (2) not residing and working in Jinan during the study period;
- (3) patients with duplicate records;
- (4) more than 1 patient admission per week.

The study was approved by the ethic committees of the 3 hospitals.

### 2.2. Data source

Data reflecting the daily average concentrations of fine particles (PM<sub>2.5</sub>), inhalable particles (PM<sub>10</sub>), sulfur dioxide (SO<sub>2</sub>), nitrogen dioxide (NO<sub>2</sub>) for 24-hours, and Ozone (O<sub>3</sub>) for 8-hours, acquired from the 14 fix-sited monitoring stations in urban areas of Jinan, during 2013 to 2015 were obtained from Jinan Environment Monitoring Center. The expectation-maximization method was used to impute missing data. Daily average air temperatures and relative humidity in the corresponding period were obtained from the Jinan Bureau of Meteorology.

### 2.3. Methods

The generalized additive model with the link function of Poisson distribution was established to assess the impact of air pollution on the hospitalization rates of individuals with AECOPD, stroke, and MI. Because of the application of link function of Poisson distribution and the emergence of extreme values in the data, the quasi-Poisson regression model was applied to reduce the influence of over-dispersion of the results.<sup>[11]</sup> Confounding factors such as day of the week, holidays, and long-term trends were included as dummy variables and subsequently adjusted.

### 2.4. Degree of freedom of basic model

The basic models of AECOPD, stroke, and MI were as follows

For AECOPD:

$$\text{Log } [E(Y_t)] = \alpha + \beta_1 C_i + \text{ns (Temperature, df = 3)} + \text{ns (Humidity, df = 5)} + \text{ns (Time, df = 6*3)} + \beta_2 \text{ factor (DOW)} + \beta_3 \text{ factor (Holiday)}$$

For stroke:

$$\text{Log } [E(Y_t)] = \alpha + \beta_1 C_i + \text{ns (Temperature, df = 4)} + \text{ns (Humidity, df = 4)} + \text{ns (Time, df = 8*3)} + \beta_2 \text{ factor (DOW)} + \beta_3 \text{ factor (Holiday)}$$

df represents degree freedom

For MI:

$$\text{Log } [E(Y_t)] = \alpha + \beta_1 C_i + \text{ns (Temperature, df = 3)} + \text{ns (Humidity, df = 6)} + \text{ns (Time, df = 7*3)} + \beta_2 \text{ factor (DOW)} + \beta_3 \text{ factor (Holiday)}$$

$Y_t$  represents the number of hospitalized patients on  $t$  days;  $E(Y_t)$  represents the expected number of hospitalized patients on  $t$  days;  $\beta_1$  represents the daily average concentration coefficient of pollutants on  $t$  days or lag  $t$  days;  $C_i$  represents the daily average concentration of pollutants on  $t$  days or lag  $t$  days; ns represents a natural spline smooth function; Time represents the long-term trend of the date; the dummy variables DOW and Holiday represent the effect of “day of the week” and the statutory holidays, respectively, and the  $\beta_2$  and  $\beta_3$  represent the coefficients of factor variables DOW and Holiday, respectively.

The basic model without pollutants was established, and the natural spline smooth function of time, daily temperature, and relative humidity were included to fit the nonlinear effects of the model. The degree of freedom for time, daily temperature, and relative humidity were determined according to the akaike information criterion.<sup>[12]</sup>

The single-pollutant composed of basic model and daily concentrations of major ambient pollutants were established to estimate the relative risk (RR) and 95% confidence interval for the associations between major ambient pollutants with an increment of 10  $\mu\text{g}/\text{m}^3$  and the risk of hospitalizations of the above diseases. A  $P < .05$  was considered statistically significant. Lag structures (from lag 0 day to lag 7 day) were defined as lag 0 to lag 7, and the multi-day moving averages lag structures (from lag 0–1 day [average] to lag 0–7 day [average]) were defined as lag 01 to lag 07. The time lag and cumulative time lag effects of major ambient pollutants were included in the model.

The 2-pollutant models of pollutants and O<sub>3</sub> with multi-day moving averages lag structures (from lag 0–1 day [average] to lag 0–7 day [average]) were established for sensitive analysis and to confirm model stability.

The stratification analyses of pollutants exposure based on gender (male or female) and age ( $< 65$  years and  $\geq 65$  years) were conducted to explore the influence of age and gender on the

**Table 1**  
**Distribution of daily data on airborne pollutants and weather parameters in Jinan, 2013–2015.**

Variable	$X \pm S$	Min	$P_{25}$	$P_{50}$	$P_{75}$	Max	IQR
Pollutants							
PM <sub>2.5</sub> , $\mu\text{g}/\text{m}^3$	94 ± 56	21	57	80	113	417	56
PM <sub>10</sub> , $\mu\text{g}/\text{m}^3$	189 ± 86	33	130	171	228	666	98
SO <sub>2</sub> , $\mu\text{g}/\text{m}^3$	79 ± 54	12	42	61	99	409	57
NO <sub>2</sub> , $\mu\text{g}/\text{m}^3$	57 ± 23	19	40	52	69	165	29
O <sub>3</sub> , $\mu\text{g}/\text{m}^3$	104 ± 57	11	54	96	150	261	96
Meteorological data							
Temperature, °C	15.2 ± 10.3	-9.4	5.8	16.6	24.1	33.7	18.3
Humidity, %	56 ± 20	15	41	55	70	100	29

IQR=interquartile range, Max=maximum value, Min=minimum value, S=standard deviation, X=mean value.

associations between air pollution and hospitalization rates of individuals with AECOPD, stroke, and MI.

### 2.5. Spearman correlation analysis

Spearman correlation was applied to describe the correlation of daily concentrations of air pollutants, temperature, and relative humidity.

## 3. Results

### 3.1. Distribution of ambient pollutants and weather data

The annual average concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, and O<sub>3</sub> were 94  $\mu\text{g}/\text{m}^3$ , 188  $\mu\text{g}/\text{m}^3$ , 79  $\mu\text{g}/\text{m}^3$ , 57  $\mu\text{g}/\text{m}^3$ , and 104  $\mu\text{g}/\text{m}^3$ , respectively, during 2013 to 2015, which were 2.7-, 2.7-, 1.3-, 1.4-, and 0.7-fold greater than the Annual Secondary National Ambient Air Quality Standards (GB3095-2013). The distribution of daily data on ambient pollutants and weather parameters is shown in Table 1. The distribution of daily ambient pollutant concentrations and temperature is shown in Figure 1.

### 3.2. Data description

Table 2 shows the demographic characteristic of patients with respiratory and cardiovascular diseases. A total of 6981 AECOPD hospitalizations (4920 males and 2061 females), 56,922 stroke hospitalizations (36,516 males and 20,406 females), and 11,583 MI hospitalizations (8085 males and 3498 females), were obtained from the databases of the primary class 3-A hospitals in Jinan from 2013 to 2015. The percentage of patients aged <65 years was 44.4% (33,492/75,486), while that of patients aged ≥65 years was 55.6% (41,994/75,486).

### 3.3. Spearman correlation analysis

Table 3 shows the Spearman correlation coefficients among daily concentrations of air pollutants, temperature, and relative humidity. Positive correlations were detected among PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, and NO<sub>2</sub>, while there was a significant negative association between these pollutants and O<sub>3</sub>, temperature, and relative humidity. The Spearman correlations between PM<sub>2.5</sub> and PM<sub>10</sub> ( $r=0.802$ ), SO<sub>2</sub> and NO<sub>2</sub> ( $r=0.783$ ), PM<sub>2.5</sub> and NO<sub>2</sub> ( $r=0.694$ ), and O<sub>3</sub> and temperature ( $r=0.840$ ) were statistically significant ( $P < .01$ ).

### 3.4. Daily hospitalizations: AECOPD, stroke, and MI

**3.4.1. AECOPD.** The strongest associations for PM<sub>2.5</sub>, PM<sub>10</sub>, and O<sub>3</sub> and the risk of AECOPD hospitalizations were observed on lag 3, lag 2, and lag 3 with an increment of 3.1%

(1.017–1.044), 1.6% (1.070–1.025), and 2.8% (1.026–1.030), respectively (Table 4). For moving averages lag structures, the strongest associations for PM<sub>2.5</sub>, PM<sub>10</sub>, and O<sub>3</sub> and the risk of AECOPD hospitalizations were observed in lag 03, lag 02, and lag 03 with an increment of 2.9% (1.018–1.040), 2.0% (1.009–1.031), and 3.5% (1.031–1.039), respectively (Table 5).

Stratified analysis based on gender and age showed the impact of concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, and O<sub>3</sub> on the hospitalization risk of AECOPD was stronger for males compared to females and stronger for older participants (≥65 years) compared to younger participants (Fig. 2).


### 3.5. Stroke

The strongest associations for PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, and NO<sub>2</sub> and the risk of stroke hospitalizations were all observed on lag 0 with an increment of 1.1% (1.010–1.012), 0.6% (1.005–1.007), 1.8% (1.017–1.020), and 2.5% (1.023–1.027), respectively. The association of O<sub>3</sub> and the risk of stroke hospitalizations were statistically significant with an increment 0.2% (1.001–1.003) (Table 6). In addition, the strongest association for O<sub>3</sub> was observed on lag 1 with an increment of 0.5% (1.004–1.006) (Table 6). For moving averages lag structures, the strongest associations for PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, and O<sub>3</sub> and the risk of stroke hospitalizations were all observed on lag 01 with an increment of 0.8% (1.007–1.009), 0.4% (1.003–1.005), 2.2% (1.020–1.023), 1.9% (1.016–1.021), and 0.7% (1.005–1.008) respectively (Table 7).

Stratified analysis based on gender and age showed that the impact of concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, and O<sub>3</sub> on stroke hospitalization was significantly stronger in patients <65 years of age compared to patients ≥65 years of age, while the impact of concentrations of SO<sub>2</sub> and NO<sub>2</sub> on stroke hospitalization was significantly stronger in patients ≥65 years of age compared to patients <65 years of age. Stratified analysis based on gender showed that the impact of concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, and SO<sub>2</sub> on stroke hospitalization was significantly stronger in males, while the impact of concentrations of NO<sub>2</sub> and O<sub>3</sub> on stroke hospitalization was significantly stronger in females (Fig. 3).

### 3.6. MI

The strongest associations between PM<sub>2.5</sub> and PM<sub>10</sub> and the risk of MI hospitalizations were both observed in lag 0 with an increment of 1.8% (1.009–1.028) and 4.7% (1.041–1.052), respectively. In addition, the strongest associations for SO<sub>2</sub> and O<sub>3</sub> and the risk of MI hospitalizations were both observed in lag 1



**Figure 1.** The temporal distribution of daily average ambient pollutant concentrations and daily mean temperature in Jinan during 2013 to 2015.

with an increment of 4.8% (1.035–1.060) and 0.8% (1.007–1.009) (Table 8). For moving averages lag structures, the strongest associations for PM<sub>10</sub>, SO<sub>2</sub>, and O<sub>3</sub> and the risk of MI hospitalizations were all observed in lag 01 with an increment of 0.3% (1.002–1.004), 3.7% (1.022–1.050), and 0.8% (1.006–1.010), respectively (Table 9).

Stratified analysis based on gender and age showed that the impact of concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, and O<sub>3</sub> and MI hospitalizations in females was stronger compared to males, and stronger in older participants ( $\geq 65$  years) compared to younger (<65 years) participants (Fig. 4).

### 3.7. Sensitive analysis

Sensitive analysis showed no significant changes in RR due to concentrations of airborne particulates for AECOPD, stroke, and MI hospitalizations after inclusion of O<sub>3</sub> in multi-day moving averages lag structures, which indicated that the effect of the single-pollutant model was robust (Figs. 5–7).

## 4. Discussion

In this epidemiological study from Jinan, China, we found that air pollution has significant adverse effects on the hospitalization rates of individuals with AECOPD, stroke, and MI. These results

**Table 2**

**Demographic characteristics of patients with respiratory and cardiovascular diseases in Jinan, 2013–2015.**

Variable		AECOPD	Stroke	MI
Total		6981	56922	11583
Gender	Male	4920	36516	8085
	Female	2061	20406	3498
Age ( $X \pm S$ )	<65	58 $\pm$ 6	55 $\pm$ 8	54 $\pm$ 8
	$\geq 65$	77 $\pm$ 7	76 $\pm$ 8	75 $\pm$ 7

AECOPD = acute exacerbation of chronic obstructive pulmonary disease, MI = myocardial infarction, S = standard deviation, X = mean value.

**Table 3****Spearman correlation coefficients among daily concentrations of air pollutants and meteorological data in Jinan, 2013–2015.**

Variable	PM <sub>2.5</sub>	PM <sub>10</sub>	SO <sub>2</sub>	NO <sub>2</sub>	O <sub>3</sub>	Temp	Hum
PM <sub>2.5</sub>	1.000	0.802**	0.554**	0.649**	−0.222**	−0.207**	0.216**
PM <sub>10</sub>	–	1.000	0.638**	0.694**	−0.253**	−0.288**	−0.104**
SO <sub>2</sub>	–	–	1.000	0.783**	−0.495**	−0.598**	−0.223**
NO <sub>2</sub>	–	–	–	1.000	−0.555**	−0.580**	−0.062*
O <sub>3</sub>	–	–	–	–	1.000	0.840**	−0.092**
Temp	–	–	–	–	–	1.000	0.156**
Hum	–	–	–	–	–	–	1.000

\*  $P < .05$ .\*\*  $P < .01$ .**Table 4****Relative risk (RR) with 95% confidence interval (CI) for the association between airborne pollutants concentrations and AECOPD hospitalizations in different lag structures in Jinan, 2013–2015.**

Variables	Lag 0	Lag 1	Lag 2	Lag 3
PM <sub>2.5</sub>				
RR	0.989	0.986	1.025*	1.031*
95% CI	0.973–1.005	0.971–1.001	1.011–1.039	1.017–1.044
PM <sub>10</sub>				
RR	0.994	0.996	1.016*	0.996
95% CI	0.987–1.001	0.983–1.008	1.070–1.025	0.997–0.998
SO <sub>2</sub>				
RR	1.018	0.987	0.988	0.984
95% CI	0.997–1.039	0.966–1.008	0.965–1.011	0.959–1.009
NO <sub>2</sub>				
RR	0.981	0.973	1.020	0.980
95% CI	0.940–1.022	0.932–1.014	0.998–1.042	0.937–1.023
O <sub>3</sub>				
RR	0.994	1.005	1.015*	1.028*
95% CI	0.985–1.005	1.002–1.007	1.012–1.017	1.026–1.030

AECOPD = acute exacerbation of chronic obstructive pulmonary disease, CI = confidence interval, RR = relative risk.

\*  $P < .05$ .**Table 5****Relative risk (RR) with 95% confidence interval (CI) for the association between airborne pollutants concentrations and AECOPD hospitalizations in different moving averages lag structures for in Jinan, 2013–2015.**

Variables	Lag 01	Lag 02	Lag 03
PM <sub>2.5</sub>			
RR	0.982	1.023*	1.029*
95% CI	0.961–1.003	1.010–1.036	1.018–1.040
PM <sub>10</sub>			
RR	0.994	1.020*	0.993
95% CI	0.978–1.010	1.009–1.031	0.985–1.001
SO <sub>2</sub>			
RR	0.993	0.986	0.975
95% CI	0.982–1.004	0.958–1.014	0.935–1.015
NO <sub>2</sub>			
RR	0.970	0.975	0.962
95% CI	0.958–0.982	0.952–1.017	0.911–1.013
O <sub>3</sub>			
RR	1.000	1.015*	1.035*
95% CI	0.997–1.004	1.011–1.018	1.031–1.039

AECOPD = acute exacerbation of chronic obstructive pulmonary disease, CI = confidence interval, RR = relative risk.


\*  $P < .05$ .

suggest that air pollution has become a major contributor to the incidence and exacerbation of acute respiratory and cardiovascular diseases in developing countries. These results highlight the importance of management of air pollution for the prevention of acute respiratory and cardiovascular diseases.

#### 4.1. AECOPD

Our findings are consistent with previous reports. Xu et al included patients with respiratory disease who visited emergency hospitals in Beijing and showed that the severest adverse effect of PM<sub>2.5</sub> on the hospitalization of patients with AECOPD was the





**Figure 2.** A stratified analysis of the effects of air pollutant concentrations at optimal lag on admission risk in patients with acute exacerbation of chronic obstructive pulmonary disease.

cumulative lag 0 to 3 (Lag 03) day with an increment of 3.15% in the emergency hospital admission rate of AECOPD.<sup>[13]</sup> Ko et al also found significant associations between hospital admissions for COPD with PM<sub>2.5</sub>; the RR for admission for every 10 mg/m<sup>3</sup> increase in PM<sub>2.5</sub> was 1.031, at a lag day ranging from lag 0 to cumulative lag 0–5.<sup>[14]</sup>

In our study, although no immediate effects were observed between PM<sub>2.5</sub>, PM<sub>10</sub>, O<sub>3</sub>, and hospitalizations of AECOPD, the effect on different lag days (lag 2, lag 3, lag 02, and lag 03) were statistically significant. As highly active oxidants with poor solubility, these airborne particulates can reach the depths of the

lungs after being inhaled and subsequently cause inflammation of the epithelial cells of the respiratory tract. The certain reaction time of airway inflammation caused by oxidants could result in a cumulative effect, rather than an immediate effect<sup>[15]</sup> as previously indicated.

As a significant risk factor for AECOPD, SO<sub>2</sub> dissolves easily in the upper respiratory tract and produces immediate stimulation to the mucosa.<sup>[16,17]</sup> However, the significant effects of SO<sub>2</sub> on hospitalization risk in patients with AECOPD were not observed in this study. The annual average concentrations of SO<sub>2</sub> in Jinan during 2013 to 2015 were 102 μg/m<sup>3</sup>, 79 μg/m<sup>3</sup>, and

**Table 6**

**Relative risk (RR) with 95% confidence interval (CI) for the association between airborne pollutants concentrations and stroke hospitalizations in different lag structures for in Jinan, 2013–2015.**

Variables	Lag 0	Lag 1	Lag 2	Lag 3
PM <sub>2.5</sub>				
RR	1.011*	1.001*	0.998	0.993
95% CI	1.010–1.012	1.000–1.002	0.995–1.001	0.984–1.002
PM <sub>10</sub>				
RR	1.006*	1.001*	0.997	0.996
95% CI	1.005–1.007	1.000–1.002	0.994–1.000	0.991–1.001
SO <sub>2</sub>				
RR	1.018*	1.014*	1.001	0.993
95% CI	1.017–1.020	1.013–1.016	1.000–1.002	0.986–1.000
NO <sub>2</sub>				
RR	1.025*	1.002*	0.986	0.984
95% CI	1.023–1.027	1.000–1.004	0.971–1.001	0.968–1.000
O <sub>3</sub>				
RR	1.002*	1.005*	1.002*	0.997
95% CI	1.001–1.003	1.004–1.006	1.001–1.003	0.993–1.001

CI=confidence interval, RR=relative risk.

\*  $P < .05$ .

**Table 7**  
**Relative risk (RR) with 95% confidence interval (CI) for the association between airborne pollutants concentrations and stroke hospitalizations in different moving averages lag structures for in Jinan, 2013–2015.**


Variables	Lag 01	Lag 02	Lag 03
PM <sub>2.5</sub>			
RR	1.008*	1.005*	1.000
95% CI	1.007–1.009	1.004–1.006	0.999–1.001
PM <sub>10</sub>			
RR	1.004*	1.002*	0.999
95% CI	1.003–1.005	1.001–1.003	0.998–1.000
SO <sub>2</sub>			
RR	1.022*	1.019*	1.014*
95% CI	1.020–1.023	1.018–1.021	1.012–1.016
NO <sub>2</sub>			
RR	1.019*	1.007*	0.996
95% CI	1.016–1.021	1.004–1.009	0.990–1.002
O <sub>3</sub>			
RR	1.007*	1.006*	1.002*
95% CI	1.005–1.008	1.005–1.008	1.001–1.004

CI=confidence interval, RR=relative risk.  
 \*P<.05.

55 µg/m<sup>3</sup>, respectively, and the average annual concentration in 2015 was below the 2-level standard limit (60 µg/m<sup>3</sup>) of the national ambient air quality standard (GB 3095-2012) for the first time. The control of coal-fired volume and the improvement of technical treatment methods, such as desulphurization and denitrification, during the “13th 5-Year Plan” period in Jinan were highly effective.

The results of stratified analysis in gender and age showed that male patients aged ≥65 years were more sensitive to air pollutants and were at a consequently higher risk of hospitalization for AECOPD compared to women aged <65 years. It has been shown that the smoking rate is 74% among men aged >35 years

in China. As China is one of the largest tobacco producing and consuming countries, the prevalence of chronic bronchitis among smokers is 2- to 8-fold higher than that of non-smokers. Meanwhile, the proportion of smoking in males has been significantly higher than that of females in underdeveloped and developing countries.<sup>[16]</sup> Moreover, a previous report revealed that male patients with a history of smoking had a higher hospital admission risk and were more sensitive to varied concentrations of pollutants.<sup>[18]</sup> Stratified analysis by age showed that patients ≥65 years of age were more sensitive to air pollutants exposure and had higher risk of hospital admission for AECOPD compared to patients <65 years of age. The weakness of the



**Figure 3.** Stratified analysis of the effects of air pollutants concentration on the risk of admission in patients with stroke.

**Table 8**  
**Relative risk (RR) with 95% confidence interval (CI) for the association between airborne pollutants concentrations and MI hospitalizations in different lag structures for in Jinan, 2013–2015.**

Variables	Lag 0	Lag 1	Lag 2	Lag 3
PM <sub>2.5</sub>				
RR	1.018*	1.004	0.994	0.997
95% CI	1.009–1.028	0.995–1.013	0.987–1.001	0.992–1.002
PM <sub>10</sub>				
RR	1.047*	1.003	0.997	1.002
95% CI	1.041–1.052	0.997–1.008	0.994–1.000	0.997–1.008
SO <sub>2</sub>				
RR	0.999	1.048*	0.998	0.992
95% CI	0.986–1.012	1.035–1.060	0.996–1.000	0.981–1.003
NO <sub>2</sub>				
RR	0.988	0.988	0.981	0.989
95% CI	0.968–1.008	0.976–1.000	0.960–1.002	0.977–1.001
O <sub>3</sub>				
RR	1.002	1.008*	0.999	0.995
95% CI	0.999–1.007	1.007–1.009	0.998–1.000	0.990–1.001

CI=confidence interval, MI=myocardial infarction, RR=relative risk.

\*  $P < .05$ .

immune system, the higher prevalence of chronic respiratory diseases, and the increased sensitivities to the particles in the elderly could be the underlying causes for the stronger association between air pollution and risk of AECOPD hospitalization in those older than 65 years of age.<sup>[19–21]</sup>

#### 4.2. Stroke

The strongest associations between particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>) and hospitalizations of stroke were found on the day of admission with an increment of 1.1% and 0.6%, respectively. Our results were supplemented by experimental studies that indicated that exposure to high concentrations of PM<sub>2.5</sub> can lead to pathophysiological changes related to the onset of cardiovascular diseases, including increased blood pressure and heart rate, reduced heart rate variability within 12 hours, changes in coagulation factors, systemic inflammation, damaged endothelial cells, and vascular dysfunction.<sup>[22]</sup> Our study also showed that

the impact of PM<sub>2.5</sub> on the admission risk of stroke patients seemed to be stronger than PM<sub>10</sub>. This is consistent with a previous meta-analysis that also indicated that stroke hospitalization increased by 1.011 (95% CI=1.010–1.012) and 1.002 (95% CI=1.000–1.003), following exposure to PM<sub>2.5</sub> and PM<sub>10</sub>, respectively.<sup>[23]</sup> Based on previous reports, it has been estimated that fine particles, ultrafine particles, and gaseous pollutants primarily cause additional systemic cerebrovascular responses,<sup>[24]</sup> compared to partial lung effects which are usually caused by particulate matter with larger aerodynamics. Moreover, particles with smaller aerodynamics can penetrate the alveolar space of the lungs, and even circulate systemically to reach other organs.<sup>[25]</sup>

Our results showed that daily exposure to concentrations of SO<sub>2</sub> and NO<sub>2</sub> may have the strongest impact on the risk of hospitalizations of stroke. A previous study by Joubert et al demonstrated that SO<sub>2</sub> can penetrate the protective blood–brain barrier via its own biochemical pathway to elicit specific

**Table 9**  
**Relative risk (RR) with 95% confidence interval (CI) for the association between airborne pollutants concentrations and MI hospitalizations in different moving averages lag structures for in Jinan, 2013–2015.**

Variables	Lag 01	Lag 02	Lag 03
PM <sub>2.5</sub>			
RR	0.998	0.997	0.996
95% CI	0.996–1.000	0.993–1.001	0.991–1.001
PM <sub>10</sub>			
RR	1.003*	1.001*	1.001*
95% CI	1.002–1.004	1.000–1.002	1.000–1.002
SO <sub>2</sub>			
RR	1.037*	1.020*	0.998
95% CI	1.022–1.050	1.003–1.040	0.996–1.000
NO <sub>2</sub>			
RR	0.994	0.998	0.998
95% CI	0.987–1.001	0.990–1.006	0.993–1.003
O <sub>3</sub>			
RR	1.008*	1.005*	1.000
95% CI	1.006–1.010	1.003–1.007	0.999–1.002

CI=confidence interval, MI=myocardial infarction, RR=relative risk.

\*  $P < .05$ .






Figure 4. Stratified analysis of the effects of air pollutants concentration on the risk of admission in patients with myocardial infarction.

pathophysiological effects and cause cerebral ischemia.<sup>[26]</sup> Results of another meta-analysis also showed that hospitalization for ischemic stroke increased by 1.79% (95% CI=1.0054–1.0306) with an incremental increase of SO<sub>2</sub>, which was in accordance with our findings.<sup>[20]</sup> In addition, epidemiological studies from Denmark, Italy, Canada, and other countries consistently showed that variations in NO<sub>2</sub> concentration were

significantly associated with the incidence of acute ischemic stroke.<sup>[27–29]</sup> All of the above evidence supports a strong association between air pollution and the risk of stroke.

Moreover, we found that the association between SO<sub>2</sub> (RR=1.018; 95% CI=1.017–1.020) and NO<sub>2</sub> (RR=1.014; 95% CI=1.013–1.016) and stroke hospitalizations were stronger than the association between PM<sub>2.5</sub> and stroke admissions (RR=1.011;




Figure 5. Comparison of cumulative lag effect between single pollutant model and sensitivity analysis model for chronic obstructive pulmonary disease.




Figure 6. Comparison of cumulative lag effect between single pollutant model and sensitivity analysis model for stroke.

95% CI=1.010–1.012) and PM<sub>10</sub> (RR=1.006; 95% CI=1.005–1.007). This finding is consistent with previous studies, which suggested that traffic pollution may cause ischemic stroke.<sup>[30–32]</sup> While the potential reasons for this association remains unclear, we

hypothesize that gaseous pollutants exhausted from traffic in Jinan may have become one of the main sources of air pollutants.

Stratified analysis by age showed that stroke patients ≥65 years of age were more sensitive to gaseous pollutants (SO<sub>2</sub> and NO<sub>2</sub>)




Figure 7. Comparison of cumulative lag effect between single pollutant model and sensitivity analysis model for myocardial infarction.

exposure and were at higher risk of stroke-related hospitalizations. Similar results were supported by a previous case-crossover study from Beijing, which also showed that the increased daily concentrations of SO<sub>2</sub> and NO<sub>2</sub> were associated with a more significant increased risk of stroke hospitalizations of patients ≥65 years of age.<sup>[33]</sup> Compared with the younger participants, the elderly are more sensitive to air pollution and have higher incidences of cardiovascular and cerebrovascular diseases. These findings are also supported by previous studies that showed that recurrent attacks of stroke were more frequent in elderly patients.<sup>[34]</sup>

Stratified analysis by gender showed that female patients were more sensitive to gaseous pollutants (NO<sub>2</sub> and O<sub>3</sub>) exposure, while male patients were more sensitive to particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>) exposure. Qin et al collected admission data of stroke patients from 3 cities in northern China in a case-crossover study and found that the significant association between air pollution and stroke hospitalization was only observed in females.<sup>[35]</sup> Moreover, a study in Beijing demonstrated that the impacts of PM<sub>2.5</sub> on the first admission due to ischemic stroke did not differ by gender.<sup>[36]</sup> In addition, a study from Shanghai showed that male stroke patients were more sensitive to changes in concentrations of PM<sub>10</sub>, SO<sub>2</sub>, and NO<sub>2</sub> than female patients.<sup>[37]</sup> In summary, the potential gender differences underlying the association between air pollution and stroke remain to be determined.

#### 4.3. MI

According to the results of this study, SO<sub>2</sub> has the strongest adverse impact on the admission of individuals with MI (RR=1.048; 95% CI=1.035–1.060), which is consistent with the result of a similar study performed in Shanghai Pudong District during 2013 to 2014.<sup>[37]</sup> This study showed that the incidence of acute MI increased by 5% with an incremental 10 μg/m<sup>3</sup> increase in SO<sub>2</sub>.<sup>[36]</sup>

However, we failed to detect a significant effect of NO<sub>2</sub> on admission rate of MI. Compared to the 4-quartile range of other pollutants (PM<sub>2.5</sub>=56 μg/m<sup>3</sup>, PM<sub>10</sub>=98 μg/m<sup>3</sup>, SO<sub>2</sub>=57 μg/m<sup>3</sup>, and O<sub>3</sub>=96 μg/m<sup>3</sup>), the quartile range of NO<sub>2</sub> only reached 29 μg/m<sup>3</sup>. The small variance in daily average concentrations of NO<sub>2</sub> in Jinan might be the potential reason for the insignificant association between NO<sub>2</sub> and admission rate of MI.

Most previous studies have shown that there is no significant association between short-term exposure to O<sub>3</sub> and morbidity and mortality of MI.<sup>[13,14,38,39]</sup> In contrast to previous studies performed in Western countries, our study demonstrated a 0.8% increased risk of hospitalizations for MI per 10 μg/m<sup>3</sup> increase in O<sub>3</sub> concentration on lag 1 and a 0.8%, 0.5% increment on lag 01 and lag 02, respectively. The relatively high concentrations of O<sub>3</sub> in Jinan could be one of the potential reasons underlying the variations between our results and the previous findings.

Although the inhalable particles (PM<sub>2.5</sub> and PM<sub>10</sub>) and O<sub>3</sub> exposure could induce inflammatory effects, the results of this study showed that the adverse effects of particulate matter (especially for PM<sub>10</sub> [RR=1.047; 95% CI=1.041–1.052]) were stronger than that of O<sub>3</sub> (RR=1.008; 95% CI=1.007–1.009). This is consistent with the study conducted by Rosenthal et al, who showed that, compared to O<sub>3</sub>, PM<sub>10</sub> plays a greater role in thrombosis.<sup>[7]</sup>

The results of the stratified analysis showed that female patients ≥65 years of age were more sensitive to air pollutants

exposure and suffered a higher risk of MI admissions than males <65 years of age. As a vulnerable group, the elderly often have atherosclerotic plaques and other cardiovascular diseases and are therefore more susceptible to the risks of air pollution exposure. Exposure to short-term air pollution is an important trigger of cardiovascular diseases, probably due to its ability to increase plaque vulnerability, platelet activation, and coagulation.<sup>[11]</sup> Some studies suggested that many factors may underlying the potential gender differences regarding the association between air pollution and MI, such secretion of hormones, the physical build, and the range of activities<sup>[36]</sup> The exact mechanisms deserve further investigation.

The limitation of our study was that we did not include patients <18 years of age.

## 5. Conclusions

In conclusion, results of our study indicate that air pollution is significantly associated with an increased risk of hospitalization of individuals with AECOPD, stroke, and MI in Jinan, China. Future studies are needed to determine the potential mechanisms involved and explore the potential importance of management of air pollution for the prevention of acute respiratory and cardiovascular diseases.

## Author contributions

**Conceptualization:** Xianfeng Wang, Wei Li, Qi Zhang.

**Data curation:** Chenguang Lv, Wei Li.

**Formal analysis:** Cai Chen, Chenguang Lv, Wei Li.

**Funding acquisition:** Cai Chen, Chenguang Lv, Leilei Dong.

**Investigation:** Cai Chen, Dedong Ma, Leilei Dong.

**Methodology:** Cai Chen, Xianfeng Wang, Dedong Ma, Leilei Dong.

**Project administration:** Xianfeng Wang, Leilei Dong.

**Resources:** Xianfeng Wang, Dedong Ma, Qi Zhang.

**Software:** Xianfeng Wang, Qi Zhang.

**Supervision:** Xianfeng Wang, Qi Zhang.

**Validation:** Dedong Ma, Qi Zhang.

**Visualization:** Qi Zhang.

Wei Li orcid: 0000-0001-7367-4178.

## References

- [1] Villeneuve PJ, Johnson JY, Pasichnyk D, et al. Short-term effects of ambient air pollution on stroke: who is most vulnerable? *Sci Total Environ* 2012;430:193–201.
- [2] Belleudi V, Faustini A, Stafoggia M, et al. Impact of fine and ultrafine particles on emergency hospital admissions for cardiac and respiratory diseases. *Epidemiology* 2010;21:414–23.
- [3] R Sciaraffa AB, Montuschi P, Gerosa GA, et al. Impact of air pollution on respiratory diseases in urban areas: a systematic review: Daniele Ignazio La Milia. *Eur J Public Health* 2017;27.
- [4] Mishra S. Is smog innocuous? Air pollution and cardiovascular disease. *Indian Heart J* 2017;69:425–9.
- [5] Mantovani KC, Nascimento LF, Moreira DS, et al. Air pollutants and hospital admissions due to cardiovascular diseases in Sao Jose do Rio Preto, Brazil. *Cien Saude Colet* 2016;21:509–15.
- [6] Lee BJ, Kim B, Lee K. Air pollution exposure and cardiovascular disease. *Toxicol Res* 2014;30:71–5.
- [7] Rosenthal FS, Kuusma M, Lanki T, et al. Association of ozone and particulate air pollution with out-of-hospital cardiac arrest in Helsinki, Finland: evidence for two different etiologies. *J Expo Sci Environ Epidemiol* 2013;23:281–8.
- [8] Xu T, Li W, Teo K, et al. Association of psychological risk factors and acute myocardial infarction in China: the INTER-HEART China study. *Chinese Med J-Peking* 2011;124:2083–8.

- [9] Wang W, Jiang B, Sun H, et al. Prevalence, incidence, and mortality of stroke in China. *Circulation* 2017;135:759.
- [10] Chan KY, Li X, Chen W, et al. Prevalence of chronic obstructive pulmonary disease (COPD) in China in 1990 and 2010. *J Glob Health* 2017;7.
- [11] Gardner B, Ling F, Hopke PK, et al. Ambient fine particulate air pollution triggers ST-elevation myocardial infarction, but not non-ST elevation myocardial infarction: a case-crossover study. Part Fibre Toxicol 2014;11:1.
- [12] Hurvich CM, Simonoff JS, Tsai CL. Smoothing parameter selection in nonparametric regression using an improved Akaike information criterion. *J R Stat Soc* 1998;60:271–93.
- [13] Xu Q, Li X, Wang S, et al. Fine particulate air pollution and hospital emergency room visits for respiratory disease in urban areas in Beijing, China, in 2013. *PLoS One* 2016;11:e0153099.
- [14] Ko FW, Tam W, Wong TW, et al. Temporal relationship between air pollutants and hospital admissions for chronic obstructive pulmonary disease in Hong Kong. *Thorax* 2007;62:780–5.
- [15] Gilmour PS, Rahman I, Donaldson K, et al. Histone acetylation regulates epithelial IL-8 release mediated by oxidative stress from environmental particles. *Am J Physiol Lung Cell Mol Physiol* 2003;284:L533–40.
- [16] Novović M, Topić V. Correlation between arterial and venous blood gas analysis parameters in patients with acute exacerbation of chronic obstructive pulmonary disease. *Srp Arh Celok Lek* 2012;140:436–40.
- [17] Wong TW, Lau TS, Yu TS, et al. Air pollution and hospital admissions for respiratory and cardiovascular diseases in Hong Kong. *Occup Environ Med* 1999;56:679–83.
- [18] Wang X, Kindzierski W, Kaul P. Air pollution and acute myocardial infarction hospital admission in Alberta, Canada: a three-step procedure case-crossover study. *PLoS One* 2015;10:e0132769.
- [19] Mustafic H, Jabre P, Caussin C, et al. Main air pollutants and myocardial infarction: a systematic review and meta-analysis. *JAMA* 2012;307:713–21.
- [20] Hurvich CMSJ, Tsai CL. Smoothing parameter selection in nonparametric regression using an improved Akaike information criterion. *J R Stat Soc* 1998;60:271–93.
- [21] Sant'Anna Neto JLda VS. Uma reflexão do espaço urbano da cidade de Campo Grande/MS na perspectiva climática. *Rev Pantaneira* 2003;3.
- [22] Brook RD, Urch B, Dvonch JT, et al. Insights into the mechanisms and mediators of the effects of air pollution exposure on blood pressure and vascular function in healthy humans. *Hypertension* 2009;54:659–67.
- [23] Andersen ZJ, Olsen TS, Andersen KK, et al. Association between short-term exposure to ultrafine particles and hospital admissions for stroke in Copenhagen, Denmark. *Eur Heart J* 2010;31:2034–40.
- [24] Shah AS, Lee KK, McAllister DA, et al. Short term exposure to air pollution and stroke: systematic review and meta-analysis. *BMJ* 2015;350:h1295.
- [25] Donaldson K, Duffin R, Langrish JP, et al. Nanoparticles and the cardiovascular system: a critical review. *Nanomedicine (Lond)* 2013;8:403–23.
- [26] Joubert J, Cumming TB, McLean AJ. Diversity of risk factors for stroke: the putative roles and mechanisms of depression and air pollution. *J Neurol Sci* 2007;262:71–6.
- [27] Kan H, London SJ, Chen G, et al. Season, sex, age, and education as modifiers of the effects of outdoor air pollution on daily mortality in Shanghai, China: the public health and air pollution in Asia (PAPA) study. *Environ Health Perspect* 2008;116:1183–8.
- [28] Miller MR, Raftis JB, Langrish JP, et al. Inhaled nanoparticles accumulate at sites of vascular disease. *ACS Nano* 2017;11:4542–52.
- [29] Andersen ZJ, Kristiansen LC, Andersen KK, et al. Stroke and long-term exposure to outdoor air pollution from nitrogen dioxide: a cohort study. *Stroke* 2012;43:320–5.
- [30] Vidale S, Bonanomi A, Guidotti M, et al. Air pollution positively correlates with daily stroke admission and in hospital mortality: a study in the urban area of Como. *Italy Neurol Sci* 2010;31:179–82.
- [31] Villeneuve PJ, Chen L, Stieb D, et al. Associations between outdoor air pollution and emergency department visits for stroke in Edmonton. *Canada Eur J Epidemiol* 2006;21:689–700.
- [32] Wellenius GA, Burger MR, Coull BA, et al. Ambient air pollution and the risk of acute ischemic stroke. *Arch Intern Med* 2012;172:229–34.
- [33] Huang F, Luo Y, Tan P, et al. Gaseous air pollution and the risk for stroke admissions: a case-crossover study in Beijing, China. *Int J Environ Res Public Health* 2017;14.
- [34] Zanobetti A, Schwartz J, Dockery DW. Airborne particles are a risk factor for hospital admissions for heart and lung disease. *Environ Health Perspect* 2000;108:1071–7.
- [35] Qin XD, Qian Z, Vaughn MG, et al. Gender-specific differences of interaction between obesity and air pollution on stroke and cardiovascular diseases in Chinese adults from a high pollution range area: a large population based cross sectional study. *Sci Total Environ* 2015;529:243–8.
- [36] Qian Y, Zhu M, Cai B, et al. Epidemiological evidence on association between ambient air pollution and stroke mortality. *J Epidemiol Community Health* 2013;67:635–40.
- [37] Wang XD, Zhang XM, Zhuang SW, et al. Short-term effects of air pollution on acute myocardial infarctions in Shanghai, China, 2013–2014. *J Geriatr Cardiol* 2016;13:132–7.
- [38] Milojevic A, Wilkinson P, Armstrong B, et al. Short-term effects of air pollution on a range of cardiovascular events in England and Wales: case-crossover analysis of the MINAP database, hospital admissions and mortality. *Heart* 2014;100:1093–8.
- [39] Clougherty JE. A growing role for gender analysis in air pollution epidemiology. *Cien Saude Colet* 2011;16:2221–38.