


Diagnostic track in amyotrophic lateral sclerosis: the Brazilian experience

O processo diagnóstico na esclerose lateral amiotrófica: a experiência brasileira

Mamede de Carvalho^{1,2} 

¹ Universidade de Lisboa, Faculdade de Medicina, Centro de Estudos Egas Moniz, Instituto de Fisiologia, Instituto de Medicina Molecular, Lisboa, Portugal.

² Centro Hospitalar Universitário de Lisboa Norte, Hospital de Santa Maria, Departamento de Neurociências e Saúde Mental, Lisboa, Portugal.

Arq. Neuropsiquiatr. 2022;80(7):661–662.

Diagnostic delay in amyotrophic lateral sclerosis (ALS) is a relevant problem that has been investigated in different countries for a long time. This delay has impact on the number of unnecessary investigations, social and personal disease economic burden, anxiety levels related to uncertain medical decisions, and increased risk of needless surgeries.¹ A wrong-surgery tends to further delay the diagnosis and to promote faster disease progression, a circumstance confirmed by neurologists dedicated to ALS and supported by scientific data.² Additionally, a diagnosis delay decreases the chances of clinical trial participation. Shortening diagnostic delay has been the rationale for establishing the Revised-El Escorial Criteria,³ the Awaji proposal (which is mainly focused on the neurophysiological features of ALS),⁴ and the Gold Coast Criteria.⁵ This demonstrates the importance of this subject, and the strong involvement of different institutions and experts on improving this constraint.

More recently, the median or mean time (variable in different sources) between the first symptoms (related to muscle weakness) and diagnosis is between 10 and 15 months, in general.⁶ Consistently, bulbar-onset disease^{7,8} and a faster observation by a neurologist has been associated with a shorter diagnostic delay.^{1,9,10} Misdiagnosis is quite common in ALS, although quite variable in different centers.⁶ Unquestionably, an initial wrong diagnosis delays ALS diagnosis, and increases the risk of inadequate medical interventions. It is uncertain if the modifications of the diagnostic criteria had a positive impact on decreasing delay of diagnosis in ALS.

The medical practice is different between countries, which depends on the health system organization, customary medical referrals routines, and other cultural influences. In Europe, medical care is first provided by general medical practitioners, who are available without costs, but in other countries this option is not so accessible. Moreover, it is reasonable to consider that the patient's salary income can influence the diagnostic track.¹⁰ For this reason, investigating the reality in each country is recommended, to endorse the appropriate interventions for each setting.

In this issue, Borthetti et al. retrospectively investigated the diagnostic track and misdiagnosis in a population of 173 ALS patients followed in Ribeirão Preto, Brazil.¹¹ They concluded that misdiagnosis was common in their population (70%), and that 7% of the patients underwent to unnecessary surgeries. In particular, limb-onset patients were more frequently observed by orthopedic surgeons, who tended to diagnose root or spinal cord lesions, and bulbar-onset patients were more commonly observed by ENT specialists. This suggests that these specialists could benefit from more information about clinical manifestations of ALS. From our experience, information about initial respiratory symptoms in ALS should be shared with pulmonologists and cardiologists, as 4% of ALS patients present with respiratory-onset and are followed by those specialists.¹²

Electromyography investigation has a critical role in the diagnosis of ALS, and is an essential step to reduce misdiagnosis and shorten diagnostic delay.¹⁰ This point was not approached in this study, but further research is welcomed.

Address for correspondence
Mamede de Carvalho
(e-mail: mamedemg@mail.
telepac.pt).

DOI [https://doi.org/
10.1055/s-0042-1755281](https://doi.org/10.1055/s-0042-1755281).
ISSN 0004-282X.

© 2022. Academia Brasileira de Neurologia. All rights reserved. This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

Finally, it should be said that ALS is a difficult clinical situation. Sometimes, neurologists decide to defer diagnosis since it is uncomfortable to communicate this dreadful disease. We need more information, and this study is important for our current knowledge.

Conflict of Interest

The author has no conflict of interests to declare.

References

- 1 Househam E, Swash M. Diagnostic delay in amyotrophic lateral sclerosis: what scope for improvement? *J Neurol Sci* 2000;180(1-2):76–81
- 2 Pinto S, Swash M, de Carvalho M. Does surgery accelerate progression of amyotrophic lateral sclerosis? *J Neurol Neurosurg Psychiatry* 2014;85(06):643–646
- 3 Brooks BR, Miller RG, Swash M, Munsat TL World Federation of Neurology Research Group on Motor Neuron Diseases. El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord* 2000;1(05):293–299
- 4 de Carvalho M, Dengler R, Eisen A, et al. Electrodiagnostic criteria for diagnosis of ALS. *Clin Neurophysiol* 2008;119(03):497–503
- 5 Shefner JM, Al-Chalabi A, Baker MR, et al. A proposal for new diagnostic criteria for ALS. *Clin Neurophysiol* 2020;131(08):1975–1978
- 6 Richards D, Morren JA, Pioro EP. Time to diagnosis and factors affecting diagnostic delay in amyotrophic lateral sclerosis. *J Neurol Sci* 2020;417:117054
- 7 Kano O, Iwamoto K, Ito H, et al. Limb-onset amyotrophic lateral sclerosis patients visiting orthopedist show a longer time-to-diagnosis since symptom onset. *BMC Neurol* 2013;13:19
- 8 Nzwalo H, de Abreu D, Swash M, Pinto S, de Carvalho M. Delayed diagnosis in ALS: the problem continues. *J Neurol Sci* 2014;343(1-2):173–175
- 9 Palese F, Sartori A, Logroscino G, Pisa FE. Predictors of diagnostic delay in amyotrophic lateral sclerosis: a cohort study based on administrative and electronic medical records data. *Amyotroph Lateral Scler Frontotemporal Degener* 2019;20(3-4):176–185
- 10 Falcão de Campos C, Gromicho M, Uysal H, et al. Delayed Diagnosis and Diagnostic Pathway of ALS Patients in Portugal: Where Can We Improve? *Front Neurol* 2021;12:761355
- 11 Borthetti VS, Cintra VP, Ramos JO, et al. Misdiagnosis in a Brazilian population with amyotrophic lateral sclerosis. *Arq Neuropsiquiatr*, In press
- 12 Pinto S, Gromicho M, Oliveira Santos MO, Swash M, De Carvalho M. Respiratory onset in amyotrophic lateral sclerosis: clinical features and spreading pattern. *Amyotroph Lateral Scler Frontotemporal Degener* 2022;•••:1–5. Doi: 10.1080/21678421.2022.2067777