



Microbial communities mediating algal detritus turnover under anaerobic conditions

Jessica M. Morrison^{1,*}, Chelsea L. Murphy^{1,*}, Kristina Baker¹, Richard M. Zamor², Steve J. Nikolai², Shawn Wilder³, Mostafa S. Elshahed¹ and Noha H. Youssef¹

¹ Department of Microbiology and Molecular Genetics, Oklahoma State University, Stillwater, OK, USA

² Grand River Dam Authority, Vinita, OK, USA

³ Department of Integrative Biology, Oklahoma State University, Stillwater, OK, USA

* These authors contributed equally to this work.

ABSTRACT

Background. Algae encompass a wide array of photosynthetic organisms that are ubiquitously distributed in aquatic and terrestrial habitats. Algal species often bloom in aquatic ecosystems, providing a significant autochthonous carbon input to the deeper anoxic layers in stratified water bodies. In addition, various algal species have been touted as promising candidates for anaerobic biogas production from biomass. Surprisingly, in spite of its ecological and economic relevance, the microbial community involved in algal detritus turnover under anaerobic conditions remains largely unexplored.

Results. Here, we characterized the microbial communities mediating the degradation of *Chlorella vulgaris* (Chlorophyta), *Chara* sp. strain IWP1 (Charophyceae), and kelp *Ascophyllum nodosum* (phylum Phaeophyceae), using sediments from an anaerobic spring (Zodlsteone spring, OK; ZDT), sludge from a secondary digester in a local wastewater treatment plant (Stillwater, OK; WWT), and deeper anoxic layers from a seasonally stratified lake (Grand Lake O' the Cherokees, OK; GL) as inoculum sources. Within all enrichments, the majority of algal biomass was metabolized within 13–16 weeks, and the process was accompanied by an increase in cell numbers and a decrease in community diversity. Community surveys based on the V4 region of the 16S rRNA gene identified different lineages belonging to the phyla Bacteroidetes, Proteobacteria (alpha, delta, gamma, and epsilon classes), Spirochaetes, and Firmicutes that were selectively abundant under various substrate and inoculum conditions. Within all kelp enrichments, the microbial communities structures at the conclusion of the experiment were highly similar regardless of the enrichment source, and were dominated by the genus *Clostridium*, or family *Veillonellaceae* within the Firmicutes. In all other enrichments the final microbial community was dependent on the inoculum source, rather than the type of algae utilized as substrate. Lineages enriched included the uncultured groups VadinBC27 and WCHB1-69 within the Bacteroidetes, genus *Spirochaeta* and the uncultured group SHA-4 within Spirochaetes, *Ruminococcaceae*, *Lachnospiraceae*, *Yongiibacter*, *Geosporobacter*, and *Acidaminobacter* within the Firmicutes, and genera *Kluyvera*, *Pantoea*, *Edwardsiella* and *Aeromonas*, and *Buttiauxella* within the Gamma-Proteobacteria order Enterobacteriales.

Submitted 15 September 2016

Accepted 18 November 2016

Published 10 January 2017

Corresponding author

Noha H. Youssef, noha@okstate.edu

Academic editor

Jonathan Eisen

Additional Information and
Declarations can be found on
page 25

DOI 10.7717/peerj.2803

© Copyright

2017 Morrison et al.

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS

Conclusions. Our results represent the first systematic survey of microbial communities mediating turnover of algal biomass under anaerobic conditions, and highlights the diversity of lineages putatively involved in the degradation process.

Subjects Microbiology

Keywords Algal detritus, Anaerobic degradation, Enrichment

INTRODUCTION

Algae represent a globally distributed group of organisms that are capable of oxygenic photosynthesis. While prevalent in aquatic marine and freshwater habitats (Cole, 1982), algal taxa are also encountered in terrestrial ecosystems such as soil, rocks, and ice/snow (Hoffmann, 1989). Collectively, algal species play an important role in global carbon, nitrogen, sulfur, and phosphorus cycling (Vanni, 2002). Taxonomically, algae are polyphyletic, and are encountered within multiple eukaryotic phyla such as the Alveolata (e.g., dinoflagellates), Stramenopiles (e.g., *Bacillariophyceae*, *Chrysophyceae*, *Eustigmatophyceae*), Viridiplanta (e.g., Chlorophyta), in addition to exclusively algal phyla such as the Euglenozoa, Cryptomonads, Haptophyta, and Rhodophyta (Amaral-Zettler, 2011). In addition to their complex evolutionary origin, these organisms exhibit a wide array of morphological diversity, pigments, ecological distribution, cellular composition, genome size, and cell wall structure.

A major characteristic of many algal taxa is their fast growth rate, enabling them to form conspicuous seasonal blooms under the appropriate environmental conditions. Such blooms are often associated with elevated nutrient (e.g., nitrogen and/or phosphorus) levels in the ecosystem, often resulting from anthropogenic inputs (e.g., sewage, industrial waste, and fertilizers) (Hallegraeff, 1993), as well as from destratification and nutrients resuspension (Wetzel, 2001). Classical examples of freshwater algal blooms involve members of the green algae (*Chlorophyceae*), whose blooms are often encountered in lakes and other freshwater habitats (Hoshaw & Mccourt, 1988), *Chara* blooms (commonly called Muskgrass), which seasonally occur in ponds and lakes and cause a strong and unpleasant musky odor (Durborow, 2014), as well as Diatoms, most commonly encountered in lakes (Sommer et al., 2012; Sommer et al., 1986). Fast growth is also a characteristic of many annual or perennial macroscopic taxa. The classical example of such taxa is the brown algae or Kelp (class *Phaeophyceae*), which is believed to be one of the most productive photosynthetic organisms and tend to attain long lengths at a very fast elongation rate (~50–60 cm/day) (Reed, Rassweiler & Arkema, 2008).

Algae provide a large input of organic carbon into aquatic ecosystems such as coastal kelp forests (Mann, 1988), meromictic and seasonally stratified lakes (Gies et al., 2014; Xia et al., 2016; Youssef et al., 2015), and coastal areas within marine environments (e.g., the North Sea (Boon et al., 1998)). When blooming subsides, the algal detritus sinks and provides a substantial organic carbon source to microbial communities within the ecosystem (Hecky & Hesslein, 1995). Algal degradation in aquatic habitats commences at or near the water

surface by the microbial phycosphere: aerobic heterotrophic bacteria that are physically attached to algal cells (Buchan et al., 2014). However, a significant fraction of algal detritus reaches the lower strata of these water bodies, providing a considerable autochthonous contribution to the carbon input in benthic layers within such ecosystems (Ask et al., 2009; Hecky & Hesslein, 1995). Sinking of algal detritus and the subsequent increase in carbon input result in the development of anoxic conditions in the lower strata and, hence, seasonal stratification. The size, intensity, and duration of these bloom-mediated anoxic zones are expected to be accentuated by future global warming trends (Paerl & Otten, 2013). Surprisingly, while a large body of research has been conducted on elucidating the microbial community composition of the algal phycosphere in the aerobic surficial marine (Amin, Parker & Armbrust, 2012; Hasegawa et al., 2007; Sapp, Wichels & Gerdt, 2007), and freshwater habitats (Bagatini et al., 2014; Cai et al., 2014; Dittami et al., 2016; Eigemann et al., 2013; Jones et al., 2013; Muylaert et al., 2002), a surprising lack of knowledge exists regarding the microbial community and patterns of algal turnover under the anoxic conditions in the lower layers of stratified water bodies.

In addition to the importance of anaerobic degradation of algal biomass to the carbon cycle in aquatic environments, the process has recently received additional attention as an integral component in algal biofuels production schemes. Direct conversion of kelp to methane (Cannell, 1990; Prabandono & Amin, 2015; Ramaraj, Unpaprom & Dussadee, 2016) has been proposed as a promising approach for biogas production (Samson & Leduy, 1982; Vanegas & Bartlett, 2013; Vergara-Fernandez et al., 2008; Wiley, Campbell & McKuin, 2011; Yen & Brune, 2007; Yuan et al., 2011). The high fat content of multiple algal taxa, e.g., *Chlorella* sp. (Guckert & Cooksey, 1990; Hu et al., 2008), and *Chara* sp. (Omer, 2013), has prompted research into their large scale production in artificial ponds, and subsequent extraction of their oil content as biodiesel (Cannell, 1990; Moazami et al., 2012; Prabandono & Amin, 2015; Ramaraj, Unpaprom & Dussadee, 2016). The economy of the process is further enhanced by anaerobic digestion of the algal detritus to produce methane as an additional source (Cannell, 1990; Prabandono & Amin, 2015; Ramaraj, Unpaprom & Dussadee, 2016) using sludge from anaerobic wastewater treatment plants as the inoculum source (Bohutskyi et al., 2015). While various aspects of the engineering and performance have been studied, there is very little documentation of the identity of the microbial community that is mediating algal detritus turnover under these anaerobic conditions.

Here, we sought to characterize patterns of algal turnover under anaerobic conditions, and identify members of the microbial community involved in the degradation of various ecologically and economically relevant algal taxa. The process was investigated in enrichments derived from three anaerobic habitats that either exhibit seasonal algal blooming, or highly eutrophic environments that receive a high input of organic compounds and previously shown to harbor a high level of microbial diversity. To our knowledge, this is the first study that systematically characterized the microbial community associated with algal degradation under anaerobic conditions.

MATERIALS AND METHODS

Algal species (substrates/carbon source)

We evaluated the microbial community mediating algal detritus degradation by setting enrichments with various types of algal biomass as the only source of carbon. Three different algal species were examined: (1) Axenic *Chlorella vulgaris* strain UTEX 2714 (phylum Chlorophyta), representing green algae that are known to bloom during summer months (June–September) in multiple freshwater bodies within the US. Strain UTEX 2714 was obtained from the University of Texas at Austin Algal Culture collection, and cultures were maintained on proteose medium plates (composition g l^{-1} : NaNO_3 , 0.25; $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.025, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.075; K_2HPO_4 , 0.075, KH_2PO_4 , 0.175, NaCl , 0.025; proteose peptone, 1, agar, 15) at 22 ± 3 °C. Fluorescent light was used (3500 lux) on a 16:8 h light to dark cycle. Cultures were harvested by scraping the growth on the surface of agar plates, and the resulting biomass was used as the carbon source. (2) *Chara* sp. strain IWP: Mats of *Chara* (class Charophyceae) were obtained from a local pond (Innovation Way Pond in Stillwater, OK, coordinates N36°6'37.75" W97°6'44.72") in August 2015, and identified using morphological and microscopic analysis as *Chara* sp. *Chara* is known to grow locally in ponds in Oklahoma and peak around June–September (B Henley, pers. comm., 2016). Samples collected were thoroughly washed and soaked in DI water for 2 h, to remove other associated biomass, before they were centrifuged and the resulting biomass was used as the carbon source to represent class Charophyceae. (3) Kelp: Due to the geographical location of the study and brown algae being common occurrences in marine environments, we were not able to obtain a fresh kelp sample to be used as a substrate. Alternatively, *Ascophyllum nodosum* (phylum Phaeophyceae) representing brown algae was obtained as a whole dried powder from a local provider (Starwest Botanicals®, Sacramento, CA, USA) and directly used as the carbon source.

It is worth noting that, with the exception of the axenic *Chlorella* culture, the *Chara* and the kelp samples were not guaranteed axenic and a minimal input of bacteria with the carbon source in these enrichment bottles could not be ruled out.

Enrichment (inoculum) sources

Grand Lake (GL)

Samples were obtained from Grand Lake O' the Cherokees (hereafter Grand Lake) to investigate the microbial community involved in algal turnover. The lake is a large (188 Km^2) man-made lake in Northeastern OK operated by the Grand River Dam Authority, which keeps continuous records of the lake water geochemistry. During summer months (starting in June), large areas of the lake become seasonally stratified, with deeper layers (June–September) becoming completely anoxic. Within these sites, surface chlorophyll-a concentration peaks in May–July, followed by algal biomass sinking to deeper anaerobic layer in September (Fig. S1). Sampling from the hypolimnion of Tree and Dream sites in GL occurred in September 2015 using a 4.0-L Van Dorn Bottle. Whole water samples were stored on ice until processed in the laboratory where the lake water was centrifuged under anaerobic conditions to collect biomass used as inoculum source.

Wastewater treatment plant (WWT)

While wastewater treatment reactors do not represent an algae rich habitat, the use of WWT material as an inoculum for biogas production from algal biomass sources has been gaining considerable attention ([Ward, Lewis & Green, 2014](#)). The process is justified by the high organic content and adaptation to organic matter turnover under anaerobic conditions, coupled to the ready availability of WWT inocula ([Sialve, Bernet & Bernard, 2009](#)). Various aspects of the engineering, kinetics, and the economy of the process, as well as optimization of the inoculum load and substrate load, and algae pretreatment methods have been investigated ([Hlavínek et al., 2016](#); [Mudhoo, 2012](#); [Nabarlatz et al., 2013](#); [Ramaraj, Unpaprom & Dussadee, 2016](#); [Samson & Leduy, 1982](#); [Vergara-Fernandez et al., 2008](#); [Ward, Lewis & Green, 2014](#)), but little research on the identity of microorganisms mediating such process has been conducted. Samples were obtained from secondary treatment sludge in the city of Stillwater, OK wastewater treatment plant in September 2015. The sample was collected anaerobically and transferred to the laboratory (5 miles away), where they were promptly centrifuged under anaerobic conditions and used as inoculum source.

Zodletone spring (ZDT)

Zodletone spring is an anaerobic surficial spring in southwestern OK (35°0'9"N 98°41'17"W). Due to the constant ejection of sulfide laden water from the spring source, the shallow spring is light exposed, yet mostly anoxic ([Buhring et al., 2011](#)). The microbial community in the spring has been extensively investigated ([Coveley, Elshahed & Youssef, 2015](#); [Youssef, Couger & Elshahed, 2010](#)), and the spring harbors a highly diverse community of phototrophs, chemolithotrophs, and heterotrophs. Samples were collected from Zodletone spring source in August 2015 in a filled mason jar (to maintain anoxic conditions), transferred to the laboratory at 4 °C and used as inoculum source on the same day of sampling.

Enrichments setup

Enrichments were prepared in 120 ml serum bottles under anaerobic conditions. Nine different treatments (three algal substrates × three different inoculum sources) were set up in triplicates. Serum bottles contained 45 ml of an anoxic solution containing (per liter): 150 ml of minerals solution I (K_2HPO_4 3 g l⁻¹), 150 ml of mineral solution II (g l⁻¹: KH_2PO_4 , 3; $(NH_4)_2SO_4$, 6; NaCl, 6; $MgSO_4 \cdot 7H_2O$, 0.6, and $CaCl_2 \cdot 2H_2O$, 0.6), 10 ml Balch vitamins solution (mg l⁻¹: biotin, 2; folic acid, 2; pyridoxine-HCl, 10; thiamine-HCl, 5; riboflavin, 5; nicotinic acid, 5; calcium pantothenate, 5; vitamin B12, 0.1; p-aminobenzoic acid, 5; lipoic acid, 5), 1 ml of Wolin's metal solution (g l⁻¹: EDTA, 0.5; $MgSO_4 \cdot 6H_2O$, 3.0; $MnSO_4 \cdot H_2O$, 0.5; NaCl, 1; $CaCl_2 \cdot 2H_2O$, 0.1; $FeSO_4 \cdot 7H_2O$, 0.1; $ZnSO_4 \cdot 7H_2O$, 0.1; $CuSO_4 \cdot 7H_2O$, 0.01; $AlK(SO_4)_2$, 0.01; $Na_2MoO_4 \cdot 2H_2O$, 0.01; boric acid, 0.01; Na_2SeO_4 , 0.005; $NiCl_2 \cdot 6H_2O$, 0.003; $CoCl_2 \cdot 6H_2O$, 0.1). The media were amended with L-cysteine hydrochloride (0.05 g/l final concentration) as a reductant, and resazurin (0.0001% final concentration) as a redox indicator, boiled under a stream of N₂ gas, dispensed in the serum bottles, autoclaved, cooled, then transferred to an anaerobic chamber (Coy Laboratory

Products Inc., Ann Arbor, MI, USA) where 5 g of sediment (Zodletone, ZDT), 5 g of sludge (wastewater treatment plant, WWT), or 5 ml of concentrated lake water (the pellet obtained after centrifugation of 1.5 L of Grand Lake water (GL)) were added as the inoculum sources. Algal biomass was added as a substrate (~0.25 g per bottle). In addition, substrate unamended controls (i.e., ZDT, WWT, and GL enrichments with no algal substrates) were included. After enrichment preparation in the anaerobic chamber, the bottles were stoppered, sealed, taken out of the chamber and the headspace in the bottles was changed by repeated flushing with 100% N₂. Samples were incubated at room temperature (22 °C) in the dark. Enrichments were periodically sampled (at 4, 7, 8, and 10 weeks) for DNA extraction by thoroughly mixing the serum bottle and anoxically withdrawing 3 ml of the enrichment. At the end of the enrichment process (13 weeks for GL, 16 weeks for ZDT and WWT), bottles were sacrificed, centrifuged and 3 ml of the pellet was used for DNA extraction. The rest of the pellet was used for chemical analysis of the remaining algal detritus.

DNA extraction, amplification, and sequencing

DNA was extracted using the PowerSoil[®] DNA Isolation Kit (MO BIO Laboratories, West Carlsbad, CA, USA) as per the manufacturer's instruction. DNA from triplicate treatments was pooled prior to amplification and sequencing. DNA from substrate-unamended controls was also extracted (hereafter pre-enrichment sample). The extracted and pooled DNA ($n = 30$; 3 inoculum sources \times 3 algal substrates \times 3 time points, plus 3 pre-enrichment samples) was quantified using Qubit fluorometer (Life technologies[®], Carlsbad, CA, USA). The genes for the V4 hypervariable region of 16S rRNA were amplified using the prokaryotic-specific primer pair 515F and 806R (Wang & Qian, 2009) to avoid amplification of eukaryotic 18S rRNA. Products were sequenced using paired-end Illumina Miseq platform, as previously described (Caporaso *et al.*, 2012). Both PCR amplification and Illumina sequencing were conducted using the services of the Genomic Sequencing and Analysis Facility (GSAF) at the University of Texas at Austin. The sequences are deposited in the SRA database under accession number [SRP083898](https://www.ncbi.nlm.nih.gov/sra/SRP083898).

Data analysis

Sequence processing

mothur software (Schloss *et al.*, 2009) was used for most of the sequence processing and operational taxonomic unit (OTU) assignments. Most of the analyses were conducted on the cowboy server, a high performance super computer housed at the Oklahoma State High Performance Computing Center (<https://hpcc.okstate.edu/>). For quality control purposes and to eliminate poor quality sequences, an average quality score of 25 was chosen as the threshold value below which sequences were considered of poor quality and removed from the dataset. In addition, sequences that contained an ambiguous base (N), sequences having a homopolymer stretch longer than 8 bases, and sequences longer than 293 bp were also removed from the datasets.

High-quality reads were aligned in mothur using the Silva alignment database as a template. Aligned sequences were then filtered to remove columns that corresponded to '.' or '-' in all sequences. Filtered alignments were then subjected to a pre-clustering

de-noising step using a pseudo-single linkage algorithm with the goal of removing sequences that are likely due to sequencing errors (Huse *et al.*, 2010). Possible chimeric sequences were identified and removed using chimera.slayer in mothur. The taxonomy of the remaining sequences was identified according to the Silva taxonomic outline (Release 123, <https://www.arb-silva.de/>). The aligned, filtered, de-noised, and chimera-free sequences were used to generate an uncorrected pair wise distance matrix. Sequences were clustered into operational taxonomic units (OTUs) at 0.03% sequence divergence cutoff using the vsearch clustering method employed through mothur. A shared file was created and was used for subsequent analyses. Most of the above steps were derived from the MiSeq SOP available from the mothur website (http://www.mothur.org/wiki/MiSeq_SOP).

Criteria used to define lineages contributing to the degradation process

Phyla considered significant to the degradation process were empirically defined as those phyla that constituted 5% or more of the community at any time during enrichment. These include phyla that were abundant prior to enrichment and remained abundant during and after enrichment, phyla that transiently increased in abundance during part of the enrichment but then decreased in abundance by the end of enrichment, and phyla that significantly and progressively increased in abundance with enrichment time. Within these abundant phyla, genera considered significant to the degradation process were also empirically defined as those whose percentage abundance represented 1% or more of the total abundance.

Diversity and community structure comparisons

Various alpha diversity indices (Shannon, Chao, Ace, Good's coverage) were performed on individual datasets in mothur. When comparing species richness across datasets (e.g., number of observed OTUs, species richness estimates using Chao and Ace estimators), numbers were reported per sample size to normalize for the differences in the number of sequences obtained between datasets. Beta diversity based on community structure was assessed by calculating pairwise Bray–Curtis dissimilarity indices and using the output to construct non-metric multidimensional scaling (NMDS) plots in mothur.

Statistical analyses

To study the significance of the effect of inoculum source versus algae type on community structure, we performed an analysis of variance using the function Adonis in the R statistical package vegan. The effect was visualized using the percentage abundances of significant phyla/classes ($\geq 5\%$ as defined above) in a constrained correspondence analysis (CCA) using the function cca in the R statistical package vegan.

Quantitative PCR

We used qPCR to quantify total Bacteria, total Archaea, as well as methanogens and sulfate-reducing bacteria in the enrichments using a MyiQ thermocycler (Bio-Rad Laboratories, Hercules, CA) and SYBR GreenERTM qPCR SuperMix for iCycler[®] Instrument (Life Technologies). Primer pair EUB-338F/UNI518R (Fierer *et al.*, 2005) was used to amplify the 16S rRNA genes from the total bacterial community, primer pair A341F/A519R (Qian

et al., 2011) was used to amplify 16S rRNA genes from the total archaeal community, primer pair mlas-ModF/mcrA-R (Angel, Claus & Conrad, 2012) was used to amplify the methyl-Coenzyme M reductase (*mcrA*) gene from the total methanogenic community, and primer pair Dsr2060F/Dsr4R (Balk, Keuskamp & Laanbroek, 2015) was used to amplify the dissimilatory sulfite reductase (*dsrB*) gene from the total sulfate reducing community. The 25- μ l PCR reaction mixtures contained 0.3 μ M of each forward and reverse primers (final concentration), 2 μ l extracted template DNA, and 12.5 μ l SYBR GreenERTM qPCR SuperMix. The reactions were heated at 95 °C for 8.5 min, followed by 40 cycles, with one cycle consisting of 30 s at 95 °C, 45 s at 50 °C (for total bacteria, total archaea, and methanogens) or 55 °C (for sulfate reducers), 30 s at 72 °C, and 15 s at 85 °C for signal reading. To calculate the total number of cells belonging to total bacteria, total archaea, methanogens, and sulfate reducers in the enrichments, a standard curve was generated using DNA from *Bacillus subtilis* strain 168 (ATCC 23857), *Haloferax sulfurifontis* strain M6 (DSM 16227), *Methanosarcina hungatei* strain JF1 (ATCC 27890), and *Desulfovibrio desulfuricans* strain G20 (ATCC BAA-1058), respectively. To account for the multiple copies of 16S rRNA genes per cell, the number of copies obtained from the standard curve was divided by an empirical value of 3.5 (average of 1–6 copies of rRNA genes in one cell). However, since the *mcrA* and *dsrB* genes are known to be present as single copies in methanogens, and sulfate reducers, respectively, no such adjustment of the total number of cells was required when calculating the total number of cells belonging to methanogens and sulfate reducers.

Chemical analysis of algal detritus

We studied the change in chemical composition of algal detritus during enrichment by quantifying the total soluble carbohydrates, total starch, total protein, and total lipid content of algal biomass pre and post enrichment. Algal detritus was dried overnight at 40 °C then weighed (DW_f; final dry weight) and ground to fine material. The ground material was first used for protein extraction using the method described previously (Rausch, 1981). Briefly, algal detritus was extracted 2–3 times with 0.5N NaOH at 80–100 °C for 10 min followed by cooling and centrifugation to collect the total protein in the supernatant. Total protein extracts were frozen at –20 °C until assayed using Qubit Protein Assay Kit (Life technologies). The pellet remaining after protein extraction was used for extraction of total soluble carbohydrates and starch. The pellets were first washed 2–3 times with 1 ml acetone to remove pigments. Total soluble carbohydrates were then extracted from the pellet using 80% ethanol according to the protocol in Maness (2010) and the total ethanol extract was dried overnight at 40 °C followed by dissolving the dried extract in water. Total starch remaining in the pellet was extracted by boiling with 1.1% HCl for 30 min followed by centrifugation. Total soluble carbohydrates, as well as total starch extracted were quantified using the anthrone method (Maness, 2010). The total crude lipids were extracted from dried algal material with chloroform and quantified with a Nile red assay modified for microplates using the protocol described previously (Higgins *et al.*, 2014).

RESULTS

Sequencing output

A total of 1,007,906 sequences were obtained from all enrichments. After implementation of all quality control criteria described above, 889,230 sequences (88.2%) were retained for further analysis. The average number of sequences per dataset was 26,946. The calculated Good's coverage for the majority of samples at putative species ($OTU_{0.03}$, 30 out of 32 samples) and family ($OTU_{0.10}$, 31 out of 32 samples) levels were always above 96 and 98.2% (average 98.4, and 99.5%, respectively), strongly indicating that the communities have been adequately sampled in all enrichments (Table S1).

Enrichment progress and diversity patterns

Multiple lines of evidence strongly indicate that in all nine treatments, algal detritus degradation occurred and was coupled to an increase in prokaryotic cell numbers and a decrease in alpha diversity both implying enrichment of specific taxa. Visual inspection of all enrichments revealed significant loss of the dried kelp powder, and the algal biomass (*Chlorella* and *Chara*) at the conclusion of the experiment. Final time point analysis demonstrated that the majority of the starting dry weight of *Chara* (86.3, 94.5, and 98.0%), *Chlorella* (96.0, 98.0, and 99.0%), and, to a relatively lower extent, kelp (56.7, and 33.6, 83.3%) in ZDT, WWT, and GL enrichments, respectively, was metabolized at the conclusion of the experiment. Analysis of the chemical composition of the remaining algal detritus in comparison to the starting material showed that the carbohydrate, lipid, and protein contents of the algal detritus were consumed to varying extents (Table 1). Quantitative PCR (qPCR) demonstrated a progressive increase in bacterial 16S rRNA gene copies/ml enrichment in all samples. An increase of 3.5–88.5 fold in total number of bacterial cells was observed by week 13–16 in all enrichments and 14.6–2,142 fold in the total number of archaeal cells was observed by week 13–16 in 6 out of 9 enrichments (Fig. 1). Finally, we followed the change in diversity estimates in the enrichments datasets as a proxy for enrichment progress. At the end of all enrichments (weeks 13 or 16), the number of observed $OTU_{0.03}$ and $OTU_{0.1}$ as well as the estimated species richness (using both Chao and ACE estimators) decreased compared to the pre-enrichment sample, hence indicating the selection for few taxa (Table 2).

Microbial community structure analysis

Bray–Curtis dissimilarity indices at $OTU_{0.03}$ coupled to non-metric multidimensional scaling (NMDS) were used to compare and visualize differences in the microbial community structure between all enrichments at all sampled data points. At first glance, it was apparent that the enriched microbial communities (week 7–10–16 in cases of WWT and ZDT microcosms, or week 4–8–13 in case of GL microcosms) within each algae type-enrichment source combination ($n = 9$, blue, green, and red shapes in Fig. 2) clustered closely together, and were distinct from the pre-enrichment microbial community (black shapes in Fig. 2). This observation strongly suggests that the abundant microbial community obtained during the first few weeks of enrichment (week 4 or 7) persisted throughout the enrichment and was responsible for the algal biomass degradation observed at the end of the enrichment

Table 1 Percentage of various algal components consumed under different enrichment conditions.^a

Algal detritus components	ZDT enrichment			WWT enrichment			GL enrichment		
	<i>Chara</i>	<i>Chlorella</i>	Kelp	<i>Chara</i>	<i>Chlorella</i>	Kelp	<i>Chara</i>	<i>Chlorella</i>	Kelp
Carbohydrate	87	96.5	52.9	99.7	98	15.7	98.3	99.2	86.2
Protein	72	92.9	60	94	96.9	70	96	97.6	82
Lipid	100	NA ^b	71	62.5	NA ^b	70	96	NA ^b	86
% Biomass lost ^c	86.3	96	56.7	94.5	98	33.6	98	99	83.3

Notes.

^aCarbohydrate, protein, and lipid contents of algal detritus were determined before and after enrichment. Percentages are calculated based on the dry weight at T_f . Original algal detritus composition was as follows (%Carbohydrate: %Protein: %Lipid): *Chara*, 88:6.5:5.5; *Chlorella*, 86.7:13.3:0; *Kelp*, 67.9:14.1:18.

^bLipids in *Chlorella* biomass were BDL.

^cBased on dry weight remaining at the end of enrichment (DW_f), and the initial dry weight used for enrichment (DW_0) using the equation: % biomass loss = $(DW_0 - DW_f)/DW_0 \times 100$. Initial dry weight for kelp was equivalent to the weight added to each enrichment bottle since it was in dry powder form. However, initial dry weight for *Chara* and *Chlorella* was determined by incubating an amount equivalent to the wet weight added to each enrichment bottle overnight at 40 °C then weighing its dry weight following moisture loss.

(Table 1). Analysis of the effect and relative contribution of algae type (*Chlorella*, *Chara*, or kelp) versus inoculum source (ZDT, WWT, and GL) on the enriched microbial communities revealed that kelp selects for a distinct and highly similar microbial community, regardless of the inoculum source (ZDT, WWT, and GL) (Fig. 2A). On the other hand, within *Chlorella* and *Chara*-derived enrichments, the inoculum source, rather than the algae type appears to be the more important factor in shaping the microbial communities (Fig. 2A). This is evident by the presence of three distinct clusters in the NMDS plot corresponding to the three sources of inoculum (ZDT, WWT, and GL) (Fig. 2A).

In addition, analysis of variance (using Adonis function) showed that both the algae type and the source of inoculum were significant in shaping the microbial community albeit to varying levels (p -value for algae type = 0.028, p -value for inoculum source = 0.001). To decipher the relative contributions of algae type versus inoculum source on the microbial community composition at the phylum/class level we employed canonical correspondence analysis (CCA) using the enriched phyla/classes relative abundances. The results (Fig. 2B) confirmed the above observation, where the algae type appears to have shaped the microbial community in case of kelp enrichments, while within *Chara* and *Chlorella* enrichments, the source of inoculum played a more important role in shaping the community (Fig. 2B).

Phylogenetic affiliation of enriched taxa in algal enrichments

In general, a handful of phyla were consistently abundant across all treatments and were considered significant to the algal degradation process (see the criteria we used for defining such phyla in 'Materials and Methods'). These phyla were: Firmicutes (in all nine enrichments), Bacteroidetes (in six enrichments), Spirochaetes (in five enrichments), and the Gamma (5 enrichments), Delta (6 enrichments), Alpha, Beta, and Epsilon (one enrichment) Proteobacteria (Figs. 3–5). However, within this limited number of phyla, the family/genus level enrichment patterns varied widely, suggesting the involvement of a wide range of bacterial lineages in the degradation process. Below, we provide a detailed




Figure 1 Total number of bacterial, archaeal, sulfate-reducing, and methanogenic cells in the pre-enrichment sample (■) versus post-enrichment samples at week 4 for GL enrichments or week 7 for ZDT and WWT enrichment (□), post-enrichment samples at week 8 for GL enrichments or week 10 for ZDT and WWT enrichment (▨), and post-enrichment samples at week 13 for GL enrichments or week 16 for ZDT and WWT enrichment (▩) as measured by quantitative PCR. The enrichment inoculum source is shown on the left, while the algae type used is shown on top. Error bars are averages \pm standard deviations from three biological replicates. Linear regression analysis was performed to examine the trend of increase in cell numbers with the weeks of enrichment, and the significance of such trend was tested by calculating the P -values of the F -statistics obtained, where “**” denotes significant P -value < 0.05 , “*” denotes p -value > 0.05 but < 0.1 , “NS” denotes non-significant P -value > 0.1 , and “ND” refers to cases where the linear regression analysis was not performed because two or more samples were below the detection level of the qPCR. In the few cases, denoted by a superscript letter a, where the total cell numbers increased initially then decreased by the last week of enrichment, the linear regression was only carried on total numbers from the first three weeks of enrichments.

analysis of the enriched families/genera across various enrichments. The detailed microbial community composition across all datasets is shown in [Table S2](#).

Chara microcosms

In *Chara* microcosms, Bacteroidetes, Firmicutes, and Delta-Proteobacteria were consistently abundant ([Table 3](#)) and, collectively, constituted the majority (40.2% to 72.7%) of the community at the end of enrichment. Spirochaetes were abundant only in WWT and ZDT enrichments, while Gamma-Proteobacteria were abundant only in GL and ZDT enrichments ([Figs. 3–5](#)).

Within the Bacteroidetes, the uncultured putative genus VadinBC27 was consistently enriched ([Table 3](#)) regardless of the inoculum source. This uncultured subgroup within the order Bacteroidales has been previously identified as a major lineage in anaerobic

Table 2 Number of OTU_{s0.03} and OTU_{s0.1} normalized to the total number of sequences, and the estimated species richness (using both Chao and ACE estimators) normalized to the total number of sequences.

Source_cutoff ^a	Weeks of enrichment	Chara			Chlorella			Kelp		
		OTUs	Chao	ACE	OTUs	Chao	ACE	OTUs	Chao	ACE
ZDT_0.03	0	0.193	0.373	0.501	0.193	0.373	0.501	0.193	0.373	0.501
	7	0.021	0.040	0.056	0.031	0.054	0.067	0.060	0.130	0.255
	10	0.030	0.051	0.068	0.022	0.043	0.058	0.069	0.148	0.265
	16	0.028	0.059	0.078	0.025	0.051	0.069	0.046	0.118	0.208
ZDT_0.1	0	0.051	0.078	0.088	0.051	0.078	0.088	0.051	0.078	0.088
	7	0.008	0.011	0.010	0.016	0.023	0.028	0.026	0.043	0.058
	10	0.012	0.017	0.016	0.009	0.013	0.015	0.026	0.042	0.053
	16	0.011	0.016	0.019	0.009	0.012	0.014	0.019	0.029	0.040
GL_0.03	0	0.043	0.091	0.140	0.043	0.091	0.140	0.043	0.091	0.140
	4	0.008	0.019	0.033	0.014	0.025	0.035	ND	ND	ND
	7	0.014	0.032	0.044	0.010	0.020	0.024	0.092	0.116	0.123
	13	0.022	0.055	0.057	0.011	0.018	0.029	0.036	0.063	0.076
GL_0.1	0	0.020	0.032	0.042	0.020	0.032	0.042	0.020	0.032	0.042
	4	0.003	0.007	0.009	0.008	0.015	0.022	ND	ND	ND
	7	0.006	0.015	0.020	0.005	0.009	0.010	0.051	0.058	0.061
	13	0.010	0.015	0.014	0.005	0.007	0.008	0.017	0.026	0.029
WWT_0.03	0	0.048	0.087	0.108	0.048	0.087	0.108	0.048	0.087	0.108
	7	0.043	0.072	0.092	0.066	0.109	0.140	0.013	0.039	0.055
	10	0.020	0.036	0.046	0.033	0.052	0.061	0.020	0.043	0.064
	16	0.030	0.051	0.068	0.021	0.036	0.043	0.012	0.025	0.035
WWT_0.1	0	0.013	0.022	0.029	0.013	0.022	0.029	0.013	0.022	0.029
	7	0.018	0.025	0.024	0.030	0.043	0.048	0.006	0.009	0.012
	10	0.006	0.009	0.009	0.013	0.019	0.019	0.006	0.010	0.014
	16	0.011	0.015	0.017	0.008	0.010	0.010	0.004	0.007	0.009

Notes.

^aSource refers to the inoculum source, while cutoff refers to the percentage divergence cutoff used to assign sequences into operational taxonomic units (OTUs). For each inoculum source, the numbers are shown for OTUs at the putative species level (0.03) and the putative order level (0.1).

ND, Not determined due to the small number of sequences obtained for this dataset.

digestors (Liu et al., 2016; Riviere et al., 2009; Xie et al., 2014; Xu et al., 2012) and was implicated as an anaerobic fermenter of sludge or other carbon sources. Other enriched Bacteroidetes members include the genera *Mangroviflexus* (ZDT microcosms), previously identified as an important *in-situ* fermenter of organic matter-rich soil (Ding et al., 2016) and anaerobic cellulolytic microcosms (Gao, Xu & Ruan, 2014), *Paludibacter* (WWT and ZDT microcosms), previously enriched from anaerobic freshwater sediment (Sanchez-Andrea et al., 2013) and shown to be an anaerobic propionate-producer (Qiu et al., 2014; Ueki et al., 2006), *Bacteroides* (WWT and GL microcosms), a well-documented complex carbohydrate degrader in a wide range of environments (Adamberg et al., 2015; Dongowski, Lorenz & Anger, 2000; Jiménez, Chaves-Moreno & Van Elsas, 2015), *Barnesiella* (GL microcosms), a known fermentative gut microbe (Wang et al., 2015), and WCHB1-69 (ZDT microcosms), a yet-uncultured Bacteroidetes family previously encountered in




Figure 2 Microbial community structure analysis in the enrichment microcosms ($n = 26$) as compared to the pre-enrichment inoculum sources ($n = 3$). The inoculum sources are denoted by shapes; ZDT (○), WWT (●), and GL (■), and the algae types are denoted by color; *Chara* (blue), *Chlorella* (green), Kelp (red), and no algae, i.e., pre-enrichment community, (black). (continued on next page...)

Figure 2 (...continued)

Each enrichment condition (inoculum source \times algae type) is represented by 3 sample points corresponding to the weeks during enrichment, except for GL-kelp enrichment where the dataset from week 4 is not shown due to the small number of sequences obtained with this dataset. (A) Non-metric multidimensional scaling plots based on Bray–Curtis dissimilarity indices at the species level (0.03). For *Chara* and *Chlorella* enrichments, communities grouped by the inoculum source, while Kelp enrichments grouped by the algae type. (B) Canonical correspondence analysis using the abundant phyla/classes relative abundances to study the effect of algae type and inoculum source on the microbial community composition. Here, the same pattern is observed at the phylum/class level, where the community structure of *Chara* and *Chlorella* enrichments were similar and grouped by inoculum source, while the microbial community of Kelp enrichments were quite distinct and grouped together regardless of the inoculum source. This pattern is reflected on the direction of the factors arrows, where the algae type is pointing in the direction of the Kelp enrichments. The CCA also depicts the abundant phyla/classes that seem to shape the microbial community in the different enrichments; Gamma-Proteobacteria in GL *Chara* and *Chlorella* enrichments, Spirochaetes and Firmicutes in ZDT-*Chlorella* enrichment, Delta-Proteobacteria and Bacteroidetes in ZDT-*Chara* enrichments and WWT *Chara* and *Chlorella* enrichments, and Epsilon-Proteobacteria and Firmicutes in Kelp enrichments regardless of the inoculum source. The constrained variables explained 57% of the variance.

organic solvent-contaminated aquifers and anaerobic digestors (Dojka et al., 1998; Xu et al., 2012) (Table 3 and Figs. 3–5).

Within the Firmicutes, all enriched taxa belonged to the order Clostridiales, a ubiquitous order of strictly anaerobic, fermentative bacteria (Xia et al., 2015). However, the profile of enriched families/genera within this order depended on the inoculum source. Members of *Ruminococcaceae* were abundant in all microcosms, while members of the families *Clostridiaceae*_1 and Family XIII were enriched only in ZDT microcosms, and members of the *Lachnospiraceae* and *Veillonellaceae* were enriched only in GL microcosms (Table 3 and Figs. 3–5).

Within the Delta-Proteobacteria, the sulfate-reducing genera *Desulfovibrio*, *Desulfobacter*, *Desulfobulbous*, and *Desulfomicrobium* were encountered as predominant members in enrichments from some or all inoculum sources. Enrichment of sulfate reducers in ZDT and WWT microcosms was accompanied by a significant decrease in the amount of sulfate in the enrichments (Fig. S2). On the other hand, sulfate concentration did not decrease in GL microcosms (Fig. S2) in spite of the apparent enrichment of SRBs (6.9% of the total enriched taxa). Similar results were previously shown for members of *Desulfovibrio* and *Desulfobulbous* in biofilms (Santegoeds et al., 1998), where not all SRBs detected by culture-independent techniques were found to be sulfidogenically active.

Members of the Spirochaetes were enriched in WWT and ZDT microcosms. The genus *Spirochaeta* and the yet uncultured family SHA-4 were identified as the major enriched Spirochaetes members in both enrichments. Both lineages appear to be widely distributed in a wide array of freshwater and marine habitats and enrichments (Bozo-Hurtado et al., 2013; Gu et al., 2004; Leschine, Paster & Canale-Parola, 2006; Wang et al., 2014).

Members of the Gamma-Proteobacteria were enriched in ZDT and GL microcosms. However, the identity of enriched families/genera differed depending on the inoculum source, where *Kluyvera* and unclassified *Enterobacteriaceae* were enriched in ZDT microcosms, while *Buttiauxella*, *Pantoea* and *Aeromonas* were enriched in GL microcosms. All such members are known carbohydrate fermenters previously encountered in microbial




Figure 3 Microbial community composition in ZDT enrichments. Abundant phyla/classes are shown as area charts for *Chara* (i), *Chlorella* (ii), and Kelp (iii) enrichments for each inoculum source. Phyla that constituted 5% or more of the community at any time during enrichment were considered significant to the degradation process and are shown in the area charts. These include phyla that were abundant prior to enrichment and remained abundant during and after enrichment (e.g., Bacteroidetes in *Chara* and *Chlorella* enrichments (i, ii)), and phyla that significantly and progressively increased in abundance with enrichment time (e.g., Firmicutes in Kelp enrichments (iii)). Bar charts depict the relative abundance of abundant genera (>1%) in each of the abundant phyla/classes shown in i-ii-iii. These include Proteobacteria (iv), Bacteroidetes (v), Firmicutes (vi), and Spirochaetes (vii). The X-axis denotes the weeks of enrichment (i-iii), or the weeks of enrichment and algae type (iv-vii). “0” denotes the community composition in the pre-enrichment inoculum source.




Figure 4 Microbial community composition in WWT enrichments. Abundant phyla/classes are shown as area charts for *Chara* (i), *Chlorella* (ii), and Kelp (iii) enrichments for each inoculum source. Phyla that constituted 5% or more of the community at any time during enrichment were considered significant to the degradation process and are shown in the area charts. These include phyla that were abundant prior to enrichment and remained abundant during and after enrichment (e.g., Bacteroidetes in *Chara* and *Chlorella* enrichments (i, ii)), phyla that transiently increased in abundance during part of the enrichment but then decreased in abundance by the end of enrichment (e.g., Delta-Proteobacteria in *Chara* and *Chlorella* enrichments (i, ii)), and phyla that significantly and progressively increased in abundance with enrichment time (e.g., Firmicutes in Kelp enrichments (iii)). Bar charts depict the relative abundance of abundant genera (>1%) in each of the abundant phyla/classes shown in i-ii-iii. These include Proteobacteria (iv), Firmicutes (v), Bacteroidetes (vi), and Spirochaetes (vii). The X-axis denotes the weeks of enrichment (i-iii), or the weeks of enrichment and algae type (iv-vii). “0” denotes the community composition in the pre-enrichment inoculum source.




Figure 5 (...continued)

Bar charts depict the relative abundance of abundant genera (>1%) in each of the abundant phyla/classes shown in i-ii-iii. These include Bacteroidetes (iv), Firmicutes (v), Delta and Epsilon-Proteobacteria (vi), Gamma-Proteobacteria (vii), Alpha and Beta Proteobacteria (viii), and Planctomycetes (ix). The X-axis denotes the weeks of enrichment (i-iii), or the weeks of enrichment and algae type (iv-ix). "0" denotes the community composition in the pre-enrichment inoculum source.

consortia degrading plant biomass (Jiménez, Chaves-Moreno & Van Elsas, 2015; Jiménez et al., 2016), in earthworm gut enrichments (Wust, Horn & Drake, 2011), and in microbial mats from bicarbonate- and ferrous-iron-rich spring (Hegler et al., 2012).

***Chlorella* microcosms**

Enrichment patterns in *Chlorella* microcosms were very similar to *Chara* enrichments; with the phyla Bacteroidetes, Firmicutes, and Delta-Proteobacteria consistently enriched in microcosms derived from all three inoculum sources (ZDT, WWT, and GL), Spirochaetes only enriched in WWT and ZDT microcosms, and Gamma-Proteobacteria enriched in GL (but not ZDT) enrichments. Similar to *Chara* enrichments, the taxa VadinBC27, *Mangroviflexus*, *Paludibacter*, *Barnesiella*, and WCHB1-69 within the Bacteroidetes; *Desulfovibrio*, *Desulfobacter*, and *Desulfomicrobium* within the Delta Proteobacteria; *Spirochaeta* and unclassified SHA-4 within the Spirochaetes were all abundant community members at the end of enrichment. Within the Firmicutes, the family *Lachnospiraceae* was abundant in all enrichments, similar to what was observed in *Chara* microcosms. However, apart from this notable exception, the enriched community of Firmicutes genera/families differed in *Chlorella* microcosms when compared to *Chara* enrichments. Within the ZDT microcosms on *Chlorella*, a wide range of Clostridiales-affiliated genera and families were encountered, with members of the genera *Geosporobacter* (family *Clostridiaceae_1*), and *Acidaminobacter* (family *Clostridiaceae_4*), *Youngiibacter* (family *Clostridiaceae_1*), and members of Clostridiales Family XIII constituting ~34% of total sequences encountered in ZDT microcosms. Further, In contrast to *Chara* enrichments where *Veillonellaceae* was only restricted to GL microcosms, *Chlorella* enrichments selected for members of this family in ZDT and WWT microcosms.

Chlorella enrichments selected for members of the Gamma-Proteobacteria only in GL microcosms where they constituted ~54% of the total taxa in these enrichments. *Buttiauxella* and *Aeromonas* were identified as major taxa in GL *Chlorella* microcosms, similar to what was observed in *Chara* enrichments. In addition, members of the genus *Edwardsiella* (family *Enterobacteriaceae*) were identified as a *Chlorella* enrichment-specific taxon (Figs. 3–5 and Table 3). Members of the genus *Edwardsiella* have been repeatedly isolated from marine and freshwater animals and some species have been linked to pathogenesis in fish (Sakazaki, 1965). This is consistent with its enrichment in microcosms from a freshwater environment such as Grand Lake.

***Kelp* enrichments**

While the microbial communities enriched on *Chara* and *Chlorella* exhibited marked similarities regardless of the inoculum source, the community enriched on kelp was quite distinct: In all kelp enrichments, Firmicutes constituted more than 70% of the

Table 3 Abundant lineages (>1%) within the abundant/enriched phyla shown in Figs. 3–5.

Phylum/Class	Class/Order	Family–genus	WWT	ZDT	GL
Chara enrichments					
Bacteroidetes	Bacteroidales	<i>Marinilabiaceaea-Mangroviflexus</i>	0.58	4.16	0
		<i>Porphyromonadaceae-Paludibacter</i>	1.47	1	0.1
		<i>Porphyromonadaceae-Bacteroides</i>	2.11	0.05	6.91
		<i>Porphyromonadaceae-Barnesiella</i>	0	0	1.58
		<i>Rikenellaceae-VadinBC27</i>	6.1	11.45	2.78
	Sphingobacteriales	WCHB1-69-unclassified	0.76	3.73	0.84
	Unclassified Bacteroidetes		14.34	4.03	0
Firmicutes	Clostridiales	<i>Clostridiaceae_1-Youngiibacter</i>	0.003	1.36	0
		Family XIII	0.41	1.85	0.1
		<i>Ruminococcaceae_Incertae_Sedis</i>	0.07	0.09	1.61
		<i>Ruminococcaceae-Ruminococcus</i>	0.02	0.03	2.38
		Other <i>Ruminococcaceae</i>	3.62	1.47	0
		<i>Lachnospiraceae_Incertae_Sedis</i>	0.16	0.45	6.87
		<i>Lachnospiraceae-Parasporobacterium-Sporobacterium</i>	0	0.04	2.51
		<i>Veillonellaceae-uncultured</i>	0	0	3.1
		Unclassified Clostridiales	3.75	3.4	0.07
	Unclassified Firmicutes		0.69	8.02	0
Spirochaetes	Spirochaetales	<i>Spirochaetaceae-Spirochaeta</i>	10.67	4.69	0
		SHA-4-unclassified	2.92	2.48	0
	Unclassified		6.91	0.29	0
Delta Proteobacteria	Desulfobacterales	<i>Desulfobacteriaceae-Desulfobacter</i>	0.36	1.39	0
		<i>Desulfobulbaceae-Desulfobulbous</i>	0.28	0.69	1.22
	Desulfovibrionales	<i>Desulfovibrionaceae-Desulfovibrio</i>	5.69	1.42	5.67
		<i>Desulfovibrionaceae-Desulfomicrobium</i>	1.89	5.69	0
Gamma Proteobacteria	Enterobacteriales	<i>Enterobacteriaceae-Kluyvera</i>	0	4.15	0.04
		<i>Enterobacteriaceae-unclassified</i>	0	4.2	0
		<i>Enterobacteriaceae-Buttiauxella</i>	0	0	44.1
		<i>Enterobacteriaceae-Pantoea</i>	0	0	11
	Aeromonadales	<i>Aeromonadaceae-Aeromonas</i>	0	0.11	1.87
Chlorella enrichments					
Bacteroidetes	Bacteroidales	<i>Marinilabiaceaea-Mangroviflexus</i>	0.16	2.27	0
		<i>Porphyromonadaceae-Paludibacter</i>	2.59	0.46	0.81
		<i>Porphyromonadaceae-Barnesiella</i>	0.002	0	1.7
		<i>Rikenellaceae-VadinBC27</i>	16.18	9.11	1.95
		Other	1.78	0.2	2.95
	Sphingobacteriales	WCHB1-69-unclassified	2.14	0.64	5.18
	Unclassified Bacteroidetes		2.69	1.18	0
Firmicutes	Clostridia/Clostridiales	<i>Clostridiaceae_1-Youngiibacter</i>	0.006	8.94	0
		<i>Clostridiaceae_4-Geosporobacter</i>	0	11.44	0

(continued on next page)

Table 3 (continued)

Phylum/Class	Class/Order	Family-genus	WWT	ZDT	GL
		Family_XII- <i>Acidaminobacter</i>	0	3.77	0
		Family XIII	0.67	6.32	0.02
		<i>Lachnospiraceae</i> - <i>Incertae_Sedis</i>	0.03	0.06	7.63
		<i>Veillonellaceae</i> -uncultured	1.56	1.69	0.94
		Unclassified Clostridiales	1.78	2.44	0
	Unclassified Firmicutes		0.82	4.87	0
Spirochaetes	Spirochaetales	<i>Spirochaetaceae</i> - <i>Spirochaeta</i>	21.65	2.62	0
		SHA-4-unclassified	5.87	2.35	0
	Unclassified		0.3	0.86	0
Delta Proteobacteria	Desulfobacterales	<i>Desulfobacteriaceae</i> - <i>Desulfobacter</i>	0.03	7.31	0
	Desulfovibrionales	<i>Desulfovibrionaceae</i> - <i>Desulfovibrio</i>	2.93	1.18	21.38
		<i>Desulfovibrionaceae</i> - <i>Desulfomicrobium</i>	0.7	6.38	0
Gamma Proteobacteria	Enterobacteriales	<i>Enterobacteriaceae</i> - <i>Buttiauxella</i>	0	0	37.75
		<i>Enterobacteriaceae</i> - <i>Edwardsiella</i>	0	0	7.64
	Aeromonadales	<i>Aeromonadaceae</i> - <i>Aeromonas</i>	0	0	8.14
Kelp enrichments					
Epsilon Proteobacteria	Campylobacterales	<i>Campylobacteraceae</i> - <i>Arcobacter</i>	8.54	0.01	0
		<i>Campylobacteraceae</i> - <i>Sulfurospirillum</i>	2.19	0.05	0.007
Gamma Proteobacteria	Aeromonadales	<i>Aeromonadaceae</i> - <i>Tolomonas</i>	5.37	0.002	0
	Enterobacteriales	<i>Enterobacteriaceae</i> - <i>Kluyvera</i>	2.95	0.006	0
		<i>Enterobacteriaceae</i> -unclassified	1.71	0	0
	Pseudomonadales	<i>Moraxellaceae</i> - <i>Acinetobacter</i>	2.41	0	0.91
	Other		2	0.992	5.09
Firmicutes	Clostridiales	<i>Clostridiaceae</i> - <i>Clostridium</i>	0.49	29.55	77.73
		<i>Lachnospiraceae</i> - <i>Incertae_Sedis</i>	0.02	19.56	0.68
		<i>Lachnospiraceae</i> - <i>Anaerospobacter</i>	0.004	8.1	0
		<i>Veillonellaceae</i> -unclassified	65.89	0.01	0
	Bacillales	<i>Paenibacillaceae</i> - <i>Paenibacillus</i>	0	2.1	0

total enriched taxa regardless of the inoculum source. In ZDT kelp enrichments, four different Firmicutes taxa were enriched; *Clostridium*, *Anaerospobacter*, *Lachnospiraceae*-*Incertae_Sedis*, and *Paenibacillus*. *Anaerospobacter*, a strictly anaerobic spore former, and other *Lachnospiraceae* members were previously isolated from cellulose and xylan-pectin enrichments of cow feces (Ziemer, 2014), and are frequently encountered within the human gut microbiota (Gagen et al., 2015; Lau et al., 2016; Martinez et al., 2013; Nava, Friedrichsen & Stappenbeck, 2011). Members of the genus *Paenibacillus* are globally distributed facultative anaerobes (Li et al., 2014), some of which are known to exhibit superior plant biomass degradation capacities (Eida et al., 2012). On the other hand, in both GL and WWT kelp microcosms, a single lineage constituted the majority of the enriched Firmicutes; Genus *Clostridium* in GL, and Family *Veillonellaceae* in WWT. Members of the genus *Clostridium* exhibit ubiquitous and global distribution in a wide range of anoxic habitats, while members of the family *Veillonellaceae* are often encountered in groundwater samples (Mosher et al., 2012), and rice paddy soil (Li et al., 2011).

In addition to Firmicutes, ZDT kelp microcosms showed an abundance of the Spirochaetes genus *Treponema* (20.2% of the total enriched taxa) previously shown to contribute to the overall cellulolytic activities in barley straw microcosms (*Kudo, Cheng & Costerton, 1987*), and WWT kelp microcosms showed an abundance of members of Epsilon (genera *Arcobacter* and *Sulfurospirillum*) and Gamma (genera *Tolumonas*, *Kluyvera*, and *Acinetobacter*) Proteobacteria, collectively comprising ~25% of the total enriched taxa. Members of these genera were previously implicated in anaerobic plant biomass degradation (*Billings et al., 2015; Caldwell et al., 2011; Cardoso et al., 2012*).

DISCUSSION

In this study we investigated the microbial community mediating algal detritus turnover under anaerobic conditions. We utilized three representative algal species: *Chlorella vulgaris* strain UTEX2714 representing the Chlorophyta, *Chara* sp. strain IWP representing the Charophyceae, and *Ascophyllum nodosum* (kelp) representing the brown algae (Phaeophyceae). We followed the turnover of these algae in enrichments that were set up using three different sources of inoculum: an anoxic freshwater sulfide- and sulfur-rich spring (Zodletone spring, OK), a wastewater treatment plant (Municipal wastewater treatment plant in Stillwater, OK), and a seasonally stratified lake that experience seasonal algal blooms (Grand Lake O' the Cherokees, OK). We identified multiple microbial lineages that were significantly enriched in these treatments. Some of these lineages appear to be substrate-specific (i.e., enriched when using a specific algal species as a substrate source regardless of the inoculum source utilized, e.g., VadinBC27 that was enriched on *Chara* and *Chlorella* regardless of the inoculum source and *Spirochaeta* that was enriched on *Chara* and *Chlorella* in ZDT and WWT microcosms), habitat-specific (i.e., enriched only when using a specific source of inoculum regardless of the algal substrate utilized, e.g., *Buttiauxella*, that was enriched in GL microcosms regardless of the algal substrate), or treatment-specific (i.e., encountered only in a specific algal substrate/inoculum source combination, e.g., *Arcobacter* in WWT microcosms on kelp, *Geosporobacter*, *Acidaminobacter*, *Anaerosporobacter*, and *Treponema* in ZDT microcosms on kelp, *Youngiibacter* in ZDT microcosms on *Chlorella*, and *Pantoea* in GL microcosms on *Chara*).

Within all nine treatments examined, a high level of diversity was invariably retained at the conclusion of the incubation process. We reason that this is a reflection of the complexity of the substrate utilized. Algal detritus harbors multiple complex macromolecules, e.g., proteins, lipids, nucleic acids, and polysaccharide, that vary considerably in structure and hence require multiple enzymes and pathways for their efficient degradation (e.g., pectin and cellulose in algal cell walls require an arsenal of degradation enzymes (*Abbott & Boraston, 2008; Doi & Kosugi, 2004*)). Such level of complexity could potentially select for a wide range of organisms, each contributing to the degradation process of a specific substrate within the algal biomass. This is in stark contrast to the selection of one/few microbial lineages in anaerobic incubations conducted using a single, chemically defined substrate (*Viggor et al., 2013; Yagi et al., 2010*).

Our results and subsequent community analysis (Figs. 2–5 and Table 3) indicate that kelp enriched for a highly similar microbial community that is mostly composed of members of the order Clostridiales; genus *Clostridium* and *Anaerosporebacter* and family *Veillonellaceae*, regardless of the inoculum source (ZDT, WWT, and GL). While only a handful of environments were examined in this study, the consistent selection for members of a specific lineage regardless of the starting inoculum suggests the ecological significance of this lineage in kelp detritus turnover in anaerobic habitats. The reason for this observed pattern of Clostridiales genera/families selection on kelp could only be speculated upon. A possible contributing factor could be the unique cell wall structure of kelp (or brown algae); multiple cellulose microfibrils layers embedded in large interfibrillar matrices that are mostly composed of alginates and fucans (Domozych, 2001; Youssef et al., 2015). Alginate (Preiss & Ashwell, 1962a; Preiss & Ashwell, 1962b) and fucans (Descamps et al., 2006; Kusaykin et al., 2016) degradation requires highly specific enzymes machineries. Organisms with alginate or fucan/fucoidan-degradation capabilities under aerobic conditions have been previously isolated (Ekborg et al., 2005; Jagtap et al., 2014; Park et al., 2012; Sakai, Kawai & Kato, 2004; Thomas et al., 2012; Yonemoto et al., 1993). On the other hand, with the exception of a few studies that used anaerobic batch-fed mixed inocula to degrade brown algae and produce methane (Moen, Horn & Østgaard, 1997a; Moen, Horn & Østgaard, 1997b; Sutherland & Varela, 2014), there is a scarcity of information on the identity of the degrading inocula under anaerobic conditions. In contrast to the number of studies on the anaerobic degradation of other common polysaccharides, e.g., cellulose and xylans, a single study by Kita et al. (2016) reported on the identity of a bacterial consortium (formed mainly of a *Clostridiaceae* bacterium and a *Porphyromonadaceae* bacterium (*Dysgonomonas capnocytophagoides*)) anaerobically degrading alginate. Based on the study by Kita et al. (2016) and the results we report here, it is possible that members of the Clostridiales represent one of very few members possessing alginate and/or fucan-degrading capabilities and that are readily enriched and propagated under laboratory incubations.

On the other hand, when using *Chara* or *Chlorella* as an algal inoculum, the final microbial community enriched was highly divergent, and the final community structure was mostly dependent on the inoculum sources (ZDT, WWT, GL), rather than the type of algal substrate provided (Figs. 2–5, Table 3). While *Chlorella* and *Chara* cell walls are quite distinct, they are both similar in being rich in fibrous cellulose and/or hemicellulose with amorphous middle layers composed mainly of pectin (homogalacturonic and rhamnogalacturonic acids polymers) in *Charophyta*, or sulfated polysaccharides in *Chlorophyceae* (Domozych, 2001; Domozych et al., 2014; Youssef et al., 2015). We reason that the relative similarity of the communities enriched on both types of algae, as well as the enrichment for multiple, rather than a single group of microbial lineages (*VadinBC27*, *Spirochaeta*, *Lachnospiraceae*, *Buttauxiella*, and *Pantoea*) is a reflection of the relative ubiquity of microbial lineages capable of the anaerobic degradation of cellulose, hemicellulose, and pectin in the algal cell walls, hence allowing ready access to the intracellular substrates within the algal cells.

Under anaerobic conditions, multiple groups of organisms and metabolic guilds are often required for the effective and complete degradation of complex organic molecules (McInerney, Sieber & Gunsalus, 2009; Morris et al., 2013). By examining the known metabolic capabilities of close relatives of lineages enriched in various treatments, one could propose a model depicting their putative involvement in the complex algal detritus degradation processes. In kelp enrichments, complex carbohydrate polymer degradation to monomers could possibly be mediated by various members of the order Clostridiales (*Clostridium*, *Anaerosporebacter*, *Lachnospiraceae* incertae sedis) as shown before (Ziemer, 2014), as well as the fermentative bacteria in the *Enterobacteriaceae* (*Kluyvera*) (*Xin & He*, 2013). Produced sugar monomers can be further fermented to various fatty acids (acetate and longer chain fatty acids e.g., butyrate, propionate, etc.) by the same members of the Clostridiales and *Enterobacteriaceae*, as well as the Epsilon Proteobacterium *Sulfurospirillum* (Stolz et al., 1999). Proteins in the initial substrate could potentially be degraded by the Epsilon Proteobacterium *Arcobacter* (Roalkvam et al., 2015). Additionally, while the majority of sequences obtained were bacterial in origin, the few archaeal sequences obtained suggest the enrichment of members of Bathyarchaeota (Table S2). Previous research using genomic sequences of different members of the Bathyarchaeota suggested their involvement in both complex carbohydrates and detrital protein degradation as well as acetate production (Lazar et al., 2016), which could explain their enrichment on kelp. Under anaerobic condition, syntrophic organisms convert the long chain fatty acids produced from the initial polymer degradation to acetate. Definitive identification of syntrophic organisms in culture-independent studies is challenging, given their close phylogenetic affiliation with fermentative lineages (Morris et al., 2013). On the other hand, saccharolytic clostridia members of the family *Lachnospiraceae* could potentially perform the initial breakdown of polymeric substances and the fermentation of the resulting sugars to acetate, hydrogen, and CO₂ (Krumholz & Bryant, 1986). While other obligate syntrophic organisms, e.g., members of the families Syntrophobacteraceae, Syntrophaceae, Syntrophomonadaceae, and Syntrophorhabdaceae, were detected in very low percentage (<0.06% of the total community in any enrichment), their role could not be ruled out. The produced acetate, hydrogen, and CO₂ would eventually be converted to methane by methanogens. The role of methanogens as the dominant terminal electron acceptor in kelp enrichment from ZDT and WWT inoculum sources is suggested by the observed increase in *mrcA* gene copy number in qPCR analysis (Fig. 1) and the identification of several sequences affiliated with known methanogens (genera *Methanosarcina*, *Methanothermococcus*, *Methanogenium*, and *Methanomicrobium*) in kelp enrichment from ZDT (Table S2). The lack of sulfate utilization in all kelp enrichments (Fig. S2) argues against the involvement of the SRBs identified in the culture-independent dataset (*Desulfovibrio*, *Desulfobacter*, *Desulfomicrobium*, and *Desulfobulbous*) and detected by qPCR (Fig. 1) in the process. Similar results were previously shown in biofilms (Santegoeds et al., 1998), where not all SRBs detected by culture-independent techniques were found to be sulfidogenically active.

Within *Chara* and *Chlorella* enrichments, complex carbohydrate (e.g., cellulose, pectin, hemicellulose) degradation to sugar monomers could be mediated by members of the

Bacteroidetes uncultured groups VadinBC27 (in all enrichments from all sources) and WCHB1-69 (in *Chara* enrichment from ZDT, and *Chlorella* enrichment from WWT and GL), as well as the Spirochaetes ([Gao, Xu & Ruan, 2014](#)) (genus *Spirochaeta* and the uncultured group SHA-4 enriched on *Chara* and *Chlorella* from ZDT and WWT sources). These lineages have been consistently enriched in anaerobic sludge digestors ([Godon et al., 1997](#); [Lee et al., 2013](#)), and microcosms with hydrocarbon or halogenated solvents ([Dojka et al., 1998](#); [Gu et al., 2004](#); [Xu et al., 2012](#)). Similarly, members of the Clostridiales (Family *Ruminococcaceae* (in *Chara* enrichments from all sources), Family *Veillonellaceae* (in *Chara* enrichments from GL), Family *Lachnospiraceae* (in *Chara* and *Chlorella* enrichments from GL), and Family *Clostridiaceae* genera *Yongiibacter*, *Geosporobacter*, *Acidaminobacter* (in *Chlorella* enrichments from ZDT)), as well as Enterobacteriales (Genera *Kluyvera* (*Chara* enrichments from ZDT), *Pantoea* (*Chara* enrichments from GL), *Edwardsiella* and *Aeromonas* (*Chlorella* enrichment from GL), and *Buttiauxella* (*Chara* and *Chlorella* enrichments from GL)) could potentially mediate complex carbohydrate degradation ([Hegler et al., 2012](#); [Jiménez, Chaves-Moreno & Van Elsas, 2015](#); [Jiménez et al., 2016](#); [Sakazaki, 1965](#); [Wust, Horn & Drake, 2011](#); [Xin & He, 2013](#); [Ziemer, 2014](#)). The monomers produced could potentially be converted to long chain volatile fatty acids, acetate, and H₂ by the Clostridiales and Enterobacteriales members above. Alternatively, long chain volatile fatty acids could be converted to acetate, H₂ and CO₂ by syntrophs, or oxidized either completely (to H₂ and CO₂) or incompletely (to acetate, H₂ and CO₂) by sulfate-reducing bacteria (e.g., the complete oxidizers (*Desulfobacter* in *Chara* and *Chlorella* enrichments from ZDT), or the incomplete oxidizers (*Desulfovibrio* in all *Chara* and *Chlorella* enrichments from all sources, *Desulfomicrobium* in *Chara* and *Chlorella* enrichments from ZDT and *Chara* enrichments from WWT, and *Desulfobulbous* in *Chara* enrichment from GL)) when sulfate is available. The produced acetate, H₂ and CO₂ could either be metabolized to methane by acetoclastic or hydrogenotrophic methanogenic lineages observed in the enrichments ([Table S2](#)) (e.g., the acetoclastic *Methanosarcina* in *Chara* and *Chlorella* enrichments from ZDT and WWT, and the hydrogenotrophic *Methanothermococcus* in *Chlorella* early (week 7) enrichments from ZDT), or metabolized by the acetoclastic autotrophic SRBs in the presence of sulfate. The increase in *dsr* copy numbers in *Chara* and *Chlorella* enrichments as measured by qPCR, the utilization of the available substrates in these enrichments (loss of sulfate ([Fig. S2](#))), as well as the presence of a large and diverse community of SRBs ([Table 3](#) and [Table S2](#)) evidenced by the culture-independent analysis, strongly argue for the co-involvement of sulfate reduction and methanogenesis as two competing terminal electron accepting processes in these enrichments. Recently, the methanogenic potential for members of the Bathyarchaeota was suggested based on genomic metabolic reconstruction ([Evans et al., 2015](#)). It is worth noting that the Bathyarchaeota phylum was enriched in ZDT *Chara* and *Chlorella* microcosms ([Table S2](#)) and could potentially be contributing to methanogenesis in these enrichments.

In conclusion, our work represents the first systematic survey of microbial communities mediating turnover of algal biomass under anaerobic conditions, and highlights the diversity of lineages putatively involved in the degradation process. The results presented here could certainly open the door for future studies that investigate the interactions

between the abundant genera identified as significant for the degradation process, as well as for targeted isolation studies for algal detritus degraders.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work has been supported by the National Science Foundation Microbial Observatories program (grant no. MCB_0240683), and the Oklahoma State University Start-up fund (NY). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Science Foundation Microbial Observatories program: MCB_0240683.

Oklahoma State University Start-up fund.

Competing Interests

The authors declare there are no competing interests.

Richard M. Zamor and Steve Nikolai are employees of Grand River Dam Authority (GRDA).

Author Contributions

- Jessica M. Morrison performed the experiments, analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Chelsea L. Murphy performed the experiments, analyzed the data, reviewed drafts of the paper.
- Kristina Baker performed the experiments, reviewed drafts of the paper.
- Richard M. Zamor and Steve J. Nikolai contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Shawn Wilder performed the experiments, contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Mostafa S. Elshahed conceived and designed the experiments, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Noha H. Youssef conceived and designed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.

DNA Deposition

The following information was supplied regarding the deposition of DNA sequences:

Genbank accession number [SRP083898](#).

Data Availability

The following information was supplied regarding data availability:

The research in this article did not generate, collect or analyse any raw data or code.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2803#supplemental-information>.

REFERENCES

- Abbott DW, Boraston AB. 2008.** Structural biology of pectin degradation by Enterobacteriaceae. *Microbiology and Molecular Biology Reviews* **72**:301–316
DOI [10.1128/MMBR.00038-07](https://doi.org/10.1128/MMBR.00038-07).
- Adamberg K, Tomson K, Talve T, Pudova K, Puurand M, Visnapuu T, Alamäe T, Adamberg S. 2015.** Levan enhances associated growth of *Bacteroides*, *Escherichia*, *Streptococcus* and *Faecalibacterium* in fecal microbiota. *PLOS ONE* **10**:e0144042
DOI [10.1371/journal.pone.0144042](https://doi.org/10.1371/journal.pone.0144042).
- Amaral-Zettler L. 2011.** Algae. In: Gargaud M, Amils R, Quintanilla JC, Cleaves HJ, Irvine WM, Pinti DL, Viso M, eds. *Encyclopedia of Astrobiology*. Berlin, Heidelberg: Springer, 27–28.
- Amin SA, Parker MS, Armbrust EV. 2012.** Interactions between Diatoms and Bacteria. *Microbiology and Molecular Biology Reviews* **76**:667–684
DOI [10.1128/MMBR.00007-12](https://doi.org/10.1128/MMBR.00007-12).
- Angel R, Claus P, Conrad R. 2012.** Methanogenic archaea are globally ubiquitous in aerated soils and become active under wet anoxic conditions. *ISME Journal* **6**:847–862 DOI [10.1038/ismej.2011.141](https://doi.org/10.1038/ismej.2011.141).
- Ask J, Karlsson J, Persson L, Ask P, Byström P, Jansson M. 2009.** Whole-lake estimates of carbon flux through algae and bacteria in benthic and pelagic habitats of clear-water lakes. *Ecology* **90**:1923–1932 DOI [10.1890/07-1855.1](https://doi.org/10.1890/07-1855.1).
- Bagatini IL, Eiler A, Bertilsson S, Klaveness D, Tessarolli LP, Vieira AAH. 2014.** Host-specificity and dynamics in bacterial communities associated with bloom-forming freshwater phytoplankton. *PLOS ONE* **9**:e85950
DOI [10.1371/journal.pone.0085950](https://doi.org/10.1371/journal.pone.0085950).
- Balk M, Keuskamp JA, Laanbroek HJ. 2015.** Potential activity, size, and structure of sulfate-reducing microbial communities in an exposed, grazed and a sheltered, non-grazed mangrove stand at the Red Sea Coast. *Frontiers in Microbiology* **6**: Article 1478 DOI [10.3389/fmicb.2015.01478](https://doi.org/10.3389/fmicb.2015.01478).
- Billings AF, Fortney JL, Hazen TC, Simmons B, Davenport KW, Goodwin L, Ivanova N, Kyrpides NC, Mavromatis K, Woyke T, DeAngelis KM. 2015.** Genome sequence and description of the anaerobic lignin-degrading bacterium *Tolomonas lignolytica* sp. nov. *Standards in Genomic Sciences* **10**: Article 106
DOI [10.1186/s40793-015-0100-3](https://doi.org/10.1186/s40793-015-0100-3).
- Bohutskyi P, Ketter B, Chow S, Adams KJ, Betenbaugh MJ, Allnut FCT, Bouwer EJ. 2015.** Anaerobic digestion of lipid-extracted *Auxenochlorella protothecoides* biomass for methane generation and nutrient recovery. *Bioresource Technology* **183**:229–239
DOI [10.1016/j.biortech.2015.02.012](https://doi.org/10.1016/j.biortech.2015.02.012).
- Boon AR, Duineveld GCA, Berghuis EM, Van der Weele JA. 1998.** Relationships between benthic activity and the annual phytopigment cycle in near-bottom water

and sediments in the southern North Sea. *Estuarine, Coastal and Shelf Science* 46:1–13.

- Bozo-Hurtado L, García-Amado MA, Chistoserdov A, Varela R, Narvaez JJ, Colwell R, Suárez P. 2013.** Identification of bacteria in enrichment cultures of sulfate reducers in the Cariaco Basin water column employing Denaturing Gradient Gel Electrophoresis of 16S ribosomal RNA gene fragments. *Aquatic Biosystems* 9:1–11 DOI 10.1186/2046-9063-9-1.
- Buchan A, LeCleir GR, Gulvik CA, Gonzalez JM. 2014.** Master recyclers: features and functions of bacteria associated with phytoplankton blooms. *Nature Reviews Microbiology* 12:686–698 DOI 10.1038/nrmicro3326.
- Buhring SI, Sievert SM, Jonkers HM, Ertefai T, Elshahed MS, Krumholz LR, Hinrichs KU. 2011.** Insights into chemotaxonomic composition and carbon cycling of phototrophic communities in an artesian sulfur-rich spring (Zodletone, Oklahoma, USA), a possible analog for ancient microbial mat systems. *Geobiology* 9:166–179.
- Cai H, Jiang H, Krumholz LR, Yang Z. 2014.** Bacterial community composition of size-fractionated aggregates within the phycosphere of cyanobacterial blooms in a eutrophic freshwater lake. *PLOS ONE* 9:e102879 DOI 10.1371/journal.pone.0102879.
- Caldwell ME, Allen TD, Lawson PA, Tanner RS. 2011.** *Tolomonas osonensis* sp. nov., isolated from anoxic freshwater sediment, and emended description of the genus *Tolomonas*. *International Journal of Systematic and Evolutionary Microbiology* 61:2659–2663 DOI 10.1099/ijs.0.023853-0.
- Cannell RJP. 1990.** Algal biotechnology. *Applied Biochemistry and Biotechnology* 26:85–105 DOI 10.1007/BF02798395.
- Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012.** Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME Journal* 6:1621–1624 DOI 10.1038/ismej.2012.8.
- Cardoso AM, Cavalcante JJV, Cantão ME, Thompson CE, Flatschart RB, Glogauer A, Scapin SMN, Sade YB, Beltrão PJMSI, Gerber AL, Martins OB, Garcia ES, De Souza W, Vasconcelos ATR. 2012.** Metagenomic Analysis of the microbiota from the crop of an invasive snail reveals a rich reservoir of novel genes. *PLOS ONE* 7:e48505 DOI 10.1371/journal.pone.0048505.
- Cole JJ. 1982.** Interactions between bacteria and algae in aquatic ecosystems. *Annual Review of Ecology and Systematics* 13:291–314 DOI 10.1146/annurev.es.13.110182.001451.
- Coveley S, Elshahed MS, Youssef NH. 2015.** Response of the rare biosphere to environmental stressors in a highly diverse ecosystem (Zodletone Spring, OK, USA). *PeerJ* 3:e1182 DOI 10.7717/peerj.1182.
- Descamps V, Colin S, Lahaye M, Jam M, Richard C, Potin P, Barbeyron T, Yvin J-C, Kloareg B. 2006.** Isolation and culture of a marine bacterium degrading the sulfated fucans from marine brown algae. *Marine Biotechnology* 8:27–39 DOI 10.1007/s10126-005-5107-0.
- Ding W, Stewart DI, Humphreys PN, Rout SP, Burke IT. 2016.** Role of an organic carbon-rich soil and Fe(III) reduction in reducing the toxicity and environmental

- mobility of chromium(VI) at a COPR disposal site. *Science of the Total Environment* **541**:1191–1199 DOI [10.1016/j.scitotenv.2015.09.150](https://doi.org/10.1016/j.scitotenv.2015.09.150).
- Dittami SM, Duboscq-Bidot L, Perennou M, Gobet A, Corre E, Boyen C, Tonon T. 2016.** Host-microbe interactions as a driver of acclimation to salinity gradients in brown algal cultures. *ISME Journal* **10**:51–63 DOI [10.1038/ismej.2015.104](https://doi.org/10.1038/ismej.2015.104).
- Doi RH, Kosugi A. 2004.** Cellulosomes: plant-cell-wall-degrading enzyme complexes. *Nature Reviews Microbiology* **2**:541–551 DOI [10.1038/nrmicro925](https://doi.org/10.1038/nrmicro925).
- Dojka MA, Hugenholtz P, Haack SK, Pace NR. 1998.** Microbial diversity in a hydrocarbon- and chlorinated-solvent-contaminated aquifer undergoing intrinsic bioremediation. *Applied and Environmental Microbiology* **64**:3869–3877.
- Domozych DS. 2001.** Algal Cell Walls. In: *Encyclopedia of Life sciences*. Hoboken: John Wiley & Sons, Ltd.
- Domozych DS, Sorensen I, Popper ZA, Ochs J, Andreas A, Fangel JU, Pielach A, Sacks C, Brechka H, Ruisi-Besares P, Willats WG, Rose JK. 2014.** Pectin metabolism and assembly in the cell wall of the charophyte green alga *Penium margaritaceum*. *Plant Physiology* **165**:105–118 DOI [10.1104/pp.114.236257](https://doi.org/10.1104/pp.114.236257).
- Dongowski G, Lorenz A, Anger H. 2000.** Degradation of pectins with different degrees of esterification by *Bacteroides thetaiotaomicron* isolated from human gut flora. *Applied and Environmental Microbiology* **66**:1321–1327 DOI [10.1128/AEM.66.4.1321-1327.2000](https://doi.org/10.1128/AEM.66.4.1321-1327.2000).
- Durborow RM. 2014.** Management of aquatic weeds. In: Chauhan BS, Mahajan G, eds. *Recent advances in weed management*. New York: Springer Science+Business Media, 281–314.
- Eida MF, Nagaoka T, Wasaki J, Kouno K. 2012.** Isolation and characterization of cellulose-decomposing bacteria inhabiting sawdust and coffee residue composts. *Microbes and Environments* **27**:226–233 DOI [10.1264/jsme2.ME11299](https://doi.org/10.1264/jsme2.ME11299).
- Eigemann F, Hilt S, Salka I, Grossart HP. 2013.** Bacterial community composition associated with freshwater algae: species specificity vs. dependency on environmental conditions and source community. *FEMS Microbiology Ecology* **83**:650–663 DOI [10.1111/1574-6941.12022](https://doi.org/10.1111/1574-6941.12022).
- Ekborg NA, Gonzalez JM, Howard MB, Taylor LE, Hutcheson SW, Weiner RM. 2005.** *Saccharophagus degradans* gen. nov., sp. nov., a versatile marine degrader of complex polysaccharides. *International Journal of Systematic and Evolutionary Microbiology* **55**:1545–1549 DOI [10.1099/ijs.0.63627-0](https://doi.org/10.1099/ijs.0.63627-0).
- Evans PN, Parks DH, Chadwick GL, Robbins SJ, Orphan VJ, Golding SD, Tyson GW. 2015.** Methane metabolism in the archaeal phylum Bathyarchaeota revealed by genome-centric metagenomics. *Science* **350**:434–438 DOI [10.1126/science.aac7745](https://doi.org/10.1126/science.aac7745).
- Fierer N, Jackson JA, Vilgalys R, Jackson RB. 2005.** Assessment of soil microbial community structure by use of taxon-specific quantitative PCR assays. *Applied and Environmental Microbiology* **71**:4117–4120 DOI [10.1128/AEM.71.7.4117-4120.2005](https://doi.org/10.1128/AEM.71.7.4117-4120.2005).
- Gagen EJ, Padmanabha J, Denman SE, McSweeney CS. 2015.** Hydrogenotrophic culture enrichment reveals rumen Lachnospiraceae and Ruminococcaceae acetogens and hydrogen-responsive *Bacteroidetes* from pasture-fed cattle. *FEMS Microbiology Letters* **362**:fnnv104 DOI [10.1093/femsle/fnnv104](https://doi.org/10.1093/femsle/fnnv104).

- Gao ZM, Xu X, Ruan LW. 2014.** Enrichment and characterization of an anaerobic cellulolytic microbial consortium SQD-1.1 from mangrove soil. *Applied Microbiology and Biotechnology* **98**:465–474 DOI [10.1007/s00253-013-4857-2](https://doi.org/10.1007/s00253-013-4857-2).
- Gies EA, Konwar KM, Beatty JT, Hallam SJ. 2014.** Illuminating microbial dark matter in meromictic Sakinaw Lake. *Applied and Environmental Microbiology* **80**:6807–6818 DOI [10.1128/AEM.01774-14](https://doi.org/10.1128/AEM.01774-14).
- Godon JJ, Zumstein E, Dabert P, Habouzit F, Moletta R. 1997.** Molecular microbial diversity of an anaerobic digester as determined by small-subunit rDNA sequence analysis. *Applied and Environmental Microbiology* **63**:2802–2813.
- Gu AZ, Hedlund BP, Staley JT, Strand SE, Stensel HD. 2004.** Analysis and comparison of the microbial community structures of two enrichment cultures capable of reductively dechlorinating TCE and cis-DCE. *Environmental Microbiology* **6**:45–54.
- Guckert JB, Cooksey KE. 1990.** Triglyceride accumulation and fatty acid profile changes in *Chlorella* (Chlorophyta) during high pH-induced cell cycle inhibition. *Journal of Phycology* **26**:72–79 DOI [10.1111/j.0022-3646.1990.00072.x](https://doi.org/10.1111/j.0022-3646.1990.00072.x).
- Hallegraeff GM. 1993.** A review of harmful algal blooms and their apparent global increase. *Phycologia* **32**:79–99 DOI [10.2216/i0031-8884-32-2-79.1](https://doi.org/10.2216/i0031-8884-32-2-79.1).
- Hasegawa Y, Martin JL, Giewat MW, Rooney-Varga JN. 2007.** Microbial community diversity in the phycosphere of natural populations of the toxic alga, *Alexandrium fundyense*. *Environmental Microbiology* **9**:3108–3121 DOI [10.1111/j.1462-2920.2007.01421.x](https://doi.org/10.1111/j.1462-2920.2007.01421.x).
- Hecky RE, Hesslein RH. 1995.** Contributions of benthic algae to lake food webs as revealed by stable isotope analysis. *Journal of the North American Benthological Society* **14**:631–653 DOI [10.2307/1467546](https://doi.org/10.2307/1467546).
- Hegler F, Lösekann-Behrens T, Hanselmann K, Behrens S, Kappler A. 2012.** Influence of seasonal and geochemical changes on the geomicrobiology of an iron carbonate mineral water spring. *Applied and Environmental Microbiology* **78**:7185–7196 DOI [10.1128/AEM.01440-12](https://doi.org/10.1128/AEM.01440-12).
- Higgins BT, Thornton-Dunwoody A, Labavitch JM, VanderGheynst JS. 2014.** Microplate assay for quantitation of neutral lipids in extracts from microalgae. *Analytical Biochemistry* **465**:81–89 DOI [10.1016/j.ab.2014.07.020](https://doi.org/10.1016/j.ab.2014.07.020).
- Hlavínek P, Strítěský L, Pešoutová R, Houdková L. 2016.** Biogas production from algal biomass from municipal wastewater treatment. *Waste Biomass Valor* **7**:747–752.
- Hoffmann L. 1989.** Algae of terrestrial habitats. *Botanical Review* **55**:77–105 DOI [10.1007/BF02858529](https://doi.org/10.1007/BF02858529).
- Hoshaw RW, Mccourt RM. 1988.** The Zygnemataceae (Chlorophyta)—a 20-year update of research. *Phycologia* **27**:511–548 DOI [10.2216/i0031-8884-27-4-511.1](https://doi.org/10.2216/i0031-8884-27-4-511.1).
- Hu Q, Sommerfeld M, Jarvis E, Ghirardi M, Posewitz M, Seibert M, Darzins A. 2008.** Microalgal triacylglycerols as feedstocks for biofuel production: perspectives and advances. *The Plant Journal* **54**:621–639 DOI [10.1111/j.1365-3113X.2008.03492.x](https://doi.org/10.1111/j.1365-3113X.2008.03492.x).
- Huse SM, Welch DM, Morrison HG, Sogin ML. 2010.** Ironing out the wrinkles in the rare biosphere through improved OTU clustering. *Environmental Microbiology* **12**:1889–1898 DOI [10.1111/j.1462-2920.2010.02193.x](https://doi.org/10.1111/j.1462-2920.2010.02193.x).

- Jagtap SS, Hehemann JH, Polz MF, Lee JK, Zhao H. 2014.** Comparative biochemical characterization of three exolytic oligoalginate lyases from *Vibrio splendidus* reveals complementary substrate scope, temperature, and pH adaptations. *Applied and Environmental Microbiology* **80**:4207–4214 DOI [10.1128/AEM.01285-14](https://doi.org/10.1128/AEM.01285-14).
- Jiménez DJ, Chaves-Moreno D, Van Elsas JD. 2015.** Unveiling the metabolic potential of two soil-derived microbial consortia selected on wheat straw. *Scientific Reports* **5**: Article 13845 DOI [10.1038/srep13845](https://doi.org/10.1038/srep13845).
- Jiménez DJ, De Lima Brossi MJ, Schückel J, Kračun SK, Willats WGT, Van Elsas JD. 2016.** Characterization of three plant biomass-degrading microbial consortia by metagenomics- and metasecretomics-based approaches. *Applied Microbiology and Biotechnology* **100**:10463–10477.
- Jones AC, Liao TSV, Najjar FZ, Roe BA, Hambright KD, Caron DA. 2013.** Seasonality and disturbance: annual pattern and response of the bacterial and microbial eukaryotic assemblages in a freshwater ecosystem. *Environmental Microbiology* **15**:2557–2572 DOI [10.1111/1462-2920.12151](https://doi.org/10.1111/1462-2920.12151).
- Kita A, Miura T, Kawata S, Yamaguchi T, Okamura Y, Aki T, Matsumura Y, Tajima T, Kato J, Nishio N, Nakashimada Y. 2016.** Bacterial community structure and predicted alginate metabolic pathway in an alginate-degrading bacterial consortium. *Journal of Bioscience and Bioengineering* **121**:286–292 DOI [10.1016/j.jbiosc.2015.06.014](https://doi.org/10.1016/j.jbiosc.2015.06.014).
- Krumholz LR, Bryant MP. 1986.** *Syntrophococcus sucromutans* sp. nov. gen. nov. uses carbohydrates as electron donors and formate, methoxybenzenoids or *Methanobrevibacter* as electron acceptor systems. *Archives of Microbiology* **143**:313–318 DOI [10.1007/BF00412795](https://doi.org/10.1007/BF00412795).
- Kudo H, Cheng KJ, Costerton JW. 1987.** Interactions between *Treponema bryantii* and cellulolytic bacteria in the *in vitro* degradation of straw cellulose. *Canadian Journal of Microbiology* **33**:244–248 DOI [10.1139/m87-041](https://doi.org/10.1139/m87-041).
- Kusaykin MI, Silchenko AS, Zakharenko AM, Zvyagintseva TN. 2016.** Fucoidanases. *Glycobiology* **26**:3–12 DOI [10.1093/glycob/cww106](https://doi.org/10.1093/glycob/cww106).
- Lau JT, Whelan FJ, Herath I, Lee CH, Collins SM, Bercik P, Surette MG. 2016.** Capturing the diversity of the human gut microbiota through culture-enriched molecular profiling. *Genome Medicine* **8**: Article 72 DOI [10.1186/s13073-016-0327-7](https://doi.org/10.1186/s13073-016-0327-7).
- Lazar CS, Baker BJ, Seitz K, Hyde AS, Dick GJ, Hinrichs K-U, Teske AP. 2016.** Genomic evidence for distinct carbon substrate preferences and ecological niches of Bathyarchaeota in estuarine sediments. *Environmental Microbiology* **18**:1200–1211 DOI [10.1111/1462-2920.13142](https://doi.org/10.1111/1462-2920.13142).
- Lee S-H, Park J-H, Kang H-J, Lee YH, Lee TJ, Park H-D. 2013.** Distribution and abundance of Spirochaetes in full-scale anaerobic digesters. *Bioresour Technol* **145**:25–32 DOI [10.1016/j.biortech.2013.02.070](https://doi.org/10.1016/j.biortech.2013.02.070).
- Leschine S, Paster BJ, Canale-Parola E. 2006.** Free-living saccharolytic Spirochetes: the genus *Spirochaeta*. In: Dworkin M, Falkow S, Rodsenberg E, Schleifer K-H, Stackebrandt E, eds. *The prokaryotes*. New York: Springer Science + Business Media, LLC, 195–210.

- Li Y-F, Calley JN, Ebert PJ, Helmes EB. 2014. *Paenibacillus lentus* sp. nov., a β -mannanolytic bacterium isolated from mixed soil samples in a selective enrichment using guar gum as the sole carbon source. *International Journal of Systematic and Evolutionary Microbiology* 64:1166–1172 DOI 10.1099/ijs.0.054726-0.
- Li H, Peng J, Weber KA, Zhu Y. 2011. Phylogenetic diversity of Fe(III)-reducing microorganisms in rice paddy soil: enrichment cultures with different short-chain fatty acids as electron donors. *Journal of Soils and Sediments* 11:1234–1242 DOI 10.1007/s11368-011-0371-2.
- Liu C, Li H, Zhang Y, Si D, Chen Q. 2016. Evolution of microbial community along with increasing solid concentration during high-solids anaerobic digestion of sewage sludge. *Bioresource Technology* 216:87–94 DOI 10.1016/j.biortech.2016.05.048.
- Maness N. 2010. Extraction and analysis of soluble carbohydrates. In: Sunkar R, ed. *Plant stress tolerance: methods and protocols*. Totowa: Humana Press, 341–370.
- Mann KH. 1988. Production and use of detritus in various fresh-water, estuarine, and coastal marine ecosystems. *Limnology and Oceanography* 33:910–930.
- Martinez I, Lattimer JM, Hubach KL, Case JA, Yang J, Weber CG, Louk JA, Rose DJ, Kyureghian G, Peterson DA, Haub MD, Walter J. 2013. Gut microbiome composition is linked to whole grain-induced immunological improvements. *ISME Journal* 7:269–280 DOI 10.1038/ismej.2012.104.
- McInerney MJ, Sieber JR, Gunsalus RP. 2009. Syntrophy in anaerobic global carbon cycles. *Current Opinion in Biotechnology* 20:623–632 DOI 10.1016/j.copbio.2009.10.001.
- Moazami N, Ashori A, Ranjbar R, Tangestani M, Eghtesadi R, Nejad AS. 2012. Large-scale biodiesel production using microalgae biomass of *Nannochloropsis*. *Biomass & Bioenergy* 39:449–453 DOI 10.1016/j.biombioe.2012.01.046.
- Moen E, Horn S, Østgaard K. 1997a. Alginate degradation during anaerobic digestion of *Laminaria hyperborea* stipes. *Journal of Applied Phycology* 9:157–166 DOI 10.1023/A:1007966126570.
- Moen E, Horn S, Østgaard K. 1997b. Biological degradation of *Ascophyllum nodosum*. *Journal of Applied Phycology* 9:347–357 DOI 10.1023/A:1007988712929.
- Morris BE, Henneberger R, Huber H, Moissl-Eichinger C. 2013. Microbial syntrophy: interaction for the common good. *FEMS Microbiology Reviews* 37:384–406 DOI 10.1111/1574-6976.12019.
- Mosher JJ, Phelps TJ, Podar M, Hurt RA, Campbell JH, Drake MM, Moberly JG, Schadt CW, Brown SD, Hazen TC, Arkin AP, Palumbo AV, Faybishenko BA, Elias DA. 2012. Microbial community succession during lactate amendment and electron acceptor limitation reveals a predominance of metal-reducing *Pelosinus* spp. *Applied and Environmental Microbiology* 78:2082–2091 DOI 10.1128/AEM.07165-11.
- Mudhoo A. 2012. Mudhoo A, ed. *Biogas production: pretreatment methods in anaerobic digestion*. Hoboken: Wiley.
- Muylaert K, Van der Gucht K, Vloemans N, Meester LD, Gillis M, Vyverman W. 2002. Relationship between bacterial community composition and bottom-up versus top-down variables in four eutrophic shallow lakes. *Applied and Environmental Microbiology* 68:4740–4750 DOI 10.1128/AEM.68.10.4740-4750.2002.

- Nabarlantz D-A, Arenas-Beltrán L-P, Herrera-Soracá D-M, Niño-Bonilla D-A. 2013.** Biogas production by anaerobic digestion of wastewater from palm oil mill industry. *Ciencia Tecnología y Futuro* 5:73–83.
- Nava GM, Friedrichsen HJ, Stappenbeck TS. 2011.** Spatial organization of intestinal microbiota in the mouse ascending colon. *ISME Journal* 5:627–638 DOI 10.1038/ismej.2010.161.
- Omer FK TFAOAT. 2013.** Oil and fatty acid composition of spirogyra and *Chara* species from Beastan SWR Spring water in Sulaimani-Kurdistan region of Iraq. *Egyptian Journal of Experimental Biology (Botany)* 9:159–162.
- Paerl HW, Otten TG. 2013.** Blooms bite the hand that feeds them. *Science* 342:433–434 DOI 10.1126/science.1245276.
- Park HH, Kam N, Lee EY, Kim HS. 2012.** Cloning and characterization of a novel oligoalginate lyase from a newly isolated bacterium *Sphingomonas* sp. MJ-3. *Marine Biotechnology* 14:189–202 DOI 10.1007/s10126-011-9402-7.
- Prabandono K, Amin S. 2015.** Production of biomethane from marine microalgae. In: Kim SK, Lee CG, eds. *Marine bioenergy: trends and developments*. Boca Raton: CRC Press, 303–323.
- Preiss J, Ashwell G. 1962a.** Alginic acid metabolism in bacteria. I. Enzymatic formation of unsaturated oligosaccharides and 4-deoxy-L-erythro-5-hexoseulose uronic acid. *Journal of Biological Chemistry* 237:309–316.
- Preiss J, Ashwell G. 1962b.** Alginic acid metabolism in bacteria. II. The enzymatic reduction of 4-deoxy-L-erythro-5-hexoseulose uronic acid to 2-keto-3-deoxy-D-gluconic acid. *Journal of Biological Chemistry* 237:317–321.
- Qian P-Y, Wang Y, Lee OO, Lau SCK, Yang J, Lafi FF, Al-Suwailem A, Wong TYH. 2011.** Vertical stratification of microbial communities in the Red Sea revealed by 16S rDNA pyrosequencing. *ISME Journal* 5:507–518 DOI 10.1038/ismej.2010.112.
- Qiu YL, Kuang XZ, Shi XS, Yuan XZ, Guo RB. 2014.** *Paludibacter jiangxiensis* sp. nov., a strictly anaerobic, propionate-producing bacterium isolated from rice paddy field. *Archives of Microbiology* 196:149–155 DOI 10.1007/s00203-013-0951-1.
- Ramaraj R, Unpaprom Y, Dussadee N. 2016.** Potential evaluation of biogas production and upgrading through algae. *International Journal of New Technology and Research* 2:128–133.
- Rausch T. 1981.** The estimation of micro-algal protein content and its meaning to the evaluation of algal biomass I. Comparison of methods for extracting protein. *Hydrobiologia* 78:237–251 DOI 10.1007/BF00008520.
- Reed DC, Rassweiler A, Arkema KK. 2008.** Biomass rather than growth rate determines variation in net primary production by giant kelp. *Ecology* 89:2493–2505 DOI 10.1890/07-1106.1.
- Riviere D, Desvignes V, Pelletier E, Chaussonnerie S, Guermazi S, Weissenbach J, Li T, Camacho P, Sghir A. 2009.** Towards the definition of a core of microorganisms involved in anaerobic digestion of sludge. *ISME Journal* 3:700–714 DOI 10.1038/ismej.2009.2.

- Roalkvam I, Drønen K, Stokke R, Daae FL, Dahle H, Steen IH. 2015.** Physiological and genomic characterization of *Arcobacter anaerophilus* IR-1 reveals new metabolic features in Epsilonproteobacteria. *Frontiers in Microbiology* **6**: Article 987.
- Sakai T, Kawai T, Kato I. 2004.** Isolation and characterization of a fucoidan-degrading marine bacterial strain and its fucoidanase. *Marine Biotechnology (NY)* **6**:335–346 DOI [10.1007/s10126-003-0033-5](https://doi.org/10.1007/s10126-003-0033-5).
- Sakazaki R. 1965.** A proposed group of the family Enterobacteriaceae, the Asakusa group. *International Journal of Systematic and Evolutionary Microbiology* **15**:45–47.
- Samson R, Leduy A. 1982.** Biogas production from anaerobic-digestion of *Spirulina-maxima* algal biomass. *Biotechnology and Bioengineering* **24**:1919–1924 DOI [10.1002/bit.260240822](https://doi.org/10.1002/bit.260240822).
- Sanchez-Andrea I, Stams AJ, Amils R, Sanz JL. 2013.** Enrichment and isolation of acidophilic sulfate-reducing bacteria from Tinto River sediments. *Environmental Microbiology Reports* **5**:672–678.
- Santegoeds CM, Ferdelman TG, Muyzer G, De Beer D. 1998.** Structural and functional dynamics of sulfate-reducing populations in bacterial biofilms. *Applied and Environmental Microbiology* **64**:3731–3739.
- Sapp M, Wichels A, Gerdts G. 2007.** Impacts of cultivation of marine diatoms on the associated bacterial community. *Applied and Environmental Microbiology* **73**:3117–3120 DOI [10.1128/AEM.02274-06](https://doi.org/10.1128/AEM.02274-06).
- Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF. 2009.** Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology* **75**:7537–7541 DOI [10.1128/AEM.01541-09](https://doi.org/10.1128/AEM.01541-09).
- Sialve B, Bernet N, Bernard O. 2009.** Anaerobic digestion of microalgae as a necessary step to make microalgal biodiesel sustainable. *Biotechnology Advances* **27**:409–416 DOI [10.1016/j.biotechadv.2009.03.001](https://doi.org/10.1016/j.biotechadv.2009.03.001).
- Sommer U, Adrian R, De Senerpont Domis L, Elser JJ, Gaedke U, Ibelings B, Jepsen E, Lürling M, Molinero JC, Mooij WM, Van Donk E, Winder M. 2012.** Beyond the Plankton Ecology Group (PEG) model: mechanisms driving plankton succession. *Annual Review of Ecology, Evolution, and Systematics* **43**:429–448 DOI [10.1146/annurev-ecolsys-110411-160251](https://doi.org/10.1146/annurev-ecolsys-110411-160251).
- Sommer U, Maciej Gliwicz Z, Lampert W, Duncan A. 1986.** The PEG model of seasonal succession of planktonic events in freshwaters. *Archiv fur Hydrobiologie* **106**:433–471.
- Stolz JF, Ellis DJ, Blum JS, Ahmann D, Lovley DR, Oremland RS. 1999.** *Sulfurospirillum barnesii* sp. nov. and *Sulfurospirillum arsenophilum* sp. nov., new members of the *Sulfurospirillum* clade of the epsilon Proteobacteria. *International Journal of Systematic and Evolutionary Microbiology* **49**(Pt 3):1177–1180 DOI [10.1099/00207713-49-3-1177](https://doi.org/10.1099/00207713-49-3-1177).
- Sutherland AD, Varela JC. 2014.** Comparison of various microbial inocula for the efficient anaerobic digestion of *Laminaria hyperborea*. *BMC Biotechnology* **14**:7–7 DOI [10.1186/1472-6750-14-7](https://doi.org/10.1186/1472-6750-14-7).

- Thomas F, Barbeyron T, Tonon T, Genicot S, Czjzek M, Michel G. 2012.** Characterization of the first alginolytic operons in a marine bacterium: from their emergence in marine Flavobacteria to their independent transfers to marine Proteobacteria and human gut Bacteroides. *Environmental Microbiology* **14**:2379–2394 DOI [10.1111/j.1462-2920.2012.02751.x](https://doi.org/10.1111/j.1462-2920.2012.02751.x).
- Ueki A, Akasaka H, Suzuki D, Ueki K. 2006.** *Paludibacter propionicigenes* gen. nov., sp. nov., a novel strictly anaerobic, Gram-negative, propionate-producing bacterium isolated from plant residue in irrigated rice-field soil in Japan. *International Journal of Systematic and Evolutionary Microbiology* **56**:39–44 DOI [10.1099/ij.s.0.63896-0](https://doi.org/10.1099/ij.s.0.63896-0).
- Vanegas CH, Bartlett J. 2013.** Green energy from marine algae: biogas production and composition from the anaerobic digestion of Irish seaweed species. *Environmental Technology* **34**:2277–2283 DOI [10.1080/09593330.2013.765922](https://doi.org/10.1080/09593330.2013.765922).
- Vanni MJ. 2002.** Nutrient cycling by animals in freshwater ecosystems. *Annual Review of Ecology and Systematics* **33**:341–370 DOI [10.1146/annurev.ecolsys.33.010802.150519](https://doi.org/10.1146/annurev.ecolsys.33.010802.150519).
- Vergara-Fernandez A, Vargas G, Alarcon N, Velasco A. 2008.** Evaluation of marine algae as a source of biogas in a two-stage anaerobic reactor system. *Biomass & Bioenergy* **32**:338–344 DOI [10.1016/j.biombioe.2007.10.005](https://doi.org/10.1016/j.biombioe.2007.10.005).
- Viggor S, Juhanson J, Jõesaar M, Mitt M, Truu J, Vedler E, Heinaru A. 2013.** Dynamic changes in the structure of microbial communities in Baltic Sea coastal seawater microcosms modified by crude oil, shale oil or diesel fuel. *Microbiological Research* **168**:415–427 DOI [10.1016/j.micres.2013.02.006](https://doi.org/10.1016/j.micres.2013.02.006).
- Wang S, Chng KR, Wilm A, Zhao S, Yang K-L, Nagarajan N, He J. 2014.** Genomic characterization of three unique *Dehalococcoides* that respire on persistent polychlorinated biphenyls. *Proceedings of the National Academy of Sciences of the United States of America* **111**:12103–12108 DOI [10.1073/pnas.1404845111](https://doi.org/10.1073/pnas.1404845111).
- Wang Y, Qian P-Y. 2009.** Conservative fragments in bacterial 16s rrna genes and primer design for 16s ribosomal DNA amplicons in metagenomic studies. *PLOS ONE* **4**:e7401 DOI [10.1371/journal.pone.0007401](https://doi.org/10.1371/journal.pone.0007401).
- Wang J, Tang H, Zhang C, Zhao Y, Derrien M, Rocher E, Van-Hylckama Vlieg JET, Strissel K, Zhao L, Obin M, Shen J. 2015.** Modulation of gut microbiota during probiotic-mediated attenuation of metabolic syndrome in high fat diet-fed mice. *ISME Journal* **9**:1–15 DOI [10.1038/ismej.2014.99](https://doi.org/10.1038/ismej.2014.99).
- Ward AJ, Lewis DM, Green FB. 2014.** Anaerobic digestion of algae biomass: a review. *Algal Research* **5**:204–214 DOI [10.1016/j.algal.2014.02.001](https://doi.org/10.1016/j.algal.2014.02.001).
- Wetzel R. 2001.** *Limnology. Lake and river ecosystems*. New York: Academic Press.
- Wiley PE, Campbell JE, McKuin B. 2011.** Production of biodiesel and biogas from algae: a review of process train options. *Water Environment Research* **83**:326–338 DOI [10.2175/106143010X12780288628615](https://doi.org/10.2175/106143010X12780288628615).
- Wust PK, Horn MA, Drake HL. 2011.** Clostridiaceae and Enterobacteriaceae as active fermenters in earthworm gut content. *ISME Journal* **5**:92–106 DOI [10.1038/ismej.2010.99](https://doi.org/10.1038/ismej.2010.99).
- Xia Y, Chin FYL, Chao Y, Zhang T. 2015.** Phylogeny-structured carbohydrate metabolism across microbiomes collected from different units in wastewater

- treatment process. *Biotechnology for Biofuels* **8**: Article 172
DOI [10.1186/s13068-014-0179-6](https://doi.org/10.1186/s13068-014-0179-6).
- Xia R, Zhang Y, Critto A, Wu J, Fan J, Zheng Z, Zhang Y. 2016.** The potential impacts of climate change factors on freshwater eutrophication: implications for research and countermeasures of water management in China. *Sustainability* **8**:229
DOI [10.3390/su8030229](https://doi.org/10.3390/su8030229).
- Xie ZF, Wang ZW, Wang QY, Zhu CW, Wu ZC. 2014.** An anaerobic dynamic membrane bioreactor (AnDMBR) for landfill leachate treatment: performance and microbial community identification. *Bioresource Technology* **161**:29–39
DOI [10.1016/j.biortech.2014.03.014](https://doi.org/10.1016/j.biortech.2014.03.014).
- Xin F, He J. 2013.** Characterization of a thermostable xylanase from a newly isolated *Kluyvera* species and its application for biobutanol production. *Bioresource Technology* **135**:309–315 DOI [10.1016/j.biortech.2012.10.002](https://doi.org/10.1016/j.biortech.2012.10.002).
- Xu M, Chen X, Qiu M, Zeng X, Xu J, Deng D, Sun G, Li X, Guo J. 2012.** Bar-Coded Pyrosequencing reveals the responses of PBDE-degrading microbial communities to electron donor amendments. *PLOS ONE* **7**:e30439
DOI [10.1371/journal.pone.0030439](https://doi.org/10.1371/journal.pone.0030439).
- Yagi JM, Suflita JM, Gieg LM, DeRito CM, Jeon C-O, Madsen EL. 2010.** Subsurface cycling of nitrogen and anaerobic aromatic hydrocarbon biodegradation revealed by nucleic acid and metabolic biomarkers. *Applied and Environmental Microbiology* **76**:3124–3134 DOI [10.1128/AEM.00172-10](https://doi.org/10.1128/AEM.00172-10).
- Yen HW, Brune DE. 2007.** Anaerobic co-digestion of algal sludge and waste paper to produce methane. *Bioresource Technology* **98**:130–134
DOI [10.1016/j.biortech.2005.11.010](https://doi.org/10.1016/j.biortech.2005.11.010).
- Yonemoto Y, Tanaka H, Hisano T, Sakaguchi K, Abe S, Yamashita T, Kimura A, Murata K. 1993.** Bacterial alginate lyase gene: nucleotide sequence and molecular route for generation of alginate lyase species. *Journal of Fermentation and Bioengineering Journal of Fermentation and Bioengineering* **75**:336–342
DOI [10.1016/0922-338X\(93\)90130-Z](https://doi.org/10.1016/0922-338X(93)90130-Z).
- Youssef NH, Couger MB, Elshahed MS. 2010.** Fine-scale bacterial beta diversity within a complex ecosystem (Zodletone Spring, OK, USA): the role of the rare biosphere. *PLOS ONE* **5**:e12414 DOI [10.1371/journal.pone.0012414](https://doi.org/10.1371/journal.pone.0012414).
- Youssef NH, Farag IF, Rinke C, Hallam SJ, Woyke T, Elshahed MS. 2015.** In silico analysis of the metabolic potential and niche specialization of candidate phylum “Latescibacteria” (WS3). *PLOS ONE* **10**:e0127499
DOI [10.1371/journal.pone.0127499](https://doi.org/10.1371/journal.pone.0127499).
- Yuan XZ, Shi XS, Zhang DL, Qiu YL, Guo RB, Wang LS. 2011.** Biogas production and microcystin biodegradation in anaerobic digestion of blue algae. *Energy & Environmental Science* **4**:1511–1515 DOI [10.1039/c0ee00452a](https://doi.org/10.1039/c0ee00452a).
- Ziemer CJ. 2014.** Newly cultured bacteria with broad diversity isolated from eight-week continuous culture enrichments of cow feces on complex polysaccharides. *Applied and Environmental Microbiology* **80**:574–585 DOI [10.1128/AEM.03016-13](https://doi.org/10.1128/AEM.03016-13).