

Letter to the Editor

Laboratory Investigations in Patients with Community Acquired Sepsis and/or Pneumonia Caused by *Burkholderia pseudomallei*

Dear Sir,

Gassiep and coauthors have made several key observations regarding associations between laboratory investigations and mortality in bacteremic melioidosis cases.¹ Though there is significant decline in the mortality rates of bacteremic melioidosis cases across various endemic areas attributed to increased awareness, with early diagnosis and optimal management the mortality of bacteremic melioidosis still remains as high as 66%.^{2,3} As pointed out rightly by the authors, there is limited information available regarding the association of various laboratory parameters at presentation with melioidosis severity and mortality. In recent years, we have witnessed efforts to examine the clinical characteristics of patients with bacteremia caused by specific pathogens.^{3,4} In this context, we elaborate on the role of laboratory-based investigations in distinguishing septicemic melioidosis cases from other causes of septicemia, in addition to their role as prognostic indicators in melioidosis.

In a prospective study at the All India Institute of Medical Sciences, Bhubaneswar, Odisha State, from July 2018 to March 2020, we identified 33 culture-confirmed cases of melioidosis among a total of 196 patients presenting with community acquired sepsis and/or pneumonia.⁵ We compared laboratory parameters between these 33 culture confirmed melioidosis cases and 163 nonmelioidosis cases presenting with community acquired sepsis and/or pneumonia. Median hemoglobin level (9.5 g/dL [interquartile range (IQR) 4.3–12]) and platelet count ($219 \times 10^3/\text{mm}^3$, (IQR [19–646]) of melioidosis cases were significantly lower compared with those of the nonmelioidosis cases (10.9 g/dL [IQR 2.4–16.3]), $P = 0.001$ and $309 \times 10^3/\text{mm}^3$, (IQR 50–856), $P = 0.003$). On admission median Sequential Organ Failure Assessment (SOFA) score (4 versus 3, $P = 0.009$) and median quick SOFA (qSOFA) score (2 versus 1, $P < 0.001$) was higher in melioidosis cases compared with nonmelioidosis cases. Out of 33 confirmed melioidosis cases, 7 died (7/33, 21%). Various hematological and biochemical test results and risk factors were compared between the survivors and nonsurvivors; on bivariate analysis, there was no significant association between specific laboratory parameters or risk factors (diabetes mellitus, alcoholism, chronic kidney disease, malignancy, or bacteremia) and mortality.

In our study, higher SOFA or qSOFA and lower median hemoglobin level and platelet count at admission were predictive of melioidosis among patients presenting with community acquired sepsis. In another recent study, median modified SOFA score was higher in patients presenting with *Burkholderia pseudomallei* bacteremia compared with that with *Escherichia coli* and *Staphylococcus aureus* bacteremia.³ In our study of a small cohort of 33 culture-confirmed melioidosis cases, we did not find association between specific hematological or biochemical test results or tested risk factors and mortality. Detailed analysis of baseline laboratory parameters stratified by pathogen group should be attempted

in large-scale studies to improve the selection of empiric antimicrobial therapy and predict patient outcomes.

BIJAYINI BEHERA
AllIMS Bhubaneswar, Microbiology
Bhubaneswar, Odisha, India
E-mail: micro_bijayini@aiimsbhubaneswar.edu.in

ANJUNA RADHAKRISHNANAN
AllIMS Jodhpur, Microbiology
Jodhpur, Rajasthan, India
E-mail: anjukrishna008@gmail.com

BAJAYANTIMALA MISHRA
AllIMS Bhubaneswar, Microbiology
Bhubaneswar, Odisha, India
E-mail: micro_bajayantimala@aiimsbhubaneswar.edu.in

PRASANTA RAGHAB MOHAPATRA
AllIMS Bhubaneswar, Pulmonary Medicine and Critical Care
Bhubaneswar, Odisha, India
E-mail: prmhapatra@hotmail.com

RAJESH KUMAR
AllIMS Bhubaneswar, General Medicine
Bhubaneswar, Odisha, India
E-mail: rrajeshdr@yahoo.co.in

Published online March 28, 2022.

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC-BY) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

REFERENCES

1. Gassiep I, Ganeshalingam V, Chatfield MD, Harris PNA, Norton RE, 2022. Melioidosis: laboratory investigations and association with patient outcomes. *Am J Trop Med Hyg* 106: 54–59.
2. Currie BJ et al., 2021. The Darwin prospective melioidosis study: a 30-year prospective, observational investigation. *Lancet Infect Dis* 21: 1737–1746.
3. Somayaji R, Hantrakun V, Teparrukkul P, Wongsuvan G, Rudd KE, Day NPJ, West TE, Limmathurotsakul D, 2021. Comparative clinical characteristics and outcomes of patients with community acquired bacteremia caused by *Escherichia coli*, *Burkholderia pseudomallei* and *Staphylococcus aureus*: a prospective observational study (Ubon-sepsis). *PLoS Negl Trop Dis* 15: e0009704.
4. Deen J, von Seidlein L, Andersen F, Elle N, White NJ, Lubell Y, 2012. Community-acquired bacterial bloodstream infections in developing countries in south and southeast Asia: a systematic review. *Lancet Infect Dis* 12: 480–487.
5. Radhakrishnan A, Behera B, Mishra B, Mohapatra PR, Kumar R, Singh AK, 2021. Clinico-microbiological description and evaluation of rapid lateral flow immunoassay and PCR for detection of *Burkholderia pseudomallei* from patients hospitalized with sepsis and pneumonia: a twenty-one months' study from Odisha, India. *Acta Trop* 221: 105994.