

Oral presentation

Open Access

OA04-03. Characterization of cell-mediated immune responses generated by recombinant modified vaccinia Ankara (rMVA)-HIV-1 in a phase I vaccine trial

JR Currier*¹, M De Souza², S Ratto-Kim¹, R Paris², V Nguay¹, J Cox³, P Earl⁴, B Moss⁴, S Sriplienchan², P Thongcharoen², J Kim¹, M Robb¹, N Michael¹ and M Marovich¹

Address: ¹Department of Vaccine Research and Development, Military HIV Research Program, Rockville, MD, USA, ²AFRIMS, Bangkok, Thailand, ³IAVI, Rockville, MD, USA and ⁴NIAID, Bethesda, MD, USA

* Corresponding author

from AIDS Vaccine 2009
Paris, France. 19–22 October 2009

Published: 22 October 2009

Retrovirology 2009, 6(Suppl 3):O27 doi:10.1186/1742-4690-6-S3-O27

This abstract is available from: <http://www.retrovirology.com/content/6/S3/O27>

© 2009 Currier et al; licensee BioMed Central Ltd.

Background

Potency of the cell-mediated immune response is now the critical metric for down-selection of candidate HIV-1 vaccines. Here we characterize the potency (magnitude and quality) of cell-mediated immunity generated in response to a multigenic rMVA-based HIV-1 (CRF01_AE-derived) vaccine.

Methods

49 healthy, vaccinia-naive volunteers were enrolled in a phase I randomized, double-blind, dose-escalation, route-comparison, placebo-controlled trial to assess the safety and immunogenicity of MVA-CMDR HIV-1 vaccine. The study was divided into Part A: low-dose 10^6 /pfu ID versus 10^7 pfu/IM and Part B: high-dose 10^7 pfu/ID vs 10^8 pfu/IM. Vaccinations were given at months 0, 1 and 3 with an active:placebo ratio of 10:2. Chromium-release CTL, IFN γ Elispot, and polyfunctional flow cytometry (IL-2/IFN γ /TNF α /MIP-1 β /CD107a), were performed on all volunteers. Synthetic peptide pools and GLP-grade MVA were used to assess insert (Gag/Pol/Env) and vector immunogenicity respectively.

Results

Vector-specific responses were robust (> 80% response rate at high-dose), durable (maintained at least 6

months), and exhibited a dose-dependent increase in both magnitude and response rate among the 4 arms of the trial. HIV-insert-specific responses were detected using all assay platforms, but were lower than the vector-specific responses in both magnitude and response rate in all arms of the trial (~60% at high-dose by CTL, Elispot and ICS assays). Specifically, polyfunctional analysis revealed a TNF α /IL-2/IFN γ bias in CD4+ T cells and a MIP-1 β /CD107a/IFN γ bias in CD8+ T cells, with CD4+ T cell responses more frequent than CD8+ T cell responses to the HIV inserts. Vector-specific immune responses showed a boosting effect from the 2nd to the 3rd immunization.

Conclusion

rMVA vaccination induces a dose-dependent, robust and durable polyfunctional cellular immune response as measured by IFN γ Elispot, CTL and intracellular cytokine stimulation assays. Although vector-specific responses tend to dominate over insert-specific responses, the data supports further exploration of MVA as a vector modality in prime-boost vaccination strategies.