

Enantioselective Synthesis of α -Quaternary Mannich Adducts by Palladium-Catalyzed Allylic Alkylation: Total Synthesis of (+)-Sibirinine

Yoshitaka Numajiri, Beau P. Pritchett, Koji Chiyoda, and Brian M. Stoltz*

Warren and Katharine Schlinger Laboratory for Chemistry and Chemical Engineering, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, California 91125, United States


S Supporting Information

ABSTRACT: A catalytic enantioselective method for the synthesis of α -quaternary Mannich-type products is reported. The two-step sequence of (1) Mannich reaction followed by (2) decarboxylative enantioselective allylic alkylation serves as a novel strategy to in effect access asymmetric Mannich-type products of “thermodynamic” enolates of substrates possessing additional enolizable positions and acidic protons. Palladium-catalyzed decarboxylative allylic alkylation enables the enantioselective synthesis of five-, six-, and seven-membered ketone, lactam, and other heterocyclic systems. The mild reaction conditions are notable given the acidic free N–H groups and high functional group tolerance in each of the substrates. The utility of this method is highlighted in the first total synthesis of (+)-sibirinine.

The Mannich reaction, first discovered in the early 20th century, is among the most robust reactions known to produce nitrogen-containing compounds. In a classic intermolecular Mannich reaction, an aldehyde, an amine, and an α -acidic carbonyl compound react to form a β -amino carbonyl compound.¹ Recent progress in this area, including modified imine donors and well-explored catalyst systems, has made available a wide variety of asymmetric α -functionalizations of carbonyl compounds.² To date, asymmetric Mannich-type reactions to establish α -quaternary carbonyl compounds have been limited to stabilized enolates³ (e.g., 1,3-dicarbonyl compounds). To our knowledge, the lone exception is a proline-catalyzed Mannich reaction with branched aldehydes.⁴ Despite the importance of the Mannich reaction, only a handful of asymmetric α -aminomethylation reactions have been reported.⁵ Enders et al.^{5a} employed enantiomerically pure α -silyl ketones for regio- and diastereoselective syntheses of Mannich bases. Córdova et al.^{5c} developed a proline-catalyzed asymmetric Mannich reaction that provides α -aminomethylated α -tertiary ketones in excellent enantioselectivities. The only example of enantioselective all-carbon quaternary center formation in this area is the palladium(II)-catalyzed aminomethylation of β -keto esters (i.e., II, R = CO₂t-Bu) performed by Sodeoka et al.^{5d} however, the reported enantioselectivities are only moderate (up to 68% ee). To date, there are no reports of enantioselective catalysis leading to α -quaternary aminomethyl “Mannich” adducts bearing only a single carbonyl moiety (i.e., II, R = alkyl).

The regioselectivity of enolate formation, potential subsequent enolization, and difficulty in controlling enantioselectivity under conditions that proceed through the thermodynamically favorable enolate all pose significant challenges to those trying to access enantioenriched α -quaternary Mannich adducts⁶ of α -alkyl-substituted ketones (Figure 1A). To overcome these challenges, we envisioned a strategy wherein the alkylation, not the aminomethylation, would be performed last (Figure 1B).

A. Asymmetric Mannich Adducts via a Traditional Approach



B. Our Re-Envisioned Strategy




Figure 1. Contrasting approaches to access α -quaternary “Mannich” products.


Over the course of the past decade, our group has demonstrated decarboxylative palladium-catalyzed asymmetric alkylation of β -keto esters as a powerful tool to access quaternary centers with high enantioselectivities.⁷ The extensive substrate scope and broad functional group compatibility of this transformation^{8,9} encouraged further exploration of palladium catalysts in the synthesis of amine-containing substrates, thereby facilitating access to enantioenriched bioactive alkaloids or pharmaceutical candidates. We therefore sought to implement our well-studied, reliable

Received: November 26, 2014

Published: January 11, 2015

alkylation chemistry in a simple yet powerful strategy for the synthesis of α -quaternary Mannich products in an enantioselective fashion. Our plan is outlined in Figure 1B. Introduction of an aminomethyl group to β -keto ester V using classical Mannich chemistry (V to VI), followed by an asymmetric allylic alkylation reaction, would provide the enantioenriched α -quaternary ketone product VII. Compound VII can be thought of as an α -aminomethylation product of the so-called "thermodynamic" enolate of compound I. We imagined that successful exploration of this inverted strategy would enable rapid, stereocontrolled total syntheses of (–)-isonitramine and (+)-sibirinine.^{10–12}

To introduce the aminomethyl moiety, we employed sulfonylmethyl carbamates (e.g., 2a) as versatile and readily available imine precursors.¹³ In the presence of Cs₂CO₃, the Boc-protected imine generated from 2 reacted with β -keto ester 1¹⁴ to smoothly afford β -aminoketone 3a, quantitatively, at ambient temperature (Scheme 1). In a similar manner, we obtained other protected aminoketones 3b–g in good to excellent yields.¹⁵


Scheme 1. Synthesis of β -Keto Ester 3a

With β -keto esters 3a–g in hand, our investigation into this substrate class commenced in the context of palladium-catalyzed allylic alkylation as shown in Table 1. We found that exposure of Boc-protected substrate 3a to a catalytic phosphinooxazoline¹⁶–palladium(0) complex in toluene at ambient temperature afforded the desired product 4a in 94% yield and 86% ee (entry 1). Cbz-protected 3b also gave an excellent yield and ee (entry 3). It is important to note that we did not detect any *N*-alkylated side products, a result that highlights the mild nature of our reaction conditions.¹⁷ Arylcarbamates 3c–e gave slightly decreased enantioselectivities in the products (entries 4–6). Changing from carbamate to benzoyl or tosyl protecting groups resulted in poor ee (entries 7, 8) presumably due to their ability to coordinate to the catalyst and/or the enhanced acidity of the N–H proton. We found that the more electron-withdrawn ligand, L1, gave higher enantioselectivity. In the case of (*S*)-*t*BuPHOX L2 (Table 1, entry 2), we observed diminished ee.

As outlined in Table 2, we have found that a broad range of ketones and amides (e.g., 5a–g) can easily be converted into enantioenriched tetrasubstituted Mannich-type products (e.g., 7a–i) with this two-step strategy. For all substrates, the first step proceeded in good to excellent yields (72–99%). In the allylic alkylation, 2-phenyl-2-propenyl-substituted 7a was obtained in high yield (91%) and excellent enantioselectivity (90% ee). Cycloheptanone 6b proved to be a good substrate and the corresponding α -quaternary cycloheptanone 7b was isolated in 93% yield and 87% ee, while cyclopentanone 6c gave a slightly lower enantioselectivity (82% ee). Vinylogous ester 6d and tetralone 6e afforded α -quaternary vinylogous ester 7d and tetralone 7e in 70% yield and 92% ee, and 74% yield and 93% ee, respectively. Heterocyclic ketone scaffolds were found to be competent substrates for this transformation, as 4-

Table 1. Optimization of the Amine Protecting Group^a

entry	R (3→4)	ligand	yield (%)	ee (%) ^b
1	Boc (3a→4a)	L1	94	86
2	Boc (3a→4a)	L2	ND ^c	80
3	Cbz (3b→4b)	L1	96	86
4	X = OMe (3c→4c)	L1	91	83
5	X = H (3d→4d)	L1	90	77
6	X = F (3e→4e)	L1	84	77
7	Bz (3f→4f)	L1	ND ^c	56
8	Ts (3g→4g)	L1	54	24



^aReaction performed with 0.2 mmol of 3, 5 mol % of Pd₂(dba)₃ (dba = dibenzylideneacetone), 12.5 mol % of ligand at 0.033 M in toluene at 23 °C. ^bDetermined by chiral SFC analysis. Absolute stereochemistry has been assigned by analogy,⁸ except in entry 3, which was assigned by conversion into (–)-isonitramine. ^cA yield was not determined.

Table 2. Two-Step Synthesis of α -Aminomethyl Carbonyl Compounds from β -Oxo Esters^a


6a 79% yield 7a 91% yield, 90% ee ^{b,c}	6b 72% yield 7b 93% yield, 87% ee	6c 86% yield 7c 98% yield, 82% ee
6d 83% yield 7d 70% yield, 92% ee ^b	6e 99% yield 7e 74% yield, 93% ee ^b	6f 80% yield 7f 78% yield, 90% ee ^b
6g 74% yield 7g 94% yield, 90% ee ^d	6h 80% yield 7h 92% yield, 99% ee ^d	6i 80% yield 7i 51% yield, 92% ee ^{b,d}

^aReaction conditions for step 2: 6 (1 equiv), Pd₂(dba)₃ (5 mol %), and L1 (12.5 mol %) in toluene (0.033 M) at 23 °C for 12–48 h. ^bPd₂(pmdba)₃ (pmdba = bis(4-methoxybenzylidene)acetone) was used instead of Pd₂(dba)₃. ^cEnantiomeric excesses were determined by chiral SFC analysis. ^dReactions were performed on 6g, 6h, and 6i at 40 °C. ^eFor individual reaction times, see Supporting Information.

piperidinone **7f** was isolated in 78% yield and 90% ee. Lastly, we were pleased to find that, under slightly elevated reaction temperatures (40 °C), the desired lactam **7g**, morpholinone **7h**, and carbazolone **7i** were obtained in moderate to excellent yields (51–94%) and excellent enantioselectivities (90–99% ee).

In order to exhibit the utility of our method for generating interesting and useful chiral building blocks, the first total synthesis of (+)-sibirinine (**12**) was carried out (Scheme 2).

Scheme 2. Natural Product Synthesis



(+)-Sibirinine is a tricyclic alkaloid featuring an *N,O*-acetal, a tertiary amine *N*-oxide, and two pairs of vicinal stereocenters, including an all-carbon quaternary center. Asymmetric allylic alkylation using 1 g of **3b** proceeded with one-half of the typical catalyst loading without any loss of enantioselectivity. Reduction of β -amino ketone **4b** with diisobutylaluminum hydride (DIBAL), followed by acetylation of the resulting alcohol, yielded carbamate **8** as a single diastereomer. Hydroboration of the terminal alkene in carbamate **8** provided primary alcohol **9** in 86% yield over three steps. Cyclization of the mesylate derived from primary alcohol **9** smoothly delivered spirocycle **10**. Removal of the acetyl and Cbz groups using potassium hydroxide furnished (–)-isonitramine (**11**) in 77% yield. Treatment of (–)-isonitramine (**11**) with excess acetaldehyde yielded the desired hemiaminal, which was smoothly oxidized by *m*-CPBA to give (+)-sibirinine (**12**) in 92% yield over two steps. Notably, conversion of (–)-isonitramine to (+)-sibirinine can be accomplished in one pot by forming the hemiaminal intermediate under an oxygen atmosphere, albeit in diminished yield. Spectral data of **11** and **12** were identical to those previously reported.^{11,12} Our synthesis of (–)-isonitramine confirms the absolute stereochemistry of **4b**.¹¹

In summary, we have developed an inverted approach to the synthesis of α -quaternary and tetrasubstituted tertiary Mannich-type products by strategic enolate formation to give products in moderate to excellent yields and good to excellent ee. This chemistry tolerates a variety of ketone, amide, and vinylogous ester functionalities even in the presence of basic tertiary amines and relatively acidic *N*–H moieties. Multiple ring sizes as well as aromatic and heteroaromatic scaffolds are also accessible via this strategy. Furthermore, this method enables the efficient construction of spirocyclic amine-containing scaffolds, as illustrated in our synthesis of the alkaloids (–)-isonitramine and (+)-sibirinine. The first total synthesis

of (+)-sibirinine was accomplished in 11 steps and 36% overall yield from commercially available diallyl pimelate. Further studies expanding the scope and applications of this two-step Mannich-like methodology are ongoing in our laboratory.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and compound characterization. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*stoltz@caltech.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This manuscript is dedicated to Professor Teruaki Mukaiyama in celebration of the 40th anniversary of the Mukaiyama Aldol reaction. The authors wish to thank NIH-NIGMS (R01GM080269), Amgen, the Gordon and Betty Moore Foundation, the Caltech Center for Catalysis and Chemical Synthesis, and Caltech for financial support. Y.N. thanks Toray Industries Inc. for a postdoctoral fellowship. B.P.P. thanks the NSF for a predoctoral fellowship. K.C. acknowledges the Japan Society for the Promotion of Science for a postdoctoral fellowship. The authors thank Mona Shahgholi and Naseem Torian for mass spectrometry assistance and Dr. Scott Virgil (Caltech) for instrumentation assistance.

■ REFERENCES

- (1) For selected reviews for Mannich reaction and other references therein, see: (a) Kleinman, E. F. In *Comprehensive Organic Synthesis*, Vol. 2; Trost, B. M., Fleming, I., Heathcock, C. H., Eds.; Pergamon: Oxford, 1991; Chapter 4.1. (b) Tramontini, M.; Angiolini, L. *Tetrahedron* **1990**, *46*, 1791. (c) Tramontini, M. *Synthesis* **1973**, 703. (d) Arend, M.; Westermann, B.; Risch, N. *Angew. Chem., Int. Ed.* **1998**, *37*, 1044.
- (2) (a) Kobayashi, S.; Mori, Y.; Fossey, J. S.; Salter, M. M. *Chem. Rev.* **2011**, *111*, 2626. (b) Córdova, A. *Acc. Chem. Res.* **2004**, *37*, 102.
- (3) For selected examples of asymmetric Mannich-type reactions to construct quaternary centers, see: (a) Hamashima, Y.; Sasamoto, N.; Hotta, D.; Somei, H.; Umabayashi, N.; Sodeoka, M. *Angew. Chem., Int. Ed.* **2005**, *44*, 1525. (b) Knudsen, K. R.; Jørgensen, K. A. *Org. Biomol. Chem.* **2005**, *3*, 1362. (c) Xue, S.; Yu, S.; Deng, Y.; Wulff, W. D. *Angew. Chem., Int. Ed.* **2001**, *40*, 2271. (d) Hatano, M.; Horibe, T.; Ishihara, K. *J. Am. Chem. Soc.* **2010**, *132*, 56. (e) Ting, A.; Lou, S.; Schaus, S. E. *Org. Lett.* **2006**, *8*, 2003.
- (4) Chowdari, N. S.; Suri, J. T.; Barbas, C. F., III. *Org. Lett.* **2004**, *6*, 2507.
- (5) (a) Enders, D.; Ward, D.; Adam, J.; Raabe, G. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 981. (b) Yamasaki, S.; Iida, T.; Shibasaki, M. *Tetrahedron* **1999**, *55*, 8857. (c) Ibrahim, I.; Casas, J.; Córdova, A. *Angew. Chem., Int. Ed.* **2004**, *43*, 6528. (d) Hamashima, Y.; Sasamoto, N.; Umabayashi, N.; Sodeoka, M. *Chem.—Asian J.* **2008**, *3*, 1443.
- (6) For nonenantioselective reactions to provide α -aminomethyl α -quaternary ketones, see: (a) Danishefsky, S.; Prisybilla, M.; Lipisko, B. *Tetrahedron Lett.* **1980**, *21*, 805. (b) Matsumoto, K.; Hashimoto, S.; Otani, S.; Atnita, F.; Osugi, J. *Synth. Commun.* **1984**, *14*, 585. (c) Desai, P.; Schildknecht, K.; Agrios, K. A.; Mossman, C.; Milligan, G. L.; Aubé, J. *J. Am. Chem. Soc.* **2000**, *122*, 7226.
- (7) Mohr, J. T.; Behenna, D. C.; Harned, A. M.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2005**, *44*, 6924.

(8) (a) Behenna, D. C.; Mohr, J. T.; Sherden, N. H.; Marinescu, S. C.; Harned, A. M.; Tani, K.; Seto, M.; Ma, S.; Novák, Z.; Krout, M. R.; McFadden, R. M.; Roizen, J. L.; Enquist, J. A., Jr.; White, D. E.; Levine, S. R.; Petrova, K. V.; Iwashita, A.; Virgil, S. C.; Stoltz, B. M. *Chem.—Eur. J.* **2011**, *17*, 14199. (b) Behenna, D. C.; Liu, Y.; Yurino, T.; Kim, J.; White, D. E.; Virgil, S. C.; Stoltz, B. M. *Nat. Chem.* **2012**, *4*, 130. (c) Bennett, N. B.; Duquette, D. C.; Kim, J.; Liu, W.-B.; Marziale, A. N.; Behenna, D. C.; Virgil, S. C.; Stoltz, B. M. *Chem.—Eur. J.* **2013**, *19*, 4414. (d) Reeves, C. M.; Eidamshaus, C.; Kim, J.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2013**, *52*, 6718.

(9) For related studies, see: (a) Nakamura, M.; Hajra, A.; Endo, K.; Nakamura, E. *Angew. Chem., Int. Ed.* **2005**, *44*, 7248. (b) Li, Z.; Zhang, S.; Wu, S.; Shen, X.; Zou, L.; Wang, F.; Li, X.; Peng, F.; Zhang, H.; Shao, Z. *Angew. Chem., Int. Ed.* **2013**, *52*, 4117. (c) Gartshore, C. J.; Lupton, D. W. *Angew. Chem., Int. Ed.* **2013**, *52*, 4113. (d) Trost, B. M.; Xu, J.; Schmidt, T. *J. Am. Chem. Soc.* **2009**, *131*, 18343. (e) Trost, B. M.; Bream, R. N.; Xu, J. *Angew. Chem., Int. Ed.* **2006**, *45*, 3109.

(10) Ibragimov, A. A.; Osmanov, Z.; Tashchodzhaev, B.; Abdullaev, N. D.; Yagudaev, M. R.; Yunusov, S. Y. *Khimiya Prir. Soedin.* **1981**, 623.

(11) For selected examples of total synthesis of isonitramine, see: (a) Deyine, A.; Poirier, J.-M.; Duhamel, L.; Duhamel, P. *Tetrahedron Lett.* **2005**, *46*, 2491. (b) François, D.; Lallemand, M.-C.; Selkti, M.; Tomas, A.; Kunesch, N.; Husson, H.-P. *Angew. Chem., Int. Ed.* **1998**, *37*, 104. (c) Pandey, G.; Kumara, C. P.; Burugu, S. K.; Puranik, V. G. *Eur. J. Org. Chem.* **2011**, 7372. (d) Park, Y.; Lee, Y. J.; Hong, S.; Lee, M.; Park, H.-G. *Org. Lett.* **2012**, *14*, 852. (e) Quirion, J.-C.; Grierson, D. S.; Royer, J.; Husson, H.-P. *Tetrahedron Lett.* **1988**, *29*, 3311.

(12) Ibragimov, A. A.; Abdullaev, N. D.; Osmanov, Z.; Yunusov, S. Y. *Khimiya Prir. Soedin.* **1987**, 685.

(13) (a) Klepacz, A.; Zwierzak, A. *Tetrahedron Lett.* **2002**, *43*, 1079. (b) Sikriwal, D.; Kant, R.; Maulik, P. R.; Dikshit, D. K. *Tetrahedron* **2010**, *66*, 6167.

(14) (a) Tsuji, J.; Nisar, M.; Shimizu, I.; Minami, I. *Synthesis* **1984**, 1009. (b) Mohr, J. T.; Krout, M. R.; Stoltz, B. M. *Org. Synth.* **2009**, *86*, 194.

(15) See Supporting Information.

(16) McDougal, N. T.; Streuff, J.; Mukherjee, H.; Virgil, S. C.; Stoltz, B. M. *Tetrahedron Lett.* **2010**, *51*, 5550.

(17) In several cases, H–N functionality acts as a good nucleophile toward π -allylpalladium complexes; see: Tsuji, J. *Palladium Reagents and Catalysts—New Perspectives for the 21st Century*; John Wiley & Sons, Ltd.: 2004.

(18) For the preparation of **5**, see Supporting Information or ref 8.