

Research article

Open Access

## Immunogenicity of a polyvalent HIV-1 candidate vaccine based on fourteen wild type gp120 proteins in golden hamsters

Ali Azizi<sup>1,2</sup>, David E Anderson<sup>2,3</sup>, Masoud Ghorbani<sup>2</sup>, Katrina Gee<sup>1,2</sup> and Francisco Diaz-Mitoma\*<sup>1,2</sup>

Address: <sup>1</sup>Infectious Disease and Vaccine Research Centre, Research Institute Children's Hospital of Eastern Ontario, 401 Smyth Road, Ottawa, ON, K1H 8L1, Canada, <sup>2</sup>Variation Biotechnologies Inc., 22 de Varennes, Suite 210, Gatineau, Quebec, J8T 8R1, Canada and <sup>3</sup>Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, 77 Avenue Louis Pasteur, Boston, MA, 02115, USA

Email: Ali Azizi - [aliazizi@variationbiotech.com](mailto:aliazizi@variationbiotech.com); David E Anderson - [deanderson@variationbiotech.com](mailto:deanderson@variationbiotech.com); Masoud Ghorbani - [mghorbani@variationbiotech.com](mailto:mghorbani@variationbiotech.com); Katrina Gee - [kgee@variationbiotech.com](mailto:kgee@variationbiotech.com); Francisco Diaz-Mitoma\* - [diaz@variationbiotech.com](mailto:diaz@variationbiotech.com)

\* Corresponding author

Published: 31 October 2006

Received: 14 July 2006

*BMC Immunology* 2006, **7**:25 doi:10.1186/1471-2172-7-25

Accepted: 31 October 2006

This article is available from: <http://www.biomedcentral.com/1471-2172/7/25>

© 2006 Azizi et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** One of the major obstacles in the design of an effective vaccine against HIV-1 is the hypervariability of the HIV-1 envelope glycoprotein. Most HIV-1 vaccine candidates have utilized envelope glycoprotein from a single virus isolate, but to date, none of them elicited broadly reactive humoral immunity. Herein, we hypothesised that a cocktail of HIV-1 gp120 proteins containing multiple epitopes may increase the breadth of immune responses against HIV-1. We compared and evaluated the immunogenicity of HIV-1 vaccines containing either gp120 protein alone or in combinations of four or fourteen gp120s from different primary HIV-1 isolates in immunized hamsters.

**Results:** We amplified and characterized 14 different gp120s from primary subtype B isolates with both syncytium and non-syncytium inducing properties, and expressed the proteins in Chinese Hamster Ovary (CHO) cell lines. Purified proteins were used either alone or in combinations of four or fourteen different gp120s to vaccinate golden hamsters. The polyvalent vaccine showed higher antibody titers to HIV-1 subtype B isolates MN and SF162 compared to the groups that received one or four gp120 proteins. However, the polyvalent vaccine was not able to show higher neutralizing antibody responses against HIV-1 primary isolates. Interestingly, the polyvalent vaccine group had the highest proliferative immune responses and showed a substantial proportion of cross-subtype CD4 reactivity to HIV-1 subtypes B, C, and A/E

**Conclusion:** Although the polyvalent approach achieved only a modest increase in the breadth of humoral and cellular immunity, the qualitative change in the vaccine (14 vs. 1 gp120) resulted in a quantitative improvement in vaccine-induced immunity.

### Background

HIV-1 gp120 is a major target for neutralizing antibodies (Nabs) and for this reason it is an important HIV immu-

nogen to include in vaccine formulations [1-3]. However, the diversity of gp120 has proven to be a significant challenge to HIV-1 vaccine development. The structure of

gp120 contains variable loops (V1-V5) which likely hide critical conserved epitope sites favoured by the Nabs. Furthermore, the crystallography structure of gp120 indicates that the protein is covered by carbohydrates which facilitates viral escape from Nabs [4,5]. Genetic variability in HIV-gp120 between groups M, N and O also affect the induction of Nabs [6,7]. These factors complicate the design of an effective candidate vaccine against HIV.

Previous vaccine studies focus on single HIV immunogens and although some of these studies show an increase in CD4/CD8+T cell immune responses, the immunogens used were not able to induce potent Nabs that mediate sterilizing immunity [8,9]. The question remains: "can a single immunogen induce a broad immune response against a diverse virus like HIV". To address this, several studies have been performed. A single and double recombinant HIV-1 gp120 protein has been used as a candidate immunogen in a phase III clinical vaccine trial. However, this vaccination was not effective to protect against HIV infection [10-12]. This lack of vaccine efficacy may be due to HIV diversity. While some single immunogens neutralize a few T-cell line adapted (TCLA) HIV-1 strains, none of the animal model or clinical studies demonstrated a broadly cross-reactive immunity against HIV-1 primary isolates [13]. Some studies demonstrated neutralizing antibody responses against HIV-1 primary isolates, however no measure of cross-reactivity was obtained as the strains of HIV virus used in the Nab assay, contained the same HIV-1 gp120 as that used for vaccination.

HIV-1 subtype B is widely distributed throughout the world and is the most common subtype in North America and Europe [14,15]. Herein, we hypothesised that immunization with several (fourteen) different wild type HIV-1 gp120 subtype B proteins would increase the breadth of specific antiviral immune responses. Fourteen wild type HIV-1 gp120 subtypes B were amplified, cloned and the recombinant gp120 proteins were expressed in mammalian cell lines. Golden hamsters were immunized with equivalent amounts of 1 vs. 4 vs. 14 distinct gp120 proteins and humoral (antibody binding and neutralization) and cellular (T helper cells) responses to HIV-1 subtypes B, C and A/E were analyzed. Although this polyvalent approach achieved only a modest increase in the breadth of humoral and cellular immunity; the qualitative change in the vaccine (14 vs. 1 gp120, same amount of total antigen) resulted in a quantitative improvement in vaccine-induced immunity.

**Results**

**Characterization and expression of HIV-1 gp120s**

Total RNA was purified from syncytium and non-syncytium inducing co-cultures of 14 HIV-1 patients (Table 1). Amplification products corresponding to the full length of

gp120 (1.6 kb) containing constant and hypervariable regions were generated by RT-PCR. The genes were completely sequenced and identified as subtype B. The V3 amino acid sequence of the amplified gp120s was compared with V3 subtypes B, C and A/E, point mutations as well as insertion and deletion mutations were detected (Fig 1). In addition, the phylogenetic relations between the 14 different gp120 sequences and HIV subtypes B, C and A/E were revealed and genetic diversity between clones was identified (Fig 2). Due to the limited number of nucleotides in the V3 region, the algorithm is applied to a longitudinal data set of the C2-V5 regions of HIV-1 gp120s using Neighbor Tree-Maker in the FASTA format. By this approach, epidemiological linkage was established between the HIV-1 gp120 clones and HIV-1 subtypes B, C and A/E. Most of the HIV-1 gp120 sequences were clustered from the HIV-1 subtype B isolate MN. This isolate was epidemiologically linked to the subtype A/E isolate 93TH975. The results revealed no particular homology between gp120 clones and HIV-1 subtype C isolate 96ZM651. Each gp120 variant was cloned into the pEF6-myc-His vector for the expression and purification of recombinant gp120 proteins *in vitro*. Stable CHO cell lines transfected with the gp120 constructs selected for using blasticidine resistance and recombinant gp120 proteins were purified by metal affinity chromatography. The expression of all recombinant HIV-1 gp120 proteins has been determined by western blotting (Fig 3).

**Immunization schedule**

Four groups of hamsters were immunized with different vaccine formulations as indicated in Table 2. In the first group (A), hamsters were immunized with MPL adjuvant alone. In the second group (B), hamsters were immunized with 100 µg of the recombinant gp120 protein purified from patient number 001 plus adjuvant. Hamsters in the

**Table 1: Viruses were isolated from the above HIV-1 seropositive patients.**

Sample	Sex	Virus isolate	Antiviral therapy
1	M	R5	Y
2	M	R5	Y
3	M	R5	Y
4	F	R5	Y
5	M	R5	Y
6	M	R5	Y
7	M	R5	Y
8	M	R5	Y
9	M	R5	Y
10	M	R5	Y
11	M	X4	Y
12	M	X4	Y
13	M	X4	Y
14	F	X4	Y

<u>Sample number</u>	<u>Length of V3 region</u>
10	TNELVLDLXLCLLHKMLSLXLYVYFFLYXCWXLXQVICTGSLNLYYDFTIVR- 54
14	TMCLS-XLLLFFLLHEMLSLGL-IYFLXYW-GGXCN-XVQNSFNLXQ-GF-NLP- 48
1	TMCLSYISWLFVLSYK-LSLGPIGTLFLVVLWGPPTVDYYRIHSVGTIRV-HWL- 52
2	TMCLSYISWLLVLSYKLLSLGPIGCLFLVVLWGPPTVDYYRIHSVGTIRV-HWL- 53
6	TMCLSYISWLFVLSYK-LSLGPICLFLSVLWFPPTVDYYRIHSVGTIRRV-HWL- 52
3	TMCLSYISCYLSYCMKCSPPW-XYMYALXYIFVVGXCTIXFYRFYQLXYYXL-SIVT 53
4	TVCFYSISYFSTCIKCPPW-XYMYASYCIGXGSCXDFYRFIQLYYYDL-SIGG 53
8	TMCLS ISCXLSCHINCSPW-SYRHXFPCXXIGXCTXDYYRFIPXYYYGL-TLFV 53
12	TMCLSYISCXLSCHINCSPW-SYRDXFPCAXIGXCTXDYYRFIPXYYYAL-TFFV 53
7	-TIWSYISFYSSCCMKCSPPWXYLVLSFFXYWCWVXYXNFYRFISCTLWFX-HCS- 52
5	TVCLSYISYFWSWLYTILSL-VLELPFFLXWXWVLYN-LLQIXSAVLNGF-SMSE 52
11	TVCLSYISYFWSWLLTILSL-VLELDFFLXWX-VLYN-LLQIXSAVLNSF-SMSE 51
9	TMCLSYXSYGLLVHTILVP-XLYVHFLFYCCWVLYX-LLXEIQLYYXGF-SXVR 52
13	----IFPFIPPVA-XALXPICILLCIWLLGLVQLISTXSFCTIMGLALFM- 48

**Figure 1**

Alignment of fourteen amino acid sequences from V3 region of HIV-1 gp120 genes. Nucleotide sequences of V3 region of HIV-1 gp120 genes were translated and aligned by using ClustalW multiple sequence alignment program. Highly variation was seen in the number and position of HIV-1 V3 regions.

third group (C) were immunized with a mixture of proteins purified from patients #001 to #004 (25 µg of each recombinant gp120 protein) and adjuvant. The polyvalent group (D) immunized with a cocktail of all 14 proteins (~7 µg of each recombinant gp120 protein). Hamsters were immunized subcutaneously (S.C.) on weeks 0, 4, and 8. Animals were rested for 8 weeks and then boosted at week 16 with the same combination of gp120 proteins. Antibody titers were determined two weeks after each immunization.


**The polyvalent candidate vaccine induces higher antibody titers against HIV-1 subtype B than other subtypes**

In order to measure antibody titers against various HIV-1 gp120 proteins, plasma from immunized hamsters was collected two weeks after each immunization and analyzed by ELISA. The plates were coated with HIV-1 gp120 subtypes B (MN and SF162), C (96ZM651), and A/E (93TH975) overnight at 4°C followed by incubation with serially diluted plasma and developed as described in the Materials and Method section. The first two immunizations elicited a low level of antibody response to gp120 proteins (data not shown). However, after the third immunization, antibodies were detectable in all groups immunized with gp120. As expected the control group

did not produce an antibody response. The highest level of antibody titer was observed in hamsters that were immunized with 14 recombinant gp120 proteins. In contrast, the control group (adjuvant alone) did not develop detectable levels of specific antibody responses. Eight weeks after the third immunization, animals were vaccinated again with respective recombinant gp120 proteins. Results similar to the third immunization were observed against gp120 subtypes between the groups. The polyvalent vaccine group induced higher levels of anti-gp120 titers to MN and SF162 (subtypes B) than the subtypes C and A/E (Fig 4). These results demonstrate that the polyvalent vaccine induces a higher antibody response to subtype B than the single gp120 candidate vaccine.

**Measurement of neutralizing antibody responses in immunized hamsters**

Two weeks after the fourth immunization, sera were collected and Nabs to HIV-1 strains were tested. Unlike many research studies, to detect the real breadth and strength of candidate vaccines, we used different HIV-1 strains, to assay for Nabs, from those, which were used in immunization. Since some background was observed in the pre-bleed samples, the Nab responses were compared with SVA-MLV (murine retrovirus Env-pseudovirus) as a back-



**Figure 2**  
Phylogenetic relationships of HIV-1 gp120 sequences with subtypes B (MN, SF162), C (96ZM651), and A/E (93TH975). The algorithm is applied to a longitudinal data set of the C2-V5 regions of HIV-1 gp120s using Neighbor Tree-Maker in the FASTA format.

ground control for non-specific activity. The group of hamsters that received all 14 gp120 proteins showed higher Nabs titres to the HIV-1 MN strain than the single gp120 ( $p < 0.05$ ), however, similar Nabs titres were observed in the group immunized with a combination of four gp120 proteins (Fig 5). Sera from immunized hamsters were further tested against other primary HIV-1 isolates but none of the sera were able to show significant neutralization.


**The polyvalent gp120 vaccine induced higher T cell proliferation in immunized hamsters**

The proliferation of CD4+T cells plays an important role in both humoral and cellular immune responses by expansion of antigen-stimulated B and CD8+T cells, respectively. A high proliferative immune response was observed to the gp120 protein isolated from patient #001 in all groups with the exception of the control group (data not shown). However, to assess and measure the strength of the candidate vaccines, splenocytes from immunized hamsters were further stimulated with HIV-1 gp120

strains and subtypes that were not included in the vaccine components. Splenocytes were cultured in the presence of HIV-1 gp120 proteins. Two days after stimulation, [<sup>3</sup>H] thymidine was added and sixteen hours later, cells were harvested and thymidine incorporation was measured. The groups B and C showed a relatively similar proliferative immune response. Interestingly, the group of hamsters that was immunized with all 14 gp120 proteins showed the highest lymphocyte proliferative immune response (Fig 6) to subtypes B (MN and SF162), C (96ZM651) and A/E (93TH975), supporting the cross-subtype activity of Th immune responses.

**Discussions and Conclusion**

The antigenic variation of HIV-1 gp120 is a major obstacle for the development of an effective vaccine [16-18]. Nabs are mainly directed against the V3 region of gp120 which plays an important function in the binding of virus to either CCR5 or CXCR4 [19]. However, antigenic hypervariability of V1-V5 regions especially V3 assists the virus in escaping neutralization [20-24]. Interestingly, a single



**Figure 3**

Western blot analysis of recombinant HIV-1 gp120 proteins. Lane 1: control gp120 protein expressed in a baculovirus expression system obtained from NIH. The lanes 2–8 are the first seven gp120 proteins (#1–7) purified from transfected CHO cells. The blot was probed with polyclonal gp120 antibody.

mutation in the V3 region, could change R5 virus into an X4 virus [25,26].

It has been shown that mono-or bi-valent HIV-1 envelope vaccines are poorly immunogenic and induce a limited breadth of reactivity. Although Nabs have been shown to control some laboratory-adapted strains, these results cannot easily be extended to primary clinical HIV-1 isolates. In addition, induced Nabs from single HIV-1 envelope studies lacked cross-subtype neutralizing activity and could not neutralize other strains of HIV except the strain contained in the vaccine preparation [27-30]. In an interesting study, Zhan et al., showed that cocktail of 50 HIV-1 envelope in the context of DNA vector, recombinant vaccinia vector and recombinant proteins is able to induce both humoral and cell-mediated immune responses in immunized macaques. Vaccinated macaques showed lower virus titers and higher amount of CD4+T cells compared to unvaccinated animals after challenge with SHIV89.6P. In addition, four out of six vaccinated macaques survived 44 week after the virus challenge, suggesting that a multi-envelope vaccine may be an effective approach to control HIV-1 infection [33].

To increase the breadth of the immune response against HIV-1 strains, ten HIV-1 strains that had a non-syncytium-inducing phenotype and four that had a syncytium-inducing phenotype were selected from a bank of clinical isolates. Although 14 gp120 proteins may not maximize the immune response and may not represent all subtype B isolates; these results may provide clues regarding the


capacity of heterologous HIV-1 subtype strains to induce antiviral immune responses that recognize other HIV-1 subtypes. The multivalent approach to increase the breadth of the immune response against heterologous strains within a subtype could provide for enhanced vaccine protection within a geographic region.

Sequence analysis from the structure of gp120s revealed that despite vivid antigenic differences between R5 and X4 isolates, the structure of their gp120 core is similar. The sequences of the HIV-1 strains were analyzed by multiple sequence ClustalW alignment program and the phylogenetic tree was constructed using the Neighbor Tree-Maker. The evolutionary history of the cohort gp120 clones was compared to other HIV-1 strains from the same HIV-1 subtype or different subtype. Phylogenetic analysis confirmed that all gp120 clones were derived from subtype B. Gp120 proteins were expressed in the CHO cells to produce a glycosylation pattern similar to the one observed in the native structural form of gp120 found on the virus

**Table 2: Immunization strategies: Groups of four hamsters were immunized four times S.C. in three different sites with one month interval between each immunization.**

<b>Group A</b>	MPL adjuvant
<b>Group B</b>	HIV-gp120 protein #1 + MPL adjuvant
<b>Group C</b>	HIV-gp120 protein #1-4 + MPL adjuvant
<b>Group D</b>	HIV-gp120 protein #1-14 + MPL adjuvant

A total of 100 µg recombinant protein in 0.2 mg of monophosphoryl lipid A was injected per animal.




**Figure 4**

Antibody titers were determined in hamsters (n = 4) two weeks after the last immunization. The 96-well plates were coated with standard HIV-1 gp120 proteins (subtypes B, C and A/E). Hamster sera were serially diluted in wells and absorbance was read at 490 nm with an ELISA plate reader. Results are shown as mean concentration. ± S.E. is shown for each group.

envelope. Gp120 proteins were poorly expressed in transiently transfected CHO cells. To solve this problem, stable CHO cell lines were created. The production of gp120 proteins were increased more than 20-fold in the stable cell lines, allowing for efficient protein purification.

It may be possible that a cocktail of HIV-1 gp120 proteins containing multiple epitopes will be able to maximize the breadth of immune response against HIV-1 compared to a single immunogen. In a recent study, several macaques were primed by DNA encoding gp120 from subtypes A, B, and C and boosted with recombinant gp120 proteins. Although strong cell-mediated immune responses were induced, a potent neutralizing antibody response was not

detected in animals [32]. So far, limited polyvalent vaccine studies have been performed and none have measured immune responses against diverse HIV strains. On the other hand, an effective HIV vaccine may have to deal with not only multiple subtypes but with many strains in each subtype. It is interesting to note that the surface envelope in each HIV subtype has featured differences. For instance, HIV-1 subtype C has enormous genetic differences compared with the recombinant subtypes A/C. The question arises whether HIV-1 immunogens from one subtype will be effective against other subtype. Leandersson et al., showed T-cell reactivity between gp160 immunogen subtype B with other subtypes in HIV individuals [33]. However, they did not detect the neutralizing cross-



**Figure 5**

Neutralizing antibody responses in immunized groups. Values are the serum dilution at which relative luminescence units (RLU) were reduced 50% compared to virus control wells. The neutralization antibody responses were compared with SVA-MLV (Murine retrovirus Env-pseudovirus) as a background control for non-specific activity. Symbol \* indicates significant difference when compared with groups A and B ( $p < 0.05$ ).

reactivity between HIV-1 subtypes. Another concern in a polyvalent vaccine approach is that there are no studies regarding the synergistic or antagonistic effects of gp120 genes. It is possible that protein components in a polyvalent vaccine cocktail interact and compete with each other and consequently may inhibit or downgrade the response to some proteins.

Due to high background in mice sera, we were not able to measure the neutralizing antibody responses in our previous vaccine studies, thus, golden hamsters were considered for measurement of the efficacy of humoral immune responses. CD8+T cell immune response was not studied due to non-existent labelled antibodies to hamster CD3 and CD8. In order to analyze the binding antibody titre, plasma from immunized hamsters was collected and HIV-1 gp120 subtypes B (MN, SF162), C (96ZM651) and A/E (93TH975) were analyzed separately by ELISA. Group D, which received all fourteen proteins, showed higher mean IgG antibody titer to HIV-1 subtype B isolates MN and SF162 in comparison to groups B and C. While results from binding assays do not predict neutralization serotypes, they suggest that antigenic subtypes may be related, but not identical, to the genetic subtypes. Theoretically, an augmented number and amount of envelope proteins may enhance the control of viremia following HIV-1 infection through a number of immunologic mechanisms, including increasing the functional activity of B

cells, promoted neutralizing antibodies, and cross-reactivity with various HIV-1 subtypes.


In this study, the candidate vaccines were able to neutralize only the HIV-1 strain MN and not any other HIV-1 subtypes. Interestingly, the immunized group with fourteen-gp120 proteins subtype B induced a response similar to the group that received four gp120 proteins. Therefore, it will be useful to evaluate the immunogenicity of each gp120 and interaction with any other gp120 protein in a non-human primate model. These results are similar to a previously published polyvalent study where the neutralization was restricted to the virus strains used for vaccination and the polyvalent vaccine was not able to neutralize any primary HIV-1 isolates [34]. Lack of cross reactivity between Nab results were also consistent to a recent study by Cavacini et al., demonstrating no quantified cross-subtype antibody reactivity and neutralization between subtype B sera against subtype C HIV virus [35]. In another study, Wang et al. showed that a polyvalent HIV-1 envelope protein approach is not able to induce neutralizing antibody responses against HIV-1. However, a polyvalent HIV-1 envelope approach in the context of DNA priming followed by protein boosting may be able to neutralize some of HIV-1 strains from different subtypes [36].

HIV-1 patients are faced with low number of CD4+Th cells in different stages of disease. An inverse correlation between gag-specific Th cells and HIV-1 RNA plasma virus load has been shown. The strongest gag specific proliferative responses were associated with lower viral load, while a high viral load was observed with weak response to gag stimulation [37]. Here, lymphocyte proliferation assay was tested to determine activation of lymphocytes after encountering with HIV-1 gp120 proteins from different subtypes. A relatively high proliferative immune response was determined to subtypes B, C and A/E in the group of hamster that immunized with fourteen gp120 proteins subtype B. This group developed stimulation indices four fold or more above baseline against all subtypes. These results were in consistent with Currier JR et al. that showed a comprehensive cross-subtype reactivity of cell-mediated immune responses between individuals with different HIV-1 subtypes [38]. In summary, the polyvalent gp120 candidate vaccine derived from subtype B was not able to neutralize primary HIV-1 isolates. However, a substantial proportion of cross-subtype CD4 reactivity was observed between the polyvalent group with HIV-1 subtypes B, C, and A/E.

## Methods

### HIV Clinical Isolates

Viruses were isolated from HIV-infected individuals during their visits to the HIV clinic in the Ottawa Hospital. Verbal consent was obtained from HIV patients for partic-



**Figure 6**

T-cell proliferation results in hamsters immunized with the candidate HIV-1 gp120 vaccines. Splenocytes were cultured and stimulated with 10 µg/ml proteins of (6A): HIV-1 gp120 subtype B (MN strain), (6B): HIV-1 gp120 subtype B (SF162 strain), (6C): HIV-1 gp120 subtype C (96ZM651 strain), and (6D): HIV-1 gp120 subtype A/E (93TH975 strain). Two days later, <sup>3</sup>H-thymidine was added and after 18 h the incorporated radioactivity was measured in harvested splenocytes. The stimulation index (SI) ± S.E. is shown for each group. Symbol \* indicates a significant difference between group D with groups B and C (p ≤ 0.05).

ipation in the study. Briefly, isolation of virus was performed by placing 2–4 × 10<sup>6</sup> peripheral blood mononuclear cells (PBMCs) in Iscove's Modified Dulbecco's Medium (IMDM: Sigma, St. Louis, MO) supplemented with 10% fetal calf serum (FCS: Life Technologies, Grand Island, NY), 2 Mm Glutamine (Sigma) and 50 µg/ml gentamicine (Sigma) with an equal number of phytohemagglutinin (PHA: Sigma) stimulated donor cells. The cells were incubated at 37°C containing 5% CO<sub>2</sub> and 20 IU/ml Interleukin-2 (IL-2: R&D systems, Minneapolis, MN) was added to increase the growth of these cells. Co-cultures were maintained for up to 4 weeks with periodic media changes. 2 × 10<sup>6</sup> freshly stimulated donor cells were added every 10 days to sustain HIV replication. Viruses were measured in the supernatant by HIV p24 antigen kit (DuPont, Wilmington, DE).

**Amplification and cloning of HIV-1 gp120 genes into pEF6-myc-His vectors**

Total RNA was purified using RNeasy extraction kit (Qiagen, Mississauga, ON) from syncytium and non-syncytium inducing HIV-1 subtype B co-cultures. The encoding

gp120 sequences (1.6 kb) were amplified by RT-PCR. The amplicons were purified using the QIAquick gel extraction kit (Qiagen, Mississauga, ON) and cloned into the PCR 2.1 TOPO-TA vector (Invitrogen, Burlington, ON). After plasmid digestion, the 1.6 kb band corresponding to the gp120 genes were sub-cloned into the pEF6-myc-His vectors (Invitrogen) which is designed for the over-production of recombinant proteins in mammalian cell lines. In this system, recombinant proteins may be purified by affinity chromatography using the c-myc epitope or poly histidine (6 × His) metal-binding tags. The expression constructs were confirmed and characterized by restriction enzymes and nucleotide sequence analysis.

**Characterization of amplified gp120s**

All fourteen HIV-1 gp120-V3 regions were aligned using "ClustalW" which is a multiple sequence alignment program for DNA or proteins [39,40]. At the end, the sequences were also manually adjusted. The phylogenetic tree was constructed by the Neighbor Tree-Maker in HIV sequence database. The program converts sequences alignment to the FASTA format.



**Chinese Hamster Ovary (CHO) culture**

CHO cells were grown at 37°C, 5% CO<sub>2</sub> in IMDM (Sigma) supplemented with 10% FCS (Life Technologies), 100 U/ml penicillin and 100 µg/ml gentamicin (Sigma).

**Expression of recombinant proteins**

CHO cells were transfected with pEF6-myc-His vectors expressing HIV-1 gp120 structural genes. Stable cell lines were established by using blasticidine-supplemented medium. Cells were harvested, sonicated and lysed in lysis buffer (25 mM Tris base, 2.5 mM Mercaptoethanol, 1% Triton-X100 and a cocktail of protease inhibitors). Recombinant viral proteins were purified with the TALON metal resin kit (Clontech, Palo Alto, CA) as per manufacturer's instructions. The recombinant proteins were confirmed by western blotting and Immunofluorescence antibody staining as previously described [41-43].

**Animal experiments**

Six-week old golden hamsters were purchased from Jackson Laboratory (Bar Harbor, ME). Hamsters were maintained in accordance with laboratory standard procedures and were housed in Animal Care in the Faculty of Medicine at University of Ottawa. A total of 100 µg recombinant protein in 0.2 mg of Monophosphoryl lipid A (MPL: Corixa, Hamilton, MT) was injected per animal. Groups of four hamsters were immunized four times subcutaneously (S.C.) in three different sites with one month interval between each immunization. Fourteen days after the last boost, the hamsters were sacrificed and their spleens and blood were collected for further testing or long-term storage in cryopreservation medium.

**Antibody measurement**

96-well ELISA plates were coated overnight at 4°C with gp120 proteins clade B (MN, SF162), C (96ZM651) and A/E (93TH975) obtained from the AIDS Research and Reference Reagent Program, National Institute of Allergy and Infectious Diseases (NIAID). The plates were washed with PBS containing 0.05% Tween 20 and then blocked for 1 h with 1% BSA in PBS. Following a second wash with PBS/0.05% Tween 20, the plates were incubated for 2 h at 37°C with serially diluted sera. The plates were washed and incubated for 2 h with peroxidase-conjugated affinity-purified rabbit anti-hamster secondary antibody (Bio-Rad). After washing, color was developed with O-phenylenediamine dihydrochloride (OPD: Sigma, St. Louise, MO). The color reaction was stopped with 1 N HCl and absorbance was read at 490 nm with an ELISA plate reader (Bio-Rad).

**Neutralizing antibody assay**

Neutralization was measured as a function of reduction in luciferase reporter gene expression after a single round of

infection in TZM-bl cells. TZM-bl cells were obtained from the NIH AIDS Research and Reference Reagent Program, as contributed by John Kappes and Xiaoyun Wu. Briefly, 200 TCID<sub>50</sub> of virus was incubated with serial 3-fold dilutions of serum sample in triplicate in a total volume of 150 µl for 1 hr at 37°C in 96-well flat-bottom culture plates. Freshly trypsinized TZM-bl cells (10,000 cells in 100 µl of growth medium containing 75 µg/ml DEAE dextran) were added to each well. Indinavir was added at a final concentration of 1 µM to prevent virus replication in the case of HIV-1 MN. One set of control wells received cells + virus (virus control) and another set received cells only (background control). After a 48 hour incubation, 100 µl of cells was transferred to 96-well black solid plates (Corning Costar, Rochester, NY) for measurement of luminescence using Bright Glo substrate solution as described by the supplier (Promega, Madison WI). An assay stock of HIV-1 MN was prepared in H9 cells. Assays stocks of Env-pseudotyped viruses were prepared by transfection in 293T cells. All virus stocks were made cell-free by 0.45-micron filtration and were stored at -70°C until use.

**Lymphocyte proliferation assay**

The assay was performed by [<sup>3</sup>H] thymidine incorporation as previously described [44]. Briefly, splenocytes from immunized hamsters were resuspended at 2 × 10<sup>6</sup> cells/ml in RPMI 1640 containing 10% FCS, 50 µM β-mercaptoethanol and 100 U/ml penicillin/streptomycin. A 100 µl aliquot containing 2 × 10<sup>5</sup> cells was added to each well of a 96 well plate. The gp120 protein subtypes B, C and A/E (100 µl at 20 µg/ml) were added to each well in triplicate. As a positive control, cells were also stimulated with phorbol 12-myristate 13-acetate and ionomycin (PMA/ION: Sigma). After 48 h of culture, 1 µCi [<sup>3</sup>H] thymidine (Amersham, Arlington Heights, IL) was added to each well. Following 16 h of incubation, cells were harvested onto glass fibre filtermats and thymidine incorporation was measured with a Microbeta beta counter (Wallac, Turku, Finland).

**Abbreviations**

Nabs: Neutralizing antibodies

CHO: Chinese hamster ovary

**Authors' contributions**

AA performed and analyzed majority of the experiments, participated in the design and wrote the manuscript. DEA participated in analyzing the data. MG helped in the cloning. KG revised the paper critically. FDM participated in the design and coordination of study and helped to draft the final manuscript. All authors read and approved the final manuscript.

## Acknowledgements

We thank Dr. David Montefiori at Duke University for the neutralizing antibody experiments. We thank Dr. William Cameron in the HIV Clinic at Ottawa Hospital for providing the blood samples. We thank D. Albert and M.C. Renaud for their technical assistance. We thank the AIDS Research and Reference Reagent program for providing some of reagents. This study was partially funded by a grant from Canadian Institutes of Health Research (CIHR) to FDM. AA is grateful to NSERC of Canada for the Industrial R&D Fellowship.

## References

- Hotzel I, Cheevers WP: **Caprine arthritis-encephalitis virus envelope surface glycoprotein regions interacting with the transmembrane glycoprotein: structural and functional parallels with human immunodeficiency virus type I gp120.** *J Virol* 2003, **77**:11578-11587.
- Kim YB, Han DP, Cao C, Cho MW: **Immunogenicity and ability of variable loop-deleted human immunodeficiency virus type I envelope glycoproteins to elicit neutralizing antibodies.** *Virology* 2003, **305**:124-137.
- Grundner C, Pancera M, Kang JM, Koch M, Sodroski J, Wyatt R: **Factors limiting the immunogenicity of HIV-1 gp120 envelope glycoproteins.** *Virology* 2004, **330**:233-248.
- Kwong PD, Wyatt R, Majeed S, Robinson J, Sweet RW, Sodroski J, Hendrickson WA: **Structures of HIV-1 gp120 envelope glycoproteins from laboratory-adapted and primary isolates.** *Structure Fold Des* 2000, **8**:1329-1339.
- Wyatt R, Kwong PD, Desjardins E, Sweet RW, Robinson J, Hendrickson WA, Sodroski JG: **The antigenic structure of the HIV gp120 envelope glycoprotein.** *Nature* 1998, **393**:705-711.
- Kandathil AJ, Ramalingam S, Kannangai R, David S, Sridharan G: **Molecular epidemiology of HIV.** *Indian J Med Res* 2005, **121**:333-344.
- Choisy M, Woelk CH, Guegan JF, Robertson DL: **Comparative study of adaptive molecular evolution in different human immunodeficiency virus groups and subtypes.** *J Virol* 2004, **78**:1962-1970.
- Ciurea A, Klenerman P, Hunziker L, Horvath E, Senn BM, Ochsenbein AF, Hengartner H, Zinkernagel RM: **Viral persistence in vivo through selection of neutralizing antibody-escape variants.** *Proc Natl Acad Sci U S A* 2000, **97**:2749-2754.
- Leung L, Srivastava IK, Kan E, Legg H, Sun Y, Greer C, Montefiori DC, zur MJ, Barnett SW: **Immunogenicity of HIV-1 Env and Gag in baboons using a DNA prime/protein boost regimen.** *AIDS* 2004, **18**:991-1001.
- Johnston R: **AIDSVAX results: an answer, or just more questions?** *AIDS Patient Care STDS* 2003, **17**:47-51.
- Billich A: **AIDSVAX. VaxGen.** *Curr Opin Investig Drugs* 2001, **2**:1203-1208.
- Billich A: **AIDSVAX VaxGen.** *Curr Opin Investig Drugs* 2004, **5**:214-221.
- Montefiori DC: **Neutralizing antibodies take a swipe at HIV in vivo.** *Nat Med* 2005, **11**:593-594.
- Sato H, Kato K, Takebe Y: **Functional complementation of the envelope hypervariable V3 loop of human immunodeficiency virus type I subtype B by the subtype E V3 loop.** *Virology* 1999, **257**:491-501.
- Hu DJ, Vanichseni S, Mastro TD, Raktham S, Young NL, Mock PA, Subbarao S, Parekh BS, Srisuwanvilai L, Sutthent R, Wasi C, Heneine W, Choopanya K: **Viral load differences in early infection with two HIV-1 subtypes.** *AIDS* 2001, **15**:683-691.
- Ferrantelli F, Rasmussen RA, Hofmann-Lehmann R, Xu W, McClure HM, Ruprecht RM: **Do not underestimate the power of antibodies--lessons from adoptive transfer of antibodies against HIV.** *Vaccine* 2002, **20 Suppl 4**:A61-A65.
- Nishimura Y, Igarashi T, Haigwood NL, Sadjadpour R, Donau OK, Buckler C, Plishka RJ, Buckler-White A, Martin MA: **Transfer of neutralizing IgG to macaques 6 h but not 24 h after SHIV infection confers sterilizing protection: implications for HIV-1 vaccine development.** *Proc Natl Acad Sci U S A* 2003, **100**:15131-15136.
- Rasmussen RA, Hofmann-Lehmann R, Li PL, Vlasak J, Schmitz JE, Reimann KA, Kuroda MJ, Letvin NL, Montefiori DC, McClure HM, Ruprecht RM: **Neutralizing antibodies as a potential secondary protective mechanism during chronic SHIV infection in CD8+ T-cell-depleted macaques.** *AIDS* 2002, **16**:829-838.
- Krachmarov CP, Kayman SC, Honnen WJ, Trochev O, Pinter A: **V3-specific polyclonal antibodies affinity purified from sera of infected humans effectively neutralize primary isolates of human immunodeficiency virus type I.** *AIDS Res Hum Retroviruses* 2001, **17**:1737-1748.
- Inouye P, Cherry E, Hsu M, Zolla-Pazner S, Wainberg MA: **Neutralizing antibodies directed against the V3 loop select for different escape variants in a virus with mutated reverse transcriptase (M184V) than in wild-type human immunodeficiency virus type I.** *AIDS Res Hum Retroviruses* 1998, **14**:735-740.
- Wei X, Decker JM, Wang S, Hui H, Kappes JC, Wu X, Salazar-Gonzalez JF, Salazar MG, Kilby JM, Saag MS, Komarova NL, Nowak MA, Hahn BH, Kwong PD, Shaw GM: **Antibody neutralization and escape by HIV-1.** *Nature* 2003, **422**:307-312.
- Wang FX, Kimura T, Nishihara K, Yoshimura K, Koito A, Matsushita S: **Emergence of autologous neutralization-resistant variants from preexisting human immunodeficiency virus (HIV) quasi species during virus rebound in HIV type I-infected patients undergoing highly active antiretroviral therapy.** *J Infect Dis* 2002, **185**:608-617.
- Chen CH, Jin L, Zhu C, Holz-Smith S, Matthews TJ: **Induction and characterization of neutralizing antibodies against a human immunodeficiency virus type I primary isolate.** *J Virol* 2001, **75**:6700-6704.
- Schreiber M, Wachsmuth C, Muller H, Odemuyiwa S, Schmitz H, Meyer S, Meyer B, Schneider-Mergener J: **The V3-directed immune response in natural human immunodeficiency virus type I infection is predominantly directed against a variable, discontinuous epitope presented by the gp120 V3 domain.** *J Virol* 1997, **71**:9198-9205.
- Yamaguchi-Kabata Y, Yamashita M, Ohkura S, Hayami M, Miura T: **Linkage of amino acid variation and evolution of human immunodeficiency virus type I gp120 envelope glycoprotein (subtype B) with usage of the second receptor.** *J Mol Evol* 2004, **58**:333-340.
- Jensen MA, Li FS, 't Wout AB, Nickle DC, Shriner D, He HX, McLaughlin S, Shankarappa R, Margolick JB, Mullins JI: **Improved coreceptor usage prediction and genotypic monitoring of R5-to-X4 transition by motif analysis of human immunodeficiency virus type I env V3 loop sequences.** *J Virol* 2003, **77**:13376-13388.
- Srivastava IK, Ulmer JB, Barnett SW: **Neutralizing antibody responses to HIV: role in protective immunity and challenges for vaccine design.** *Expert Rev Vaccines* 2004, **3**:533-552.
- Li M, Gao F, Mascola JR, Stamatatos L, Polonis VR, Koutsoukos M, Voss G, Goepfert P, Gilbert P, Greene KM, Bilska M, Kothe DL, Salazar-Gonzalez JF, Wei X, Decker JM, Hahn BH, Montefiori DC: **Human Immunodeficiency Virus Type I env Clones from Acute and Early Subtype B Infections for Standardized Assessments of Vaccine-Elicited Neutralizing Antibodies.** *J Virol* 2005, **79**:10108-10125.
- Sarmati L, d'Ettorre G, Nicastrì E, Ercoli L, Uccella I, Massetti P, Parisi SG, Vullo V, Andreoni M: **Neutralizing antibodies against autologous human immunodeficiency virus Type I isolates in patients with increasing CD4 cell counts despite incomplete virus suppression during antiretroviral treatment.** *Clin Diagn Lab Immunol* 2001, **8**:822-824.
- Locher CP, Grant RM, Collisson EA, Reyes-Teran G, Elbeik T, Kahn JO, Levy JA: **Antibody and cellular immune responses in breakthrough infection subjects after HIV type I glycoprotein 120 vaccination.** *AIDS Res Hum Retroviruses* 1999, **15**:1685-1689.
- Zhan X, Martin LN, Slobod KS, Coleclough C, Lockett TD, Brown SA, Stambas J, Bonsignori M, Sealy RE, Blanchard JL, Hurwitz JL: **Multi-envelope HIV-1 vaccine devoid of SIV components controls disease in macaques challenged with heterologous pathogenic SHIV.** *Vaccine* 2005, **23**:5306-5320.
- Pal R, Wang S, Kalyanaraman VS, Nair BC, Whitney S, Keen T, Hocker L, Hudacik L, Rose N, Cristillo A, Mboudjeka I, Shen S, Wu-Chou TH, Montefiori D, Mascola J, Lu S, Markham P: **Polyvalent DNA prime and envelope protein boost HIV-1 vaccine elicits humoral and cellular responses and controls plasma viremia in rhesus macaques following rectal challenge with an R5 SHIV isolate.** *J Med Primatol* 2005, **34**:226-236.

33. Leandersson AC, Gilljam G, Fredriksson M, Hinkula J, Alaeus A, Lidman K, Albert J, Bratt G, Sandstrom E, Wahren B: **Cross-reactive T-helper responses in patients infected with different subtypes of human immunodeficiency virus type 1.** *J Virol* 2000, **74**:4888-4890.
34. Cho MW, Kim YB, Lee MK, Gupta KC, Ross W, Plishka R, Buckler-White A, Igarashi T, Theodore T, Byrum R, Kemp C, Montefiori DC, Martin MA: **Polyvalent envelope glycoprotein vaccine elicits a broader neutralizing antibody response but is unable to provide sterilizing protection against heterologous Simian/human immunodeficiency virus infection in pigtailed macaques.** *J Virol* 2001, **75**:2224-2234.
35. Cavacini LA, Duval M, Patil A, Wood C, Mayer KH, Ruprecht RM, Posner MR: **Dichotomy in cross-clade reactivity and neutralization by HIV-1 sera: Implications for active and passive immunotherapy.** *J Med Virol* 2005, **76**:146-152.
36. Wang S, Pal R, Mascola JR, Chou TH, Mboudjeka I, Shen S, Liu Q, Whitney S, Keen T, Nair BC, Kalyanaraman VS, Markham P, Lu S: **Polyvalent HIV-1 Env vaccine formulations delivered by the DNA priming plus protein boosting approach are effective in generating neutralizing antibodies against primary human immunodeficiency virus type 1 isolates from subtypes A, B, C, D and E.** *Virology* 2006, **350**:34-47.
37. Klein MR, Veenstra J, Holwerda AM, Roos MT, Gow I, Patou G, Coutinho RA, De Wolf F, Miedema F: **Gag-specific immune responses after immunization with p17/p24:Ty virus-like particles in HIV type 1-seropositive individuals.** *AIDS Res Hum Retroviruses* 1997, **13**:393-399.
38. Currier JR, Dowling WE, Wasunna KM, Alam U, Mason CJ, Robb ML, Carr JK, McCutchan FE, Birx DL, Cox JH: **Detection of high frequencies of HIV-1 cross-subtype reactive CD8 T lymphocytes in the peripheral blood of HIV-1-infected Kenyans.** *AIDS* 2003, **17**:2149-2157.
39. Pavlovic-Lazetic GM, Mitic NS, Beljanski MV: **Bioinformatics analysis of SARS coronavirus genome polymorphism.** *BMC Bioinformatics* 2004, **5**:65.
40. Li KB: **ClustalW-MPI: ClustalW analysis using distributed and parallel computing.** *Bioinformatics* 2003, **19**:1585-1586.
41. Ghorbani M, Nass T, Azizi A, Soare C, Aucoin S, Giulivi A, Anderson DE, Diaz-Mitoma F: **Comparison of antibody- and cell-mediated immune responses after intramuscular hepatitis C immunizations of BALB/c mice.** *Viral Immunol* 2005, **18**:637-648.
42. Azizi A, Aucoin S, Tadesse H, Frost R, Ghorbani M, Soare C, Naas T, Diaz-Mitoma F: **A combined nucleocapsid vaccine induces vigorous SARS-CD8+ T-cell immune responses.** *Genet Vaccines Ther* 2005, **3**:7.
43. Naas T, Ghorbani M, Alvarez-Maya I, Lapner M, Kothary R, De Repentigny Y, Gomes S, Babiuk L, Giulivi A, Soare C, Azizi A, Diaz-Mitoma F: **Characterization of liver histopathology in a transgenic mouse model expressing genotype 1a hepatitis C virus core and envelope proteins 1 and 2.** *J Gen Virol* 2005, **86**:2185-2196.
44. Azizi A, Ghorbani M, Kryworuchko M, Aucoin S, Diaz-Mitoma F: **Potency of cell-mediated immune responses to different combined HIV-1 immunogens in a humanized murine model.** *Hum.Vaccin* 2005, **1**(4):170-176.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

