

Article

Venomic Analysis of the Poorly Studied Desert Coral Snake, *Micrurus tschudii tschudii*, Supports the 3FTx/PLA₂ Dichotomy across *Micrurus* Venoms

Libia Sanz ¹, Davinia Pla ¹, Alicia Pérez ¹, Yania Rodríguez ¹, Alfonso Zavaleta ^{2,3}, Maria Salas ², Bruno Lomonte ⁴ and Juan J. Calvete ^{1,*}

¹ Laboratorio de Venómica Estructural y Funcional, Instituto de Biomedicina de Valencia, CSIC, Jaime Roig 11, Valencia 46010, Spain; libia.sanz@ibv.csic.es (L.S.); dpla@ibv.csic.es (D.P.); aperez@ibv.csic.es (A.P.); yrodriguez@ibv.csic.es (Y.R.)

² Departamento Academico de Ciencias Celulares y Moleculares, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima 31, Perú; alfonso.zavaleta@upch.pe (A.Z.); maria.salas@upch.pe (M.S.)

³ Instituto Nacional de Salud, Ministerio de Salud, Lima 11, Perú

⁴ Instituto Clodomiro Picado, Facultad de Microbiología, Universidad de Costa Rica, San Jose 11501, Costa Rica; bruno.lomonte@ucr.ac.cr

* Correspondence: jcalvete@ibv.csic.es; Tel.: +34-96-339-1778; Fax: +34-96-369-0800

Academic Editor: Stephen P. Mackessy

Received: 23 March 2016; Accepted: 1 June 2016; Published: 7 June 2016

Abstract: The venom proteome of the poorly studied desert coral snake *Micrurus tschudii tschudii* was unveiled using a venomic approach, which identified ≥ 38 proteins belonging to only four snake venom protein families. The three-finger toxins (3FTxs) constitute, both in number of isoforms (~30) and total abundance (93.6% of the venom proteome), the major protein family of the desert coral snake venom. Phospholipases A₂ (PLA₂s; seven isoforms, 4.1% of the venom proteome), 1–3 Kunitz-type proteins (1.6%), and 1–2 L-amino acid oxidases (LAO, 0.7%) complete the toxin arsenal of *M. t. tschudii*. Our results add to the growing evidence that the occurrence of two divergent venom phenotypes, *i.e.*, 3FTx- and PLA₂-predominant venom proteomes, may constitute a general trend across the cladogenesis of *Micrurus*. The occurrence of a similar pattern of venom phenotypic variability among true sea snake (Hydrophiinae) venoms suggests that the 3FTx/PLA₂ dichotomy may be widely distributed among Elapidae venoms.

Keywords: *Micrurus tschudii tschudii* venom; venomics; snake venom proteome; three-finger toxin; snake venom phospholipase A₂; mass spectrometry

1. Introduction

New World genus *Micrurus* (Elapidae) (Wagler, 1824) [1] represents a monophyletic clade of some 80 currently recognized species of venomous coral snakes [2–8], although the topology of the tree is still unresolved. Coral snakes are widely distributed in tropical and subtropical regions from the southern United States to northeastern Argentina, including several continental islands inhabited by endemic forms [2,3]. Coral snakes are considered by herpetologists to be among the most beautiful snakes of the planet, as they are adorned with unique combinations of red-, black-, and yellow-colored banding. Based on their body ring color pattern characteristics, tail proportion with respect to body length, and hemipenial morphology, coral snakes are traditionally arranged in four species groups: a tricolored monadal group, a bicolored group, a Central American tricolored triadal group, and a South American triadal group [3,9–12]. The monadal group comprises long-tailed species with a single black ring separating the white and red rings; the bicolored group contains short-tailed species

with white or red rings separated by black rings; the Central American triadal group taxa are long-tailed species with three black rings separated by white rings between red rings; the South American triadal group is represented by short-tailed species with the same color pattern as the Central American triadal group species but is restricted geographically to South America, extending from Panama to southern Argentina.

In the last decade, the increasing application of omics techniques to the study of snake venoms has greatly enhanced our knowledge on their composition, evolution, biological activities, and clinical effects [13–17]. However, the venoms of only a handful of the vast number (~130) of species and subspecies that constitute the genus *Micrurus* have been the subject of proteomic studies (consult Table 2 of [18]). To understand this fact it should be taken into consideration that, despite producing among the most potent neurotoxic venoms of any New World snake, bites and fatalities by coral snakes are very rare. This is facilitated by the fact that coral snakes are not aggressive; when confronted by humans, coral snakes will almost always attempt to flee, and bite only as a last resort. On the other hand, coral snakes generally inhabit sparsely populated areas and are thus infrequently encountered, and they have short fangs that cannot penetrate thick leather clothing. Consequently, envenomings by coral snakes are less frequent than those produced by sympatric pitvipers, by far the most dangerous snakes of South America, accounting for less than 3% of the snakebites recorded in the Americas [19–21].

The desert coral snake *M. tschudii tschudii* (January 1858) [22], one of the lesser-studied species of the genus *Micrurus*, belongs to the South American triad coral snakes [11,12,23]. Named after the Swiss naturalist, explorer and diplomat Johann Jakob von Tschudi (1818–1889), who traveled extensively in South America and published important works in herpetology, *M. t. tschudii* is a small (adults average 45 to 55 cm in length), tri-colored (Figure 1), mainly diurnal and terrestrial coral snake, found in tropical deciduous forest, dry tropical forest, and thorn scrub, mainly along watercourses, from near sea level to 1450 m elevation, in the western slopes of the Andes on the semi-arid Pacific coast of South America, from southern Ecuador to southwestern Peru [2,3,24]. The desert coral snake feeds on geckkonid lizards (*Phyllodactylus* spp.), Amphisbaenids (*A. occidentalis*), and colubrids of the species *Mastigodryas haathi* [3]. However, available literature about its reproduction and activity is scarce, and in particular virtually nothing (not even a single entry in PubMed) is known about its venom. Here, we have applied a venomomics approach [13,14] to gain an insight into the spectrum of toxins that make up the venom proteome of *M. t. tschudii*.

2. Results and Discussion

2.1. The Venom Proteome of the Desert Coral Snake

The venom of *M. t. tschudii* was fractionated into 26 RP-HPLC fractions (Figure 1A). Each chromatographic fraction was analyzed by SDS-PAGE (Figure 1B), and the protein bands were excised and submitted to mass spectrometric analysis. The MS/MS data, listed in Supplementary Table S1, resulted in the identification of ≥ 38 proteins belonging to only four snake venom protein families: three-finger toxin (3FTx), phospholipase A₂ (PLA₂), Kunitz-type, and L-amino acid oxidase (LAO) (Table 1). The relative abundances of *M. t. tschudii* venom proteins and toxin families are displayed, respectively, in Table S1 and Figure 1C. The 3FTxs constitute, by far, both in number of isoforms (~30) and total abundance (93.6% of the total venom proteins), the major protein family of the desert coral snake venom. Seven PLA₂ isoforms, 1–3 Kunitz-type proteins, and 1–2 LAO molecules (accounting, respectively, for 4.1%, 1.6%, and 0.7% of the venom proteome) complete the toxin arsenal of the desert coral snake.




Figure 1. Panel (A) reverse-phase HPLC separation of the venom proteins from *M. t. tschudii*. Photo credit: Dr. Med. Vet. Gualberto Marcas Cáceres, Centro Nacional de Productos Biológicos, Instituto Nacional de Salud, Ministerio de Salud, Perú; Panel (B) SDS-PAGE of the isolated chromatographic fractions run under reduced conditions; Panel (C) displays the relative abundance (in % of the total venom proteins) of the toxin families found in *M. t. tschudii* venom.

The 3FTx and PLA₂ molecules are hallmark components of the venoms of Elapidae. Catalytically active PLA₂ molecules typically exhibit presynaptic neurotoxic activity, myotoxic activity, or both, although some forms exhibit antiplatelet activity [25,26]. Despite their pronounced structural similarity, members of the 3FTx family exhibit a wide variety of pharmacological effects including postsynaptic neurotoxicity, cytotoxicity, cardiotoxicity, and anticoagulant, and antiplatelet activities [27,28]. In addition, L-type Ca²⁺ channel antagonists of the 3FTx family may act synergistically with muscarinic three-finger toxins to promote hypotension [29]. Kunitz-type serine protease inhibitor isolated from elapid venoms blocked the activity of a range of serine proteases, producing an antihemorrhagic effect [30]. Non-covalent PLA₂-KUN complexes have been characterized from venom of some *Micrurus* species [18 and references cited]. These complexes, originally discovered in the venom of the Texas coral snake *M. tener*, are target acid-sensing receptors ASIC1a/2 evoking pain [31,32]. There are many examples of venom-derived toxins that elicit notoriously intense pain [33]. Pain response may cause paralysis and serve as a warning signal for discouraging potentially threatening predators by triggering lasting acute physiological distress. The L-amino acid oxidases are flavoenzymes that catalyze oxidative deamination of L-amino acids to form corresponding α -keto acids, hydrogen peroxide and ammonia. LAOs are thought to contribute to the toxicity of the venom due to the production of hydrogen peroxide during the oxidation reaction [34,35], although their contribution to the envenoming process remains elusive.

The low yield of venom and the challenge of long-term maintenance of the desert coral snake in captivity precluded a detailed toxicovenomics analysis of the venom components of this poorly studied *Micrurus* species.

2.2. 3FTx- and PLA₂-Predominant Venom Proteomes among Coral Snakes

Despite their extremely adaptive ecological radiation, coral and sea snakes show virtually identical rates of body size evolution to other elapids, suggesting that their diversification may be correlated with phenotypic traits other than body size [36]. Given the central role that diet has played in the adaptive radiation of snakes [37], venom represents a key adaptation that has played an important role in the diversification of these animals [38,39]. Growing evidence, including the venom analysis of the desert coral snake here reported, supports the view that the occurrence of two divergent venom phenotypes, *i.e.*, 3FTx- and PLA₂-predominant venom proteomes, may constitute a general trend across *Micrurus* cladogenesis [18,40]. Thus, *M. alleni* (77% 3FTx *vs.* 11% PLA₂) and *M. mosquitensis* (56% PLA₂, 22% 3FTx) [18]; *M. corallinus* (82% 3FTx, 12% PLA₂) [41]; *M. nigrocinctus* (38% 3FTx, 48% PLA₂) [42]; *M. fulvius* (21% 3FTx, 65% PLA₂) [43,44]; *M. tener* (38% 3FTx, 46% PLA₂) [45]; *M. dumerilii* (28% 3FTx, 52% PLA₂) [40]; and *M. clarki* (48.2% 3FTx, 36.5% PLA₂) [46] all are species included in the well-supported monophyletic monadal clade [3,10,47,48]. The 22 valid species of coral snakes classified in the South American triadal group, to which *M. t. tschudii* belongs, also form a strongly supported clade [3,11,12,23]. Biochemical and/or proteomic analyses have been reported for the venoms of *M. altirostris* (80% 3FTx, 14% PLA₂) [44], *M. surinamensis* (3FTx-predominant venom) [49], *M. frontalis* (3FTx-predominant venom) [50], *M. pyrrhocryptus* (3FTx-rich venom proteome) [51], and *M. clarki* (moderately 3FTx-predominant venom) [46]. On the other hand, the venom proteomes of *M. multifasciatus* and *M. mipartitus* from the bicolor species group are both dominated by 3FTxs [52].

Table 1. Comparison of reported median lethal doses (LD₅₀, µg venom/g mouse) for mice of 3FTx- and PLA₂-predominant *Micrurus* venoms. Route of venom administration: *i.v.*, intravenous; *i.p.*, intraperitoneal; *s.c.*, sub-cutaneous; *tw*, this work.

3FTx-Rich Venom	LD ₅₀	Reference	PLA ₂ -Rich Venom	LD ₅₀	References
<i>M. altirostris</i>	<i>i.p.</i> 0.26–0.65	[41]	<i>M. fulvius</i>	<i>i.v.</i> 0.32 ± 0.12 <i>i.p.</i> 2.60–4.40	[43] [53]
<i>M. corallinus</i>	<i>i.p.</i> 0.25–1.35	[41]	<i>M. tener</i>	<i>s.c.</i> 4.4 <i>i.v.</i> 0.78 ± 0.14	[45] [54,55]
<i>M. mipartitus</i>	<i>i.p.</i> 0.47	[52]	<i>M. nigrocinctus</i>	<i>i.v.</i> 0.3–0.5 <i>i.p.</i> 0.4–1.2 <i>s.c.</i> 1.7–2.5	[45]
<i>M. multifasciatus</i>	<i>i.p.</i> 1.35	[52]	<i>M. mosquitensis</i>	<i>i.v.</i> 0.20–0.61	[18]
<i>M. alleni</i>	<i>i.v.</i> 0.23–0.55 <i>i.v.</i> 0.74 ± 0.16	[18]	<i>M. dumerilii</i>	<i>i.p.</i> 0.8–1.9	[40]
<i>M. tschudii</i>	<i>i.p.</i> 0.44–0.81	[<i>tw</i>]			
<i>M. frontalis</i>	<i>i.p.</i> 0.20–1.45	[53]			
<i>M. pyrrhocryptus</i>	<i>i.p.</i> 1.10 ± 0.10	[51]			
<i>M. surinamensis</i>	<i>i.p.</i> 2.15–4.35	[53]			
<i>M. clarki</i>	<i>i.v.</i> 0.42–1.38	[46]			

Experiments in mice determined the Median Lethal Dose (LD₅₀) for *M. t. tschudii* venom at 10.5 µg/mouse (7.4–13.8 µg/mouse, 95% confidence limits) by the *i.p.* route. This figure corresponds to 0.62 µg venom/g of mouse body weight (0.44–0.81 µg/g, 95% confidence limits) (Table 1). Comparison of reported median lethal doses (LD₅₀) clearly shows that 3FTx-rich and PLA₂-rich *Micrurus* venoms are not statistically different in terms of lethality to mice (Table 1). Although a more meaningful correlation would require comparing LD₅₀ values for their natural prey, current data suggest that *Micrurus* venoms may have evolved under the action of balancing selection [56].

Margres and coworkers [44] have detected strong evidence of positive selection for the 3FTx and PLA₂ toxin families of *M. fulvius*. This accelerated evolution is most likely due to their direct involvement in fitness. Figure 2 shows a clear geographical distribution of PLA₂- and

3FTx-predominant *Micrurus* venoms along their north-south dispersal axis, further supporting the adaptive nature of this phenotypic dichotomy.




Figure 2. Geographic distribution of *Micrurus* species for which quantitative estimation of proteome compositions have been reported in the literature (Table 1). Distribution ranges were adapted from [3] and The Reptile Database (<http://www.reptile-database.org>), and are color-coded: green, PLA₂-rich venom phenotype; red, 3FTx-predominant venom composition. The arrow highlights the trend towards diverging venom phenotypes along the *Micrurus* north-south dispersal, suggesting the epicenter of the divergence in Mesoamerica.

2.3. Tracing the Evolutionary Origin of the 3FTx/PLA₂ Dichotomy across Elapidae

Elapid snakes are a relatively young group. The crown elapid radiation is approximately 38 million years (My) old, yet elapid snakes exhibit some of the highest diversification rates in reptiles [57]. In particular, two clades, *Hydrophis* and *Micrurus*, show anomalously high rates of diversification within Elapidae [36]. The Asian coral snake genus *Calliophis* emerges as monophyletic and the

sister group to all other American and Asian coral snakes [47], suggesting an Asiatic origin for the common ancestor of these elapids [58]. Coral snakes diversified around 30–25 My ago, and sea snakes (Hydrophiini) are approximately 16 My old [36]. Estimated divergence times suggest that Hydrophiini is a young and rapidly speciating clade that had a common ancestor ~8 My ago, although the majority of extant lineages diversified more recently, over the last ~1.5–3.5 My [36,59].

A similar pattern of venom phenotypic variability has been documented by the proteomic analysis of true sea snake (Hydrophiinae) venoms. Hence, the venom of *A. laevis* comprises mainly PLA₂ (71%) and 3FTxs (25%) [60], whereas the venoms of *H. schistosus* [61], *H. cyanocinctus* [62], and *P. platura* [63] all share 3FTx-rich phenotypes (*i.e.*, 70.5% 3FTx/26.5% PLA₂; 81% 3FTx/19% PLA₂; and 50% 3FTx/33% PLA₂, respectively). Transcripts for 3FTx and PLA₂ molecules also represent the two main constituents of the venom gland toxin-encoding transcriptomes of *Acalyptophis peronii* (77.5% 3FTx/17.5 PLA₂) and *Lapemis curtus* (92.4% 3FTx; 7.5% PLA₂) [64].

The evolutionary origin and adaptive relevance of the puzzling 3FTx/PLA₂ dichotomy remains elusive since 3FTx- and PLA₂-predominant venoms are scattered through the phylogenetic tree of *Micrurus*. The dominant protein families in the venom proteome of *C. bivirgata flaviceps* are PLA₂ (41%), 3FTx (22.6%) and SVMP (18.7%) [65], suggesting that the ancestor venom phenotype of coral snakes might have been of the PLA₂-predominant type. Alternatively, 3FTx- and PLA₂-rich venom proteomes may represent pedomorphic and ontogenetic traits, as has been documented in *Crotalus*. Rattlesnake venoms belong to one of two distinct phenotypes, which broadly correspond to type I (high levels of SVMPs and low toxicity, LD₅₀ > 1 mg/g mouse body weight) and type II (low metalloproteinase activity and high toxicity, LD₅₀ < 1 mg/g mouse body weight) [66]. In Neotropical rattlesnakes, the adaptive pressure for type I to type II transition was the gain of neurotoxicity and lethality to rodents, and this transition represents a miRNA-modulated pedomorphic trait that correlates with the increased concentration of crotoxin along the axis of *Crotalus* radiation in South America [67,68]. In Nearctic species, such as *C. s. scutulatus* (Css), the venom phenotype changes geographically from SVMP-rich to Mojave toxin-rich (type-II) as one moves from south central to southeastern Arizona, with a transitional zone between the SVMP and Mojave toxin phenotypes [69]. Understanding the phylogenetic origin of the 3FTx-rich and PLA₂-predominant venom phenotypes across *Micrurus*, and whether there is parallelism between the 3FTx/PLA₂ (*Micrurus*) and the type I/type II (*Crotalus*) venom dichotomies, requires genus-wide profiling of the venom proteomes and the venom gland transcriptomes of adult and juvenile *Crotalus* and *Micrurus* congeneric specimens.

3. Concluding Remarks and Perspectives

While genomic studies on model organisms have been widely applied to identify traits involved in maintaining functional genetic variation, few studies have drawn the links between genotype, phenotype, and fitness, and the environmental pressures that act to maintain variation that affects organismal phenotypes [70,71]. Venom represents a key adaptive trophic trait that has played an important role in the radiation of advanced snakes, and thus could be a particularly powerful model system to investigate players and mechanisms of adaptive variation at the phenotype level [17,70,72]. In this regard, structural and functional studies of *Micrurus* venoms point to balancing selection as the mechanism acting to maintain a 3FTx/PLA₂ venom dichotomy. The realization that this 3FTx/PLA₂ dichotomy may have deep roots in the evolution of coral snakes may also have important translational implications. Thus, both types of venoms should be part of immunization mixtures aimed at generating broad-spectrum antivenoms.

4. Materials and Methods

4.1. Venom

Adult *M. t. tschudii* specimens were caught in the Peruvian Pacific coast regions of La Libertad, Ancash, and Lima, and kept in captivity at the serpentarium of the Instituto Nacional de Salud, Lima,

Perú. Venom from nine adult *M. t. tschudii* specimens was collected and pooled during the first three months of captivity. Venom was lyophilized and stored at -20°C until used.

4.2. Isolation and Characterization of Venom Proteins

First 0.5 milligrams of crude, lyophilized venom were dissolved in 200 μL of 5% acetonitrile in water containing 0.1% trifluoroacetic acid (TFA), centrifuged to remove debris, and separated by reverse-phase HPLC using a Teknokroma Europa Protein 300 C18 (0.4 cm \times 25 cm, 5 μm particle size, 300 \AA pore size) column and an LC 1100 High Pressure Gradient System (Agilent Technologies, Santa Clara, CA, USA) equipped with DAD detector and micro-Auto-sampler [73]. The flow rate was set to 1 mL/min and the column was developed with a linear gradient of 0.1% TFA in water (solution A) and acetonitrile (solution B) using the following column elution conditions: isocratically (5% B) for 5 min, followed by 5%–25% B for 10 min, 25%–45% B for 60 min, and 45%–70% for 10 min. Protein detection was carried out at 215 nm with a reference wavelength of 400 nm. Fractions were collected manually, dried in a vacuum centrifuge (Savant), and redissolved in water, and submitted to molecular mass determination using a SYNAPT[®] G2 High Definition Mass Spectrometry System (Waters Corp., Milford, MA, USA), and SDS-PAGE analysis in 15% polyacrylamide gels, under reducing and non-reducing conditions. Gels were stained with Coomassie Brilliant Blue R-250 (Sigma-Aldrich, St. Louis, MO, USA).

4.3. Characterization of the Venom Peptidome and Proteome

Electrophoretic protein bands were excised from a Coomassie Brilliant Blue-stained SDS-PAGE gel and subjected to in-gel reduction (10 mM dithiothreitol) and alkylation (50 mM iodoacetamide), followed by overnight sequencing-grade trypsin digestion (66 ng/ μL in 25 mM ammonium bicarbonate, 10% acetonitrile; 0.25 μg /sample) in an automated processor (ProGest Protein Digestion Workstation, Genomic Solution Ltd., Cambridgeshire, UK) following the manufacturer's instructions. Tryptic digests were dried in a SpeedVac (Savant[™], Thermo Scientific Inc., West Palm Beach, FL, USA), redissolved in 15 μL of 0.1% formic acid in water, and submitted to LC-MS/MS. To this end, tryptic peptides were separated by nano-Acquity UltraPerformance LC[®] (UPLC[®], Waters Corporation, Milford, MA, USA) using BEH130 C18 (100 μm \times 100 mm, 1.7 μm particle size) column in-line with a SYNAPT[®] G2 High Definition Mass Spectrometry System (Waters Corp., Milford, MA, USA). The flow rate was set to 0.6 μL /min and the column was developed with a linear gradient of 0.1% formic acid in water (solution A) and 0.1% formic acid in acetonitrile (solution B), isocratically 1% B for 1 min, followed by 1%–12% B for 1 min, 12%–40% B for 15 min, 40%–85% B for 2 min. Doubly and triply charged ions were selected for collision-induced dissociation (CID) MS/MS. Fragmentation spectra were interpreted (a) manually (*de novo* sequencing); (b) using the on-line form of the MASCOT program at <http://www.matrixscience.com> against NCBIInr database, a comprehensive, non-identical protein database compiled from GenBank CDS translations, PIR, SwissProt, PRF, and PDB; and (c) processed in Waters Corporation's (Milford, MA, USA) ProteinLynx Global SERVER 2013 version 2.5.2. (with Expression version 2.0) and the generated .pkl peak list files were exported to MASCOT for protein identification against the NCBIInr database. MS/MS mass tolerance was set to ± 0.6 Da. Carbamidomethyl cysteine and oxidation of methionine were selected as fixed and variable modifications, respectively. Cut-off for MASCOT reporting was set to top 10 hits. All MASCOT identifications were manually verified. Amino acid sequence similarity searches were performed against the NCBIInr and UniProtKB databases using the BLASTP program implemented in the WU-BLAST2 search engine at <http://www.bork.embl-heidelberg.de>.

The relative abundances (expressed as percentage of the total venom proteins) of the different protein families were calculated as the ratio of the sum of the areas of the reverse-phase chromatographic peaks containing proteins from the same family to the total area of venom protein peaks in the reverse-phase chromatogram [14,74]. When more than one protein band was present in a reverse-phase fraction, their proportions were estimated by densitometry of Coomassie-stained

SDS-polyacrylamide gels using ImageJ version 1.47 (Free Software Foundation, Boston, MA, USA) (<http://rsbweb.nih.gov/ij>). Conversely, the relative abundances of different proteins contained in the same SDS-PAGE band were estimated based on the relative ion intensities of the three more abundant peptide ions associated with each protein by MS/MS analysis. Finally, protein family abundances were estimated as the percentages of the total venom proteome.

4.4. Determination of LD₅₀ for Mice

Animal experiments were performed in accordance with protocols approved by the Institutional Committee for the Care and Use of Laboratory Animals of the University of Costa Rica (CICUA 041-15, 21 October 2015), using CD-1 mice of either sex, provided by Instituto Clodomiro Picado. To evaluate the lethal activity, various doses of *M. t. tschudii* venom, dissolved in 200 µL of PBS, were injected into groups of five CD-1 mice (16–18 g body weight), by the intraperitoneal (i.p.) route. Deaths were scored over a 48 h period and the median lethal dose (LD₅₀) was calculated by probits [75].

Supplementary Materials: The following are available online at www.mdpi.com/2072-6651/8/6/178/s1. Table S1: Summary of the MS/MS identification of the SDS-PAGE protein bands of *M. t. tschudii* venom.

Acknowledgments: This study was conducted as part of an agreement signed between the Instituto Nacional de Salud (Perú) and the Universidad Peruana Cayetano Heredia. Venomic analyses were supported by Grant BFU2013-42833-P from the Ministerio de Economía y Competitividad, Madrid (Spain). Authors gratefully acknowledge Dr. Med. Vet. Gualberto Marcas Cáceres, Centro Nacional de Productos Biológicos, Instituto Nacional de Salud, Ministerio de Salud, Perú, for kindly providing pictures of *M. t. tschudii*.

Author Contributions: L.S., D.P., B.L., and J.J.C. conceived and designed the experiments; L.S., D.P., and B.L. performed the experiments; A.Z., B.L., and J.J.C. contributed reagents/materials/analysis tools; L.S., D.P., B.L. and J.J.C. analyzed the data. J.J.C. wrote the paper. All authors revised the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Wagler, J. Serpentina Brasiliensium species novae, ou histoire naturelle des espèces nouvelles de serpens. In *Animalia Nova Sive Species Novae*; de Spix, J., Ed.; Franc. Seraph. Hübschmanni: Munich, Germany, 1824.
2. Roze, J.A. *Coral Snakes of the Americas. Biology, Identification and Venoms*; Krieger Publishing Company: Malabar, FL, USA, 1996; p. 328.
3. Campbell, J.A.; Lamar, W.W. *The Venomous Reptiles of the Western Hemisphere*; Cornell University Press: Ithaca, NY, USA, 2004; p. 476.
4. The Reptile Database. Available online: <http://www.reptile-database.org> (accessed on 12 February 2016).
5. Pires, M.G.; da Silva, N.J., Jr.; Feitosa, D.T.; Prudente, A.L.; Filho, G.A.; Zaher, H. A new species of triadal coral snake of the genus *Micrurus* Wagler, 1824 (Serpentes: Elapidae) from northeastern Brazil. *Zootaxa* **2014**, *3811*, 569–584. [[CrossRef](#)] [[PubMed](#)]
6. Feitosa, D.T.; da Silva, N.J.; Pires, M.G.; Zaher, H.; Prudente, A.L. A new species of monadal coral snake of the genus *Micrurus* (Serpentes, Elapidae) from western Amazon. *Zootaxa* **2015**, *3974*, 538–554. [[CrossRef](#)] [[PubMed](#)]
7. Slowinski, J.B.; Keogh, J.S. Phylogenetic relationship of coral snakes based on cytochrome b mtDNA sequences. *Mol. Phylogenet. Evol.* **2005**, *15*, 157–164. [[CrossRef](#)] [[PubMed](#)]
8. Harvey, M.B.; Gutberlet, R.L. The evolution of New World venomous snakes. In *The Venomous Reptiles of the Western Hemisphere*; Campbell, J.A., Lamar, W.W., Eds.; Cornell University Press: Ithaca, NY, USA, 2004; pp. 634–682.
9. Campbell, J.A.; Lamar, W.W. *The Venomous Reptiles of Latin America*; Cornell University Press: Ithaca, NY, USA; London, UK, 1989; p. 425.
10. Savage, J.M.; Slowinski, J.B. The coloration of the venomous coral snakes (family Elapidae) and their mimics (families Aniliidae and Colubridae). *Biol. J. Linn. Soc.* **1992**, *45*, 235–254. [[CrossRef](#)]
11. Slowinski, J.B. A phylogenetic analysis of the New World coral snakes (Elapidae: *Leptomicrurus*, *Micruroides* and *Micrurus*) based on allozymic and morphological characters. *J. Herpetol.* **1995**, *29*, 325–338. [[CrossRef](#)]

12. Silva, N.J., Jr.; Sites, J.W., Jr. Phylogeny of South American triad coral snakes (Elapidae: *Micrurus*) based on molecular characters. *Herpetologica* **2001**, *57*, 1–22.
13. Calvete, J.J. Snake venomomics: From the inventory of toxins to biology. *Toxicon* **2013**, *75*, 44–62. [[CrossRef](#)] [[PubMed](#)]
14. Calvete, J.J. Next-generation snake venomomics: Protein-locus resolution through venom proteome decomplexation. *Expert Rev. Proteom.* **2014**, *11*, 315–329. [[CrossRef](#)] [[PubMed](#)]
15. Brahma, R.K.; McCleary, R.J.; Kini, R.M.; Doley, R. Venom gland transcriptomics for identifying, cataloging, and characterizing venom proteins in snakes. *Toxicon* **2015**, *93*, 1–10. [[CrossRef](#)] [[PubMed](#)]
16. Calvete, J.J.; Lomonte, B. A bright future for integrative venomomics. *Toxicon* **2015**, *107*, 159–162. [[CrossRef](#)] [[PubMed](#)]
17. Sunagar, K.; Morgenstern, D.; Reitzel, A.M.; Moran, Y. Ecological venomomics: How genomics, transcriptomics and proteomics can shed new light on the ecology and evolution of venom. *J. Proteom.* **2016**, *135*, 62–72. [[CrossRef](#)] [[PubMed](#)]
18. Fernández, J.; Vargas-Vargas, N.; Pla, D.; Sasa, M.; Rey-Suárez, P.; Sanz, L.; Gutiérrez, J.M.; Calvete, J.J.; Lomonte, B. Snake venomomics of *Micrurus alleni* and *Micrurus mosquitensis* from the Caribbean region of Costa Rica reveals two divergent compositional patterns in New World elapids. *Toxicon* **2015**, *107*, 217–233. [[CrossRef](#)] [[PubMed](#)]
19. Warrell, D.A. Snakebites in central and South America: Epidemiology, clinical features, clinical management. In *The Venomous Reptiles of the Western Hemisphere*; Campbell, J.A., Lamar, W.W., Eds.; Cornell University Press: Ithaca, NY, USA, 2004; pp. 709–761.
20. Bucarechi, F.; de Capitani, E.M.; Vieira, R.J.; Rodrigues, C.K.; Zannin, M.; da Silva, N.J., Jr.; Casais-E-Silva, L.L.; Hyslop, S. Coral snake bites (*Micrurus* spp.) in Brazil: A review of literature reports. *Clin. Toxicol.* **2016**, *25*, 1–13.
21. Norris, R.L.; Pfalzgraf, R.R.; Laing, G. Death following coral snake bite in the United States—First documented case (with ELISA confirmation of envenomation) in over 40 years. *Toxicon* **2009**, *53*, 693–697. [[CrossRef](#)] [[PubMed](#)]
22. Jan, G. Plan d’une iconographie descriptive des ophidiens et description sommaire de nouvelles espèces deserpents. *Rev. Mag. Zool. Pure Appliquée (Paris)* **1858**, *2*, 438–439 and 514–527.
23. Silva, N.J., Jr.; Sites, J.W., Jr. Revision of the *Micrurus frontalis* complex (Serpentes: Elapidae). *Herpetol. Monogr.* **1999**, *13*, 142–194. [[CrossRef](#)]
24. Peters, J.A.; Orejas-Miranda, B. Catalogue of the neotropical Squamata: Part I. Snakes. *US Natl. Mus. Bull.* **1970**, *297*, 1–347. [[CrossRef](#)]
25. Mackessy, S.P.; Tu, A.T. Biology of the sea snakes and biochemistry of their venoms. In *Toxin-Related Diseases: Poisons Originating from Plants, Animals and Spoilage*; Tu, A.T., Ed.; Oxford & IBH Publishing Co.: New Delhi, India, 1993; pp. 305–351.
26. Jackson, T.N.; Sunagar, K.; Undheim, E.A.; Koludarov, I.; Chan, A.H.; Sanders, K.; Ali, S.A.; Hendrikx, I.; Dunstan, N.; Fry, B.G. Venom down under: Dynamic evolution of Australian elapid snake toxins. *Toxins* **2013**, *5*, 2621–2655. [[CrossRef](#)] [[PubMed](#)]
27. Kini, R.M.; Doley, R. Structure, function and evolution of three-finger toxins: Mini proteins with multiple targets. *Toxicon* **2010**, *56*, 855–867. [[CrossRef](#)] [[PubMed](#)]
28. Utkin, Y.N. Three-finger toxins, a deadly weapon of elapid venom-milestones of discovery. *Toxicon* **2013**, *62*, 50–55. [[CrossRef](#)] [[PubMed](#)]
29. Du, X.-Y.; Clemetson, K.J. Snake venom L-amino acid oxidases. *Toxicon* **2002**, *40*, 659–665. [[CrossRef](#)]
30. Filippovich, I.; Sorokina, N.; Masci, P.P.; de Jersey, J.; Whitaker, A.N.; Winzor, D.J.; Gaffney, P.J.; Lavin, M.F. A family of textilinin genes, two of which encode proteins with antihemorrhagic properties. *Br. J. Haematol.* **2002**, *119*, 376–384. [[CrossRef](#)] [[PubMed](#)]
31. Bohlen, C.J.; Chesler, A.T.; Sharif-Naeini, R.; Medzihradzsky, K.F.; Zhou, S.; King, D.; Sánchez, E.E.; Burlingame, A.L.; Basbaum, A.I.; Julius, D. A heteromeric Texas coral snake toxin targets acid-sensing ion channels to produce pain. *Nature* **2011**, *479*, 410–414. [[CrossRef](#)] [[PubMed](#)]
32. Bacongus, I.; Bohlen, C.J.; Goehring, A.; Julius, D.; Gouaux, E. X-ray structure of acid-sensing ion channel 1-snake toxin complex reveals open state of a Na(+)-selective channel. *Cell* **2014**, *156*, 717–729. [[CrossRef](#)] [[PubMed](#)]

33. Bohlen, C.J.; Julius, D. Receptor-targeting mechanisms of pain-causing toxins: How ow? *Toxicon* **2012**, *60*, 254–264. [[CrossRef](#)] [[PubMed](#)]
34. Aird, S.D. Ophidian envenomation strategies and the role of purines. *Toxicon* **2002**, *40*, 335–393. [[CrossRef](#)]
35. Guo, C.; Liu, S.; Yao, Y.; Zhang, Q.; Sun, M.-Z. Past decade study of snake venom L-amino acid oxidase. *Toxicon* **2012**, *60*, 302–311. [[CrossRef](#)] [[PubMed](#)]
36. Lee, M.S.Y.; Sanders, K.L.; King, B.; Palci, A. Diversification rates and phenotypic evolution in venomous snakes (Elapidae). *R. Soc. Open Sci.* **2016**, *3*, 150277. [[CrossRef](#)] [[PubMed](#)]
37. Greene, H.W. Dietary correlates of the origin and radiation of snakes. *Am. Zool.* **1983**, *23*, 431–441. [[CrossRef](#)]
38. Daltry, J.C.; Wüster, W.; Thorpe, R.S. Diet and snake venom evolution. *Nature* **1996**, *379*, 537–540. [[CrossRef](#)] [[PubMed](#)]
39. Barlow, A.; Pook, C.E.; Harrison, R.A.; Wüster, W. Coevolution of diet and prey-specific venom activity supports the role of selection in snake venom evolution. *Proc. Biol. Sci.* **2009**, *276*, 2443–2449. [[CrossRef](#)] [[PubMed](#)]
40. Rey-Suárez, P.; Núñez, V.; Fernández, J.; Lomonte, B. Integrative characterization of the venom of the coral snake *Micrurus dumerilii* (Elapidae) from Colombia: Proteome, toxicity, and cross-neutralization by antivenom. *J. Proteom.* **2016**, *136*, 262–273. [[CrossRef](#)] [[PubMed](#)]
41. Corrêa-Netto, C.; Junqueira-de-Azevedo, I.L.; Silva, D.A.; Ho, P.L.; Leitão-de-Araújo, M.; Alves, M.L.; Sanz, L.; Foguel, D.; Zingali, R.B.; Calvete, J.J. Snake venomomics and venom gland transcriptomic analysis of Brazilian coral snakes, *Micrurus altirostris* and *M. corallinus*. *J. Proteom.* **2011**, *74*, 1795–1809. [[CrossRef](#)] [[PubMed](#)]
42. Fernández, J.; Alape-Girón, A.; Angulo, Y.; Sanz, L.; Gutiérrez, J.M.; Calvete, J.J.; Lomonte, B. Venomic and antivenomic analyses of the Central American coral snake, *Micrurus nigrocinctus* (Elapidae). *J. Proteome Res.* **2011**, *10*, 1816–1827. [[CrossRef](#)] [[PubMed](#)]
43. Vergara, I.; Pedraza-Escalona, M.; Paniagua, D.; Restano-Cassulini, R.; Zamudio, F.; Batista, C.V.; Possani, L.D.; Alagón, A. Eastern coral snake *Micrurus fulvius* venom toxicity in mice is mainly determined by neurotoxic phospholipases A₂. *J. Proteom.* **2014**, *105*, 295–306. [[CrossRef](#)] [[PubMed](#)]
44. Margres, M.J.; Aronow, K.; Loyacano, J.; Rokyta, D.R. The venom-gland transcriptome of the eastern coral snake (*Micrurus fulvius*) reveals high venom complexity in the intragenomic evolution of venoms. *BMC Genom.* **2013**, *14*, 531. [[CrossRef](#)] [[PubMed](#)]
45. Bénard-Valle, M.; Carbajal-Saucedo, A.; de Roodt, A.; López-Vera, E.; Alagón, A. Biochemical characterization of the venom of the coral snake *Micrurus tener* and comparative biological activities in the mouse and a reptile model. *Toxicon* **2014**, *77*, 6–15. [[CrossRef](#)] [[PubMed](#)]
46. Lomonte, B.; Sasa, M.; Rey-Suárez, P.; Bryan, W.; Gutiérrez, J.M. Venom of the coral snake *Micrurus clarki*: Proteomic profile, toxicity, immunological cross-neutralization, and characterization of a three-finger toxin. *Toxins* **2016**, *8*, 138. [[CrossRef](#)] [[PubMed](#)]
47. Castoe, T.A.; Smith, E.N.; Brown, R.M.; Parkinson, K.L. Higher-level phylogeny of Asian and American coralsnakes, their placement within the Elapidae (Squamata), and the systematic affinities at the enigmatic Asian coralsnake *Hemibungarus calligaster* (Weigmann, 1834). *Zool. J. Linn. Soc.* **2007**, *151*, 809–831. [[CrossRef](#)]
48. Renjifo, C.; Smith, E.N.; Hodgson, W.C.; Renjifo, J.M.; Sanchez, A.; Acosta, R.; Maldonado, J.H.; Rivero, A. Neuromuscular activity of the venoms of the Colombian coral snakes *Micrurus dissoleucus* and *Micrurus mipartitus*: An evolutionary perspective. *Toxicon* **2012**, *59*, 132–142. [[CrossRef](#)] [[PubMed](#)]
49. Olamendi-Portugal, T.; Batista, C.; Restano-Cassulini, R.; Pando, V.; Villa-Hernandez, O.; Zavaleta-Martínez-Vargas, A.; Salas-Arruz, M.C.; Rodríguez de la Vega, R.C.; Becerril, B.; Possani, L. Proteomic analysis of the venom from the fish eating coral snake *Micrurus surinamensis*: Novel toxins, their function and phylogeny. *Proteomics* **2008**, *8*, 1919–1932. [[CrossRef](#)] [[PubMed](#)]
50. Ciscotto, P.H.; Rates, B.; Silva, D.A.; Richardson, M.; Silva, L.P.; Andrade, H.; Donato, M.F.; Cotta, G.A.; Maria, W.S.; Rodrigues, R.J.; *et al.* Venomic analysis and evaluation of antivenom cross-reactivity of South American *Micrurus* species. *J. Proteom.* **2011**, *74*, 1810–1825. [[CrossRef](#)] [[PubMed](#)]
51. Dokmetjian, J.C.; Del Canto, S.; Vinzón, S.; de Jiménez Bonino, M.B. Biochemical characterization of the *Micrurus pyrrhocryptus* venom. *Toxicon* **2009**, *53*, 375–382. [[CrossRef](#)] [[PubMed](#)]
52. Rey-Suárez, P.; Núñez, V.; Gutiérrez, J.M.; Lomonte, B. Proteomic and biological characterization of the venom of the redtail coral snake, *Micrurus mipartitus* (Elapidae), from Colombia and Costa Rica. *J. Proteom.* **2011**, *75*, 655–667. [[CrossRef](#)] [[PubMed](#)]

53. Tanaka, G.D.; Furtado, M.F.; Portaro, F.C.; Sant'Anna, O.A.; Tambourgi, D.V. Diversity of *Micrurus* snake species related to their venom toxic effects and the prospective of antivenom neutralization. *PLoS Negl. Trop. Dis.* **2010**, *4*, e622. [[CrossRef](#)] [[PubMed](#)]
54. Salazar, A.M.; Vivas, J.; Sánchez, E.E.; Rodríguez-Acosta, A.; Ibarra, C.; Gil, A.; Carvajal, Z.; Girón, M.E.; Estrella, A.; Navarrete, L.F.; *et al.* Hemostatic and toxinological diversities in venom of *Micrurus tener tener*, *Micrurus fulvius fulvius* and *Micrurus isozonus* coral snakes. *Toxicon* **2011**, *58*, 35–45. [[CrossRef](#)] [[PubMed](#)]
55. Aguilar, I.; Sánchez, E.E.; Girón, M.E.; Estrella, A.; Guerrero, B.; Rodríguez-Acosta, F.A. Coral snake antivenom produced in chickens (*Gallus domesticus*). *Rev. Inst. Med. Trop. São Paulo* **2014**, *56*, 61–66. [[CrossRef](#)] [[PubMed](#)]
56. Gloss, A.D.; Whiteman, N.K. Balancing Selection: Walking a Tightrope. *Curr. Biol.* **2016**, *26*, R73–R76. [[CrossRef](#)] [[PubMed](#)]
57. Sanders, K.L.; Lee, M.S.Y. Molecular evidence for a rapid late-Miocene radiation of Australasian venomous snakes. *Mol. Phylogenet. Evol.* **2008**, *46*, 1165–1173. [[CrossRef](#)] [[PubMed](#)]
58. Pyron, R.A.; Burbrink, F.T.; Wiens, J.J. A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC Evol. Biol.* **2013**, *13*, 93. [[CrossRef](#)] [[PubMed](#)]
59. Sanders, K.L.; Lee, M.S.Y.; Mumpuni, B.T.; Rasmussen, A.R. Multilocus phylogeny and recent rapid radiation of the viviparous sea snakes (Elapidae: Hydrophiinae). *Mol. Phylogenet. Evol.* **2013**, *66*, 575–591. [[CrossRef](#)] [[PubMed](#)]
60. Laustsen, A.H.; Gutiérrez, J.M.; Rasmussen, A.R.; Engmark, M.; Gravlund, P.; Sanders, K.L.; Lohse, B.; Lomonte, B. Danger in the reef: Proteome, toxicity, and neutralization of the venom of the olive sea snake, *Aipysurus laevis*. *Toxicon* **2015**, *107*, 187–196. [[CrossRef](#)] [[PubMed](#)]
61. Tan, C.H.; Tan, K.Y.; Lim, S.E.; Tan, N.H. Venomics of the beaked sea snake, *Hydrophis schistosus*: A minimalist toxin arsenal and its cross-neutralization by heterologous antivenoms. *J. Proteom.* **2015**, *126*, 121–130. [[CrossRef](#)] [[PubMed](#)]
62. Calvete, J.J.; Ghezellou, P.; Paiva, O.; Matainaho, T.; Ghassempour, A.; Goudarzi, H.; Kraus, F.; Sanz, L.; Williams, D.J. Snake venomics of two poorly known Hydrophiinae: Comparative proteomics of the venoms of terrestrial *Toxicocalamus longissimus* and marine *Hydrophis cyanocinctus*. *J. Proteom.* **2012**, *75*, 4091–4101. [[CrossRef](#)] [[PubMed](#)]
63. Lomonte, B.; Pla, D.; Sasa, M.; Tsai, W.C.; Solórzano, A.; Ureña-Díaz, J.M.; Fernández-Montes, M.L.; Mora-Obando, D.; Sanz, L.; Gutiérrez, J.M.; *et al.* Two color morphs of the pelagic yellow-bellied sea snake, *Pelamis platura*, from different locations of Costa Rica: Snake venomics, toxicity, and neutralization by antivenom. *J. Proteom.* **2014**, *103*, 137–152. [[CrossRef](#)] [[PubMed](#)]
64. Pahari, S.; Bickford, D.; Fry, B.G.; Kini, R.M. Expression pattern of three-finger toxin and phospholipase A₂ genes in the venom glands of two sea snakes, *Lapemis curtus* and *Acalyptophis peronii*: Comparison of evolution of these toxins in land snakes, sea kraits and sea snakes. *BMC Evol. Biol.* **2007**, *7*, 175. [[CrossRef](#)] [[PubMed](#)]
65. Tan, C.H.; Fung, S.Y.; Yap, M.K.; Leong, P.K.; Liew, J.L.; Tan, N.H. Unveiling the elusive and exotic: Venomics of the Malayan blue coral snake (*Calliophis bivirgata flaviceps*). *J. Proteom.* **2016**, *132*, 1–12. [[CrossRef](#)] [[PubMed](#)]
66. Mackessy, S.P. Venom composition in rattlesnakes: trends and biological significance. In *The Biology of Rattlesnakes*; Hayes, W.K., Beaman, K.R., Cardwell, M.D., Bush, S.P., Eds.; Loma Linda University Press: Loma Linda, CA, USA, 2008; pp. 495–510.
67. Calvete, J.J.; Sanz, L.; Cid, P.; de la Torre, P.; Flores-Díaz, M.; Dos Santos, M.C.; Borges, A.; Bremo, A.; Angulo, Y.; Lomonte, B.; *et al.* Snake venomics of the Central American rattlesnake *Crotalus simus* and the South American *Crotalus durissus* complex points to neurotoxicity as an adaptive paedomorphic trend along *Crotalus* dispersal in South America. *J. Proteome Res.* **2010**, *9*, 528–544. [[CrossRef](#)] [[PubMed](#)]
68. Durban, J.; Pérez, A.; Sanz, L.; Gómez, A.; Bonilla, F.; Rodríguez, S.; Chacón, D.; Sasa, M.; Angulo, Y.; Gutiérrez, J.M.; *et al.* Integrated “omics” profiling indicates that miRNAs are modulators of the ontogenetic venom composition shift in the Central American rattlesnake, *Crotalus simus simus*. *BMC Genom.* **2013**, *14*, 234. [[CrossRef](#)] [[PubMed](#)]
69. Massey, D.J.; Calvete, J.J.; Sánchez, E.E.; Sanz, L.; Richards, K.; Curtis, R.; Boesen, K. Venom variability and envenoming severity outcomes of the *Crotalus scutulatus scutulatus* (Mojave rattlesnake) from Southern Arizona. *J. Proteom.* **2012**, *75*, 2576–2587. [[CrossRef](#)] [[PubMed](#)]

70. Diz, A.P.; Martínez-Fernández, M.; Rolán-Alvarez, E. Proteomics in evolutionary ecology: Linking the genotype with the phenotype. *Mol. Ecol.* **2012**, *21*, 1060–1080. [[CrossRef](#)] [[PubMed](#)]
71. Baer, B.; Millar, A.H. Proteomics in evolutionary ecology. *J. Proteom.* **2016**, *135*, 4–11. [[CrossRef](#)] [[PubMed](#)]
72. Diz, A.P.; Calvete, J.J. Ecological proteomics: Is the field ripe for integrating proteomics into evolutionary ecology research? *J. Proteom.* **2016**, *135*, 1–3. [[CrossRef](#)] [[PubMed](#)]
73. Eichberg, S.; Sanz, L.; Calvete, J.J.; Pla, D. Constructing comprehensive venom proteome reference maps for integrative venomics. *Expert Rev. Proteom.* **2015**, *12*, 557–573. [[CrossRef](#)] [[PubMed](#)]
74. Calvete, J.J.; Juárez, P.; Sanz, L. Snake venomics, strategy and applications. *J. Mass Spectrom.* **2007**, *42*, 1405–1414. [[CrossRef](#)] [[PubMed](#)]
75. Finney, D.J. *Statistical Methods in Biological Assay*; Charles Griffin and Company Ltd.: London, UK, 1971.



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).