





The Effects of Quinone Imine, a New Potent Nitrification Inhibitor, Dicyandiamide, and Nitrapyrin on Target and Off-Target Soil Microbiota

 Evangelia S. Papadopoulou,^{a,b} Eleftheria Bachtsevani,^a Christina V. Papazlatani,^a Constantina Rousidou,^a Antonios Brouziotis,^a Eleni Lampronikou,^a Myrto Tsiknia,^c Sotirios Vasileiadis,^a Ioannis Ipsilantis,^d Urania Menkissoglu-Spiroudi,^e Constantinou Ehaliotis,^c  Laurent Philippot,^f Graeme W. Nicol,^g Dimitrios G. Karpouzas^a

^aLaboratory of Plant and Environmental Biotechnology, Department of Biochemistry and Biotechnology, University of Thessaly, Larissa, Greece

^bLaboratory of Environmental Microbiology, Department of Environmental Sciences, University of Thessaly, Larissa, Greece

^cLaboratory of Soils and Agricultural Chemistry, Agricultural University of Athens, Athens, Greece

^dLaboratory of Soil Sciences, School of Agriculture, Forestry and Environment, Faculty of Agriculture, Aristotle University of Thessaloniki, Thessaloniki, Greece

^ePesticide Science Laboratory, School of Agriculture, Forestry and Environment, Faculty of Agriculture, Aristotle University of Thessaloniki, Thessaloniki, Greece

^fUniversité Bourgogne Franche-Comté, INRAE, AgroSup Dijon, Agroécologie, Dijon, France

^gEnvironmental Microbial Genomics Group, Laboratoire Ampère, École Centrale de Lyon, CNRS UMR 5005, Université de Lyon, Lyon, France

ABSTRACT Dicyandiamide (DCD) and nitrapyrin (NP) are nitrification inhibitors (NIs) used in agriculture for over 40 years. Recently, ethoxyquin (EQ) was proposed as a novel potential NI, acting through its derivative quinone imine (QI). Still, the specific activity of these NIs on the different groups of ammonia-oxidizing microorganisms (AOM), and mostly their effects on other soil microbiota remain unknown. We determined the impact of QI, and comparatively of DCD and NP, applied at two doses (regular versus high), on the function, diversity, and dynamics of target (AOM), functionally associated (nitrite-oxidizing bacteria-NOB), and off-target prokaryotic and fungal communities in two soils mainly differing in pH (5.4 versus 7.9). QI was equally effective to DCD but more effective than NP in inhibiting nitrification in the acidic soil, while in the alkaline soil QI was less efficient than DCD and NP. This was attributed to the higher activity of QI toward AOA prevailing in the acidic soil. All NIs induced significant effects on the composition of the AOB community in both soils, unlike AOA, which were less responsive. Beyond on-target effects, we noted an inhibitory effect of all NIs on the abundance of NOB in the alkaline soil, with *Nitrobacter* being more sensitive than *Nitrospira*. QI, unlike the other NIs, induced significant changes in the composition of the bacterial and fungal communities in both soils. Our findings have serious implications for the efficiency and future use of NIs on agriculture and provide unprecedented evidence for the potential off-target effects of NIs on soil microbiota.

IMPORTANCE NIs could improve N use efficiency and decelerate N cycling. Still, we know little about their activity on the distinct AOM groups and about their effects on off-target soil microorganisms. Here, we studied the behavior of a new potent NI, QI, compared to established NIs. We show that (i) the variable efficacy of NIs across soils with different pH reflects differences in the inherent specific activity of the NIs to AOA and AOB; (ii) beyond AOM, NIs exhibit negative effects on other nitrifiers, like NOB; (iii) QI was the sole NI that significantly affected prokaryotic and fungal diversity. Our findings (i) highlight the need for novel NI strategies that consider the variable sensitivity of AOM groups to the different NIs (ii) identify QI as a potent AOA inhibitor, and (iii) stress the need for monitoring NIs' impact on off-target soil microorganisms to ensure sustainable N fertilizers use and soil ecosystem functioning.

KEYWORDS quinone imine, DCD, nitrapyrin, ammonia-oxidizing microorganisms, nitrite-oxidizing bacteria, soil microbial toxicity

Editor Konstantinos Aristomenis Kormas, University of Thessaly

Copyright © 2022 Papadopoulou et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Evangelia S. Papadopoulou, evapapadopoulou@uth.gr.

Received 14 February 2022

Accepted 24 June 2022

Published 20 July 2022

Nitrification, the microbial conversion of ammonia (NH_3) to nitrate (NO_3^-), is an essential process of the global nitrogen (N) cycle with paramount importance for plant nutrition and productivity (1). In agricultural soil ecosystems, where N is typically added through the application of inorganic fertilizers, nitrification can result in a significant loss of N, through NO_3^- leaching and N oxides (N_xO) or elemental N_2 emissions, raising agricultural production costs and contributing to environmental pollution and climate change (2). It is therefore essential to regulate nitrification in agricultural soils, particularly the first and often rate-limiting step of ammonia oxidation, to prolong the presence of ammonium (NH_4^+) in the root zone and facilitate uptake of N by plants (3). In this first step of nitrification, ammonia is oxidized to hydroxylamine by ammonia monooxygenase and further to nitrite (NO_2^-) via nitric oxide (4), by canonical ammonia-oxidizing bacteria (AOB), ammonia-oxidizing archaea (AOA), and the recently discovered complete ammonia-oxidizing (comammox) bacteria (5).

The direct inhibition of soil ammonia-oxidizing microorganisms (AOM) using commercial nitrification inhibitors (NIs) is an established strategy for improving N use efficiency in intensive farming systems (6). Hundreds of compounds have been identified as potential NIs (7); however, only three have been widely used in agricultural practice: dicyandiamide (DCD), nitrapyrin (NP; 2-chloro-6-[trichloromethyl] pyridine), and 3,4-dimethylpyrazole phosphate (DMPP) (8). While the underlying inhibition mechanisms of these NIs have been associated with their presumed capacity to interfere with the activation of ammonia monooxygenase, an enzyme carried by all chemolithoautotrophic AOM (9), their efficacy in regulating soil N transformations is highly variable across soils (10, 11). Soil properties such as pH, clay content, and organic matter content (12, 13) and environmental variables such as temperature (14) and moisture (15) are known to affect the performance of NIs in soil. For example, the inhibitory efficiency of DCD and NP in soils with variable physicochemical properties varied from 5.5 to 83.8% for the former (13) and from -39.8 to 135.3% for the latter (16).

To date, the impact of NIs on the indigenous soil microbial communities has been typically considered for target AOM, while little is known about their off-target effects on other functionally associated microbial groups, such as nitrite-oxidizing bacteria (NOB), fueled by substrates produced during ammonia oxidation. Soil studies have shown a trend of selective inhibition of commercial NIs toward AOB, which is further modulated by the application rates and the composition of the metabolically active AOM in soil (17–19). The variable sensitivity of soil AOM to different NIs may arise from fundamental differences in their biochemistry and physiology (1) and their well-documented niche specialization (20) and implies a potential suboptimal efficiency of the NIs currently used in agriculture. This stresses the need for the development of novel nitrification inhibition strategies that could rely on new broad-range NIs, or perhaps on mixtures of NIs with complementary activity against the different AOM groups.

Besides the effects of NIs on target microorganisms, studies on the effects of NIs on nontarget microorganisms are limited. The studies available have explored the effects of DCD- (or DMPP-) fortified fertilizers on the abundance of bacteria or fungi or on microbial community structure determined by low-resolution methods (e.g., PLFAs and DGGE), with contradictory results (21, 22). On the other hand, the potential effects of NP on nontarget soil microorganisms are less studied, with recent reports suggesting possible effects on the overall soil microbial community within and beyond N cycling (23, 24). Careful evaluation of the off-target effects of NIs on the soil microbiota is essential to define potential risks associated with their use in agricultural settings that might compromise soil ecosystem diversity and functioning, beyond nitrification and N cycling.

Ethoxyquin (EQ; 1,2-dihydro-6-ethoxy-2,2,4-trimethylquinoline) is an antioxidant used as a preservative in fruit-packaging plants (25). Previous soil studies have shown that it is rapidly transformed to 2,6-dihydro-2,2,4-trimethyl-6-quinone imine (QI) and suggested a strong inhibitory effect on both AOA and AOB, without, however, affecting soil microbial activity endpoints and PLFA composition (26). A follow-up analysis of the spectrum of the inhibitory activity of EQ and its derivatives, compared to DCD and NP, against a range of soil AOB and AOA strains revealed a selective inhibitory activity

TABLE 1 Properties of the soils used in this study

Property	Acidic soil	Alkaline soil
Site name	Livadi, Ellasona	Rhodia, Ampelonas
Latitude	40°08'29.7N	39°46'57.6N
Longitude	22°10'13.5'E	22°19'36.6E
Soil order	Cambisol	Cambisol
Soil texture	loam	loam
Sand/silt/clay (%)	40.8/47.8/11.4	45.8/36.0/18.2
Land use	Arable	Arable
Crops/Plants	Lettuce-potato-fallow rotation (One harvest per yr)	Peach (5 sequential growing seasons)
Soil pH (H ₂ O)	5.40	7.90
Water Holding Capacity (WHC) (%)	50.32	60.20
Cation-exchange capacity (cmol kg ⁻¹)	9.3	8.5
Organic C content (%)	2.1	1.8
Organic matter content (%)	4.3	3.6
Total N content (%)	0.13	0.08
Exch. NH ₄ ⁺ -N (mg kg ⁻¹)	54.01	72.27
NO ₃ ⁻ -N (mg kg ⁻¹)	33.40	32.61
Ca (mg kg ⁻¹)	1182	1164
K (mg kg ⁻¹)	90	89
Mg (mg kg ⁻¹)	271	224
Na (mg kg ⁻¹)	21	18
Olsen-P (mg kg ⁻¹)	39.1	43.5

of DCD on AOB, whereas NP showed equivalent inhibitory activity against AOB and AOA. EQ showed inhibitory activity against AOA and AOB, comparable to that of NP and DCD, respectively, with its major transformation product QI being the active NI (27). Considering the low cost of EQ (25), and its ability to be transformed in soil to a highly potent NI like QI, there is a major potential for its use in agricultural settings.

The main objective of this study was to determine the efficacy and off-target effects of QI, the active NI component of EQ, in comparison with two established NIs, such as DCD and NP, on the soil microbiota, and to define how these effects are modulated by soil pH and NI application rate. To minimize the involvement of other confounding soil factors on NIs performance, we selectively determined NI effects in two soils that largely differed in their pH (acidic versus alkaline) but had otherwise similar main physicochemical properties (Table 1). We hypothesized that (i) the efficacy of the tested NIs toward nitrification would depend on the functionally dominant AOM groups/phylogenotypes in each soil, in accord with the pH-driven niche specialization of AOM (20), and the different sensitivity of the distinct AOM groups to the tested NIs (27); (ii) the inhibitory effects of NIs go beyond ammonia oxidation and extend to other soil microbiota components within and beyond N cycling; (iii) high NI dose rates would minimize differences in NIs efficacy and maximize potential off-target effects on soil microbiota. To test these hypotheses, we compared the impact of QI, DCD, and NP, applied at two dose rates, a low one equivalent to the agriculturally recommended application (low dose, *LD*) and a high dose aiming at universal AOM inhibition (high dose, *HD*), on (i) the activity of AOM by monitoring N-pools, potential nitrification rates, and changes in *amoA* gene and transcripts abundance; (ii) the diversity and dynamics of AOM communities and functionally linked NOB, respectively; and (iii) the diversity and dynamics of the total prokaryotic and fungal soil communities. To relate our results to the level and duration of the microbial exposure to these compounds, the dissipation of the NIs in soil was also determined.

RESULTS

Dissipation of NIs. The dissipation patterns of DCD, NP, and QI in the two soils, as described by the single first order (SFO) kinetic model (28), are shown in Fig. S1. DCD showed both a dose-dependent and soil-specific dissipation pattern with dissipation

TABLE 2 The kinetic parameters describing the dissipation of dicyandiamide (DCD), nitrapyrin (NP), and quinone imine (QI) applied at low (LD) and high (HD) dose rates in the two studied soils (acidic and alkaline)^a

Treatment	Soil	K (d ⁻¹)	DT ₅₀ (d)	DT ₉₀ (d)	χ ^{2α} (%)
DCD LD	Acidic	0.030	23.06	76.61	5.88
	Alkaline	0.012	57.20	190.0	3.74
DCD HD	Acidic	0.024	28.93	96.09	4.05
	Alkaline	0.006	110.20	366.2	3.64
NP LD	Acidic	0.246	2.82	9.38	22.37
	Alkaline	0.133	5.21	17.31	17.13
NP HD	Acidic	0.147	4.73	15.71	7.08
	Alkaline	0.178	3.90	12.96	8.81
QI LD	Acidic	0.282	2.46	8.17	3.14
	Alkaline	0.210	3.30	10.96	2.68
QI HD	Acidic	0.385	1.80	5.98	4.13
	Alkaline	0.102	6.83	22.68	7.95

^aThe error χ² provides an indication of how accurately the kinetic model describes the dissipation pattern. Error values of 15% suggest acceptable fit to the observed data. Dissipation kinetic parameters were calculated with the single first order (SFO) kinetic model which provided adequate fit to the dissipation data in all cases.

time 50% (DT₅₀) values of 23.1 days (acidic soil) and 57.2 days (alkaline soil) in the samples treated with the low dose, and 28.9 days (acidic soil) and 110.2 (alkaline soil) days in the samples treated with the high dose (Table 2). The dissipation of NP and QI did not show a clear dose-dependent pattern. For example, the DT₅₀ values of NP in samples treated with the low and the high dose rates were 2.8 and 5.2 days in the acidic and 4.7 and 3.9 days in the alkaline soil, respectively. Similarly, QI DT₅₀ values in the acidic and alkaline soil were 2.5 and 3.3 days, respectively, at the low dose rate and 1.8 and 6.8 days at the high dose rate.

The impact of NIs on the dynamics of soil inorganic nitrogen. We first determined the effects of NIs on inorganic nitrogen pools and potential nitrification as measures of potential effects on nitrification activity. In both soils, the low dose rate of the NIs did not induce a clear temporal pattern in NH₄⁺-N levels, unlike their high dose rate, which resulted in a significant ($P < 0.001$) and persistent increase of NH₄⁺-N levels only in the acidic soil (Fig. 1a).

NO₃⁻-N, DCD, and QI low dose rates induced a significant reduction at days 14 and 35 in the acidic soil, whereas at high dose rates all NIs induced a significant ($P < 0.001$) and persistent decrease, with NP showing the lowest and least persistent effect (Fig. 1b). In the alkaline soil, all tested NIs at both dose rates significantly ($P < 0.05$) decreased NO₃⁻-N formation from day 7 onward (Fig. 1b).

DCD and QI exerted a significant and persistent inhibitory effect ($P < 0.001$) on potential nitrification in the acidic soil, whereas the effect of NP was weaker and less persistent, especially at the low dose rate (Fig. 1c). In the alkaline soil, both NI doses induced a significant decrease ($P < 0.001$) in potential nitrification, with recovery observed only at the low dose rates at 35 days (Fig. 1c). QI at the high dose rate showed a weaker inhibitory effect ($P < 0.05$) compared to DCD and NP, still being significantly lower than the control (Fig. 1c).

The impact of NIs on the abundance and transcriptional activity of AOs. In parallel we monitored, via q-PCR and RT-q-PCR, the effects of NIs on the abundance and transcriptional activity of AOB, AOA, and comammox bacteria. In the acidic soil, regardless of the treatments, the abundance of the AOA *amoA* genes (2.2×10^6 to 3.0×10^7 g⁻¹ d.w. soil) was significantly higher than that of AOB (9.2×10^4 to 8.8×10^5 g⁻¹ d.w. soil). Conversely, in the alkaline soil the AOB *amoA* genes (3.5×10^7 to 2.4×10^8 g⁻¹ d.w. soil) were more abundant ($P < 0.05$) than those of AOA (7.3×10^5 to 2.1×10^7 g⁻¹ d.w. soil). Comammox clade A and B *amoA* genes were not detected in either soil.

In the acidic soil, QI induced a significant reduction ($P < 0.05$) in AOB abundance compared to the control on days 35 (low dose) and 14 (high dose). DCD and NP induced a significant ($P < 0.05$) and persistent reduction in AOB abundance only at the




FIG 1 The effect of dicyandiamide (DCD), nitrapyrin (NP), and quinone imine (QI), applied at two dose rates (low and high), on ammonium concentration (a), nitrate concentration (b), and potential nitrification (PN) (c) in an acidic and alkaline soil. Each value is the mean of three replicates \pm standard error. At each time point, groups designated by the same letter are not significantly different at the selected P -value levels.

high dose rate (Fig. 2a). Regarding AOA, both doses of NIs significantly decreased ($P < 0.05$) their abundance at day 35, with the high NP dose showing significantly lower inhibition ($P < 0.05$) compared to DCD and QI (Fig. 2b). Bacterial *amoA* gene transcripts were below the limit of detection (<48 gene transcripts g^{-1} d.w. soil). Regarding AOA, both doses of DCD and QI induced a significant and persistent reduction ($P < 0.05$) of *amoA* gene transcripts, while NP showed a weaker and temporary inhibition of *amoA* transcription ($P < 0.05$) (Fig. 2d).

In the alkaline soil, all NIs at both dose rates significantly and persistently reduced ($P < 0.05$) the abundance of AOB (Fig. 2a). Regarding AOA, we observed a reduction in their abundance ($P < 0.05$) in the samples treated with the high NI dose rates only at 35 days (Fig. 2b). At the transcription level, NP and QI reduced ($P < 0.05$) AOB *amoA* gene transcript abundance at both dose rates, with the impact of the QI low dose only being significant ($P < 0.05$) on day 35. DCD induced a significant reduction ($P < 0.05$) of AOB *amoA* gene transcripts only at the high dose rate (Fig. 2c). Regarding AOA, the application of NIs at both dose rates induced a significant and persistent reduction ($P < 0.05$) of *amoA* gene transcripts ($P < 0.001$) (Fig. 2d).

The impact of NIs on the dynamics of NOB. We expanded our tests to NOB, functionally associated with AOM. In the acidic soil, all tested NIs induced a significant reduction on the abundance of the *nxrB* gene of *Nitrobacter* and *Nitrospira* at day 35 ($P < 0.05$) at the low dose rate. In the alkaline soil, all NIs significantly ($P < 0.05$) decreased the abundance of the *nxrB* gene of *Nitrobacter* at both dose rates, compared to *Nitrospira*, whose abundance was negatively affected only at the high dose rates of DCD and QI (Fig. S2).

The impact of NIs on the abundance of bacteria, fungi, and Thaumarchaeota. Beyond N cycling, we investigated, via q-PCR, the impact of NIs on the abundance of total bacteria, fungi, and Thaumarchaeota. DCD, NP, and QI, at both rates, significantly and persistently decreased ($P < 0.05$) the abundance of Thaumarchaeota in the acidic soil but not of bacteria and fungi. In the alkaline soil, NIs did not show a clear temporal dose-dependent effect on the abundance of any of the studied microbial groups (Fig. S3). The ratio of AOA *amoA* to thaumarchaeotal 16S rRNA gene abundance ranged from 0.5 to 192.6 in the acidic soil (median 16.8), and from 0.02 to 113.9 in the alkaline soil




FIG 2 The effect of dicyandiamide (DCD), nitrapyrin (NP), and quinone imine (QI), applied at two dose rates (low and high), on the abundance of the *amoA* genes and transcripts of AOB (a, c) and AOA (b, d), in an acidic and alkaline soil. Each value is mean of three replicates \pm standard error. All values are expressed in log scale. At each time point, groups designated by the same letter are not significantly different at the selected *P*-value levels. In the acidic soil bacterial *amoA* gene transcripts were below the limit of detection.

(median 5.7). The resulting ratios of *amoA*: thaumarcheotal 16S rRNA did not vary significantly between NI treatments ($P > 0.05$) and increased only in the acidic soil at day 14 upon application of NP (low dose rate) and QI (high dose rate) (Table S1).

The impact of NIs on the diversity of the different soil microbial groups. Eventually we determined the impact of NIs on the diversity of (i) target microorganisms (AOM) via amplicon sequencing of the *amoA* gene of AOA and AOB and (ii) the total prokaryotic and fungal community via amplicon sequencing of the 16S rRNA and ITS, respectively. The per-sample average sequence numbers obtained and passing quality control screening are presented in Table S2. Our sequencing effort provided adequate coverage of the microbial diversity as evidenced by rarefaction curves that reached a plateau in all samples and for all microbial domains (Fig. S4).

The two tested soils supported similar bacterial and fungal communities at the phylum level. The soil bacterial communities were composed of Proteobacteria (39.5 and 44.6%, in the acidic and alkaline soil, respectively), Actinobacteria (16.6 and 15.0%), Gemmatimonadota (9.7 and 7.5%), Acidobacteriota (10.7 and 9.0%), and Chloroflexi (7.4 and 5.3%) (Fig. S5). NI-driven changes in the relative abundance of the bacterial phyla were evident with QI (both dose rates) significantly increasing the relative abundance of Proteobacteria ($P < 0.001$) compared to the control, while other phyla like Gemmatimonadota were significantly decreased upon application of QI ($P < 0.001$). The fungal community was dominated by Ascomycetes (84.7 and 85.6%) (orders Sordariomycetes (38.7 and 56.7%), Dothideomycetes (9.5 and 17.3%), Eurotiomycetes (15.0 and 4.5%), Leotiomycetes (15.2 and 0.4%), and Basidiomycetes (11.3 and 9.6%) (Fig. S6). In the acidic and alkaline soil AOB *amoA* amplicon sequencing analysis

showed a dominance of *Nitrosospira* (89.2 and 43.4%, respectively) and unclassified *amoA* sequences (10.8 and 56.6%, respectively), with *N. briensis* (11.3 and 34.3%, respectively) identified as the only characterized *Nitrosospira* cluster, according to Abell et al. (29) (Fig. S7, Table S3). Regarding AOA, the two soils supported distinct communities, with the acidic soil dominated by *Nitrososphaerales* γ clade (γ -2.2. [69.1%] and γ -2. [26.1%]), and the alkaline soil dominated by *Nitrososphaerales* α , γ , and ε clades (γ -2.1. [34.3%], ε -2.2. [22.8%], α -3.1. [10.2%], γ -1.1. [8.8%]) (Fig. S8, Table S4).

(i) Effects on α -diversity. The diversity of the bacterial community was only affected by the high dose rate of QI in the alkaline soil (Inverse Simpson index and Pielou's evenness, $P < 0.05$), whereas QI had a significant effect at both doses on the diversity of the fungal community in the acidic soil only. DCD applied at the high dose rate also significantly decreased ($P < 0.05$) the Shannon index for fungi in the acidic soil. The observed richness of AOB was only affected by the low dose rate of QI ($P < 0.05$) in the acidic soil, whereas QI at both dose rates and NP at the high dose rate significantly reduced the observed richness of AOB ($P < 0.05$) in the alkaline soil. NP and QI, at both dose rates, reduced the α -diversity of AOB in the acidic (Inverse Simpson, $P < 0.05$) and the alkaline (Shannon, Inverse Simpson and Pielou's evenness, $P < 0.05$) soil. In contrast, DCD significantly reduced the α -diversity of AOB only at the high dose rate in the acidic (Inverse Simpson, $P < 0.05$) and the alkaline (Inverse Simpson and Pielou's evenness, $P < 0.05$) soil. For AOA, we noted a significant reduction at Pielou's evenness values in the alkaline soil by the high dose rate of all NIs (Table S5).

(ii) Effects on β -diversity. We employed multivariate analysis to determine the effects of NIs on the β -diversity of the different microbial groups studied. Canonical correspondence analysis (CCA) or redundancy analysis (RDA) (not considering the time factor) coupled with pairwise permutational multivariate analysis of variance (PERMANOVA) showed that, for the NIs tested, only QI induced a significant ($P \leq 0.006$) and universal (dose-wise) effect on the structure of the bacterial and fungal communities in both soils (Fig. 3 and 4). Regarding AOB, all NIs significantly affected their community structure ($P < 0.05$), except for the low dose rate of DCD and the high dose rate of QI in the acidic soil (Fig. 3 and 4). Conversely, the AOA community was less responsive with significant changes only induced by the high dose rates of NP in both soils and QI only in the alkaline soil (Fig. 3 and 4).

We further looked for amplicon sequencing variants (ASVs) (30) that were associated with certain NI treatments. We first looked for universal responses to NIs across soils. *Sphingomonas* ASVs were significantly reduced in the QI-treated samples from both soils (Fig. 5, S9 to S12). We also identified taxa exhibiting soil-specific responses to QI like *Rhodanobacter* (Fig. 5, S9, S10) and *Lysobacter* (Fig. 5, S11, S12) that significantly increased in the QI-treated samples of the acidic and alkaline soil, respectively. Regarding fungi, in both soils ASVs belonging to *Fusarium*, *Trichoderma*, and *Solicoccozyma* showed a significant increase in their relative abundance upon exposure to QI; conversely ASVs belonging to *Coniochaeta* were significantly reduced in soils treated with QI (Fig. 6, S13 to S16).

For AOM, QI significantly affected the relative abundance of AOB belonging to *Nitrosospira* sp. Nsp5 group in the acidic soil (Fig. 7a, S17), whereas DCD and NP induced a significant reduction on ASVs belonging to *Nitrosospira* sp. Nsp65 and *Nitrosospira briensis* groups at high dose rates (Fig. 7b, S18). In the alkaline soil, the majority of ASVs significantly affected by NIs belonged to the *N. briensis* group (Fig. 7d and e, S19, S20). Regarding AOA, NP applied in the alkaline soil suppressed ASVs of the *Nitrososphaerales* ε -2.2 clade but favored ASVs belonging to clade γ . In the same soil, ASVs belonging to *Nitrososphaerales* α -3 and ε -2.2. clades decreased with the high dose rate of QI (Fig. 7, S22-23).

DISCUSSION

NIs soil dissipation. QI and NP did not persist in both soils, in line with previous studies (26, 31), unlike DCD, which was the most persistent NI, showing a dose-




FIG 3 Canonical correspondence analysis (CCA) or redundancy analysis (RDA) of bacterial (a, e), fungal (b, f), AOB (c, g), and AOA (d, h) community structures in samples of the acidic soil treated with the low (a, b, c, d) or the high (e, f, g, h) dose rates of the three NIs (DCD, QI, NP) and in untreated samples (CTL). The percentages of the plotted canonical variance and *P*-values of the multilevel pairwise PERMANOVA are provided next to the axes and at the top-right tables, respectively. Ellipses encompass all samples of each treatment.

dependent increase in its persistence (32, 33). The DT_{50} values of DCD in the acidic soil was within the range reported in studies for acidic soils (6.2 to 39 days, at 20–25°C) (14, 33–35), whereas there is a lack of studies on its persistence in alkaline soils, where higher DT_{50} values were recorded in our study. The different persistence of DCD in the two soils might be attributed to their different pH, since the differences in other properties of the two soils known to affect the environmental fate of organic compounds (e.g., organic matter content and soil texture) were small to induce such a distinct effect on the persistence of DCD. pH is a major factor governing the degradation activity of soil microbes (36) but also the polarity of chemicals and hence their sorption and bioavailability (37).

On-target effects of NIs on the function and diversity of AOM. An accumulation of NH_4^+ was observed only at the high dose rates of the tested NIs and only in the acidic soil. Previous studies looking at the effect of DCD and NP on NH_4^+ in various soils have reported contrasting results with accumulation (17, 38, 39) or no increase in NH_4^+ retention (17, 40). This is perhaps not surprising considering that, in addition to nitrification, the NH_4^+ soil equilibrium is affected by numerous other soil-specific processes, including sorption and NH_3 volatilization, both of which can be greater in alkaline soils (41). A clear inhibitory effect by all NIs was evident when potential nitrification and NO_3^- levels were considered, and it was more consistent in the alkaline soil. Comprehensive mechanistic studies with stable isotope labeling are required to elucidate the effects of NIs on gross N transformation rates and to provide best insights into N processes other than nitrification (42).

QI reduced the abundance and the transcriptional activity of AOA and AOB in both soils, although effects on the transcriptional activity of AOB were weaker. Earlier studies by Papadopoulou et al. (26) showed a clear and persistent inhibition of the transcription of *amoA* genes of both AOA and AOB by QI applied at rate of 16.1 mg kg⁻¹ d.w. soil




FIG 4 Canonical correspondence analysis (CCA) or redundancy analysis (RDA) of bacterial (a, e), fungal (b, f), AOB (c, g), and AOA (d, h) community structures, in samples of the alkaline soil treated with the low (a, b, c, d) or the high (e, f, g, h) dose rates of the three NIs (DCD, QI, NP) and in untreated samples (CTL). The percentages of the plotted canonical variance and the *P*-values of the multilevel pairwise PERMANOVA are provided next to the axes and at the top-right tables, respectively. Ellipses encompass all samples of each treatment.

(i.e., approximately half of the high QI dose in the current study) in an alkaline soil (pH 7.9). The weaker effect of QI on AOB in the current study is probably a function of the lower dose rate of QI used in this study ($10 \text{ mg kg}^{-1} \text{ d.w. soil}$) compared to our earlier study and reflects the lower intrinsic sensitivity of AOB to QI compared to AOA (27).

The impact of DCD on the abundance and activity of the different AOM groups varied in the two studied soils. In the acidic soil DCD showed a more consistent inhibitory effect on AOA, compared to the alkaline soil where DCD was more inhibitory on AOB. At transcriptional level DCD showed again a soil-dependent effect on AOA and AOB. In the acidic soil, it reduced the transcriptional activity of AOA at both dose rates, compared to the alkaline soil, where DCD had a more consistent effect on AOA than on AOB transcripts. Most studies to date have investigated the effects of DCD on the abundance of *amoA* genes but not on the *amoA* transcripts, an indicator of AOM ammonia oxidation activity. These studies have shown that DCD significantly decreased the abundance of AOB, while AOA showed no or little response (19, 43). The few studies that have reported a significant inhibition of AOA by DCD (44, 45) were all conducted in acidic soils (pH 4.2 to 5.4), often dominated by AOA. Robinson et al. (12) investigated the effect of DCD on AOB and AOA abundance in a field site where soil pH was maintained at native (pH 6.0) or adjusted to alkaline or acidic levels. They found that DCD inhibited AOB at alkaline and native pH and AOA in acidic pH and concluded that DCD could effectively inhibit AOB or AOA under soil conditions favoring their growth. Our data concur with these findings: in the acidic soil, where AOA prevail functionally and numerically, DCD inhibits AOA more effectively, whereas in the alkaline soil, where AOB are numerically dominant, DCD reduces AOB abundance more effectively than AOA.

NP showed a similar inhibitory pattern with DCD in the two soils. It reduced more effectively the abundance of AOA and AOB in the acidic and alkaline soil, respectively. When looking for effects at *amoA* transcript levels, NP strongly inhibited the expression




FIG 5 Heatmaps of differentially abundant bacterial ASVs (Kruskal-Wallis and Wilcoxon rank sum analysis, test significance is indicated in each ASV annotation * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$), scaled from 0 to 1 (blue to red) in the acidic (a, b) and alkaline (c, d) soil untreated (CTL) or treated with a low (a, c) or high (b, d) dose rate of NIs (DCD, QI, NP). The mean relative abundance of each ASV is indicated by the white-to green column and key. Hierarchical clustering was performed for depicting treatment and ASV groupings. Per ASV, barplots with the corresponding P -values/ASV taxonomies/treatment relative abundances are provided in Fig. S9 to S12 in the supplemental material.

of AOB and AOA *amoA* gene in the alkaline soil, as opposed to the acidic soil, where the transcriptional activity of the functionally dominant AOA was only temporarily affected by NP. Only a few studies have examined the effect of NP on AOA and AOB dynamics, with contrasting results. In studies performed in acidic soils where AOA functionally and numerically prevailed, NP applied at the recommended dose rates reduced the abundance of AOA but had no effect on AOB (18, 46), while others reported no effect of NP on AOB in soils numerically and functionally dominated by AOB (38). Recently, Hayden et al. (24) reported an increase in AOA abundance in response to NP application in a range of soils, attributed to reduced competition by AOB, whose population was reduced by NP.

Besides their impact on the functional activity of AOM, all NIs induced significant effects in the composition of AOB, unlike AOA, whose community was marginally affected only by the high dose rates of NP and QI. Previous studies (43, 47) suggested a selective effect of DCD on the diversity of AOB, while AOA were not affected, in line with our findings. Recently Schmidt et al. (23) showed that NP, when applied at rates similar to our high dose rate, induced a significant shift on the AOA soil community, in accord with our results. Our study provides first evidence for the effects of NP and QI on the diversity of AOB and AOA, indicating a greater impact of both compounds on AOB compared to AOA, even in soils numerically and functionally dominated by AOA. When looking for specific AOB and AOA that were responsive to the tested NIs, a clear phylogenetic-based inhibition trend was not evident. However, we should note that the sets of primers used in our study may not cover the full range of the AOA harboring *amoA* gene (48, 49).




FIG 6 Heatmaps of differentially abundant fungal ASVs (Kruskal-Wallis and Wilcoxon rank sum analysis, test significance is indicated in each ASV annotation * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$), scaled from 0 to 1 (blue to red)) in the acidic (a, b) and alkaline (c, d) soil for samples treated with the low (a, c) or the high (b, d) dose rate of NIs (DCD, QI, NP) or untreated (CTL). The mean relative abundance of each ASV is indicated by the white-to green column and key. Hierarchical clustering was performed for depicting treatment and ASV groupings. Per ASV barplots with the corresponding P -values/ASV taxonomies/treatment relative abundances are provided in Fig. S13 to S16 in the supplemental material.

Off-target effects of NIs on the abundance and diversity of microbial groups beyond AOM. Consistent effects on NOB, were observed only in the alkaline soil, where all NIs suppressed *Nitrobacter* at both dose rates, unlike *Nitrospira*, which were inhibited only by the high dose rates of DCD and QI. Previous studies looking at the effect of DCD on NOB in various soils (pH 5.2 to 7.2) reported contrasting results leading to either a decrease (21) or no effect on the abundance of *Nitrobacter* and *Nitrospira* (19, 50). We suggest that the higher inhibitory effect of NIs on *Nitrobacter* over *Nitrospira* NOB most probably reflects the stronger inhibitory activity of the NIs on ammonia oxidation in the alkaline versus the acidic soil. This has resulted in lower substrate (NO_2^- -N) availability in the former, a condition expected to favor *Nitrospira*, characterized as k-strategists thriving in soils with nitrite limitations, over *Nitrobacter* which are superior competitors at high resources availability (51). Our hypothesis fits well with the theory that nonrandom associations exist between different AOM and NOB lineages that, along with edaphic factors, shape field-scale spatial patterns of nitrifying communities (52). Still a higher intrinsic tolerance of *Nitrospira* over *Nitrobacter* to NIs cannot be excluded, although data from relevant *in vitro* assays are only available for *Nitrobacter* (27).

The NIs tested did not induce significant alterations in the composition of nontarget prokaryotes and fungal communities, with QI being the sole exception. Little is known




FIG 7 Heatmaps of differentially abundant ASVs of AOB (a, b, d, e) and AOA (c, f) (Kruskal-Wallis and Wilcoxon rank sum analysis, test significance is indicated in each ASV annotation * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$) in the acidic (a, b, c) and the alkaline (d, e, f) soil for samples treated with the low (a, d) or the high (b, c, e, f) dose rate of NIs (DCD, QI, NP) or untreated (CTL). The mean relative abundance of each ASV is indicated by the white-to green column and key. Hierarchical clustering was performed for depicting treatment and ASV groupings. Per ASV barplots with the corresponding P -values/ASV taxonomies/treatment relative abundances are provided in Fig. S17 to S23 in the supplemental material.

regarding the effects of NIs on the overall microbial diversity in soil, with previous studies typically using lower resolution methods that are not directly comparable to our findings (21, 26). Recent studies, using amplicon sequencing, showed that DCD, applied at a dose similar to the low dose rate used in our study, did not alter the composition of the soil bacterial community (50, 53), in line with our findings. We further examined which bacteria and fungi were benefited or suppressed by QI. We noted a soil independent negative effect of QI on *Sphingomonas*. Bacteria of this genus possess multifaceted functions ranging from the biodegradation of environmental contaminants to the production of beneficial phytohormones (54). In contrast, QI stimulated *Lysobacter*, which are considered to be facultative predators (55) or biocontrol microbes, producing secondary metabolites involved in the suppression of plant pathogens (56), pointing to potential impacts of QI on microbial predator-prey interactions in the soil food web. Beyond bacteria, QI favored fungi belonging to *Fusarium*, *Trichoderma*, and *Solicoccozyma* encompassing plant pathogens (e.g., *Fusarium*) (57), plant growth promoting agents (e.g., *Trichoderma*, *Fusarium*) (58, 59), and soil lignocellulose decomposers (e.g., *Solicoccozyma*) (60). These effects could be attributed either (i) to the direct toxicity of QI on target and off-target soil microorganisms that have reciprocal and cascading effects on the proliferation of other soil microorganisms (release from competition, cellular content release as nutrient sources for copiotrophs) or (ii) to the use of QI as a growth substrate favoring the proliferation of the fraction of the soil microbiota being capable of catabolizing and using QI as an energy source. Further studies using shotgun metagenomic and metatranscriptomic approaches are needed to provide insights into the causality of the effects of QI on nontarget soil

bacteria and fungi and identify how these diversity changes translate into effects on agricultural soil ecosystem functioning.

Conclusions. Using a combination of molecular, biochemical, and analytical tools we provide a comprehensive assessment of the on-target activity (AOM) and off-target effects (NOB, bacteria, fungi) of QI, a new potent NI compound, and two currently used in agriculture NIs, DCD, and NP, on the soil microbiota of two soils mostly varying in pH known to affect microbial community structure, AOM niche differentiation, and chemicals persistence. Our study (i) highlights QI as an alternative, highly efficient NI against AOA, showing, however, significant effects on off-target soil microbiota, within and beyond N cycling, that might reciprocate broader effects on soil ecosystem functioning; (ii) verifies the different sensitivity of the distinct AOM groups and phylotypes to the different NIs that should be considered along with soil microbial dynamics and community composition data for the design of novel, more efficient NI strategies; and (iii) provides first insights into the potential impact of NIs on the wider soil microbial community, beyond the target microbial groups.

MATERIALS AND METHODS

Soils and chemicals. Two arable loamy soils, one acidic vegetable (pH 5.4) and one alkaline (pH 7.9) orchard soil, characterized as Cambisols, were collected in 2019 from two field sites in Thessaly, Greece (61). The two soils were selected to confer very similar physicochemical properties and texture but largely differed in pH (Table 1). Topsoil samples were obtained from 5 selected points (0 to 10 cm depth), according to ISO (62, 63). For each soil, individual samples were mixed thoroughly to provide a single bulk sample (2 kg). After collection, the soils were partially air dried and sieved to pass through a 2 mm mesh sieve. For the preparation of NIs, working solutions for soil treatment and analytical purposes, analytical standards of DCD (99% purity), and NP ($\geq 98\%$), purchased from Sigma-Aldrich (Germany), were used. The oxidation derivative of ethoxyquin QI was synthesized as described by Thorisson et al. (64).

Experimental setup. Each bulk soil was treated with 80 mL of 0.42 M urea (corresponding to 0.47 g N kg⁻¹ d.w. soil) to stimulate nitrification and left to equilibrate for 5 days at room temperature, to ensure the ammonification of urea residues in both soils. The final concentration of urea was expected to result in NH₄⁺-N levels, which have been demonstrated to support the growth of both AOA and AOB (65). Each bulk soil sample was then divided into two sets of four subsamples (0.5 kg each). The first set of subsamples was treated with aqueous suspensions (DCD) or acetonitrile solutions (QI and NP) of NIs, resulting in nominal concentrations in soil of 15.4, 10, and 0.86 mg kg⁻¹ d.w. soil of DCD, QI, and NP, respectively. These dose rates were recommended for agricultural use for DCD and NP, assuming that (i) NIs are distributed through the top 5 cm of the soil profile of a field site of 0.1 ha and (ii) soil bulk density was equal to 1.3 g cm⁻³. As QI has no established recommended dose for use as an NI, the nominal concentration of 10 mg kg⁻¹ d.w. soil was used, based on previous observations from soil microcosms and liquid batch culture assays (26, 27). A subsample from the first set of each soil was treated with an equivalent amount of acetonitrile and water to serve as an untreated control. DCD-amended samples were also treated with 5 mL acetonitrile to ensure that all treatments received the same volume of water and acetonitrile. Control samples used in the present study were the same as those used in a previous study of our group (61) that was running in parallel, following a similar experimental setup. The second set of subsamples was treated in the same way with the three NIs resulting in concentrations of 200, 30, and 5 mg kg⁻¹ d.w. soil for DCD, QI, and NP, respectively. These high dose rates were selected based on the concentrations previously found to suppress the activity of both AOB and AOA strains in liquid culture assays (27). Soil samples were left to equilibrate for 1 h, and sterile water was added to adjust moisture to 40% of the water holding capacity. Soil samples were then split into ~30-g portions and placed in aerated plastic bags, which were incubated in the dark at 20°C for 35 days. Triplicate samples of each treatment were removed from the incubator at 3, 7, 14, 21, and 35 days and stored at -20°C for NI dissipation measurements or at -80°C for molecular analyses. Fresh soil samples were used for NH₄⁺-N, NO₃⁻-N, and potential nitrification measurements. The moisture content of all soil samples was kept constant with regular addition of deionized water.

Analysis of NI residues in soil samples. DCD residues were extracted from soil according to Kim et al. (66) with slight modifications. Briefly, 5 g of soil was mixed with 10 mL dH₂O in 50 mL tubes and shaken on a horizontal shaker for 1 h, at 250 rpm, at room temperature. The tubes were centrifuged (7232 × g, 5 min), and 2 mL of the soil extract were acidified with 0.2 mL of 0.66 M H₂SO₄ and were left to settle for 30 min before centrifugation (16200 × g, 10 min). For the extraction of NP, 5 g of soil was mixed with 10 mL acetonitrile. Samples were shaken for 90 min, at 250 rpm, sonicated for 30 min, and centrifuged at 7232 × g, for 5 min. Residues of QI were extracted from soil as described by Papadopoulou et al. (26). Before analysis, extracts were passed through 0.45 μm nylon syringe filters. Analysis was performed in a Shimadzu LC-20AD HPLC-PDA system and a Shimadzu GVP-ODS (4.6 mm × 10 mm) precolumn, connected to a RP Shimadzu VP-ODS (4.6 mm × 150 mm, 5 μm) column as described by Papadopoulou et al. (27).

Analysis of inorganic nitrogen. To determine NH₄⁺-N and NO₃⁻-N levels in soil samples, soils (2 g) were extracted with 20 mL of 1 M KCl, shaken on a horizontal shaker, at 300 rpm, at room temperature

for 30 min, and filtered gravimetrically through Whatman filter paper. NH_4^+ -N and NO_3^- -N were determined in KCl extracts using a modified indophenol method based on the classical Berthelot reaction (67), and on the VCl_3 /Griess method (68), respectively. Potential nitrification was determined with the method of Kandelner (69).

Nucleic acid extraction. RNA was extracted from 2 g of soil using the RNeasy PowerSoil Total RNA isolation kit (Qiagen, Germany) with DNA co-eluted using the RNeasy PowerSoil DNA elution accessory kit (Qiagen, Germany). DNA and RNA purity was determined spectrophotometrically with integrity (shearing and degradation) checked by agarose gel electrophoresis. Nucleic acids were quantified using a Qubit fluorometer (Invitrogen).

qPCR analysis of AOB, AOA, Comammox bacteria, NOB, total bacteria, fungi, and Thaumarcheota. The effect of NIs on the abundance of these microbial groups was assessed in samples collected at 14- and 35-days post application. Primers and thermocycling conditions for each target gene are presented in Table S5. qPCR amplification efficiencies were between 82 and 110%, with R^2 values of ≥ 0.993 . Standard curves were obtained using serial dilutions of plasmid vectors containing amplicons of nearly full-length target genes.

RT-qPCR analysis of AOB and AOA *amoA* gene transcripts. The effect of NIs on the transcriptional activity of AOB and AOA was further tested via reverse transcription-qPCR (RT-qPCR) of *amoA* gene transcripts. Soil RNA (~300 ng) was treated with DNase I (1 U/ μL) (amplification grade; Invitrogen), reverse transcribed, using 200 U Superscript II (Invitrogen) and a 0.2 μM *amoA*2R and Arch-*amoA*R primers for AOB and AOA, respectively, according to the manufacturer's instructions. Amplification of cDNAs for both AOB and AOA was performed following the same procedure as for DNA.

Amplicon sequencing and bioinformatic analysis. Microbial diversity analysis for total bacteria, fungi, AOB, and AOA populations were performed on samples of all treatments collected at days 14 and 35, via multiplex amplicon sequencing. A detailed description of the multiplexing method can be found elsewhere (70). Sequencing was carried out at an Illumina HiSeq 2500 sequencer at the Brigham Young University sequencing facilities (Provo, UT, USA), generating 250 bp paired-end reads (bacteria, fungi, AOA) and an Illumina MiSeq sequencer at the Macrogen facilities (Seoul, South Korea), generating 300 bp paired-end reads (AOB). Primers and thermal cycling conditions are shown in Table S6, while primer-indexing sequences are given in Table S7. The read-pairs were assembled and demultiplexed with Flexbar, version 3.0.3 (71). Sequence quality control, generation of ASVs matrices, and sequence classification were performed with dada2 version 1.18.0 (72) using R v3.3.2, software (73) as described in (74). Bacterial 16S rRNA gene amplicon sequence annotations were obtained through assembly of paired reads and subsequently compared with the Silva database v138 (75). Fungal ITS amplicon sequences were classified using UNITE general fasta release version 8.2 (76). AOB and AOA *amoA* amplicons were compared with the databases of Abell et al. (29) and Alves et al. (77), respectively, for classifying taxonomically the generated ASVs. Unclassified AOB *amoA* sequences were subjected to blastx against the nr database (78) and those that were assigned as unclassified or untargeted products were excluded from further analysis.

Data analysis. The SFO kinetic model provided adequate fit to the data, and it was used to calculate soil dissipation kinetics of the NIs. The SFO kinetic model is based on the assumption that the change in a given NI concentration with time (dC/dt) is directly proportional to the actual concentration of the NI at this time (28). The goodness of fit was assessed using a χ^2 test (<15%, for an α of 0.05), visual inspection, and the distribution of residuals. Data from NH_4^+ -N, NO_3^- -N, potential nitrification, qPCR, and RT-qPCR measurements were subjected to two-way-ANOVA, where treatment and time constituted the two main factors. In case significant interactions between these two factors were observed, significant differences between treatments within each time point were determined via Tukey's *post hoc* test.

The ASV matrices of bacteria, fungi, AOA, and AOB were used to assess the impact of NIs on the α - and β -diversity values. α -diversity indices used included the Shannon index (79), the Inverse Simpson index (80), observed richness (S), and the Pielou's evenness index (81). Microbial diversity coverage was assessed through rarefaction curves (82). ANOVA and a Tukey's *post hoc* analysis or their nonparametric equivalents (Kruskal-Wallis test followed by the Wilcoxon rank-sum test) in case the ANOVA conditions were not met, were implemented for assessing statistically significant differences of each α -diversity index between the treatments. This was performed using the Agricolae v1.3.3 package (83) implemented in R. β -diversity analysis was performed using multivariate and canonical multivariate analysis tests. Detrended correspondence analysis (DCA) on Hellinger-transformed matrices (84) was used to assess the type of ASV responses to the environmental gradients as denoted by the first axis length (unimodal for values higher than 2.5 standard deviations or linear for lower values). Then, CCA or RDA was preferred depending on the first axis value: CCA if higher and RDA if lower than 2.5 standard deviations (85). The PERMANOVA (86), accompanying CCA or RDA, was performed using the pairwise Adonis, v0.0.1 package (87) with the default Bray-Curtis dissimilarity estimation. The nonparametric Kruskal-Wallis test was used for identifying differentially abundant ASVs among the different treatments to assess potential effect at an ASV level (88). All statistical analyses were performed with R.

Data availability. The data were submitted to the NCBI Sequence Read Archive with bioproject accession number [PRJNA733873](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA733873).

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 5.2 MB.

ACKNOWLEDGMENTS

This work is part of the project “NITRIC–Looking up for Novel nITRification Inhibitors: New Stories with Old Compounds” which has received funding from the Hellenic Foundation for Research and Innovation (HFRI) and the General Secretariat for Research and Technology (GSRT), under Grant Agreement No. 1229.

REFERENCES

- Schleper C, Nicol GW. 2010. Ammonia-oxidizing archaea-physiology, ecology, and evolution. *Adv Microb* 57:1–41. <https://doi.org/10.1016/B978-0-12-381045-8.00001-1>.
- Sutton M, Oenema O, Erisman J, Leip A, van Grinsven H, Winiwarter W. 2011. Too much of a good thing. *Nature* 472:159–161. <https://doi.org/10.1038/472159a>.
- Norton J, Ouyang Y. 2019. Controls and adaptive management of nitrification in agricultural soils. *Front Microbiol* 10:1931. <https://doi.org/10.3389/fmicb.2019.01931>.
- Caranto JD, Lancaster KM. 2017. Nitric oxide is an obligate bacterial nitrification intermediate produced by hydroxylamine oxidoreductase. *Proc Natl Acad Sci USA* 114:8217–8222. <https://doi.org/10.1073/pnas.1704504114>.
- Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M, Jehmlich N, Palatinszky M, Vierheilig J, Bulaev A, Kirkegaard RH, von Bergen M, Rattei M, Bendinger B, Nielsen PH, Wagner M. 2015. Complete nitrification by *Nitrospira* bacteria. *Nature* 528:504–509. <https://doi.org/10.1038/nature16461>.
- Abbasi MK, Adams WA. 1998. Loss of nitrogen in compacted grassland soil by simultaneous nitrification and denitrification. *Plant Soil* 200: 265–277. <https://doi.org/10.1023/A:1004398520150>.
- McCarty GW. 1999. Modes of action of nitrification inhibitors. *Biol Fertil Soils* 29:1–9. <https://doi.org/10.1007/s003740050518>.
- Zerulla W, Barth T, Dressel J, Erhardt K, von Locquenghien KH, Pasda G, Rädle M, Wissemeyer A. 2001. 3,4-Dimethylpyrazole phosphate (DMPP)—a new nitrification inhibitor for agriculture and horticulture: an introduction. *Biol Fertil Soils* 34:79–84. <https://doi.org/10.1007/s003740100380>.
- Ruser R, Schulz R. 2015. The effect of nitrification inhibitors on the nitrous oxide (N₂O) release from agricultural soils—a review. *J Plant Nutr Soil Sci* 178:171–188. <https://doi.org/10.1002/jpln.201400251>.
- Wakelin S, Williams E, O’Sullivan CA, Cameron KC, Di HJ, Cave V, O’Callaghan M. 2014. Predicting the efficacy of the nitrification inhibitor dicyandiamide in pastoral soils. *Plant Soil* 381:35–43. <https://doi.org/10.1007/s11104-014-2107-8>.
- Shi X, Hu H-W, Müller C, He J-Z, Chen D, Sute HC. 2016. Effects of the nitrification inhibitor 3,4-dimethylpyrazole phosphate on nitrification and nitrifiers in two contrasting agricultural soils. *Appl Environ Microbiol* 82: 5236–5248. <https://doi.org/10.1128/AEM.01031-16>.
- Robinson A, Di HJ, Cameron KC, Podolyan A, He J. 2014. The effect of soil pH and dicyandiamide (DCD) on N₂O emissions and ammonia oxidiser abundance in a stimulated grazed pasture soil. *J Soils Sediments* 14: 1434–1444. <https://doi.org/10.1007/s11368-014-0888-2>.
- McGeough KL, Watson CJ, Müller C, Laughlin RJ, Chadwick DR. 2016. Evidence that the efficacy of the nitrification inhibitor dicyandiamide (DCD) is affected by soil properties in UK soils. *Soil Biol Biochem* 94:222–232. <https://doi.org/10.1016/j.soilbio.2015.11.017>.
- Di HJ, Cameron KC. 2004a. Effects of temperature and application rate of a nitrification inhibitor, dicyandiamide (DCD) on nitrification rate and microbial biomass in a grazed pasture soil. *Soil Res* 42:927–932. <https://doi.org/10.1071/SR04050>.
- Di HJ, Cameron KC, Podolyan A, Robinson A. 2014. Effect of soil moisture status and a nitrification inhibitor, dicyandiamide, on ammonia oxidizer and denitrifier growth and nitrous oxide emissions in a grassland soil. *Soil Biol Biochem* 73:59–68. <https://doi.org/10.1016/j.soilbio.2014.02.011>.
- Wolt JD. 2004. A meta-evaluation of nitrapyrin agronomic and environmental effectiveness with emphasis on corn production in Midwestern USA. *Nutr Cycl Agroecosystems* 69:23–41. <https://doi.org/10.1023/B:FRES.0000025287.52565.99>.
- Cui P, Fan F, Yin C, Li Z, Song A, Wan Y, Liang Y. 2013. Urea- and nitrapyrin-affected N₂O emission is coupled mainly with ammonia oxidizing bacteria growth in microcosms of three typical Chinese arable soils. *Soil Biol Biochem* 66:214–221. <https://doi.org/10.1016/j.soilbio.2013.08.001>.
- Lehtovirta-Morley LE, Verhamme DT, Nicol GW, Prosser JI. 2013. Effect of nitrification inhibitors on the growth and activity of *Nitrosotalea devanattera* in culture and soil. *Soil Biol Biochem* 62:129–133. <https://doi.org/10.1016/j.soilbio.2013.01.020>.
- Fu Q, Clark IM, Zhu J, Hu H, Hirsch PR. 2018. The short-term effects of nitrification inhibitors on the abundance and expression of ammonia and nitrite oxidizers in a long-term field experiment comparing land management. *Biol Fertil Soils* 54:163–172. <https://doi.org/10.1007/s00374-017-1249-2>.
- Prosser JI, Hink L, Gubry-Rangin C, Nicol GW. 2020. Nitrous oxide production by ammonia oxidizers: physiological diversity, niche differentiation and potential mitigation strategies. *Glob Chang Biol* 26:103–118. <https://doi.org/10.1111/gcb.14877>.
- Carneiro J, Cardenas LM, Hatch DJ, Trindade H, Scholefield D, Clegg CD, Hobbs P. 2010. Effect of the nitrification inhibitor dicyandiamide on microbial communities and N₂O from an arable soil fertilized with ammonium sulphate. *Environ Chem Lett* 8:237–246. <https://doi.org/10.1007/s10311-009-0212-3>.
- O’Callaghan M, Gerard EM, Carter PE, Lardner R, Sarathchandra U, Burch G, Ghani A, Bell N. 2010. Effect of the nitrification inhibitor dicyandiamide (DCD) on microbial communities in a pasture soil amended with bovine urine. *Soil Biol Biochem* 42:1425–1436. <https://doi.org/10.1016/j.soilbio.2010.05.003>.
- Schmidt R, Wang X-B, Garbeva P, Yergeau É. 2020. Nitrapyrin has far reaching effects on the soil microbial community structure, composition, diversity and functions. *BioRxiv Preprint* <https://doi.org/10.1101/2020.07.21.205765>.
- Hayden HL, Phillips LA, Marshall AJ, Condon JR, Doran GS, Wells GS, Mele PM. 2021. Nitrapyrin reduced ammonia oxidation with different impacts on the abundance of bacterial and archaeal ammonia oxidisers in four agricultural soils. *Appl Soil Ecol* 157:103759. <https://doi.org/10.1016/j.apsoil.2020.103759>.
- Błaszczak AB, Augustyniak A, Skolimowski J. 2013. Ethoxyquin: an antioxidant used in animal feed. *Int J Food Sci* 2013:585931. <https://doi.org/10.1155/2013/585931>.
- Papadopoulou E, Tsachidou B, Sulowic S, Menkissoglu-Spiroudi U, Karpouzou DG. 2016. Land spreading of wastewaters from the fruit packaging industry and potential effects on soil microbes: effects of the antioxidant ethoxyquin and its metabolites on ammonia oxidizers. *Appl Environ Microbiol* 82:747–755. <https://doi.org/10.1128/AEM.03437-15>.
- Papadopoulou ES, Bachtsevani E, Lampronikou E, Adamou E, Katsaouni A, Thion C, Vasileiadis S, Menkissoglu-Spiroudi U, Nicol GW, Karpouzou DG. 2020. Comparison of novel and established nitrification inhibitors relevant to agriculture on soil ammonia- and nitrite oxidizing isolates. *Front Microbiol* 11:581283. <https://doi.org/10.3389/fmicb.2020.581283>.
- FOCUS. 2006. Guidance document on estimating persistence and degradation kinetics from environmental fate Studies on pesticides in EU registration. Report of the FOCUS Work Group on Degradation Kinetics, EC Document Reference Sanco/10058/2005 Version, 2.0, 434. https://esdac.jrc.ec.europa.eu/public_path/projects_data/focus/dk/docs/finalreportFOCDegKinetics.pdf.
- Abell GC, Robert SS, Frampton MF, Volkman JK, Rizwi F, Csontos J, Bodrossy L. 2012. High-throughput analysis of ammonia oxidiser community composition via a novel, amoA-based functional gene array. *PLoS One* 7:e51542. <https://doi.org/10.1371/journal.pone.0051542>.
- Callahan B, McMurdie P, Holmes S. 2017. Exact sequence variants should replace operational taxonomic units in marker-gene data analysis. *ISME J* 11:2639–2643. <https://doi.org/10.1038/ismej.2017.119>.
- Woodward EE, Edwards TM, Givens CE, Kolpin DW, Hladik ML. 2021. Widespread use of the nitrification inhibitor nitrapyrin: assessing benefits and costs to agriculture, ecosystems, and environmental health. *Environ Sci Technol* 55:1345–1353. <https://doi.org/10.1021/acs.est.0c05732>.
- Kelliher FM, Clough TJ, Clark H, Rys G, Sedcole JR. 2008. The temperature dependence of dicyandiamide (DCD) degradation in soils: a data synthesis. *Soil Biol Biochem* 40:1878–1882. <https://doi.org/10.1016/j.soilbio.2008.03.013>.
- Singh J, Saggari S, Giltrap DL, Bolan NS. 2008. Decomposition of dicyandiamide (DCD) in three contrasting soils and its effect on nitrous oxide

- emission, soil respiratory activity, and microbial biomass – an incubation study. *Soil Res* 46:517–525. <https://doi.org/10.1071/SR07204>.
34. Williamson JC, Menneer JC, Torrens RS. 1996. Impact of dicyandiamide on the internal nitrogen cycle of a volcanic, silt loam soil receiving effluent. *Appl Soil Ecol* 4:39–48. [https://doi.org/10.1016/0929-1393\(96\)00100-X](https://doi.org/10.1016/0929-1393(96)00100-X).
 35. Bronson KF, Touchton JT, Hauck RD. 1989. Decomposition rate of dicyandiamide and nitrification inhibition. *Commun Soil Sci Plant Anal* 20: 2067–2078. <https://doi.org/10.1080/00103628909368201>.
 36. Rousidou C, Karaiskos D, Myti D, Karanasios E, Karas PA, Tourna M, Tzortzakakis EA, Karpouzias DG. 2017. Distribution and function of carbamate hydrolase genes *cehA* and *mcd* in soils: the distinct role of soil pH. *FEMS Microbiol Ecol* 93:fiw219. <https://doi.org/10.1093/femsec/fiw219>.
 37. Franco A, Fu W, Trapp S. 2009. Influence of soil pH on the sorption of ionizable chemicals: modeling advances. *Environ Toxicol Chem* 28:458–464. <https://doi.org/10.1897/08-178.1>.
 38. Di HJ, Cameron KC. 2011. Inhibition of ammonium oxidation by a liquid formulation of 3,4-Dimethylpyrazole phosphate (DMPP) compared with a dicyandiamide (DCD) solution in six New Zealand grazed grassland soils. *J Soils Sediments* 11:1032–1039. <https://doi.org/10.1007/s11368-011-0372-1>.
 39. Fisk LM, Maccarone LD, Barton L, Murphy DV. 2015. Nitrapyrin decreased nitrification of nitrogen released from soil organic matter but not *amoA* gene abundance at high soil temperature. *Soil Biol Biochem* 88:214–223. <https://doi.org/10.1016/j.soilbio.2015.05.029>.
 40. Di HJ, Cameron KC. 2004b. Treating grazed pasture soil with a nitrification inhibitor, *eco-n*TM, to decrease nitrate leaching in a deep sandy soil under spray irrigation—a lysimeter study. *N Z J Agric Res* 47:351–361. <https://doi.org/10.1080/00288233.2004.9513604>.
 41. Kim D-G, Saggat S, Roudier P. 2012. The effect of nitrification inhibitors on soil ammonia emissions in nitrogen managed soils: a meta-analysis. *Nutr Cycl Agroecosyst* 93:51–64. <https://doi.org/10.1007/s10705-012-9498-9>.
 42. Zhu G, Ju X, Zhang J, Müller C, Rees RM, Thorman RE, Sylvester-Bradley R. 2019. Effects of the nitrification inhibitor DMPP (3,4-dimethylpyrazole phosphate) on gross N transformation rates and N₂O emissions. *Biol Fertil Soils* 55:603–615. <https://doi.org/10.1007/s00374-019-01375-6>.
 43. Di HJ, Cameron KC, Shen J-P, Winefield CS, O'Callaghan M, Bowatte S, He J-Z. 2010. Ammonia-oxidizing bacteria and archaea grow under contrasting soil nitrogen conditions. *FEMS Microbiol Ecol* 72:386–394. <https://doi.org/10.1111/j.1574-6941.2010.00861.x>.
 44. Yang W, Wang Y, Tago K, Tokuda S, Hayatsu M. 2017. Comparison of the effects of phenylhydrazine hydrochloride and dicyandiamide on ammonia-oxidizing bacteria and archaea in andosols. *Front Microbiol* 8:2226. <https://doi.org/10.3389/fmicb.2017.02226>.
 45. Zhang LM, Hu HW, Shen JP, He JZ. 2012. Ammonia-oxidizing archaea have more important role than ammonia-oxidizing bacteria in ammonia oxidation of strongly acidic soils. *ISME J* 6:1032–1045. <https://doi.org/10.1038/ismej.2011.168>.
 46. Gu Y, Mi W, Xie Y, Ma Q, Wu L, Hu Z, Dai F. 2019. Nitrapyrin affects the abundance of ammonia oxidizers rather than community structure in a yellow clay paddy soil. *J Soils Sediments* 19:872–882. <https://doi.org/10.1007/s11368-018-2075-3>.
 47. Cassman NA, Soares JR, Pijl A, Lourenço KS, van Veen JA, Cantarella H, Kuramae EE. 2019. Nitrification inhibitors effectively target N₂O-producing *Nitrosospora* spp. in tropical soil. *Environ Microbiol* 21:1241–1254. <https://doi.org/10.1111/1462-2920.14557>.
 48. Pester M, Rattei T, Flechl S, Gröngroft A, Richter A, Overmann J, Reinhold-Hurek B, Loy A, Wagner M. 2012. *amoA*-based consensus phylogeny of ammonia-oxidizing archaea and deep sequencing of *amoA* genes from soils of four different geographic regions. *Environ Microbiol* 14:525–539. <https://doi.org/10.1111/j.1462-2920.2011.02666.x>.
 49. Meinhardt KA, Bertagnolli A, Pannu MW, Strand SE, Brown SL, Stahl DA. 2015. Evaluation of revised polymerase chain reaction primers for more inclusive quantification of ammonia-oxidizing archaea and bacteria. *Environ Microbiol Rep* 7:354–363. <https://doi.org/10.1111/1758-2229.12259>.
 50. Suleiman AKA, Gonzatto R, Aita C, Lupatini M, Jacques RJS, Kuramae EE, Antoniolli ZI, Roesch LFW. 2016. Temporal variability of soil microbial communities after application of dicyandiamide-treated swine slurry and mineral fertilizers. *Soil Biol Biochem* 97:71–82. <https://doi.org/10.1016/j.soilbio.2016.03.002>.
 51. Daims H, Lückner S, Wagner M. 2016. A new perspective on microbes formerly known as nitrite-oxidizing bacteria. *Trends Microbiol* 24:699–712. <https://doi.org/10.1016/j.tim.2016.05.004>.
 52. Jones CM, Hallin S. 2019. Geospatial variation in co-occurrence networks of nitrifying microbial guilds. *Mol Ecol* 28:293–306. <https://doi.org/10.1111/mec.14893>.
 53. Morales SE, Jha N, Saggat S. 2015. Impact of urine and the application of the nitrification inhibitor DCD on microbial communities in dairy-grazed pasture soils. *Soil Biol Biochem* 88:344–353. <https://doi.org/10.1016/j.soilbio.2015.06.009>.
 54. Asaf S, Numan M, Khan AL, Al-Harrasi A. 2020. *Sphingomonas*: from diversity and genomics to functional role in environmental remediation and plant growth. *Crit Rev Biotechnol* 40:138–152. <https://doi.org/10.1080/07388551.2019.1709793>.
 55. Seccareccia I, Kost C, Nett M. 2015. Quantitative analysis of *Lysobacter* predation. *Appl Environ Microbiol* 81:7098–7105. <https://doi.org/10.1128/AEM.01781-15>.
 56. de Bruijn I, Cheng X, de Jager V, Exposito RG, Watrous J, Patel N, Postma J, Dorrestein PC, Kobayashi D, Raaijmakers JM. 2015. Comparative genomics and metabolic profiling of the genus *Lysobacter*. *BMC Genomics* 16:1–17. <https://doi.org/10.1186/s12864-015-2191-z>.
 57. Dean R, Van Kan JAL, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD, Rudd JJ, Dickman M, Kahmann R, Ellis J, Foster GD. 2012. The top 10 fungal pathogens in molecular plant pathology. *Mol Plant Pathol* 13:414–430. <https://doi.org/10.1111/j.1364-3703.2011.00783.x>.
 58. Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM, Monte E, Mukherjee PK, Zeilinger S, Grigoriev IV, Kubicek CP. 2011. Trichoderma: the genomics of opportunistic success. *Nat Rev Microbiol* 9: 749–759. <https://doi.org/10.1038/nrmicro2637>.
 59. Pappas ML, Liapoura M, Papantoniou D, Avramidou M, Kavroulakis N, Weinhold A, Broufas GD, Papadopoulou KK. 2018. The beneficial endophytic fungus *Fusarium solani* strain K alters tomato responses against spider mites to the benefit of the plant. *Front Plant Sci* 9:1603. <https://doi.org/10.3389/fpls.2018.01603>.
 60. Mašínová T, Bahnmann BD, Větrovský T, Tomšovský M, Merunková K, Baldrian P. 2017. Drivers of yeast community composition in the litter and soil of a temperate forest. *FEMS Microbiol Ecol* 93:fiw223. <https://doi.org/10.1093/femsec/fiw223>.
 61. Bachtsevani E, Papazlatani CV, Rousidou C, Lampronikou E, Menkissoglou-Spiroudi U, Nicol GW, Karpouzias DG, Papadopoulou ES. 2021. Effects of the nitrification inhibitor 3,4-dimethylpyrazole phosphate (DMPP) on the activity and diversity of the soil microbial community under contrasting soil pH. *Biol Fertil Soils* 57:1117–1135. <https://doi.org/10.1007/s00374-021-01602-z>.
 62. International Organization for Standardization. 2002. Soil quality—sampling. Part 1: guidance on the design of sampling programmes. ISO 10381–1. International Organization for Standardization, Geneva, Switzerland.
 63. International Organization for Standardization. 2002. Soil quality—sampling. Part 2: guidance on sampling techniques. ISO 10381–2. International Organization for Standardization, Geneva, Switzerland.
 64. Thorisson S, Gunstone FD, Hardy R. 1992. Some oxidation products of ethoxyquin including those found in autoxidising systems. *Chem Phys Lipids* 60:263–271. [https://doi.org/10.1016/0009-3084\(92\)90078-4](https://doi.org/10.1016/0009-3084(92)90078-4).
 65. Hink L, Gubry-Rangin C, Nicol GV, Prosser JL. 2018. The consequences of niche and physiological differentiation of archaeal and bacterial ammonia oxidisers for nitrous oxide emissions. *ISME J* 12:1084–1093. <https://doi.org/10.1038/s41396-017-0025-5>.
 66. Kim D-G, Giltrap DL, Saggat S, Hanly JA. 2014. Field studies assessing the effect of dicyandiamide (DCD) on N transformations. *N Z J Agric Res* 57: 271–293. <https://doi.org/10.1080/00288233.2013.855244>.
 67. Kandeler E, Gerber H. 1988. Short-term assay of soil urease activity using colorimetric determination of ammonium. *Biol Fertil Soils* 6:68–72. <https://doi.org/10.1007/BF00257924>.
 68. Doane TA, Horwath WR. 2003. Spectrophotometric determination of nitrate with a single reagent. *Anal Lett* 36:2713–2722. <https://doi.org/10.1081/AL-120024647>.
 69. Kandeler E. 1995. Potential nitrification, p 146–149. In Schinner F, Ohlinger R, Kandeler E, Margesin R (ed), *Methods in soil biology*. Springer, Heidelberg, Germany.
 70. Katsoula A, Vasileiadis S, Karamanolis K, Vokou D, Karpouzias DG. 2021. Factors structuring the epiphytic archaeal and fungal communities in a semi-arid Mediterranean ecosystem. *Microb Ecol* 82:638–651. <https://doi.org/10.1007/s00248-021-01712-z>.
 71. Dodt M, Roehr JT, Ahmed R, Dieterich C. 2012. FLEXBAR—Flexible barcode and adapter processing for next-generation sequencing platforms. *Biology (Basel)* 1:895–905. <https://doi.org/10.3390/biology1030895>.
 72. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016a. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods* 13:581–583. <https://doi.org/10.1038/nmeth.3869>.

73. R Core Team. 2020. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
74. Callahan B, Sankaran K, Fukuyama J, McMurdie P, Holmes S. 2016b. Bioconductor Workflow for Microbiome Data Analysis: from raw reads to community analyses [version 2; peer review: 3 approved]. *F1000Res* 5: 1492. <https://doi.org/10.12688/f1000research.8986.2>.
75. McLaren MR. 2020. Silva SSU taxonomic training data formatted for DADA2 (Silva version 138). Zenodo. <https://doi.org/10.5281/zenodo.3986799>.
76. Abarenkov K, Zirk A, Piirmann T, Pöhönen R, Ivanov F, Henrik NR, Kõljalg U. 2020. UNITE General FASTA Release for Fungi. Application/gzip. UNITE Community.
77. Alves RJE, Minh BQ, Urich T, von Haeseler A, Schleper C. 2018. Unifying the global phylogeny and environmental distribution of ammonia-oxidising archaea based on amoA genes. *Nat Commun* 9:1517. <https://doi.org/10.1038/s41467-018-03861-1>.
78. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).
79. Spellerberg I, Fedor P. 2003. A tribute to Claude Shannon (1916–2001) and a plea for more rigorous use of species richness, species diversity and the “Shannon–Wiener” index. *Glob Ecol Biogeogr* 12:177–179. <https://doi.org/10.1046/j.1466-822X.2003.00015.x>.
80. Hill MO. 1973. Diversity and evenness: a unifying notation and its consequences. *Ecology* 54:427–432. <https://doi.org/10.2307/1934352>.
81. Pielou EC. 1966. The measurement of diversity in different types of biological collections. *J Theor* 13:131–144. [https://doi.org/10.1016/0022-5193\(66\)90013-0](https://doi.org/10.1016/0022-5193(66)90013-0).
82. Sanders HL. 1968. Marine benthic diversity: a comparative study. *Am Nat* 102:243–282. <https://doi.org/10.1086/282541>.
83. de Mendiburu F. 2020. *Agricolae*: Statistical procedures for agricultural research (version 1.3–3). <https://CRAN.R-project.org/package=agricolae>.
84. Legendre P, Gallagher ED. 2001. Ecologically meaningful transformations for ordination of species data. *Oecologia* 129:271–280. <https://doi.org/10.1007/s004420100716>.
85. Lepš J, Šmilauer P. 2003. *Multivariate analysis of ecological data using CANOCO*. Cambridge University Press, New York.
86. Anderson MJ. 2017. *Permutational multivariate analysis of variance (PERMANOVA)*. In: *Wiley Stats Ref: Statistics Reference Online*.
87. Martinez Arbizu P. 2020. *PairwiseAdonis*: pairwiseAdonis: pairwise multi-level comparison using Adonis. Version 0.0.1. <https://github.com/pm-artinezarbizu/pairwiseAdonis>.
88. Weiss S, Xu ZZ, Peddada S, Amir A, Bittinger K, Gonzalez A, Lozupone C, Zaneveld JR, Vázquez-Baeza Y, Birmingham A, Hyde ER, Knight R. 2017. Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome* 5:27. <https://doi.org/10.1186/s40168-017-0237-y>.