

RESEARCH ARTICLE

Open Access

Prevalence of metabolic syndrome in Murcia Region, a southern European Mediterranean area with low cardiovascular risk and high obesity

Diana Gavrila^{1,2*}, Diego Salmerón^{2,1,4}, José-Manuel Egea-Caparrós³, José M Huerta^{2,1}, Alfonso Pérez-Martínez³, Carmen Navarro^{1,2,4} and María-José Tormo^{1,2,4}

Abstract

Background: Metabolic syndrome (MS) is associated with subsequent appearance of diabetes and cardiovascular disease. As compared to other Spanish regions, Murcia (southern Spain) registers increased obesity as well as cardiovascular morbidity and mortality. The aim of this study was to assess the prevalence of MS and its components, awareness of obesity as a health risk and associated lifestyles.

Methods: A population-based, cross-sectional study was conducted in 2003, covering a sample of 1555 individuals 20 years and over. MS was defined according to the Revised National Cholesterol Education Program Adult Treatment Panel III (R-ATPIII), International Diabetes Federation (IDF) and Joint Interim Statement (JIS) criteria. Both low (94/80) and high (102/88) waist circumference (WC) thresholds were considered.

Results: Prevalence of MS was 27.2% (95%CI: 25.2-29.2), 32.2% (95%CI: 30.1-34.3) and 33.2% (95%CI: 31.2-35.3) according to the R-ATPIII, IDF and JIS94/80 respectively. It increased with age until reaching 52.6% (R-ATPIII) or 60.3% (JIS94/80) among persons aged 70 years and over, and was higher in persons with little or no formal education (51.7% R-ATPIII, 57.3% JIS94/80). The most common risk factors were hypertension (46.6%) and central obesity (40.7% and 66.1% according to high and low WC cut-off points respectively). Although most persons were aware that obesity increased health risks, regular exercise was very unusual (13.0% centrally obese, 27.2% non-centrally obese). Adherence to dietary recommendations was similar among centrally obese and non-centrally obese subjects.

Conclusions: Prevalence of MS is high in our population, is comparable to that found in northern Europe and varies with the definition used. Adherence to preventive recommendations and to adequate weight promotion is very low. In the absence of a specific treatment for MS, integrated intervention based on a sustained increase in physical activity and changes in diet should be reinforced.

Background

Metabolic syndrome (MS), a cluster of metabolically related cardiovascular risk factors, is associated with increased risk of death, cardiovascular events or diabetes [1]. Nowadays, estimates suggest that the population-attributable fraction for MS is approximately 6%-7% for all-cause mortality, 12%-17% for cardiovascular disease and 30%-52% for diabetes, depending on the criteria

used [1]. Insulin resistance and central obesity seem to be closely involved in its aetiology [2,3].

Although the Murcia Region registers low cardiovascular mortality [4] and low incidence of acute myocardial infarction [5] at a European level, its myocardial infarction incidence and mortality rates are nevertheless among the highest in Spain [5,6]. Prevalence of diabetes in Murcia is considered to be at the upper limit of the moderate diabetes prevalence range [7], despite a notably high adherence to the Mediterranean dietary pattern and its components [8-10], something that has been shown to have a beneficial effect on insulin sensitivity

* Correspondence: diana.gavrila@carm.es

¹Department of Epidemiology, Regional Authority for Health & Consumer Affairs (Consejería de Sanidad y Consumo), Ronda de Levante 11, 30008 Murcia, Spain

Full list of author information is available at the end of the article

[11]. Overweight and general obesity are very frequent and affect two out of every three adults [12]. Abdominal obesity is one of the highest recorded among European populations [13].

Bearing in mind the important role of central obesity in its appearance, a high prevalence of MS is expected in the adult population of the Murcia Region, thus justifying our interest in the matter. The main difficulty hindering its assessment is the lack of consensus on a definition. Currently, there are several definitions in more or less widespread use. There are: a) the World Health Organization definition [14] including a criterion of albuminuria, hardly available in large population-based surveys; b) the European Group for the Study of Insulin Resistance definition [15] focusing on insulin resistance with the expensive determination of insulinaemia; c) the revised National Cholesterol Education Program Adult Treatment Panel III (R-ATPIII) definition [16] and d) the International Diabetes Federation (IDF) definition [3], focusing on central obesity and using different cut-off points for waist circumference with the aim of being adaptable to different population. Recently, the major organizations involved in the clinical characterisation of MS (the IDF Task Force on Epidemiology and Prevention, the National Heart, Lung and Blood Institute, the American Heart Association, the World Heart Federation, the International Atherosclerosis Society and the International Association for the Study of Obesity) attempted to unify criteria by issuing a Joint Interim Statement (JIS)[17]. Accordingly, our study sought to assess the prevalence of the MS according to the R-ATPIII, IDF and JIS definitions, as well as the prevalence of each of its components, awareness of obesity as a health risk, and associated lifestyles.

Methods

Study Population

Murcia, a Mediterranean area lying in the south-east of Spain, is one of the 17 Autonomous Regions (*comunidades autónomas*) into which the country is divided. The framework for this study was the region's entire adult population aged 20 years and over, a total of 901,920 persons (2001 Census). Two thousand, five hundred and sixty-two subjects were selected by randomised sampling, stratified by age, sex, health area and type of residence (urban or rural), and invited to participate from July 2001 to June 2003. Exclusion criteria were pregnancy and/or any serious physical or mental disease. Subjects were invited to a primary health care centre, where they answered a lifestyle questionnaire administered by trained interviewers, underwent a physical examination, and provided a blood sample for biochemical determinations including fasting plasma glucose, HDL-cholesterol and triglycerides. Details on sampling,

study design, biochemical determinations and physical examination have been published elsewhere [7]. The study was approved by the Ethics Committee of the Virgen de la Arrixaca University Teaching Hospital, the main regional hospital, and all subjects gave their written informed consent.

Lifestyle questionnaire

To assess awareness of the pathogenic role of obesity, subjects were asked whether they considered it a disease, by answering the questions "In your opinion, is obesity a disease?" and "Do you know that obesity is associated with diabetes and heart attacks?" Questions on leisure-time physical activity were based on a previously validated questionnaire [18]. The number of hours of leisure-time physical activity per week was recorded. The American College of Sports Medicine and the American Heart Association recommend that adults should do physical activity for a minimum of 30 minutes a day, five days a week, in order to promote and maintain good health [19]. In line with these recommendations, moderate to vigorous physical activity for two and a half hours per week was considered adequate. Intake of the main dietary components (cereals, fruit, vegetables, legumes, fish, red meat, olive oil and wine) was measured using a food-frequency questionnaire. According to the Mediterranean diet pyramid [20], using olive oil daily, eating cereals, fruit, vegetables more than once daily, legumes and fish more than once per week and red meat less than twice per week was considered adequate. Insofar as alcohol was concerned, wine consumption of one glass per week to three per day for men and one glass per week to one per day for women was deemed to be the recommended moderate intake. Smoking was classified as never, ex- and current smoker. Educational level was defined as little or no formal education, primary, secondary and university.

Physical examination

The participants underwent a physical examination, in which weight and height were measured using a stadiometer with an accuracy of 1 cm (Año-Sayol model 1SB, Barcelona, Spain) and a digital scale with an accuracy of 100 g (TEFAL model SC-3301, GROUPE SEB Iberica, S.A., Barcelona, Spain). Body mass index (BMI) was calculated as weight divided by the square height, in kg/m². Additionally, waist and hip circumferences were measured according to WHO recommendations [21]. Arterial blood pressure was measured according to the MONICA protocol [22], using a digital sphygmomanometer (BOSO oscillomat®). Systolic and diastolic arterial blood pressure was calculated as the mean of two consecutive measures.

Definitions of the metabolic syndrome, impaired fasting glucose and diabetes

Participants were classified as having or not having MS according to the R-ATPIII [16], IDF [3] and JIS definitions. Under the R-ATPIII, a diagnosis of MS, requires three or more of the following criteria: 1) central obesity, defined as elevated waist circumference ≥ 102 cm for males or ≥ 88 cm for females; 2) elevated triglycerides ≥ 150 mg/dL; 3) reduced HDL-cholesterol < 40 mg/dL in men or < 50 mg/dL in women; 4) elevated blood pressure $\geq 130/85$ mmHg or use of an antihypertensive agent; and 5) elevated fasting glucose ≥ 100 mg/dL or previous diagnosis of diabetes or impaired glucose tolerance. Diagnosis of MS according to the IDF definition requires the presence of abdominal obesity defined as elevated waist circumference ≥ 94 cm among men or ≥ 80 cm among women for European populations, together with at least two of the other four criteria of the R-ATPIII definition using the same categorical cut-off points. In much the same way, the JIS definition requires the presence of at least three of the five components and considers two levels of abdominal obesity for European populations, namely: a high cut-off point (102/88) for waist circumference proposed by European Cardiovascular Societies[23]; and a low cut-off point (94/80) as proposed by the IDF. As the JIS high cut-off point definition is identical to the R-ATPIII definition, only data for the JIS low cut-off point definition differ from those for the other former definitions of MS. Individuals on antihypertensive, diabetes and/or lipid-lowering medication were included in the elevated blood pressure, elevated fasting glucose and/or elevated triglyceride groups respectively.

Participants with fasting glucose ≥ 100 mg/dL and < 126 mg/dL, not meeting the criteria for diabetes, were identified as having impaired fasting glucose [24]. Criteria for diabetes were the following: fasting glucose concentration ≥ 126 mg/dL; treatment for diabetes; or plasma HbA1c $\geq 7\%$ [25].

Statistical analysis

Prevalences and their 95% confidence intervals (CIs) were calculated for MS and each of its components. Prevalence estimates were standardised by age using the European standard population. A Wald test was used to test the significance of the differences observed between centrally obese and non-centrally obese subjects in terms of awareness of obesity as a health risk and associated lifestyles. For this purpose central obesity was defined with the high waist-circumference cut-off points (≥ 102 cm for men and ≥ 88 cm for women). All statistical analyses were performed using the STATA 10.0 software package (Stata Corporation, College Station, TX, USA).

Results

Of the 2562 people invited to participate in the study, 61% agreed to give a fasting blood sample, yielding a final sample of 1555 (718 men and 837 women). The age and sex distribution of the sample was similar to that of the general population of the Murcia Region.

The main characteristics of the study population are shown in Table 1. While men displayed a worse profile than did women in terms of glucose status, lipids and smoking, women had a lower educational level and engaged in less physical activity.

Table 2 shows the prevalence of MS and each of its components across categories of sex, age, residence, education and glucose status. Prevalence of MS varied from 27.2% (95%CI: 25.2-29.2) to 32.2% (95%CI: 30.1-34.3) and 33.2% (95%CI: 31.2-35.3), according to the R-ATPIII, IDF and JIS94/80 respectively. Whereas the prevalence of MS according to the IDF and JIS94/80 definitions increased with age to a figure of approximately 60% in persons aged 70 years and over, prevalence of R-ATPIII MS seemed to increase until age 60 years and then stabilise at around 53% in persons aged 60 years and over. The most common cardiovascular risk factors and components of MS in our population were hypertension (46.6%) and central obesity (40.7% or 66.1%, according to high or low waist circumference cut-off points respectively). Prevalence of MS and each of its components increased markedly with age (p for trend < 0.001). Prevalence of MS based on any of the three definitions, and prevalence of each of its components were both higher in persons with little or no formal education (51.7% R-ATPIII, 56.4% IDF and 57.3% JIS94/80) and glucose disorders.

As can be seen from Figure 1, regardless of the definition used, prevalence of MS increased with age, except for men aged 70 years and over, who registered a lower estimated prevalence than did their counterparts in the 60-69 age group. As the IDF and JIS94/80 definitions use lower waist-circumference cut-offs, the estimates of MS prevalence with these definitions were higher in each age and sex group than were the corresponding R-ATPIII/JIS102/88 estimates, with these differences being greater among men.

Figure 2 depicts the degree of awareness of the pathogenic role of obesity and adherence to preventive recommendations and adequate weight control. Although most persons were aware that obesity increased health risks, regular exercise was very unusual (13.0% centrally obese, 27.2% non-centrally obese). The proportion of women doing the recommended two and a half hours of moderate/vigorous physical exercise weekly was extremely low, both for centrally obese (9.9%) and non-centrally obese subjects (19.8%), and amounted to half the corresponding proportions among

Table 1 Characteristics of study subjects

	Men (n = 718)	Women (n = 837)	Total (n = 1555)	p
Age (years), mean ± SD	47.2 ± 17.4	47.6 ± 18.0	47.4 ± 17.7	0.622
20-29%	18.8	18.8	18.8	0.799
30-39	21.6	22.5	22.1	
40-49	17.8	16.5	17.1	
50-59	13.7	13.9	13.8	
60-69	14.2	12.5	13.3	
70+	13.9	15.9	15.0	
Residence, %				
Urban	74.0	76.9	75.6	0.172
Rural	26.0	23.1	24.4	
Education, %				
Little or no formal education	21.1	29.1	25.4	0.004
Primary	20.3	17.0	18.5	
Secondary	39.1	35.6	37.2	
University	19.6	18.3	18.9	
Waist circumference (cm), mean ± SD	97.5 ± 10.8	87.3 ± 13.4	92.1 ± 13.3	<0.001
Systolic BP (mmHg), mean ± SD	134.4 ± 20.6	122.5 ± 24.0	128.0 ± 23.3	<0.001
Diastolic BP (mmHg), mean ± SD	80.4 ± 11.2	74.2 ± 10.3	77.0 ± 11.2	<0.001
Fasting glucose (mg/dl), mean ± SD	99.5 ± 29.8	93.4 ± 25.3	96.2 ± 27.6	<0.001
HDL-Cholesterol (mg/dl), mean ± SD	49.3 ± 11.2	59.6 ± 13.5	54.9 ± 13.5	<0.001
Triglycerides (mg/dl), mean ± SD	128.9 ± 111.2	92.5 ± 51.4	109.3 ± 86.4	<0.001
Glucose status, %				
Normoglycaemia	68.0	78.7	73.8	<0.001
Impaired fasting glucose	19.5	11.7	15.3	
Diabetes	12.5	9.6	10.9	
Smoking, %				
Non-smoker	46.3	65.4	56.6	<0.001
Ex-smoker	15.0	4.1	9.1	
Current smoker	38.7	30.5	34.3	
Leisure-time physical activity				
Household chores (hours/week), mean ± SD	3.8 ± 6.4	22.9 ± 17.3	14.1 ± 16.5	<0.001
Walking (hours/week), mean ± SD	6.9 ± 7.3	5.9 ± 6.1	6.4 ± 6.7	0.003
Moderate/vigorous PA (hours/week), mean ± SD	1.9 ± 3.8	0.9 ± 2.2	1.4 ± 3.1	<0.001

SD: standard deviation. BP: Blood pressure. HDL: High-density lipoprotein. PA: Physical activity

the men. Only 17% of men and 10% of women reported walking at least two hours per week, with practically no difference between centrally obese and non-centrally obese subjects. Frequency of eating the main dietary components was similar among centrally obese and non-centrally obese persons. Only 21% of men and 31% of women reported eating vegetables more than once

daily, and 27% of men and 35% of women consumed red meat less than twice per week. The proportion of women following the recommended frequency of eating fruit, fish and red meat proved slightly higher among centrally obese than non-centrally obese persons. At the opposite extreme, about 95% of persons used olive oil daily, irrespective of sex or obese status.

Table 2 Prevalence (%) of metabolic syndrome and its components according to the R-ATPIII, IDF and JIS94/80 criteria

	Elevated waist circumference 102/88 ¹	Elevated waist circumference 94/80 ²	Elevated triglycerides	Reduced HDL-cholesterol	Elevated blood pressure	Elevated fasting glucose	Metabolic syndrome R-ATPIII	Metabolic syndrome IDF	Metabolic syndrome JIS94/80
	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)	% (95%CI)
Total unadjusted	40.7 (38.6-42.7)	66.1 (64.1-68.2)	22.8 (20.8-24.7)	27.3 (25.1-29.4)	46.6 (44.6-48.6)	26.0 (24.1-28.0)	27.2 (25.2-29.2)	32.2 (30.1-34.3)	33.2 (31.2-35.3)
Total age-adjusted	39.9 (37.8-42.0)	65.8 (63.7-67.9)	22.7 (20.7-24.7)	27.0 (24.8-29.2)	44.9 (42.8-47.0)	25.0 (23.1-27.0)	26.1 (24.1-28.2)	31.1 (28.9-33.2)	32.1 (29.9-34.2)
European population									
Sex									
Men	33.8 (30.6-37.0)	63.7 (60.5-66.9)	29.0 (25.7-32.2)	23.7 (20.6-26.8)	57.7 (54.4-61.0)	32.0 (29.0-35.1)	28.2 (25.1-31.2)	36.9 (33.6-40.2)	38.9 (35.6-42.2)
Women	46.6 (44.0-49.2)	68.3 (65.7-70.9)	17.4 (15.1-19.8)	30.3 (27.3-33.4)	37.1 (34.8-39.5)	20.9 (18.5-23.3)	26.3 (23.8-28.9)	28.1 (25.5-30.8)	28.4 (25.8-31.0)
Age, years									
20-29	10.7 (7.1-14.2)	27.8 (22.8-32.9)	8.2 (5.1-11.3)	16.8 (12.6-21.0)	16.5 (12.3-20.6)	3.8 (1.6-5.9)	3.4 (1.3-5.5)	6.2 (3.4-9.0)	6.2 (3.4-9.0)
30-39	22.3 (17.9-26.7)	53.7 (48.4-58.9)	20.7 (16.8-24.6)	25.4 (20.8-30.0)	23.1 (19.0-27.2)	8.7 (5.8-11.7)	12.5 (9.1-15.9)	18.1 (14.2-22.1)	19.2 (15.2-23.3)
40-49	33.8 (28.1-39.5)	69.6 (63.9-75.2)	22.2 (17.4-27.0)	26.3 (21.1-31.5)	38.3 (33.0-43.5)	22.9 (18.2-27.7)	21.1 (16.3-26.0)	29.5 (24.3-34.8)	30.2 (24.9-35.5)
50-59	59.2 (53.4-64.9)	85.0 (80.4-89.6)	31.3 (25.0-37.6)	30.4 (24.1-36.7)	58.0 (51.5-64.6)	35.5 (29.2-41.9)	37.4 (30.8-43.9)	39.6 (33.0-46.3)	42.1 (35.4-48.7)
60-69	71.4 (65.4-77.3)	92.2 (88.7-95.7)	32.4 (26.1-38.6)	37.7 (31.4-44.0)	80.2 (74.9-85.5)	49.8 (42.9-56.6)	53.6 (46.9-60.3)	57.8 (51.0-64.5)	58.9 (52.3-65.6)
70+	69.3 (64.2-74.4)	88.6 (84.6-92.6)	28.3 (22.7-34.0)	32.2 (26.4-38.0)	88.7 (84.6-92.8)	53.2 (47.0-59.5)	52.6 (46.4-58.7)	59.6 (53.5-65.8)	60.3 (54.2-66.5)
Residence									
Urban	39.7 (37.4-42.1)	64.9 (62.5-67.3)	21.7 (19.5-23.9)	26.9 (24.4-29.4)	45.0 (42.8-47.3)	25.3 (23.1-27.5)	26.1 (23.9-28.4)	30.7 (28.4-33.1)	31.7 (29.4-34.1)
Rural	43.5 (39.5-47.6)	69.9 (66.0-73.8)	26.1 (21.9-30.2)	28.4 (24.1-32.7)	51.6 (47.5-55.7)	28.4 (24.3-32.5)	30.5 (26.5-34.6)	36.8 (32.4-41.1)	37.9 (33.5-42.3)
Education									
Little or no formal education	72.8 (68.7-76.8)	90.7 (88.0-93.5)	33.0 (28.4-37.5)	37.3 (32.7-41.9)	79.1 (75.2-83.0)	45.4 (40.6-50.3)	51.7 (46.8-56.5)	56.4 (51.6-61.3)	57.3 (52.4-62.1)
Primary	49.8 (44.1-55.5)	79.3 (74.6-84.0)	23.7 (18.8-28.6)	26.5 (21.4-31.5)	55.8 (50.2-61.4)	33.8 (28.5-39.1)	34.1 (28.7-39.6)	38.2 (32.7-43.8)	39.4 (33.8-44.9)
Secondary	25.2 (21.7-28.7)	53.4 (49.4-57.4)	18.4 (15.3-21.4)	25.1 (21.6-28.7)	29.4 (25.8-33.0)	13.7 (11.0-16.4)	14.4 (11.6-17.2)	20.0 (16.8-23.2)	21.2 (17.9-24.4)
University	19.6 (15.1-24.1)	45.7 (40.2-51.2)	16.7 (12.6-20.9)	18.4 (14.0-22.9)	28.5 (23.5-33.5)	16.7 (12.6-20.8)	12.7 (8.9-16.4)	18.6 (14.2-22.9)	18.8 (14.5-23.2)
Glucose status									
Normoglycaemia	31.8 (29.4-34.2)	58.2 (55.6-60.8)	16.9 (14.8-19.0)	24.0 (21.5-26.4)	35.6 (33.1-38.1)	0.0 (0.0-13.6)	11.8 (10.0-13.6)	15.9 (13.9-18.0)	16.1 (14.1-18.2)
Impaired fasting glucose	61.3 (55.2-67.3)	87.7 (83.5-91.9)	34.5 (28.5-40.4)	31.5 (25.6-37.5)	73.3 (67.7-78.9)	100 (60.3-72.2)	66.2 (71.1-81.9)	76.5 (73.8-84.0)	78.9 (73.8-84.0)

Table 2 Prevalence (%) of metabolic syndrome and its components according to the R-ATPIII, IDF and JIS94/80 criteria (Continued)

	72.3 (65.5-79.1)	90.4 (85.9-94.8)	45.9 (38.4-53.3)	43.5 (36.2-50.9)	84.0 (78.5-89.5)	98.2 (96.3-100)	76.9 (70.7-83.2)	81.9 (76.1-87.8)	85.2 (79.9-90.5)
--	---------------------	---------------------	---------------------	---------------------	---------------------	--------------------	---------------------	---------------------	---------------------

R-ATPIII: Revised National Cholesterol Education Program Adult Treatment Panel III. IDF: International Diabetes Federation.

JIS 94/80: Joint Interim Statement with waist-circumference cut-off points of ≥ 94 cm for men and ≥ 80 cm for women

¹Waist circumference ≥ 102 cm for men and ≥ 88 cm for women

²Waist circumference ≥ 94 cm for men and ≥ 80 cm for women

Discussion

In the Murcia Region, one-third of the general population aged 20 years and over met the IDF and JIS94/80 definitions for MS, which was slightly higher than the proportion yielded by the R-ATPIII definition. The crude and European population age-adjusted MS prevalence figures were very similar, namely, 27.2% vs. 26.1%, 32.2% vs. 31.1% and 33.2% vs. 32.1% according to the R-ATPIII, IDF and JIS94/80 definitions respectively.

Our findings on the prevalence of MS are comparable to those reported both by Qiao's study [26] based on collaboratively analysed data from nine European cohort studies, and by other studies undertaken in Northern European countries [27,28]. The prevalence in our study was higher than that encountered by other studies conducted in Australia [29], Russia [30], Italy [31] or even other Spanish regions [32-34] in which it ranged from 15 to 22%. Even so, the prevalence estimated by us was lower than that for the US population [35]. To a certain extent, such differences in MS prevalence among studies are due to differences in methodology and, more particularly, in

the measures of obesity used. A recent study by Vega et al. [36] showed that prevalence of MS was influenced by the measure of obesity used, and went on to question the inclusion of any measure of obesity among the diagnostic components of MS. The so-called abridged ATPIII MS - defined as 3 of 4 independent risk factors after obesity is removed from the diagnostic criteria - successfully identified a subgroup of patients who were at higher risk of cardiovascular disease and were therefore candidates for intensive cardiovascular risk reduction. Moreover, regardless of the definition used, MS identifies a group of subjects at significantly greater coronary heart disease risk, and the proportion of such higher-risk subjects is comparable regardless of whether the high or low waist-circumference cut-off point is used, as was recently shown by Alkerwi et al. [37]. In contrast, overweight and obesity measured by reference to body mass index failed to detect any increased cardiovascular risk in our population [6].

In agreement with Martínez-Candela's study [34], conducted on a similar population, and with the studies undertaken by Halldin in Sweden [27], and Alkerwi in Luxembourg [37], prevalence of MS was higher in men than in women. This finding differs from other studies which reported similar prevalences in both sexes [28,35,38] or, on the contrary, a higher prevalence in women than in men [30,31,33,39]. As with other studies, prevalence of MS and of each of its components was observed to increase with age and be higher among the population with a low education level. The results of our study are in line with several other studies [29,40] that have reported a levelling-off or even a decline in the prevalence of MS among men after the sixth or seventh decade of life versus a linear increase with age among women. This difference may be due to selective survival with increased mortality rates among men in general, and among those with metabolic risk factors in particular.

Although most persons were aware of the pathogenic role of obesity, only a small proportion reported physical activity and dietary habits consistent with the preventive recommendations for weight control. While the lowest adherence to the dietary recommendations was found for vegetables and red meat, the highest adherence was found for olive oil. The food consumption patterns reported by centrally obese and non-centrally obese




Figure 1 Prevalence and 95%CI of metabolic syndrome according to the R-ATPIII, IDF and JIS94/80 definitions, by sex and age groups. R-ATPIII: Revised National Cholesterol Education Program Adult Treatment Panel III. JIS 94/80: Joint Interim Statement with waist-circumference cut-off points of ≥ 94 cm for men and ≥ 80 cm for women. IDF: International Diabetes Federation.




Figure 2 Awareness of the pathogenic role of obesity and adherence to preventive recommendations for adequate weight control.
 Central obesity was defined with the high waist-circumference cut-off points (≥ 102 cm for men and ≥ 88 cm for women). ^aWine intake from a minimum of 1/week until a maximum of 3/day in men and 1/day in women. * $p < 0.05$. ** $p < 0.01$.

persons were very similar, except for the higher frequency of fruit and fish intake and lower frequency of red meat intake among centrally obese women.

Substantial differences in physical activity between the centrally obese and non-centrally obese were observed for men and women alike, with the proportion of non-centrally obese persons doing the recommended physical activity being twice as high as the corresponding proportion of centrally obese persons. This is in agreement with Halldin et al's study [27], which observed a strong inverse dose-response association between leisure-time physical activity and MS, and stressed the role of physical activity in the prevention and treatment of MS. Even in the non-centrally obese population, the proportion of persons with an adequate frequency of

physical activity is excessively low, especially among women. A possible explanation for the extremely low proportion of women doing the recommended physical activity could be the long hours that women devote to household chores (an average of 23 hours per week for women vs. 4 hours per week for men), which thus competes with the time available for physical activity.

This study has several limitations. First, the 61% blood-extraction response rate was less than desirable. Nevertheless, as the sex- and age-distribution of the sample was similar to that of the general population, selection bias can be assumed to be minimal. Secondly, while the use of a food-frequency questionnaire without any quantitative or semi-quantitative measure hinders the assessment of energy intake, an important factor in

the appearance and maintenance of obesity, it nevertheless allows for comparison with the Mediterranean diet pyramid [20]. Thirdly, due to the study's cross-sectional design, the dietary pattern observed in the centrally obese population may, to a certain extent, reflect changes in diet introduced as a result of medical advice. Finally, a misreporting bias cannot be ruled out, since obese subjects are known to be prone to overestimate their adherence to healthy lifestyles, e.g., by under-reporting their energy intake and overreporting their intake of fruit and vegetables [41].

The strengths of our study include the fact that it benefits from being a population-based study covering a wide age range, and that prevalence of MS was estimated in accordance with the latest, most widely used definitions. Moreover, by examining both degree of awareness of the pathogenic role of obesity and adherence to preventive recommendations and adequate weight control, we were able to identify the discrepancy between the high degree of awareness and its low impact on improving adherence to healthier lifestyles. The study was based on the most recent data on cardiovascular risk factors available in our region and provides valuable information for decision-makers with regard to areas of health promotion that should be reinforced. Measures targeted at improving adherence to preventive recommendations and adequate weight promotion are required in both sexes, with special stress on dietary recommendations in men and increased physical activity in women.

Further studies are called for to assess trends in cardiovascular risk factor prevalence and changes in lifestyles in our region, and their impact on risk of coronary heart disease. Using a low cut-off point for waist circumference identifies more persons who need intervention; and, as this might prove excessive, further studies are needed to ascertain which waist-circumference cut-off point would be more appropriate for our population. Similarly, longitudinal studies are warranted to assess the impact of MS and abdominal obesity on cardiovascular risk among the local population.

Conclusions

The prevalence of MS is high in our population, is comparable to that found in northern Europe, and varies with the definition used. Adherence to preventive recommendations and adequate weight promotion is very low. In the absence of a specific treatment for MS, integrated intervention based on a sustained increase of physical activity and changes in diet should be reinforced.

Acknowledgements

This study was supported by grants from the Spanish Health Research Fund (Fondo de Investigación Sanitaria Ref 01/0711) and Carlos III Health Institute

(C03/09). The authors would like to thank Mónica Ballesta for her assistance with the graphs.

Author details

¹Department of Epidemiology, Regional Authority for Health & Consumer Affairs (Consejería de Sanidad y Consumo), Ronda de Levante 11, 30008 Murcia, Spain. ²CIBER Epidemiología y Salud Pública (CIBERESP), Spain.

³Clinical Laboratory, Hospital Morales Meseguer, Avda. Marqués de los Vélez s/n, 30008 Murcia, Spain. ⁴Department of Socio-health Sciences, University of Murcia School of Medicine, Spain.

Authors' contributions

DG performed the statistical analyses and drafted the manuscript. DS and JMH conducted a critical review of the manuscript and helped with the statistical analyses. JMEC and APM performed the clinical analysis and contributed to data interpretation and the critical review of the manuscript. CN was involved in the study design, critically reviewing and revising the manuscript, made substantial contributions to its final content, and obtained funding. MUT was involved in study conception and design, co-ordinated the field work, participated in the analysis and in the drafting of the manuscript, and obtained funding. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 16 February 2011 Accepted: 14 July 2011

Published: 14 July 2011

References

1. Ford ES: Risks for all-cause mortality, cardiovascular disease, and diabetes associated with the metabolic syndrome: a summary of the evidence. *Diabetes Care* 2005, 28:1769-1778.
2. Reaven GM: Banting lecture 1988. Role of insulin resistance in human disease. *Diabetes* 1988, 37:1595-1607.
3. Alberti KG, Zimmet P, Shaw J: Metabolic syndrome—a new world-wide definition. A Consensus Statement from the International Diabetes Federation. *Diabet Med* 2006, 23:469-480.
4. Regidor E, Gutiérrez-Fisac JL, Alfaro M: Indicadores de Salud 2009. Evolución de los indicadores del estado de salud en España y su magnitud en el contexto de la Unión Europea. Madrid: Ministerio de Sanidad y Política Social; 2009, 4-6-2010.
5. Marrugat J, Elosua R, Aldasoro E, Tormo MJ, Vanaclocha H, Segura A, et al: Regional variability in population acute myocardial infarction cumulative incidence and mortality rates in Spain 1997 and 1998. *Eur J Epidemiol* 2004, 19:831-839.
6. Huerta JM, Tormo MJ, Gavrilà D, Navarro C: Cardiovascular risk estimated after 13 years of follow-up in a low-incidence Mediterranean region with high-prevalence of cardiovascular risk factors. *BMC Public Health* 2010, 10:640.
7. Valverde JC, Tormo MJ, Navarro C, Rodriguez-Barranco M, Marco R, Egea JM, et al: Prevalence of diabetes in Murcia (Spain): a Mediterranean area characterised by obesity. *Diabetes Res Clin Pract* 2006, 71:202-209.
8. Agudo A, Slimani N, Ocke MC, Naska A, Miller AB, Kroke A, et al: Consumption of vegetables, fruit and other plant foods in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts from 10 European countries. *Public Health Nutr* 2002, 5:1179-1196.
9. Linseisen J, Bergstrom E, Gafa L, Gonzalez CA, Thiebaut A, Trichopoulou A, et al: Consumption of added fats and oils in the European Prospective Investigation into Cancer and Nutrition (EPIC) centres across 10 European countries as assessed by 24-hour dietary recalls. *Public Health Nutr* 2002, 5:1227-1242.
10. Sieri S, Agudo A, Kesse E, Klipstein-Grobusch K, San-Jose B, Welch AA, et al: Patterns of alcohol consumption in 10 European countries participating in the European Prospective Investigation into Cancer and Nutrition (EPIC) project. *Public Health Nutr* 2002, 5:1287-1296.
11. Ryan M, McInerney D, Owens D, Collins P, Johnson A, Tomkin GH: Diabetes and the Mediterranean diet: a beneficial effect of oleic acid on insulin sensitivity, adipocyte glucose transport and endothelium-dependent vasoreactivity. *QJM* 2000, 93:85-91.

12. Martinez-Ros MT, Tormo MJ, Navarro C, Chirlaque MD, Perez-Flores D: Extremely high prevalence of overweight and obesity in Murcia, a Mediterranean region in south-east Spain. *Int J Obes Relat Metab Disord* 2001, 25:1372-1380.
13. Haftenberger M, Lahmann PH, Panico S, Gonzalez CA, Seidell JC, Boeing H, et al: Overweight, obesity and fat distribution in 50- to 64-year-old participants in the European Prospective Investigation into Cancer and Nutrition (EPIC). *Public Health Nutr* 2002, 5:1147-1162.
14. Alberti KG, Zimmet PZ: Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation [see comments]. *Diabet Med* 1998, 15:539-553.
15. Balkau B, Charles MA, Drivsholm T, Borch-Johnsen K, Wareham N, Yudkin JS, et al: Frequency of the WHO metabolic syndrome in European cohorts, and an alternative definition of an insulin resistance syndrome. *Diabetes Metab* 2002, 28:364-376.
16. Grundy SM, Cleeman JL, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al: Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 2005, 112:2735-2752.
17. Alberti KG, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JL, Donato KA, et al: Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation* 2009, 120:1640-1645.
18. Pols MA, Peeters PH, Ocke MC, Slimani N, Bueno-de-Mesquita HB, Collette HJ: Estimation of reproducibility and relative validity of the questions included in the EPIC Physical Activity Questionnaire. *Int J Epidemiol* 1997, 26(Suppl 1):S181-9, S181-S189.
19. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al: Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation* 2007, 116:1081-1093.
20. Willett WC, Sacks F, Trichopoulou A, Drescher G, Ferro-Luzzi A, Helsing E, et al: Mediterranean diet pyramid: a cultural model for healthy eating. *Am J Clin Nutr* 1995, 61:1402S-1406S.
21. WHO Regional Office for Europe: Measuring obesity. Classification and description of anthropometric data. Report on a WHO Consultation on the Epidemiology of Obesity Copenhagen; 1988, Ref Type: Report.
22. Keil U, Kuulasmaa K: WHO MONICA Project: risk factors [published erratum appears in Int J Epidemiol 1990 Sep;19(3):following 775]. *Int J Epidemiol* 1989, 18:S46-S55.
23. Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, et al: European guidelines on cardiovascular disease prevention in clinical practice: executive summary. *Atherosclerosis* 2007, 194:1-45.
24. Diagnosis and classification of diabetes mellitus. *Diabetes Care* 2008, 31(Suppl 1):S55-S60.
25. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 1997, 20:1183-1197.
26. Qiao Q: Comparison of different definitions of the metabolic syndrome in relation to cardiovascular mortality in European men and women. *Diabetologia* 2006, 49:2837-2846.
27. Halldin M, Rosell M, de FU, Hellenius ML: The metabolic syndrome: prevalence and association to leisure-time and work-related physical activity in 60-year-old men and women. *Nutr Metab Cardiovasc Dis* 2007, 17:349-357.
28. Hildrum B, Myklebust A, Hole T, Midtjell K, Dahl AA: Age-specific prevalence of the metabolic syndrome defined by the International Diabetes Federation and the National Cholesterol Education Program: the Norwegian HUNT 2 study. *BMC Public Health* 2007, 7:220.
29. Adams RJ, Appleton S, Wilson DH, Taylor AW, Dal GE, Chittleborough C, et al: Population comparison of two clinical approaches to the metabolic syndrome: implications of the new International Diabetes Federation consensus definition. *Diabetes Care* 2005, 28:2777-2779.
30. Sidorenkov O, Nilssen O, Brenn T, Martiushev S, Arkhipovsky VL, Grjibovski AM: Prevalence of the metabolic syndrome and its components in Northwest Russia: the Arkhangelsk study. *BMC Public Health* 2010, 10:23.
31. Miccoli R, Bianchi C, Odoguardi L, Penno G, Caricato F, Giovannitti MG, et al: Prevalence of the metabolic syndrome among Italian adults according to ATP III definition. *Nutr Metab Cardiovasc Dis* 2005, 15:250-254.
32. Boronat M, Chirino R, Varillas VF, Saavedra P, Marrero D, Fabregas M, et al: Prevalence of the metabolic syndrome in the island of Gran Canaria: comparison of three major diagnostic proposals. *Diabet Med* 2005, 22:1751-1756.
33. Martinez-Larrad MT, Fernandez-Perez C, Gonzalez-Sanchez JL, Lopez A, Fernandez-Alvarez J, Riviriego J, et al: [Prevalence of the metabolic syndrome (ATP-III criteria). Population-based study of rural and urban areas in the Spanish province of Segovia]. *Med Clin (Barc)* 2005, 125:481-486.
34. Martinez CJ, Franch NJ, Romero OJ, Canovas DC, Gallardo MA, Lopez Yepes ML: [Predictive capacity of the diagnostic criteria of metabolic syndrome on the insulin-resistance and the coronary risk]. *Med Clin (Barc)* 2007, 129:601-606.
35. Ford ES: Prevalence of the metabolic syndrome defined by the International Diabetes Federation among adults in the US. *Diabetes Care* 2005, 28:2745-2749.
36. Vega GL, Barlow CE, Grundy SM: Prevalence of the metabolic syndrome as influenced by the measure of obesity employed. *Am J Cardiol* 2010, 105:1306-1312.
37. Alkenwi A, Donneau AF, Sauvageot N, Lair ML, Scheen A, Albert A, et al: Prevalence of the metabolic syndrome in Luxembourg according to the Joint Interim Statement definition estimated from the ORISCAV-LUX study. *BMC Public Health* 2011, 11:4.
38. Boronat M, Saavedra P, Varillas VF, Wagner AM, Lopez-Plasencia Y, Alberice MP, et al: Differences in traditional and emerging cardiovascular risk factors of subjects discordantly classified by metabolic syndrome definitions of the International Diabetes Federation and the National Cholesterol Education Program. *Nutr Metab Cardiovasc Dis* 2009, 19:417-422.
39. Xu WH, Ruan XN, Fu XJ, Zhu QL, Zhang H, Bai Y, et al: Prevalence of the metabolic syndrome in Pudong New Area of Shanghai using three proposed definitions among Chinese adults. *BMC Public Health* 2010, 10:246.
40. Al-Lawati JA, Mohammed AJ, Al-Hinai HQ, Jousilahti P: Prevalence of the metabolic syndrome among Omani adults. *Diabetes Care* 2003, 26:1781-1785.
41. Livingstone MB, Black AE: Markers of the validity of reported energy intake. *J Nutr* 2003, 133(Suppl 3):895S-920S.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1471-2458/11/562/prepub>

doi:10.1186/1471-2458-11-562

Cite this article as: Gavrila et al.: Prevalence of metabolic syndrome in Murcia Region, a southern European Mediterranean area with low cardiovascular risk and high obesity. *BMC Public Health* 2011 11:562.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

