

Chicken Juice Enhances Surface Attachment and Biofilm Formation of *Campylobacter jejuni*

Helen L. Brown,^a Mark Reuter,^a Louise J. Salt,^a Kathryn L. Cross,^a Roy P. Betts,^b Arnoud H. M. van Vliet^a

Institute of Food Research, Norwich, United Kingdom^a; Campden BRI, Chipping Campden, Gloucestershire, United Kingdom^b

The bacterial pathogen *Campylobacter jejuni* is primarily transmitted via the consumption of contaminated foodstuffs, especially poultry meat. In food processing environments, *C. jejuni* is required to survive a multitude of stresses and requires the use of specific survival mechanisms, such as biofilms. An initial step in biofilm formation is bacterial attachment to a surface. Here, we investigated the effects of a chicken meat exudate (chicken juice) on *C. jejuni* surface attachment and biofilm formation. Supplementation of brucella broth with $\geq 5\%$ chicken juice resulted in increased biofilm formation on glass, polystyrene, and stainless steel surfaces with four *C. jejuni* isolates and one *C. coli* isolate in both microaerobic and aerobic conditions. When incubated with chicken juice, *C. jejuni* was both able to grow and form biofilms in static cultures in aerobic conditions. Electron microscopy showed that *C. jejuni* cells were associated with chicken juice particulates attached to the abiotic surface rather than the surface itself. This suggests that chicken juice contributes to *C. jejuni* biofilm formation by covering and conditioning the abiotic surface and is a source of nutrients. Chicken juice was able to complement the reduction in biofilm formation of an aflagellated mutant of *C. jejuni*, indicating that chicken juice may support food chain transmission of isolates with lowered motility. We provide here a useful model for studying the interaction of *C. jejuni* biofilms in food chain-relevant conditions and also show a possible mechanism for *C. jejuni* cell attachment and biofilm initiation on abiotic surfaces within the food chain.

Infection by *Campylobacter* species is a global public health concern, estimated to affect 1% of the population in the developed world annually (1). *Campylobacter jejuni* is the most common cause of human *Campylobacter* infection, representing up to 90% of isolates from clinical cases (2). Infection with *C. jejuni* is also linked to severe postinfectious sequelae, such as Guillain-Barré syndrome and reactive arthritis (3–6). This combination of high disease load and severe postinfectious complications makes *C. jejuni* infection a significant economic and disease burden in many countries worldwide.

The major transmission route for *C. jejuni* is thought to be via contaminated food stuffs, with poultry meat being the main source of infection in urban cases. Sampling of chicken meat from supermarkets showed that up to 70% of meat is contaminated with *C. jejuni* (7). In laboratory conditions, *C. jejuni* is a fastidious organism that requires a temperature of 34 to 44°C and microaerobic conditions for growth. However, during transmission through the food chain it encounters stresses, such as changes in temperature, exposure to aerobic conditions, and lack of nutrients. Significant advances have been made in the understanding of *C. jejuni* stress responses; however, there is still a lack of understanding of how these work together to allow survival of *C. jejuni* in the human food chain. One possible contributor to this survival is the ability of *C. jejuni* to form biofilms (8–11).

Biofilms are commonly defined as attached bacterial colonies of either single or multiple species, encased in an extracellular matrix (11). Biofilms support the survival of bacteria in suboptimal conditions and increase resistance to disinfectants, antimicrobials, and antibiotics (10, 12). To date, it is estimated that 99% of bacteria can grow in biofilms, and it has been suggested that for the majority of bacteria, biofilms are the normal mode of existence (13). *C. jejuni* has been shown to form a monospecies biofilm (8–11, 14, 15) and can also integrate into preexisting biofilms (16).

A serious problem in food processing areas is insufficient or ineffective removal of organic material. Spilled foodstuffs or run-

off from carcass eviscerations contain a complex blend of carbohydrates, proteins, lipids, and sugars (17), providing an ideal medium for bacteria to thrive and survive. A build-up of these organic materials on a surface is here referred to as a conditioning layer. Conditioning layers assist bacterial attachment to surfaces by altering the surface physicochemical properties and attracting the bacteria to the surface due to the increased nutrient availability (18, 19). One well-studied example of a conditioning layer is the oral pellicle, which assists in the attachment of bacterial species such as *Streptococcus mutans* to the tooth surface and contributes to subsequent periodontal disease (20). Surface conditioning layers have also been shown to be important for the initial attachment of food-borne pathogens; for example, *Listeria monocytogenes* survival rates increase when biological soil is present on stainless steel surfaces (21), and milk proteins are able to increase the attachment of *Escherichia coli*, *L. monocytogenes*, and *Staphylococcus aureus* to stainless steel (22).

To date, most studies on *C. jejuni* biofilms have been performed in laboratory conditions, which do not mimic the conditions encountered in the processing environment. It is important to ensure that studies are designed to allow accurate interpretation and extrapolation of laboratory-obtained results to the food industry (23). Various experimental systems have been used to mimic the conditions encountered by *C. jejuni* in the food chain. These models typically include the use of cooked or raw meat (24),

Received 8 August 2014 Accepted 1 September 2014

Published ahead of print 5 September 2014

Editor: M. W. Griffiths

Address correspondence to Arnoud H. M. van Vliet, arnoud.vanvliet@ifr.ac.uk.

Copyright © 2014 Brown et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

doi:10.1128/AEM.02614-14

modeling relevant packaging conditions (23), or the use materials relevant to the food chain such as stainless steel (25). One such model system is the “chicken juice” model (26). This model is based on the collection of exudate from defrosted, commercially obtained chicken carcasses, followed by supplementation or replacement of standard laboratory media with this sterile-filtered liquid. Supplementation of brucella broth with chicken juice resulted in increased survival of planktonic cells of *C. jejuni* following both chilled and frozen storage (26, 27).

We investigated here the effect of chicken juice on the attachment of *C. jejuni* to surfaces and subsequent biofilm formation. We show that in the presence of chicken juice, *C. jejuni* biofilm formation is increased and that this increase in biofilm levels is not simply due to increased cell numbers within the suspensions but to an increase in attachment to abiotic surfaces. We show that this increase in attachment is due to the ability of chicken juice to condition abiotic surfaces relevant to food processing environments.

MATERIALS AND METHODS

***C. jejuni* strains and growth conditions.** *C. jejuni* reference strains NCTC 11168 (28), 81116 (29), 81-176 (30), and RM1221 (31), an NCTC 11168 nonmotile (aflagellate) mutant (NCTC 11168 Δ *flaAB*) (10), and *C. coli* clinical isolate 15-537360 (32) were routinely cultured in a MACS-MG-1000 controlled atmosphere cabinet (Don Whitley Scientific) under microaerobic conditions (85% N₂, 5% O₂, 10% CO₂) at 37°C. For growth on plates, strains were either grown on brucella agar or blood agar base (Becton Dickinson) with Skirrow supplements (10 µg/ml vancomycin, 5 µg/ml trimethoprim, 2.5 IU of polymyxin B). Broth culture was carried out in brucella broth (Becton Dickinson). An Innova 4230 (New Brunswick Scientific) incubator was used for aerobic culture at 37°C.

Preparation of chicken juice. Chicken juice (meat exudate) was prepared as described previously (26). Briefly, frozen whole chickens were purchased from four different United Kingdom supermarkets, with no significant differences observed between different supermarkets and different whole chicken (data not shown). The whole chickens were thawed overnight at room temperature, and the exudate was collected, centrifuged to remove debris, and sterilized by using a 0.2-µm-pore-size sterile polyethersulfone syringe filter (Millipore). Chicken juice was divided into aliquots and stored at -20°C until use. Chicken juice was diluted to various percentages (vol/vol) in brucella broth unless stated otherwise.

Precoating of abiotic surfaces. Sterile stainless steel coupons (stainless steel type 1.4301 according to European Standard EN 10088-1, with a type 2B finish according to European Standard EN 10088-2) were placed in a six-well polystyrene tissue culture plate (Corning) and incubated with 4 ml of brucella broth, brucella broth containing chicken juice, or 100% chicken juice. Likewise, sterile borosilicate glass test tubes (Corning) were incubated with 1 ml of brucella broth, brucella broth containing chicken juice, or 100% chicken juice. Samples were incubated overnight at 37°C in aerobic conditions to allow precoating. The medium was subsequently removed, and surfaces were washed with an equal volume of phosphate-buffered saline (PBS; 1 ml for test tubes, 4 ml for six-well plates with stainless steel coupons) and immediately used for biofilm assays using brucella broth.

***Campylobacter* growth for biofilm assay.** Single-use glycerol stocks of *C. jejuni* were thawed, inoculated onto Skirrow plates and grown overnight at 37°C in microaerobic conditions (5% O₂, 10% CO₂, 85% N₂). Cells from the Skirrow plate were used to inoculate brucella broth and incubated overnight shaking (37°C, microaerobic conditions). After overnight growth, cell cultures were adjusted to an A₆₀₀ of 0.05 in brucella broth, brucella broth supplemented with 5% (vol/vol) chicken juice or 100% chicken juice. To allow biofilm formation, 1 ml of this solution was added to either a sterile borosilicate glass test tube (Corning) or a 24-well polystyrene tissue culture plate (Corning), or 3 ml was added to a six-well

polystyrene tissue culture plate (Corning) containing a sterile stainless steel coupon. Tubes were incubated at 37°C under either microaerobic or atmospheric air conditions for 48 h before staining.

Congo red staining. A 0.1% (vol/vol) concentration of Congo red was added to brucella broth, brucella broth supplemented with 5% (vol/vol) chicken juice, or 100% chicken juice with or without *C. jejuni*, before static incubation at 37°C in microaerobic conditions for 48 h. At the end of the incubation period, the medium was removed from the tube before washing with 1 ml of PBS and drying at 37°C. Bound Congo red dye was dissolved by adding 20% acetone–80% ethanol, followed by incubation on a rocking platform for 15 min at room temperature. The level of dissolved dye was measured at a wavelength of 500 nm using a Biomate 5 spectrophotometer (Thermo Scientific).

Crystal violet staining. Biofilms were formed in brucella broth, brucella broth supplemented with 5% (vol/vol) chicken juice, or 100% chicken juice. After biofilm formation, the medium was removed from the test tubes before the samples were washed with water and dried at 60°C for 30 min. Next, 1 ml of 1% (wt/vol) crystal violet solution was added, and the tubes were further incubated on a rocker at room temperature for 30 min. The unbound dye was removed from the tubes by thorough washing in water, followed by drying at 37°C. Bound crystal violet was dissolved by adding 20% acetone–80% ethanol, followed by incubation on a rocking platform for 15 min at room temperature. The resulting dissolved dye was measured at a wavelength of 590 nm by using a Biomate 5 spectrophotometer (Thermo Scientific).

TTC staining. 2,3,5-Triphenyltetrazolium chloride (TTC) staining was carried out as described previously (33). Briefly, cell suspensions were removed after 48 h of incubation, and the tubes were washed twice with 1 ml of sterile PBS. Then, 1.2 ml of brucella broth supplemented with 0.05% (wt/vol) TTC was added to each test tube before incubation at 37°C under microaerobic conditions for 72 h. After incubation, the TTC solution was removed, and the test tubes were air dried. Bound TTC dye was dissolved as described above using 20% acetone–80% ethanol, and the A₅₀₀ of the solution was measured.

Assessment of cell viability by culture. To determine the number of viable cells in the planktonic fraction, the medium of biofilm experiments was 10-fold serially diluted in PBS, and 5 µl of each dilution was spotted onto brucella agar plates. After 2 days of growth, the dilution resulting in two or more colonies was recorded. Cell viability in biofilm assays was assessed upon initial addition of cultures into static culture, before washing and the addition of TTC-containing media and after incubation with TTC-containing media. The cell viability in growth assays was assessed every 2 h in the first 8 h of the experiment and every 24 h thereafter.

Use of TTC as a growth indicator. *C. jejuni* was grown as described above and diluted to an A₆₀₀ of 0.05 in brucella broth supplemented with 0.05% TTC before incubation at 37°C under microaerobic conditions for 48 h (33). Formazan crystals were then dissolved by adding an equal volume of 20% acetone–80% ethanol and incubating the samples at room temperature for 30 min before centrifugation (20,000 × g, 10 min at room temperature). The A₅₀₀ of the supernatant was then measured.

SEM. The biofilms were collected on Thermanox coverslips (Agar Scientific, Stansted, United Kingdom) and fixed with 2.5% glutaraldehyde in 0.1 M PIPES buffer (pH 7.4) for 1 h. The fixative was then replaced with three changes of 0.1 M PIPES buffer. The cells, supported by the coverslips, were then dehydrated in a series of ethanol solutions (at 30, 40, 50, 60, 70, 80, and 90% and then three times at 100%) for at least 20 min for each step. Samples were critical point dried in a Polaron E3000 critical point drier using liquid carbon dioxide as the transition fluid. The coverslips were then mounted with the cell layer facing upward on aluminum scanning electron microscopy (SEM) stubs using sticky tabs. The samples were coated with gold in an Agar high resolution sputter-coater apparatus. SEM was carried out using a Zeiss Supra 55 VP FEG SEM operating at 3 kV.

Statistics. Statistical analysis was carried out using both GraphPad Prism and SPSS software. At least three biological replicates (each with




FIG 1 Biofilm formation and growth of *C. jejuni* NCTC 11168 in the presence of chicken juice. (A) Static incubation of *C. jejuni* in brucella broth supplemented with chicken juice results in increased biofilm formation, as shown by using a TTC biofilm assay. (B) Growth of *C. jejuni* in media supplemented with 5% chicken juice is not significantly different from unsupplemented brucella broth. White bars represent unsupplemented brucella broth, and black bars represent brucella broth supplemented with 5% (vol/vol) chicken juice. Error bars show the standard errors of the mean, and significance was measured by using the Bonferroni post test following ANOVA (**, $P < 0.01$; ***, $P < 0.001$).

three technical replicates) were used to calculate means and the standard errors of the mean. Significance was measured using either Mann-Whitney U test or Bonferroni post test values following analysis of variance (ANOVA).

RESULTS

***C. jejuni* forms increased levels of biofilm in the presence of chicken juice.** Meat and meat exudates have been previously reported to allow for an increase in survival of *C. jejuni* (23, 26, 27). To assess whether meat exudates affect *C. jejuni* biofilm formation, we measured the biofilm levels in static *C. jejuni* NCTC 11168 cultures supplemented with meat exudates recovered from defrosted chicken carcasses (chicken juice) and from pork steaks. Since dyes such as crystal violet and Congo red aspecifically bind to meat exudate components (33), we measured biofilm formation by *C. jejuni* via conversion of the respiratory dye TTC, which relies on detecting redox activity from adhered bacterial cells. Supplementation of brucella broth with chicken juice resulted in an increase in biofilm formation compared to brucella broth alone in both microaerobic and aerobic conditions (Fig. 1A). Replacement of medium by 100% chicken juice gave the highest level of biofilm

formation, and this was not due to differences in viability, since cultures incubated in brucella broth, brucella broth with 5% chicken juice, and 100% chicken juice had similar levels of viable planktonic cells. Likewise, addition of pork exudate resulted in a 2-fold increase in biofilm formation in both microaerobic and aerobic conditions (data not shown).

To differentiate between growth and biofilm formation, we assessed growth of *C. jejuni* NCTC 11168 in brucella broth, brucella broth supplemented with 5% chicken juice, and 100% chicken juice in shaking cultures. There was no statistical difference between growth in brucella broth and media supplemented with 5% chicken juice over a 24 h period (Fig. 1B), and thus the increase in biofilm formation in the presence of chicken juice is likely to be solely due to increased attachment of *Campylobacter* to the abiotic surface. In 100% chicken juice, the mean A_{500} value of the 24 h sample was significantly higher than that of the unsupplemented brucella control (data not shown), suggesting that the increased biofilm formation present in 100% chicken juice could in part be due to enhanced growth of *C. jejuni*. These results also show that chicken juice supports *C. jejuni* growth.

Chicken juice increases biofilm formation in different *Campylobacter* isolates and on different abiotic surfaces. In order to ensure that the effect observed in the glass test tubes was present on other abiotic surfaces and not specific for strain NCTC 11168, we repeated the previous assay using polystyrene plates and stainless steel coupons and extended the assay to three other *C. jejuni* reference isolates (81116, 81-176, and RM1221) and one *C. coli* clinical isolate (15-537360). Stainless steel is a commonly used material within the food chain and so is an important surface for bacterial attachment and subsequent biofilm formation and survival. All *C. jejuni* and *C. coli* strains showed a significant increase in biofilm formation when brucella broth was supplemented with 5% chicken juice in borosilicate test tubes and 24-well polystyrene wells under both microaerobic and aerobic conditions (Fig. 2A to D). The chicken juice-dependent increase in biofilm formation was particularly clear in *C. jejuni* RM1221 and *C. coli* 15-537360, since these strains showed very low levels of biofilm formation in brucella broth alone (Fig. 2A to D). Biofilm formation was also significantly increased in the presence of chicken juice on food grade stainless steel coupons (Fig. 2E and F). Hence, chicken juice is able to promote biofilm formation, independently of *Campylobacter* isolate or abiotic surface.

***C. jejuni* preferentially attaches to chicken juice particulates.** Since biofilm formation was increased by chicken juice on different surfaces, we investigated the effect of chicken juice on an abiotic surface in the absence of *C. jejuni*. Brucella broth with or without 5% chicken juice and also 100% chicken juice were incubated in static glass tubes under the standard assay conditions and stained with TTC, crystal violet, or Congo red (Fig. 3A). There was a significant increase in crystal violet and Congo red staining in the presence of chicken juice, while staining with TTC (measuring bacterial respiration) was negative, demonstrating that components of chicken juice bind to the abiotic surface but do not interfere with TTC staining. Since the formation of precipitates (particulates) was also observed, we hypothesized that chicken juice components may form a conditioning layer on the abiotic surface, facilitating bacterial attachment.

In order to further investigate this phenomenon, *C. jejuni* NCTC 11168 biofilms obtained with brucella broth, brucella broth supplemented with 5% chicken juice, or 100% chicken juice




FIG 2 Static incubation of four strains of *C. jejuni* and one strain of *C. coli* in the presence of chicken juice leads to increased biofilm formation in various abiotic materials. Graphs A, C, and E show data for biofilms incubated in atmospheric conditions, and graphs B, D, and F show data from a comparable treatment in microaerobic conditions. The materials tested included borosilicate glass (A and B), polystyrene (C and D), and stainless steel (E and F). There was an increase in TTC conversion when brucella broth was supplemented or replaced by chicken juice. The figure shows the quantity of biofilm formation measured by TTC conversion. Error bars show the standard errors of the mean, and significance was measured by using the Bonferroni post test (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$).

were analyzed by SEM (Fig. 3B to D). In the presence of chicken juice, *C. jejuni* cells preferentially bind to the particulates rather than directly to the abiotic surface (Fig. 3C and D). This is especially apparent in the 5% chicken juice image (Fig. 3C), where only the chicken juice particulates, but not the abiotic surface, are bound by *C. jejuni* cells. Figure 3D also visually supports the observations in Fig. 1B that the total number of cells within the biofilm is increased in 100% chicken juice. Hence, chicken juice provides a highly adhesive environment supporting subsequent formation of a *C. jejuni* biofilm.

Precoating assay tubes with chicken juice increases biofilm formation. All previous experiments were performed with simultaneous addition of *C. jejuni* and chicken juice, and therefore we investigated whether precoating surfaces with chicken juice also

enhanced biofilm formation. A range of chicken juice concentrations was tested during the precoating stage from brucella broth supplemented with 10 to 90% chicken juice and with 100% chicken juice for 24 h at 37°C. Subsequent replacement of precoating medium with *C. jejuni* NCTC 11168 in unsupplemented brucella broth resulted in a significant increase in levels of biofilm formation with all concentrations of chicken juice compared to brucella broth under both aerobic and microaerobic conditions (Fig. 4). There was no significant increase in levels of biofilm formation with increasing concentrations of chicken juice. This was also observed by precoating stainless steel coupons with chicken juice.

Chicken juice complements reduced biofilm formation by aflagellated *C. jejuni*. Flagella are known to contribute to attach-




FIG 3 Chicken juice facilitates binding of *C. jejuni* via modification of abiotic surfaces. (A) Chicken juice components bind to abiotic surfaces such as glass tubes, as shown by crystal violet staining (top row) and Congo red staining (middle row). TTC staining (bottom row) shows that the material bound is




FIG 4 Precoating of test tubes with chicken juice increases biofilm formation by *C. jejuni* NCTC 11168. Tubes were precoated with a range of chicken juice concentrations before being used in the standard TTC biofilm assay under both aerobic (A) and microaerobic (B) conditions, using unsupplemented brucella broth. Error bars show the standard errors of the mean, and significance was assessed by using the Mann-Whitney U test (*, $P < 0.05$).

ment and biofilm formation in several bacterial pathogens (34, 35), and an aflagellated *C. jejuni* Δ *flaAB* mutant produces significantly less biofilm than the wild-type NCTC 11168 strain (10, 36). Incubation with chicken juice or precoating of tubes with chicken juice both resulted in a significant increase of biofilm formation with the *C. jejuni* Δ *flaAB* mutant compared to incubation in brucella broth alone (Fig. 5). In the presence of chicken juice, biofilm levels were similar to that of wild-type *C. jejuni* NCTC 11168 (Fig. 5), showing that chicken juice can complement the lack of flagella and support biofilm formation by aflagellated strains. This sup-

not redox reactive. The left column shows results for brucella broth only, the middle column shows results for brucella broth supplemented with 5% chicken juice, and the right column shows results for 100% chicken juice. (B to D) Representative SEM images of *C. jejuni* biofilms grown in brucella broth supplemented with 0% (B), 5% (C), or 100% (D) chicken juice on coverslips. In chicken juice-containing samples (C and D), *C. jejuni* can be seen to adhere to the juice particulates rather than the abiotic surface. A large chicken juice particulate can be seen adhered to the slide surface in panel C, with *C. jejuni* attached to it in preference to the slide surface. In panel D, particulates are densely packed and so cover the field of view. Scale bar, 10 μ m.




FIG 5 Chicken juice increases the ability of *C. jejuni* NCTC 11168 Δ *flaAB* mutants to form biofilms in static culture. Static suspensions of *C. jejuni* Δ *flaAB* mutants were incubated for 48 h to allow biofilm formation in various types of media before TTC staining. A bar chart shows (from left to right) results for *C. jejuni* Δ *flaAB* mutants in brucella broth (with no pretreatment of the test tubes), *C. jejuni* Δ *flaAB* mutants in 100% chicken juice (with no pretreatment of the test tubes), *C. jejuni* Δ *flaAB* mutants in brucella broth (with a 24-h brucella broth pretreatment of the test tubes), *C. jejuni* Δ *flaAB* mutants in brucella broth (with a 24-h 100% chicken juice pretreatment of the test tubes), and a *C. jejuni* NCTC 11168 wild-type (WT) culture (with no pretreatment of the test tubes). Error bars show standard errors of the mean, and images above the bar chart are representative of the TTC staining observed for each condition. Significance was measured using a Bonferroni post test following ANOVA (**, $P < 0.01$).

ports our hypothesis that the effect of chicken juice is mediated through facilitating attachment and not via chemotactic motility.

DISCUSSION

In this study we investigated the effect of meat exudates on *C. jejuni* biofilm formation and show that chicken juice is able to enhance biofilm formation compared to brucella broth. Our data show that this is mediated by the ability of chicken juice to provide a conditioning layer on abiotic surfaces, providing an adhesive foundation onto which a *C. jejuni* biofilm can establish itself and grow. This is observed in both isolates capable of forming biofilms in brucella broth and isolates that are otherwise poor biofilm formers (Fig. 2). In an industrial food setting, this means that the presence of meat exudates can aggravate the problem of contamination by food-borne pathogens such as *C. jejuni*. Conditioning is defined as the development of absorbed layers onto a surface (22) and can also be seen as biofouling if it is in a undesirable area, for example, the food chain or pipelines (37). Within the food chain, biofouling is an important area of study since it contributes to increased biofilm formation, the loss of heat transfer efficiency, and reduced liquid flow in pipes (37). Our findings add another dimension to the conclusions in a recent literature review on *C. jejuni* biofilms (38), which concluded that attachment and survival on surfaces and in existing biofilms of other species is the

most likely mechanism for *C. jejuni* to persist in the food chain, rather than *de novo* biofilm formation. Although the aforementioned mechanisms indeed contribute to persistence, meat exudates can enhance survival of *C. jejuni* by increasing surface adhesion and by providing a scaffold with nutrients and materials the bacterium can use to form a biofilm.

Biofilms are frequently found in the food chain and support bacterial persistence in suboptimal conditions. They are also frequently detected in many different areas of poultry processing plants, from conveyor belts (39) and stainless steel surfaces (25) to floor sealant (40). The food chain is very complex and dynamic, containing varied bacterial contamination sources, environmental conditions, and nutrient sources (41). *In vitro* laboratory studies allow for a reductionist approach, controlling variables to assess the effect of specific conditions, material, or genes on biofilm formation; however, a middle ground must be found in which experimental setup allows control but reflects the complexity of the food chain. The chicken juice system (26) is one method of experimenting with food chain relevance in a laboratory setting. Chicken juice more accurately reflects the conditions in the food chain but is easy to manipulate and reproducible. Several food relevant compounds have been identified to be able to form conditioning layers by their ability to increase biofilm formation in various food relevant bacteria. Bacterial soil increases *L. monocytogenes* survival on surfaces (21), while milk residues and chicken fillet suspension increase the survival of planktonic *Salmonella enterica* serovar Enteritidis and *C. jejuni* on stainless steel (42). Although the notion of conditioning layers within the food chain is not novel, to our knowledge this is the first study proposing a mechanism for the effect of chicken juice on *C. jejuni* and *C. coli* biofilm formation, as well as investigating the capacity of chicken juice to condition food chain-relevant abiotic surfaces.

Many animal macromolecules have been reported to be able to form a conditioning film, but they are not always able to enhance biofilm formation. For instance, bovine serum albumin reduces biofilm formation in *S. aureus* (43) and *Burkholderia cepacia* (18). Conversely, whey protein and casein are important for *Cronobacter* biofilm formation (44), although skimmed milk and milk albumin had the opposite effect, inhibiting biofilm formation (22). Additional factors, such as surface roughness and hydrophobicity, will also affect bacterial attachment and biofilm formation. Hydrophilic surfaces, such as stainless steel and glass, increase the time required for bacterial attachment and biofilm formation (37). Surface microstructure is also capable of affecting protein absorption (45), again leading to variability in surface conditioning and subsequent biofilm formation. We have demonstrated that biofilm formation in *C. jejuni* NCTC 11168 Δ *flaAB* mutants is also enhanced following the precoating of test tubes with chicken juice (Fig. 5), thus complementing the reduced biofilm phenotype of the mutant. This mutant is aflagellated, unable to swarm, and thus unable to migrate toward food sources (46). This means that an increase in attachment and subsequent biofilm formation must be due to alteration of the glass surface properties by the conditioning layer from the chicken juice rather than due to increased chemotactic or energy taxis-directed motility toward a food source.

Since poultry is the most important source of *Campylobacter* infection in the Western world, we have limited this investigation to the effect that chicken juice has on biofilm formation. Both *C. jejuni* and *C. coli* are able to contaminate not only chicken but also

turkey, pork, and beef (47). For instance, 49.3% of chicken samples tested were positive for *Campylobacter* species, along with turkey (37.5% of samples), duck (45.8% of samples), beef (3.2% of samples), pork (5.1% of samples), lamb (11.8% of samples), oysters (2.3% of samples), and milk (1.6% of samples) (48). Subsequent speciation suggested that *C. jejuni* and *C. coli* accounted for 83.4 and 16.6% of the isolates, respectively. In our SEM images (Fig. 3B to D), *C. jejuni* can be observed preferentially attaching to the adhered chicken juice components rather than the surface of the slide. This highlights the need for future studies to not only investigate the link between chicken or pork soil and surface conditioning but also assess the effect of other meat exudates on biofilm formation.

In conclusion, chicken juice allows increased attachment of *C. jejuni* as it attaches to the surface of the test tubes, providing a conditioned surface for the bacteria to adhere to. This conditioning surface is still present following a simple washing procedure and able to increase biofilm formation if the subsequent incubation with bacteria lacks chicken juice in the broth. Chicken juice also provides a suitable laboratory model for the study of *C. jejuni* biofilm formation in the food chain, allowing investigators to more closely mimic the food chain conditions that lead to *C. jejuni* spread and cross contamination of carcasses. Furthermore, identification of the chicken juice components involved in surface conditioning and bacterial attachment may give the opportunity for targeted intervention and prevention strategies to reduce transmission of *C. jejuni* and *C. coli* through the food chain.

ACKNOWLEDGMENTS

We thank the members of the IFR *Campylobacter* Research Group and Gary Barker for helpful discussions. We also thank Maddy Houchen for microbiology medium support and Lawrence Staniforth for the gift of food-grade stainless steel coupons.

We gratefully acknowledge the support of the Biotechnology and Biological Sciences Research Council (BBSRC) via the BBSRC Institute Strategic Programme (BB/J004529/1) and a BBSRC CASE studentship (BB/I15321/1) with CASE funding from Campden BRI.

REFERENCES

- Altekruse SF, Stern NJ, Fields PI, Swerdlow DL. 1999. *Campylobacter jejuni*: an emerging foodborne pathogen. *Emerg. Infect. Dis.* 5:28–35. <http://dx.doi.org/10.3201/eid0501.990104>.
- Cody AJ, McCarthy ND, Jansen van Rensburg M, Isinkaye T, Bentley SD, Parkhill J, Dingle KE, Bowler IC, Jolley KA, Maiden MC. 2013. Real-time genomic epidemiological evaluation of human *Campylobacter* isolates by use of whole-genome multilocus sequence typing. *J. Clin. Microbiol.* 51:2526–2534. <http://dx.doi.org/10.1128/JCM.00066-13>.
- Nichols GL, Richardson JF, Sheppard SK, Lane C, Sarran C. 2012. *Campylobacter* epidemiology: a descriptive study reviewing 1 million cases in England and Wales between 1989 and 2011. *BMJ Open* 2:e001179. <http://dx.doi.org/10.1136/bmjopen-2012-001179>.
- Tam CC, Rodrigues LC, Viviani L, Dodds JP, Evans MR, Hunter PR, Gray JJ, Letley LH, Rait G, Tompkins DS, O'Brien SJ. 2012. Longitudinal study of infectious intestinal disease in the UK (IID2 study): incidence in the community and presenting to general practice. *Gut* 61:69–77. <http://dx.doi.org/10.1136/gut.2011.238386>.
- Wakerley BR, Yuki N. 2013. Infectious and noninfectious triggers in Guillain-Barre syndrome. *Exp. Rev. Clin. Immunol.* 9:627–639. <http://dx.doi.org/10.1586/1744666X.2013.811119>.
- Zautner AE, Johann C, Strubel A, Busse C, Tareen AM, Masanta WO, Lugert R, Schmidt-Ott R, Gross U. 2014. Seroprevalence of campylobacteriosis and relevant postinfectious sequelae. *Eur. J. Clin. Microbiol. Infect. Dis.* 33:1019–1027. <http://dx.doi.org/10.1007/s10096-013-2040-4>.
- Jorgensen F, Ellis-Iversen J, Rushton S, Bull SA, Harris SA, Bryan SJ, Gonzalez A, Humphrey TJ. 2011. Influence of season and geography on *Campylobacter jejuni* and *C. coli* subtypes in housed broiler flocks reared in Great Britain. *Appl. Environ. Microbiol.* 77:3741–3748. <http://dx.doi.org/10.1128/AEM.02444-10>.
- Ica T, Caner V, Istanbul O, Nguyen HD, Ahmed B, Call DR, Beyenal H. 2011. Characterization of mono- and mixed-culture *Campylobacter jejuni* biofilms. *Appl. Environ. Microbiol.* 78:1033–1038. <http://dx.doi.org/10.1128/AEM.07364-11>.
- Reeser RJ, Medler RT, Billington SJ, Jost BH, Joens LA. 2007. Characterization of *Campylobacter jejuni* biofilms under defined growth conditions. *Appl. Environ. Microbiol.* 73:1908–1913. <http://dx.doi.org/10.1128/AEM.00740-06>.
- Reuter M, Mallett A, Pearson BM, van Vliet AHM. 2010. Biofilm formation by *Campylobacter jejuni* is increased under aerobic conditions. *Appl. Environ. Microbiol.* 76:2122–2128. <http://dx.doi.org/10.1128/AEM.01878-09>.
- Siringan P, Connerton PL, Payne RJ, Connerton IF. 2011. Bacteriophage-mediated dispersal of *Campylobacter jejuni* biofilms. *Appl. Environ. Microbiol.* 77:3320–3326. <http://dx.doi.org/10.1128/AEM.02704-10>.
- Sofos JN, Geornaras I. 2010. Overview of current meat hygiene and safety risks and summary of recent studies on biofilms, and control of *Escherichia coli* O157:H7 in nonintact, and *Listeria monocytogenes* in ready-to-eat, meat products. *Meat Sci.* 86:2–14. <http://dx.doi.org/10.1016/j.meatsci.2010.04.015>.
- Costerton JW, Stewart PS, Greenberg EP. 1999. Bacterial biofilms: a common cause of persistent infections. *Science* 284:1318–1322. <http://dx.doi.org/10.1126/science.284.5418.1318>.
- Teh KH, Flint S, French N. 2010. Biofilm formation by *Campylobacter jejuni* in controlled mixed-microbial populations. *Int. J. Food Microbiol.* 143:118–124. <http://dx.doi.org/10.1016/j.ijfoodmicro.2010.07.037>.
- Sulaeman S, Le Bihan G, Rossero A, Federighi M, De E, Tresse O. 2010. Comparison between the biofilm initiation of *Campylobacter jejuni* and *Campylobacter coli* strains to an inert surface using BioFilm ring test. *J. Appl. Microbiol.* 108:1303–1312. <http://dx.doi.org/10.1111/j.1365-2672.2009.04534.x>.
- Hanning I, Jarquin R, Slavik M. 2008. *Campylobacter jejuni* as a secondary colonizer of poultry biofilms. *J. Appl. Microbiol.* 105:1199–1208. <http://dx.doi.org/10.1111/j.1365-2672.2008.03853.x>.
- Chmielewski RAN, Frank JF. 2007. Inactivation of *Listeria monocytogenes* biofilms using chemical sanitizers and heat, p 73–104. In *Biofilms in the food environment*. Blackwell Publishing, Ltd., London, United Kingdom.
- Hwang G, Kang S, El-Din MG, Liu Y. 2012. Impact of conditioning films on the initial adhesion of *Burkholderia cepacia*. *Colloids Surf. B Biointerfaces* 91:181–188. <http://dx.doi.org/10.1016/j.colsurfb.2011.10.059>.
- Dat NM, Hamanaka D, Tanaka F, Uchino T. 2010. Surface conditioning of stainless steel coupons with skim milk solutions at different pH values and its effect on bacterial adherence. *Food Control* 21:1769–1773. <http://dx.doi.org/10.1016/j.foodcont.2010.06.012>.
- Di Giulio M, Di Bartolomeo S, Di Campli E, Sancilio S, Marsich E, Travan A, Cataldi A, Cellini L. 2013. The effect of a silver nanoparticle polysaccharide system on streptococcal and saliva-derived biofilms. *Int. J. Mol. Sci.* 14:13615–13625. <http://dx.doi.org/10.3390/ijms140713615>.
- Van Houdt R, Michiels CW. 2010. Biofilm formation and the food industry, a focus on the bacterial outer surface. *J. Appl. Microbiol.* 109:1117–1131. <http://dx.doi.org/10.1111/j.1365-2672.2010.04756.x>.
- Barnes LM, Lo MF, Adams MR, Chamberlain AH. 1999. Effect of milk proteins on adhesion of bacteria to stainless steel surfaces. *Appl. Environ. Microbiol.* 65:4543–4548.
- Balamurugan S, Nattress FM, Baker LP, Dilts BD. 2011. Survival of *Campylobacter jejuni* on beef and pork under vacuum packaged and retail storage conditions: examination of the role of natural meat microflora on *C. jejuni* survival. *Food Microbiol.* 28:1003–1010. <http://dx.doi.org/10.1016/j.fm.2011.01.012>.
- Yoon KS, Burnette CN, Oscar TP. 2004. Development of predictive models for the survival of *Campylobacter jejuni* (ATCC 43051) on cooked chicken breast patties and in broth as a function of temperature. *J. Food Prot.* 67:64–70.
- Sanders SQ, Frank JF, Arnold JW. 2008. Temperature and nutrient effects on *Campylobacter jejuni* attachment on multispecies biofilms on stainless steel. *J. Food Prot.* 71:271–278.
- Birk T, Ingmer H, Andersen MT, Jorgensen K, Brondsted L. 2004. Chicken juice, a food-based model system suitable to study survival of *Campylobacter jejuni*. *Lett. Appl. Microbiol.* 38:66–71. <http://dx.doi.org/10.1046/j.1472-765X.2003.01446.x>.

27. Birk T, Rosenquist H, Brondsted L, Ingmer H, Bysted A, Christensen BB. 2006. A comparative study of two food model systems to test the survival of *Campylobacter jejuni* at -18°C . *J. Food Prot.* 69:2635–2639.
28. Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, Basham D, Chillingworth T, Davies RM, Feltwell T, Holroyd S, Jagels K, Karlyshev AV, Moule S, Pallen MJ, Penn CW, Quail MA, Rajandream MA, Rutherford KM, van Vliet AH, Whitehead S, Barrell BG. 2000. The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature* 403:665–668. <http://dx.doi.org/10.1038/35001088>.
29. Pearson BM, Gaskin DJH, Segers RPAM, Wells JM, Nuijten PJM, van Vliet AHM. 2007. The complete genome sequence of *Campylobacter jejuni* strain 81116 (NCTC11828). *J. Bacteriol.* 189:8402–8403. <http://dx.doi.org/10.1128/JB.01404-07>.
30. Hofreuter D, Tsai J, Watson RO, Novik V, Altman B, Benitez M, Clark C, Perbost C, Jarvie T, Du L, Galan JE. 2006. Unique features of a highly pathogenic *Campylobacter jejuni* strain. *Infect. Immun.* 74:4694–4707. <http://dx.doi.org/10.1128/IAI.00210-06>.
31. Fouts DE, Mongodin EF, Mandrell RE, Miller WG, Rasko DA, Ravel J, Brinkac LM, DeBoy RT, Parker CT, Daugherty SC, Dodson RJ, Durkin AS, Madupu R, Sullivan SA, Shetty JU, Ayodeji MA, Shvartsbeyn A, Schatz MC, Badger JH, Fraser CM, Nelson KE. 2005. Major structural differences and novel potential virulence mechanisms from the genomes of multiple *Campylobacter* species. *PLoS Biol.* 3:e15. <http://dx.doi.org/10.1371/journal.pbio.0030015>.
32. Pearson BM, Rokney A, Crossman LC, Miller WG, Wain J, van Vliet AHM. 2013. Complete genome sequence of the *Campylobacter coli* clinical isolate 15-537360. *Genome Announc.* 1:e01056-13. <http://dx.doi.org/10.1128/genomeA.01056-13>.
33. Brown HL, van Vliet AHM, Betts RP, Reuter M. 2013. Tetrazolium reduction allows assessment of biofilm formation by *Campylobacter jejuni* in a food matrix model. *J. Appl. Microbiol.* 115:1212–1221. <http://dx.doi.org/10.1111/jam.12316>.
34. Lemon KP, Higgins DE, Kolter R. 2007. Flagellar motility is critical for *Listeria monocytogenes* biofilm formation. *J. Bacteriol.* 189:4418–4424. <http://dx.doi.org/10.1128/JB.01967-06>.
35. O'Toole GA, Kolter R. 1998. Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development. *Mol. Microbiol.* 30:295–304. <http://dx.doi.org/10.1046/j.1365-2958.1998.01062.x>.
36. Kalmokoff M, Lanthier P, Tremblay TL, Foss M, Lau PC, Sanders G, Austin J, Kelly J, Szymanski CM. 2006. Proteomic analysis of *Campylobacter jejuni* 11168 biofilms reveals a role for the motility complex in biofilm formation. *J. Bacteriol.* 188:4312–4320. <http://dx.doi.org/10.1128/JB.01975-05>.
37. Agle ME. 2007. Biofilms in the food industry, p 3–18. *Biofilms in the food environment*. Blackwell Publishing, Ltd., London, United Kingdom.
38. Teh AH, Lee SM, Dykes GA. 13 June 2014. Do *Campylobacter jejuni* form biofilms in food-related environments? *Appl. Environ. Microbiol.* <http://dx.doi.org/10.1128/AEM.01493-14>.
39. Lindsay D, Geornaras I, von Holy A. 1996. Biofilms associated with poultry processing equipment. *Microbios* 86:105–116.
40. Blackman IC, Frank JF. 1996. Growth of *Listeria monocytogenes* as a biofilm on various food-processing surfaces. *J. Food Prot.* 59:827–831.
41. Habimana O, Heir E, Langsrud S, Aringsli AW, Møretro T. 2010. Enhanced surface colonization by *Escherichia coli* O157:H7 in biofilms formed by an *Acinetobacter calcoaceticus* isolate from meat-processing environments. *Appl. Environ. Microbiol.* 76:4557–4559. <http://dx.doi.org/10.1128/AEM.02707-09>.
42. Kusumaningrum HD, Riboldi G, Hazeleger WC, Beumer RR. 2003. Survival of foodborne pathogens on stainless steel surfaces and cross-contamination to foods. *Int. J. Food Microbiol.* 85:227–236. [http://dx.doi.org/10.1016/S0168-1605\(02\)00540-8](http://dx.doi.org/10.1016/S0168-1605(02)00540-8).
43. Xu C-P, Boks NP, de Vries J, Kaper HJ, Norde W, Busscher HJ, van der Mei HC. 2008. *Staphylococcus aureus*-fibronectin interactions with and without fibronectin-binding proteins and their role in adhesion and desorption. *Appl. Environ. Microbiol.* 74:7522–7528. <http://dx.doi.org/10.1128/AEM.00948-08>.
44. Healy B, Cooney S, O'Brien S, Iversen C, Whyte P, Nally J, Callanan JJ, Fanning S. 2010. *Cronobacter (Enterobacter sakazakii)*: an opportunistic foodborne pathogen. *Foodborne Pathog. Dis.* 7:339–350. <http://dx.doi.org/10.1089/fpd.2009.0379>.
45. Singh AV, Vyas V, Patil R, Sharma V, Scopelliti PE, Bongiorno G, Podesta A, Lenardi C, Gade WN, Milani P. 2011. Quantitative characterization of the influence of the nanoscale morphology of nanostructured surfaces on bacterial adhesion and biofilm formation. *PLoS One* 6:e25029. <http://dx.doi.org/10.1371/journal.pone.0025029>.
46. Reuter M, van Vliet AHM. 2013. Signal balancing by the CetABC and CetZ chemoreceptors controls energy taxis in *Campylobacter jejuni*. *PLoS One* 8:e54390. <http://dx.doi.org/10.1371/journal.pone.0054390>.
47. Zhao C, Ge B, De Villena J, Sudler R, Yeh E, Zhao S, White DG, Wagner D, Meng J. 2001. Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* serovars in retail chicken, turkey, pork, and beef from the greater Washington, DC, area. *Appl. Environ. Microbiol.* 67:5431–5436. <http://dx.doi.org/10.1128/AEM.67.12.5431-5436.2001>.
48. Whyte P, McGill K, Cowley D, Madden RH, Moran L, Scates P, Carroll C, O'Leary A, Fanning S, Collins JD, McNamara E, Moore JE, Cormican M. 2004. Occurrence of *Campylobacter* in retail foods in Ireland. *Int. J. Food Microbiol.* 95:111–118. <http://dx.doi.org/10.1016/j.ijfoodmicro.2003.10.018>.