

# The Indoor Fungus *Cladosporium halotolerans* Survives Humidity Dynamics Markedly Better than *Aspergillus niger* and *Penicillium rubens* despite Less Growth at Lowered Steady-State Water Activity

Frank J. J. Segers,<sup>a</sup> Karel A. van Laarhoven,<sup>b</sup> Hendrik P. Huinink,<sup>b</sup> Olaf C. G. Adan,<sup>b</sup> Han A. B. Wösten,<sup>c</sup> Jan Dijksterhuis<sup>a</sup>

CBS-KNAW Fungal Biodiversity Centre, Applied and Industrial Mycology, Utrecht, The Netherlands<sup>a</sup>; Eindhoven University of Technology, Department of Applied Physics, Eindhoven, The Netherlands<sup>b</sup>; Utrecht University, Microbiology, Department of Biology, Utrecht, The Netherlands<sup>c</sup>

## ABSTRACT

Indoor fungi cause damage in houses and are a potential threat to human health. Indoor fungal growth requires water, for which the terms water activity ( $a_w$ ) and relative humidity (RH) are used. The ability of the fungi *Aspergillus niger*, *Cladosporium halotolerans*, and *Penicillium rubens* at different developmental stages to survive changes in  $a_w$  dynamics was studied. Fungi grown on media with high  $a_w$  were transferred to a controlled environment with low RH and incubated for 1 week. Growth of all developmental stages was halted during incubation at RHs below 75%, while growth continued at 84% RH. Swollen conidia, germ-lings, and microcolonies of *A. niger* and *P. rubens* could not reinitiate growth when retransferred from an RH below 75% to a medium with high  $a_w$ . All developmental stages of *C. halotolerans* showed growth after retransfer from 75% RH. Dormant conidia survived retransfer to medium with high  $a_w$  in all cases. In addition, retransfer from 84% RH to medium with high  $a_w$  resulted in burst hyphal tips for *Aspergillus* and *Penicillium*. Cell damage of hyphae of these fungi after incubation at 75% RH was already visible after 2 h, as observed by staining with the fluorescent dye TOTO-1. Thus, *C. halotolerans* is more resistant to  $a_w$  dynamics than *A. niger* and *P. rubens*, despite its limited growth compared to that of these fungi at a lowered steady-state  $a_w$ . The survival strategy of this phylloplane fungus in response to the dynamics of  $a_w$  is discussed in relation to its morphology as studied by cryo-scanning electron microscopy (cryo-SEM).

## IMPORTANCE

Indoor fungi cause structural and cosmetic damage in houses and are a potential threat to human health. Growth depends on water, which is available only at certain periods of the day (e.g., during cooking or showering). Knowing why fungi can or cannot survive indoors is important for finding novel ways of prevention. Until now, the ability of fungi to grow on media with little available water at steady state (unchanging conditions) has been important for evaluating whether a fungus can grow indoors. In the present study, we found that the fungus *Cladosporium halotolerans*, a common indoor fungus, is more resistant to changes in available water than the fungi *Aspergillus niger* and *Penicillium rubens*, despite the fact that the latter fungi can grow on media with low water availability. We concluded that the ability of fungi to deal with changes in humidity is at least as important as the ability to grow on low-water media.

People in Europe spend only 1.6 h a day outdoors (1), emphasizing the need for a healthy indoor environment (2). Indoor fungal growth represents a global problem. It has been estimated that about 25% of social housing in the European Union shows fungal growth. This not only causes disfigurement of the building materials but also poses a health threat (3, 4). Asthmatic and allergic patients are particularly at risk due to activation of the immune system by airborne fungal structures that are released from surface-grown fungal colonies (5–9). Mycotoxins produced by indoor fungi may also contribute to the health risk, but it is still unknown to what extent these toxic compounds are produced and released into the indoor environment (10, 11).

Highly diverse fungal species are present in the outdoor air, in particular species of *Aspergillus*, *Penicillium*, and *Cladosporium* (12). Their abundance in temperate climates is influenced by the seasons. For instance, *Cladosporium* is particularly prevalent during fall (13, 14). The abundance of fungal species in the indoor environment is affected by their predominance in the outdoor environment and by the indoor conditions (13–17). *Penicillium chrysogenum* and *Aspergillus versicolor* are particularly abundant after water damage or direct moistening, while *Cladosporium* oc-

curs predominantly in indoor environments without water-related incidents (12, 18–22).

A lot of studies have been done on fungal growth in relation to water activity ( $a_w$ ) (23–25) and relative humidity (RH) (26, 27). Both  $a_w$  and RH express water's chemical potential, and therefore the availability of water. In this study, we use both terms, with the following relationship in equilibrium:  $RH = a_w \times 100\%$ .

Earlier studies which were done in an attempt to predict indoor

Received 17 February 2016 Accepted 6 May 2016

Accepted manuscript posted online 17 June 2016

Citation Segers FJJ, van Laarhoven KA, Huinink HP, Adan OCG, Wösten HAB, Dijksterhuis J. 2016. The indoor fungus *Cladosporium halotolerans* survives humidity dynamics markedly better than *Aspergillus niger* and *Penicillium rubens* despite less growth at lowered steady-state water activity. *Appl Environ Microbiol* 82:5089–5098. doi:10.1128/AEM.00510-16.

Editor: A. A. Brakhage, HKI and University of Jena

Address correspondence to Jan Dijksterhuis, j.dijksterhuis@cbs.knaw.nl.

Copyright © 2016 Segers et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

**TABLE 1** Developmental stages exposed to changes in  $a_w$ 

Developmental stage no. (description)	Species	Culture time (h) <sup>a</sup>
I (dormant conidia)	<i>A. niger</i> , <i>C. halotolerans</i> , and <i>P. rubens</i>	0
II (swollen conidia)	<i>A. niger</i> , <i>C. halotolerans</i> , and <i>P. rubens</i>	7–8
III (germlings)	<i>A. niger</i> and <i>P. rubens</i>	18
	<i>C. halotolerans</i>	24
IV (microcolonies without aerial hyphae)	<i>A. niger</i> and <i>P. rubens</i>	24
	<i>C. halotolerans</i>	32–36
V (microcolonies with aerial hyphae)	<i>A. niger</i> and <i>P. rubens</i>	32–36
	<i>C. halotolerans</i>	44–48
VI (sporulating microcolonies)	<i>A. niger</i> and <i>P. rubens</i>	48–50
	<i>C. halotolerans</i>	48–60

<sup>a</sup> Time needed to reach the indicated developmental stage on DG18 agar.

fungal growth focused on the effects of RH and fungal growth on building materials, such as gypsum (28–31), wood (32–35), and concrete (36). Most fungi show growth in the range of 90 to 100% RH (12), but a restricted subset of fungi, such as *Cladosporium sphaerospermum*, *P. chrysogenum*, and *A. versicolor*, also grow on building materials, such as plywood, pine sapwood, and gypsum board, at a static RH as low as 75 to 80% (33). However, dynamic humidity of around 80% RH resulted in less-than-expected growth in the case of some materials, such as gypsum board (33, 37). This value of 80% RH is stated as the lower limit for indoor fungal growth. Indoor RH is dynamically averaged to be around 50%, thus raising the question of how fungal cells respond to periods of lower RH (38). Survival of fully developed colonies of *Penicillium rubens* in response to  $a_w$  dynamics was reported by Bekker (29). The effect of dynamic RH on the fungi *Penicillium brevicompactum*, *Alternaria tenuissima*, and *Trichoderma harzia-*


*num* in wood samples was studied by Li and Wadsö (35). They measured more heat release due to increased metabolic activity of fungal cells following an increase of RH by using an isothermal calorimeter. The impact of dynamic water availability on fungal growth was also studied in a model system of moistened building material that was dried and moistened again (22). The fungal growth seen in that study was due to the moisture content in the building material. The relevance of moisture content was shown in a study done by van Laarhoven et al. (31). Gypsum tablets equilibrated at a certain RH or soaked in a glycerol solution with the corresponding  $a_w$  were inoculated with *P. rubens*. The soaked tablets that had a higher moisture content showed markedly faster elongation of hyphae. However, there is a scarcity of knowledge on the effects of humidity changes on the cellular level.

The purpose of this study was to compare the responses of indoor fungi, namely, *Aspergillus niger*, *Cladosporium halotolerans*, and *P. rubens*, at different developmental stages to steady-state and dynamic water activity. *C. halotolerans* had markedly better survival at both increasing and decreasing  $a_w$  than that of *P. rubens* and *A. niger*. This difference in survival at dynamic  $a_w$  was despite the growth limit of *C. halotolerans* at a higher steady-state  $a_w$  than that for *P. rubens* and *A. niger*.

## MATERIALS AND METHODS

**Organisms and growth conditions.** *A. niger* N402 (39), *C. halotolerans* CBS 139586 (40), and *P. rubens* CBS 401.92 (28) were used in this study. The *P. rubens* strain was first identified as *Penicillium chrysogenum* but was later reclassified (41). Strains were routinely grown at 25°C on dichloran-18% glycerol agar (DG18 agar; Oxoid) ( $a_w = 0.96$ ) (42, 43).

**Growth at steady-state  $a_w$ .** Measurements of the growth of fungi at lower steady-state  $a_w$  were done as described by Segers et al. (40). Fungi were grown on malt extract agar (MEA) complemented with 0 to 50% glycerol ( $a_w$  of 0.99 to 0.75) to assess the lower limits of growth with respect to  $a_w$ . The  $a_w$  values for the glycerol-agar mixtures were deter-



**FIG 1** Schematic overview of the developmental stages of the indoor fungi used in this study, i.e., dormant conidia (I), swollen conidia (II), germlings (III), and mycelia without aerial hyphae (IV), with aerial hyphae (V), and with conidium-forming conidiophores (VI).




FIG 2 Flowchart of the experimental setup to study survival of indoor fungi after periods of lowered relative humidity.

mined before and after growth experiments by using a Novasina labmaster- $a_w$  instrument (Novasina, Lachen, Switzerland) as also described by Segers et al. (40). Cultures were inoculated with  $3 \mu\text{l}$  of a spore solution containing  $1 \times 10^6$  conidia  $\text{ml}^{-1}$ . These conidia were harvested from a 7-day-old culture by use of a T spatula (VWR, Amsterdam, The Nether-




FIG 3 Growth (change in colony diameter) of *A. niger* (green triangles), *C. halotolerans* (gray squares) (data derived from the work of Segers et al. [40]), and *P. rubens* (blue diamonds) on MEA with 0 to 50% glycerol.

TABLE 2 Survival of developmental stages of *A. niger*, *C. halotolerans*, and *P. rubens* after 1 week of incubation at 33, 58, 75, or 84% RH and subsequent 1 to 2 days of incubation on DG18 agar

RH (%) during 1 week of incubation	Developmental stage <sup>a</sup>	Survival <sup>b</sup>		
		<i>A. niger</i>	<i>C. halotolerans</i>	<i>P. rubens</i>
33	I	++	++	++
	II	-	-	-
	III	-	-	-
	IV	-	-	-
	V	-	-	-
	VI	+++	+++	+++
58	I	++++	++++	++++
	II	-	-	-
	III	-	-	-
	IV	-	-	-
	V	-	+	-
	VI	+++	+++	+++
75	I	++++	++++	++++
	II	-	++++	-
	III	-	++++	-
	IV	-	++++	-
	V	+	++++	-
	VI	++++	++++	++++
84	I	++++	++++	++++
	II	++++	++++	+++
	III	+++	++++	+++
	IV	+++	++++	+++
	V	++++	++++	++++
	VI	++++	++++	++++


<sup>a</sup> The developmental stages comprised dormant conidia (I), swollen conidia (II), germlings (III), and microcolonies without aerial hyphae (IV), with aerial hyphae (V), and with conidium-forming conidiophores (VI).

<sup>b</sup> Survival was qualified as no survival (-), <5% survival (+), <50% survival (++), >50% survival (+++), and >95% survival (++++).

lands), using ice-cold sterile 10 mM *N*-(2-acetamido)-2-aminoethanesulfonic acid, 0.02% Tween 80 (ACES; pH 6.8). The colony diameter was measured 3 times a week for 3 weeks. The growth speed was determined from the regression coefficients based on graphs of each colony.

**Survival during periods of dynamic  $a_w$ .** To study survival of fungal cells during changes in  $a_w$ , DG18 plates were inoculated with  $50 \mu\text{l}$  of spore solution ( $1 \times 10^6$  conidia  $\text{ml}^{-1}$ ). The inoculum was homogeneously spread on the agar surface by use of a T spatula and left to dry. A polycarbonate (PC) filter (47-mm diameter, 0.1- $\mu\text{m}$  pore size; GE Water & Process Technologies) was placed on the surface of the inoculated agar to pick up the spores. Filters were immediately transferred to a new DG18 plate, with the side containing the spores oriented upwards. This "stamping" method resulted in 100 to 500 dispersed conidia per filter. The filters containing conidia were incubated in the dark on DG18 agar for 0 to 50 h, until the developmental stage of interest (Table 1; Fig. 1), as confirmed by light microscopy. Filters were removed from the agar plate and transferred to glass desiccators (6 liters) containing 150 to 300 ml saturated potassium chloride (RH,  $84.3\% \pm 0.2\%$ ), sodium chloride (RH,  $75.3\% \pm 0.1\%$ ), sodium bromide (RH,  $57.6\% \pm 0.4\%$ ), or magnesium chloride (RH,  $32.8\% \pm 0.2\%$ ) to control relative humidity (44). The filters were transferred to and from the desiccators within a humid environment (RH of >85%) to prevent exposure to low RH during transfer. The RH was quantified using a Testo 174H hygrometer (Testo, Lenzkirch, Germany). After 1 week of incubation within the desiccator, filters were transferred to DG18 agar plates, and incubation was prolonged at an  $a_w$  of 0.96 for up to






**FIG 4** Branching hyphae of *P. rubens* (stage IV) (A, B, E, and F) and *C. halotolerans* (C, D, G, and H) immediately after incubation for 1 week at 75% RH (A and C) or 84% RH (E and G) and after subsequent incubation on DG18 agar for 1 day (B, D, F, and H). Bar = 100  $\mu$ m.

1 month. Survival of fungal developmental structures, growth, and bursting of hyphal tips were evaluated and documented using a stereomicroscope (Zoom AZ-100; Nikon, Amsterdam, The Netherlands) within 30 min after transfer and after 1 and 2 days of incubation. The fungi were considered to have survived if hyphae reinitiated growth or the conidia germinated. These conidia could be originating from stage I and stage II or be newly formed at stage VI. A flowchart of this experimental setup is

depicted in Fig. 2. Experiments were performed using at least two technical replicates and three biological replicates.

**Fluorescence microscopy.** A Zeiss Axioplan II microscope, a Zeiss Plan NeoFluar 40 $\times$ /0.75 objective, and a blue BP450-490 (FT510, LP520) excitation filter were used for fluorescence microscopy. Images were captured with a Zeiss AxioCam MRc digital camera run by Zeiss Axiovision 4. A 0.1 mM solution of the fluorescent dye quinolinium, 1-1'-[1,3-pro-





**FIG 5** TOTO-1 staining, an indication of dead fungal structures, of *A. niger* (A, D, G, and J), *C. halotolerans* (B, E, H, and K), and *P. rubens* (C, F, I, and L) after treatment with steam for 20 min (A to C) or after exposure to 70% alcohol (D to F), to 75% RH for 45 min (G to I), or to 75% RH for 105 min (J to L). Bar = 20  $\mu\text{m}$ .


panediybis[(dimethyliminio)-3,1-propanediy]]bis[4-[(3-methyl-2(3H)-benzothiazolydene)methyl]-, tetraiodide (TOTO-1; Molecular Probes, Breda, The Netherlands) in dimethyl sulfoxide (DMSO) was used as a stain of dead colonies. Colonies without aerial hyphae, grown on top of PC filters on DG18 agar (see above), were placed in a desiccator containing a saturated NaCl solution (75% RH) for 45 or 105 min. Filters were placed back on DG18 agar and stained for 25 min by using 2  $\mu\text{M}$  TOTO-1 in ACES buffer (45, 46). A 1-cm<sup>2</sup> square of the filter and the underlying DG18 agar was placed on top of an objective glass and covered with a coverslip. Colonies grown on a PC filter and killed with 70% alcohol or a 20-min steam treatment (47) served as a control.

**Cryo-SEM.** Small rectangular 5- by 5-mm blocks of agar medium topped with a filter were transferred to copper cups for snap-freezing in nitrogen slush. They were glued to the copper surface with frozen tissue medium (KP-Cryoblock; Klinipath, Duiven, The Netherlands) and sputter coated 3 times for 1 min each by using a gold target. Cryo-scanning electron microscopy (cryo-SEM) was done with a JEOL 5600LV scanning

electron microscope (JEOL, Tokyo, Japan) equipped with an Oxford CT1500 Cryostation. Electron micrographs were taken at an acceleration voltage of 5 kV.

## RESULTS

**Minimal and optimal water activities for fungal growth.** The minimal and optimal  $a_w$  values for growth of *A. niger*, *C. halotolerans*, and *P. rubens* were determined. Colony diameters of *A. niger*, *C. halotolerans*, and *P. rubens* colonies were monitored 3 times a week during a 3-week period on MEA plates with  $a_w$  values of 0.75 to 0.99 (Fig. 3). *A. niger* was the fastest-growing species, with an optimal growth speed of 14.4 mm day<sup>-1</sup>, while the optimal growth speed of *P. rubens* was 7.9 mm day<sup>-1</sup>. *C. halotolerans* has a growth speed of 4.1 mm day<sup>-1</sup> as derived from the work of Segers et al. (40). These optimal growth speeds were observed at  $a_w$  values of 0.96 to 0.98. The minimal  $a_w$  values needed to support



**FIG 6** (A and B) Leading hyphae of *P. rubens* growing at 84% RH had burst after transfer to DG18 agar ( $a_w = 0.96$ ). (B) New hyphae formed that originated from the center of the colony (white arrows) when the colony was left to grow for another 24 h after being transferred to DG18 agar. Bars = 100  $\mu\text{m}$  (A) and 200  $\mu\text{m}$  (B).

growth of *A. niger*, *C. halotolerans*, and *P. rubens* were 0.80, 0.82 (40), and 0.82, respectively.

**Survival during periods of dynamic  $a_w$ .** *A. niger*, *C. halotolerans*, and *P. rubens* were tested for the ability to survive during periods of defined lowered  $a_w$ , which were obtained using closed environments containing a saturated salt solution. Filters overlying DG18 agar medium were inoculated with freshly harvested conidia (stage I) and cultured at an  $a_w$  of 0.96 until conidia were swollen (stage II), conidia formed germ tubes (stage III), or microcolonies formed without aerial hyphae (stage IV), with aerial hyphae (stage V), or with conidium-forming conidiophores (stage VI) (Table 1; Fig. 1). The filters with these developmental stages were removed from agar and transferred to 33%, 58%, 75%, and 84% RH. No growth was observed during incubation in the desiccator at RHs of 33 to 75%, while growth continued in the desiccator at RHs above 84% for all developmental stages of the 3 species.

After 1 week of incubation, survival was assessed by removing the filters from the desiccators and placing them on DG18 agar with an  $a_w$  of 0.96. The response of the fungi was monitored for up to 1 month. Incubation for 2 weeks to up to 1 month resulted in no more survival as determined 2 days after removal from the desiccator. Developmental stages II to V did not reinitiate growth at an  $a_w$  of 0.96 after a 1-week incubation at 33% RH. In contrast, about 50% of the conidia (stage I) of the 3 species germinated after transfer to medium with an  $a_w$  of 0.96. Similarly, germination of newly formed conidia was observed in the case of conidium-producing colonies (stage VI). Similar results were obtained after incubation at 58% RH. However, in this case, more conidia from stages I and VI germinated than those after incubation at 33% RH. Notably, stage V colonies of *C. halotolerans* also showed some growth at an  $a_w$  of 0.96 after exposure to 58% RH. All developmental stages of *C. halotolerans* incubated for 1 week at 75% RH showed survival after transfer of the filters to medium with an  $a_w$  of 0.96. Incubation of *A. niger* and *P. rubens* at 75% RH gave results similar to those obtained after incubation at 33% and 58% RH, except that more conidia of stages I and VI germinated. Thus, *A. niger* and *P. rubens* hyphae were unable to reinitiate growth after incubation at 75% RH (Table 2). Figure 4 shows examples of

*C. halotolerans* and *P. rubens* after 1 week at 75% or 84% RH. *P. rubens* (stage IV) did not survive for 1 week at 75% RH (Fig. 4A and B), but in contrast, *C. halotolerans* did (Fig. 4C and D). Both fungi continued to grow during the period of 1 week at 84% RH, and both formed conidiophores (Fig. 4E and G). They showed strongly increased growth after 1 day of being rehydrated on DG18 agar (Fig. 4F and H).

Staining with TOTO-1 showed that some and most hyphae of *A. niger* and *P. rubens* were killed after 45 min (Fig. 5G and I) and 105 min (Fig. 5J and L), respectively, of exposure to 75% RH. In contrast, *C. halotolerans* showed no or little fluorescence after these exposures (Fig. 5H and K), indicating that most hyphae survived the treatment.

As mentioned above, growth of the colonies of the 3 species continued, albeit at a lower rate, during incubation at 84% RH and subsequent transfer to DG18 agar. However, 14.7% and 7.5% of all hyphal tips of microcolonies, including small branching hyphae, burst directly after the transfer for *A. niger* and *P. rubens*, respectively (Fig. 6). In contrast, hyphal tips of *C. halotolerans* hardly (0.2%) showed rupture. Most bursting hyphae were found at the periphery of the colony (Fig. 6A), while hyphae at the center of the microcolony resumed growth, as indicated with arrows (Fig. 6B).


## DISCUSSION

Relative humidity (RH) can change considerably during and between days in indoor situations. As a result, indoor fungi not only have to be able to survive periods of low RH but also have to be able to resume growth within the time frame of favorable RH. In this study, we compared the capacities of *A. niger*, *C. halotolerans*, and *P. rubens* to grow during static or dynamic water activity ( $a_w$ ) regimens.

*A. niger* grew at an  $a_w$  of 0.80 at 25°C, which was lower than the minimal  $a_w$  of 0.82 for *C. halotolerans* and *P. rubens*. The latter data confirm earlier findings (23, 40, 48, 49), although *A. niger* has been reported to grow even at an  $a_w$  of 0.78. The latter value was obtained at the optimal growth temperature of 30°C (23).

A controlled decrease in  $a_w$  was used to address the responses of *A. niger*, *C. halotolerans*, and *P. rubens* to dynamic  $a_w$  conditions.





**FIG 7** Cryo-SEM images of *C. halotolerans*, showing a 1-day-old colony growing on DG18 agar on top of a 0.1- $\mu\text{m}$ -pore-size polycarbonate filter. The microcolony (A) can be seen with strengthened cells in the center (B, D, and E), while bundles of hyphae stretch out as leading hyphae (B and C). Bars = 100  $\mu\text{m}$  (A), 10  $\mu\text{m}$  (B to D), and 5  $\mu\text{m}$  (E).



To this end, conidia were inoculated onto filters overlying DG18 medium ( $a_w = 0.96$ ) and cultured until conidia had swollen or germinated or until microcolonies had formed without aerial hyphae, with aerial hyphae, or with conidium-forming conidiophores. The filters were then transferred to 84%, 75%, 58%, and 33% RH ( $a_w = 0.84, 0.75, 0.58, \text{ and } 0.33$ ). Conidia germinated at 84% RH but not at lower RHs, as could be expected according to the data presented above. Yet some or even all spores that had been exposed to the lower-RH conditions germinated when they were retransferred to DG18 agar after 1 week. The percentage of conidia that germinated increased with increasing RHs used during the 1-week incubation. This is interesting because conidia of *Aspergillus fumigatus* that had been dried slowly at room temperature survived for a year, with full germination (50). In our study, some of the conidia of the different fungal species lost their viability after a relatively quick shift from 96% to 33% RH and a week-long incubation, which indicates that the speed of change in RH is an important factor in survival even for survival structures, such as conidia. Alternatively, the extent of drying may be a factor that improves survival of spores. Ascospores of *Talaromyces macrosporus* and *Neosartorya fischeri* survive better when stringently dried (down to 0% RH) than when dried in ambient air (which is typically 40 to 60% RH as measured with a hygrometer) (51). This can be explained by the large amounts of compatible solutes that result in a glassy state at water contents below 2 to 3%. The mobility of molecules is very low in this state, thus suppressing the incidence of detrimental chemical reactions (e.g., with reactive oxygen species) (52). Environmental conditions during conidium formation are also important for survival of asexual spores. In preliminary experiments, we used MEA instead of DG18 agar, and this indicated less survival after a 1-week incubation period at 75% RH for *C. halotolerans*. It may well be that colonies grown on DG18 agar had adapted to the lower  $a_w$  of this medium and transferred this adaptation to the spores (53). It may also be that the glycerol in DG18 medium is taken up and serves as a compatible solute that helps the fungus and its conidia to survive during periods of low RH (24, 51).

This study showed that the capability to grow at low  $a_w$  does not always reflect survival during changes in RH. All developmental stages of *C. halotolerans* were able to survive for 1 week at 75% RH. This was not the case for *A. niger*, despite its ability to grow at a lower steady-state  $a_w$ . Notably, increasing water availability also affected survival. *C. halotolerans* hyphae had a much lower bursting incidence after transfer from 85% RH to medium with an  $a_w$  of 0.96 than those of *A. niger* and *P. rubens*. This was probably due to the suddenly increased  $a_w$  causing water influx into the hyphae. This may be due to differences in accumulation of compatible solutes between the species. Alternatively, the structure of the cell wall at the hyphal tips may differ. A more rigid cell wall at tips of *C. halotolerans* hyphae would more easily overcome a transfer from low to high water availability. At the same time, such a cell wall would restrict growth speed (54), which agrees with the lower growth speed of this species than those of *A. niger* and *P. rubens*. The rigidity of the *C. halotolerans* cell wall may be due to the presence of melanin. Nonlinear spectral imaging (NLSI) microscopy indicates that the cell wall of vegetative hyphae of *C. halotolerans* contains melanin, while melanin is absent in *P. rubens* (our unpublished results). Formation of enlarged cells with strengthened pigmented cell walls in the center of the colony (55), as well

as formation of bundles of hyphae, may also help cells to overcome sudden changes in humidity (Fig. 7).

The ability of *C. halotolerans* to cope with dynamic water availability is probably related to the ecological niche of this fungus. *Cladosporium* species grow on leaves and are therefore called phylloplane fungi (56, 57). The available water for fungi growing on leaves is highly dynamic and is influenced by changing temperature, dew formation, sunlight, and rain. It is interesting that the indoor environment is also characterized by changes in humidity during the day. It has been shown that phylloplane fungi can restore growth after minutes to hours of rehydration after drying for 2 to 3 weeks (56). Furthermore, these fungi have to withstand large amounts of UV radiation from direct sunlight. Damage from UV radiation is prevented by melanin within the cell walls of conidia and hyphae (58–60). The protection against UV radiation is of less concern in indoor environments, which are generally darker aside from the small amount of UV passing through open windows. However, melanin can help in protection against reactive oxygen species or other reactive molecules present in building materials.

This study shows for the first time, to our knowledge, that steady-state  $a_w$  measurements, for so long the hallmark to determine the xerotolerance of a fungus, are not enough to predict the response of a fungus to humidity dynamics. The indoor environment is characterized by periods of lowered  $a_w$  as well as sudden increases in water availability, and fungal cells have to deal with both types of change. It has to be expected that the adaptability of a fungal species to deal with humidity dynamics is important for the predominance of the species on the relevant building material. Fungi may even be able to influence the (micro)environment by using enzymes or other excretions. For example, germinating conidia of *P. rubens* are known to produce an extracellular matrix on gypsum (61).

## ACKNOWLEDGMENTS

This research was supported by the Dutch Technology Foundation STW, which is part of the Netherlands Organization for Scientific Research (NWO) and is partly funded by the Ministry of Economic Affairs.

We thank Joey van den Ende for technical assistance and Bart Theelen and Sylvia Klaubauf for critical readings of the manuscript. Elke van Nieuwenhuijzen is thanked for fruitful discussions.

## FUNDING INFORMATION

This work, including the efforts of Frank J. J. Segers, Karel A. van Laarhoven, Hendrik P. Huinink, Olaf C. G. Adan, Han A. B. Wösten, and Jan Dijksterhuis, was funded by Dutch Technology Foundation STW (11117).

## REFERENCES

- Schweizer C, Edwards RD, Bayer-Oglesby L, Gauderman WJ, Ilacqua V, Juhani Jantunen M, Lai HK, Nieuwenhuijzen M, Kunzli N. 2006. Indoor time-microenvironment-activity patterns in seven regions of Europe. *J Expos Sci Environ Epidemiol* 17:170–181.
- Adan OCG, Ng-A-Tham J, Hanke W, Sigsgaard T, van den Hazel P, Wu F. 2007. In search of a common European approach to a healthy indoor environment. *Environ Health Perspect* 115:983–988. <http://dx.doi.org/10.1289/ehp.8991>.
- Bonnefoy X. 2007. Inadequate housing and health: an overview. *Int J Environ Pollut* 30:411–429. <http://dx.doi.org/10.1504/IJEP.2007.014819>.
- Bonnefoy XR, Braubach M, Moissonnier B, Monollbaev K, Robbel N. 2003. Housing and health in Europe: preliminary results of a pan-European study. *Am J Public Health* 93:1559–1563. <http://dx.doi.org/10.2105/AJPH.93.9.1559>.
- Denning DW, O'Driscoll BR, Hogaboam CM, Bowyer P, Niven RM. 2006. The link between fungi and severe asthma: a summary of the evi-



- dence. *Eur Respir J* 27:615–626. <http://dx.doi.org/10.1183/09031936.06.00074705>.
6. Reboux G, Bellanger AP, Roussel S, Grenouillet F, Millon L. 2010. Moulds in dwellings: health risks and involved species. *Rev Fr D Allergol* 50:611–620. <http://dx.doi.org/10.1016/j.reval.2010.03.009>.
  7. Miller JD, Sun M, Gilyan A, Roy J, Rand TG. 2010. Inflammation-associated gene transcription and expression in mouse lungs induced by low molecular weight compounds from fungi from the built environment. *Chem Biol Interact* 183:113–124. <http://dx.doi.org/10.1016/j.cbi.2009.09.023>.
  8. Rand TG, DiPenta J, Robbins C, Miller JD. 2011. Effects of low molecular weight fungal compounds on inflammatory gene transcription and expression in mouse alveolar macrophages. *Chem Biol Interact* 190:139–147. <http://dx.doi.org/10.1016/j.cbi.2011.02.017>.
  9. Green BJ, Schmechel D, Summerbell RC. 2011. Aerosolized fungal fragments, p 211–243. In Adan OCG, Samson RA (ed), *Fundamentals of mold growth in indoor environments and strategies for healthy living*, vol 1. Wageningen Academic Publishers, Wageningen, The Netherlands.
  10. Miller JD, McMullin DR. 2014. Fungal secondary metabolites as harmful indoor air contaminants: 10 years on. *Appl Microbiol Biotechnol* 98:9953–9966. <http://dx.doi.org/10.1007/s00253-014-6178-5>.
  11. Miller JD. 2011. Health effects from mold and dampness in housing in Western societies: early epidemiology studies and barriers to further progress, p 183–210. In Adan OCG, Samson RA (ed), *Fundamentals of mold growth in indoor environments and strategies for healthy living*, vol 1. Wageningen Academic Publishers, Wageningen, The Netherlands.
  12. Nevalainen A, Taubel M, Hyvarinen A. 2015. Indoor fungi: companions and contaminants. *Indoor Air* 25:125–156. <http://dx.doi.org/10.1111/ina.12182>.
  13. de Ana SG, Torres-Rodriguez JM, Ramirez EA, Garcia SM, Belmonte-Soler J. 2006. Seasonal distribution of *Alternaria*, *Aspergillus*, *Cladosporium* and *Penicillium* species isolated in homes of fungal allergic patients. *J Investig Allergol Clin Immunol* 16:357–363.
  14. Crawford JA, Rosenbaum PF, Anagnost SE, Hunt A, Abraham JL. 2015. Indicators of airborne fungal concentrations in urban homes: understanding the conditions that affect indoor fungal exposures. *Sci Total Environ* 517:113–124. <http://dx.doi.org/10.1016/j.scitotenv.2015.02.060>.
  15. Adams RI, Miletto M, Taylor JW, Bruns TD. 2013. Dispersal in microbes: fungi in indoor air are dominated by outdoor air and show dispersal limitation at short distances. *ISME J* 7:1262–1273. <http://dx.doi.org/10.1038/ismej.2013.28>.
  16. Horner WE, Worthan AG, Morey PR. 2004. Air- and dustborne mycoflora in houses free of water damage and fungal growth. *Appl Environ Microbiol* 70:6394–6400. <http://dx.doi.org/10.1128/AEM.70.11.6394-6400.2004>.
  17. McGinnis MR. 2007. Indoor mold development and dispersal. *Med Mycol* 45:1–9. <http://dx.doi.org/10.1080/13693780600928495>.
  18. Andersen B, Frisvad JC, Sondergaard I, Rasmussen IS, Larsen LS. 2011. Associations between fungal species and water-damaged building materials. *Appl Environ Microbiol* 77:4180–4188. <http://dx.doi.org/10.1128/AEM.02513-10>.
  19. Fradkin A, Tarlo SM, Tobin RS, Tucicporretta M, Malloch D. 1987. Species identification of airborne molds and its significance for the detection of indoor pollution. *JAPCA* 37:51–53. <http://dx.doi.org/10.1080/08940630.1987.10466201>.
  20. Pasanen P, Korpi A, Kalliokoski P, Pasanen AL. 1997. Growth and volatile metabolite production of *Aspergillus versicolor* in house dust. *Environ Int* 23:425–432. [http://dx.doi.org/10.1016/S0160-4120\(97\)00027-5](http://dx.doi.org/10.1016/S0160-4120(97)00027-5).
  21. Polizzi V, Delmulle B, Adams A, Moretti A, Susca A, Picco AM, Rosseel Y, Kindt R, Van Bocxlaer J, De Kimpe N, Van Peteghem C, De Saeger S. 2009. Jem spotlight: fungi, mycotoxins and microbial volatile organic compounds in mouldy interiors from water-damaged buildings. *J Environ Monit* 11:1849–1858. <http://dx.doi.org/10.1039/b906856b>.
  22. Pasanen AL, Kasanen JP, Rautiala S, Ikaheimo M, Rantamaki J, Kaariainen H, Kalliokoski P. 2000. Fungal growth and survival in building materials under fluctuating moisture and temperature conditions. *Int Biodeterior Biodegradation* 46:117–127. [http://dx.doi.org/10.1016/S0964-8305\(00\)00093-7](http://dx.doi.org/10.1016/S0964-8305(00)00093-7).
  23. Ayerst G. 1969. The effects of moisture and temperature on growth and spore germination in some fungi. *J Stored Prod Res* 5:127–141. [http://dx.doi.org/10.1016/0022-474X\(69\)90055-1](http://dx.doi.org/10.1016/0022-474X(69)90055-1).
  24. Hallsworth JE, Magan N. 1995. Manipulation of intracellular glycerol and erythritol enhances germination of conidia at low water availability. *Microbiology* 141:1109–1115. <http://dx.doi.org/10.1099/13500872-141-5-1109>.
  25. Magan N, Lacey J. 1984. Effect of temperature and pH on water relations of field and storage fungi. *Trans Br Mycol Soc* 82:71–81. [http://dx.doi.org/10.1016/S0007-1536\(84\)80213-2](http://dx.doi.org/10.1016/S0007-1536(84)80213-2).
  26. Grant C, Hunter CA, Flannigan B, Bravery AF. 1989. The moisture requirements of molds isolated from domestic dwellings. *Int Biodeterior* 25:259–284. [http://dx.doi.org/10.1016/0265-3036\(89\)90002-X](http://dx.doi.org/10.1016/0265-3036(89)90002-X).
  27. Nielsen KF, Holm G, Uttrup LP, Nielsen PA. 2004. Mould growth on building materials under low water activities. Influence of humidity and temperature on fungal growth and secondary metabolism. *Int Biodeterior Biodegradation* 54:325–336.
  28. Adan OCG. 1994. On the fungal defacement of interior finishes. Eindhoven University of Technology, Eindhoven, The Netherlands.
  29. Bekker M. 2014. Growth of *Penicillium rubens* after desiccation. Technical University of Eindhoven, Eindhoven, The Netherlands.
  30. Bekker M, Erich SJF, Hermanns SPM, van Maris MPFHL, Huinink HP, Adan OCG. 2015. Quantifying discoloration caused by the indoor fungus *Penicillium rubens* on building material at controlled humidity. *Build Environ* 90:60–70. <http://dx.doi.org/10.1016/j.buildenv.2015.03.020>.
  31. van Laarhoven KA, Huinink HP, Segers FJJ, Dijksterhuis J, Adan OCG. 2015. Separate effects of moisture content and water activity on the hyphal extension of *Penicillium rubens* on porous media. *Environ Microbiol* 17:5089–5099. <http://dx.doi.org/10.1111/1462-2920.13012>.
  32. Johansson P, Bok G, Ekstrand-Tobin A. 2013. The effect of cyclic moisture and temperature on mould growth on wood compared to steady state conditions. *Build Environ* 65:178–184. <http://dx.doi.org/10.1016/j.buildenv.2013.04.004>.
  33. Johansson P, Ekstrand-Tobin A, Svensson T, Bok G. 2012. Laboratory study to determine the critical moisture level for mould growth on building materials. *Int Biodeterior Biodegradation* 73:23–32. <http://dx.doi.org/10.1016/j.ibiod.2012.05.014>.
  34. Viitanen HA. 1997. Modelling the time factor in the development of mould fungi—the effect of critical humidity and temperature conditions on pine and spruce sapwood. *Int J Biol Chem Phys Technol Wood* 51:6–14.
  35. Li Y, Wadsö L. 2013. Fungal activities of indoor moulds on wood as a function of relative humidity during desorption and adsorption processes. *Eng Life Sci* 13:528–535. <http://dx.doi.org/10.1002/elsc.201200100>.
  36. Viitanen H, Ojanen T. 2007. Improved model to predict mold growth in building materials. In *Abstr Thermal Performance of the Exterior Envelopes of Whole Buildings X*, Clearwater Beach, FL, 2 to 7 December 2007.
  37. Johansson P, Svensson T, Ekstrand-Tobin A. 2013. Validation of critical moisture conditions for mould growth on building materials. *Build Environ* 62:201–209. <http://dx.doi.org/10.1016/j.buildenv.2013.01.012>.
  38. Adan OCG, Samson RA. 2011. Introduction, p 15–36. In Adan OCG, Samson RA (ed), *Fundamentals of mold growth in indoor environments and strategies for healthy living*. Wageningen Academic Publishers, Wageningen, The Netherlands.
  39. Bos CJ, Debets AJM, Swart K, Huybers A, Kobus G, Slakhorst SM. 1988. Genetic-analysis and the construction of master strains for assignment of genes to 6 linkage groups in *Aspergillus niger*. *Curr Genet* 14:437–443. <http://dx.doi.org/10.1007/BF00521266>.
  40. Segers FJJ, Meijer M, Houbraken J, Samson RA, Wösten HAB, Dijksterhuis J. 2015. Xerotolerant *Cladosporium sphaerospermum* are predominant on indoor surfaces compared to other *Cladosporium* species. *PLoS One* 10:e0145415. <http://dx.doi.org/10.1371/journal.pone.0145415>.
  41. Houbraken J, Frisvad JC, Samson RA. 2011. Fleming's penicillin producing strain is not *Penicillium chrysogenum* but *P. rubens*. *IMA Fungus* 2:87–95. <http://dx.doi.org/10.5598/ima fungus.2011.02.01.12>.
  42. Beuchat LR, Hwang CA. 1996. Evaluation of modified dichloran 18% glycerol (DG18) agar for enumerating fungi in wheat flour: a collaborative study. *Int J Food Microbiol* 29:161–166. [http://dx.doi.org/10.1016/0168-1605\(95\)00042-9](http://dx.doi.org/10.1016/0168-1605(95)00042-9).
  43. Hocking AD, Pitt JL. 1980. Dichloran-glycerol medium for enumeration of xerophilic fungi from low-moisture foods. *Appl Environ Microbiol* 39:488–492.
  44. Greenspan L. 1977. Humidity fixed points of binary saturated aqueous solutions. *J Res Natl Bureau Stand* 81:89–96.
  45. Van Leeuwen MR, Golovina EA, Dijksterhuis J. 2009. The polyene antimycotics nystatin and filipin disrupt the plasma membrane, whereas natamycin inhibits endocytosis in germinating conidia of *Penicillium dis-*

- color. *J Appl Microbiol* 106:1908–1918. <http://dx.doi.org/10.1111/j.1365-2672.2009.04165.x>.
46. Van Leeuwen MR, Van Doorn TM, Golovina EA, Stark J, Dijksterhuis J. 2010. Water- and air-distributed conidia differ in sterol content and cytoplasmic microviscosity. *Appl Environ Microbiol* 76:366–369. <http://dx.doi.org/10.1128/AEM.01632-09>.
  47. Fritsche W, Laplace F. 1999. *Mikrobiologie*. Spektrum Akademischer Verlag, Heidelberg, Germany.
  48. Samson RA, Houbraeken J, Thrane U, Frisvad JC, Andersen B. 2010. Food and indoor fungi. CBS-KNAW Fungal Biodiversity Centre, Utrecht, The Netherlands.
  49. Sautour M, Dantigny P, Divies C, Bensoussan M. 2001. A temperature-type model for describing the relationship between fungal growth and water activity. *Int J Food Microbiol* 67:63–69. [http://dx.doi.org/10.1016/S0168-1605\(01\)00471-8](http://dx.doi.org/10.1016/S0168-1605(01)00471-8).
  50. Lamarre C, Sokol S, Debeaupuis JP, Henry C, Lacroix C, Glaser P, Coppee JY, Francois JM, Latge JP. 2008. Transcriptomic analysis of the exit from dormancy of *Aspergillus fumigatus* conidia. *BMC Genomics* 9:417. <http://dx.doi.org/10.1186/1471-2164-9-417>.
  51. Wyatt TT, van Leeuwen MR, Golovina EA, Hoekstra FA, Kuenstner EJ, Palumbo EA, Snyder NL, Visagie C, Verkennis A, Hallsworth JE, Wosten HAB, Dijksterhuis J. 2015. Functionality and prevalence of trehalose-based oligosaccharides as novel compatible solutes in ascospores of *Neosartorya fischeri* (*Aspergillus fischeri*) and other fungi. *Environ Microbiol* 17:395–411. <http://dx.doi.org/10.1111/1462-2920.12558>.
  52. Hoekstra FA, Golovina EA, Buitink J. 2001. Mechanisms of plant desiccation tolerance. *Trends Plant Sci* 6:431–438. [http://dx.doi.org/10.1016/S1360-1385\(01\)02052-0](http://dx.doi.org/10.1016/S1360-1385(01)02052-0).
  53. Dantigny P, Nanguy SPM. 2009. Significance of the physiological state of fungal spores. *Int J Food Microbiol* 134:16–20. <http://dx.doi.org/10.1016/j.ijfoodmicro.2009.02.005>.
  54. Wessels JG. 1993. Tansley review no. 45. Wall growth, protein excretion and morphogenesis in fungi. *New Phytol* 123:397–413.
  55. Dijksterhuis J. 2011. The fungal cell, p 83–100. *In* Adan OCG, Samson RA (ed), *Fundamentals of mold growth in indoor environments and strategies for healthy living*, vol 1. Wageningen Academic Publishers, Wageningen, The Netherlands.
  56. Park D. 1982. Phylloplane fungi: tolerance of hyphal tips to drying. *Trans Br Mycol Soc* 79:174–178. [http://dx.doi.org/10.1016/S0007-1536\(82\)80212-X](http://dx.doi.org/10.1016/S0007-1536(82)80212-X).
  57. Moody SA, Newsham KK, Ayres PG, Paul ND. 1999. Variation in the responses of litter and phylloplane fungi to UV-B radiation (290–315 nm). *Mycol Res* 103:1469–1477. <http://dx.doi.org/10.1017/S0953756299008783>.
  58. Dadachova E, Bryan RA, Howell RC, Schweitzer AD, Aisen P, Nosanchuk JD, Casadevall A. 2008. The radioprotective properties of fungal melanin are a function of its chemical composition, stable radical presence and spatial arrangement. *Pigment Cell Melanoma Res* 21:192–199. <http://dx.doi.org/10.1111/j.1755-148X.2007.00430.x>.
  59. Feofilova EP. 2010. The fungal cell wall: modern concepts of its composition and biological function. *Microbiology* 79:711–720. <http://dx.doi.org/10.1134/S0026261710060019>.
  60. Gostincar C, Grube M, de Hoog S, Zalar P, Gunde-Cimerman N. 2010. Extremotolerance in fungi: evolution on the edge. *FEMS Microbiol Ecol* 71:2–11. <http://dx.doi.org/10.1111/j.1574-6941.2009.00794.x>.
  61. Bekker M, Huinink HP, Adan OCG, Samson RA, Wyatt T, Dijksterhuis J. 2012. Production of an extracellular matrix as an isotropic growth phase of *Penicillium rubens* on gypsum. *Appl Environ Microbiol* 78:6930–6937. <http://dx.doi.org/10.1128/AEM.01506-12>.