

## N-(2-Benzoylethyl)propan-2-aminium chloride

Abdullah Aydin,<sup>a\*</sup> Mehmet Akkurt,<sup>b</sup> Halise Inci Gul,<sup>c</sup> Ebru Mete<sup>d</sup> and Ertan Sahin<sup>d</sup>

<sup>a</sup>Department of Science Education, Faculty of Education, Kastamonu University, 37200 Kastamonu, Turkey, <sup>b</sup>Department of Physics, Faculty of Sciences, Erciyes University, 38039 Kayseri, Turkey, <sup>c</sup>Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Ataturk University, 25240 Erzurum, Turkey, and <sup>d</sup>Department of Chemistry, Faculty of Sciences, Ataturk University, 25240 Erzurum, Turkey  
Correspondence e-mail: aaydin@kastamonu.edu.tr


Received 31 July 2012; accepted 8 August 2012

Key indicators: single-crystal X-ray study;  $T = 294\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.008\text{ \AA}$ ;  $R$  factor = 0.075;  $wR$  factor = 0.224; data-to-parameter ratio = 16.1.

In the title salt,  $\text{C}_{12}\text{H}_{18}\text{NO}^+\cdot\text{Cl}^-$ , N—H···Cl interactions between the free chloride anions and the organic cations connect the molecules into hydrogen-bond dimers, forming a  $R_2^2(8)$  motif. The dimers are linked by C—H···O hydrogen bonds into chains extending along [301]. The carbonyl group is co-planar with the phenyl ring [ $\text{C}=\text{C}=\text{O}$  torsion angle =  $-3.3(7)^\circ$ ]. The side chain has an *E* conformation.

### Related literature

For the details of the pharmacological effects of Mannich bases and for their synthesis, see: Dimmock & Kumar (1997); Gul *et al.* (2004, 2005a,b, 2009); Mete *et al.* (2011a,b); Kucukoglu *et al.* (2011); Canturk *et al.* (2008); Chen *et al.* (1991); Gul (2005); Suleyman *et al.* (2007); Plati *et al.* (1949). For bond-length data, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995); Etter (1990). For some related structures, see: Abonia *et al.* (2011); Tuzina *et al.* (2006).



### Experimental

#### Crystal data

$\text{C}_{12}\text{H}_{18}\text{NO}^+\cdot\text{Cl}^-$   
 $M_r = 227.72$   
Monoclinic,  $P2_1/n$

$a = 8.036(5)\text{ \AA}$   
 $b = 8.656(5)\text{ \AA}$   
 $c = 18.403(5)\text{ \AA}$

$\beta = 97.174(5)^\circ$   
 $V = 1270.1(11)\text{ \AA}^3$   
 $Z = 4$   
Mo  $K\alpha$  radiation

$\mu = 0.28\text{ mm}^{-1}$   
 $T = 294\text{ K}$   
 $0.16 \times 0.13 \times 0.12\text{ mm}$

#### Data collection

Rigaku R-AXIS RAPID-S diffractometer  
Absorption correction: multi-scan (Blessing, 1995)  
 $T_{\min} = 0.958$ ,  $T_{\max} = 0.967$

15799 measured reflections  
2334 independent reflections  
1204 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.116$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.075$   
 $wR(F^2) = 0.224$   
 $S = 1.05$   
2334 reflections  
145 parameters  
2 restraints

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.49\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.19\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N···Cl1 <sup>i</sup>	0.87 (4)	2.30 (4)	3.141 (4)	163 (4)
N1—H2N···Cl1	0.87 (3)	2.27 (3)	3.131 (4)	169 (4)
C10—H10···O1 <sup>ii</sup>	0.98	2.56	3.284 (7)	130

Symmetry codes: (i)  $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{3}{2}$ , (ii)  $-x + 1, -y + 1, -z + 2$ .

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are indebted to the Department of Chemistry, Ataturk University, Erzurum, Turkey, for use of the X-ray diffractometer purchased under grant No. 2003/219 of the University Research Fund.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: QM2079).

### References

- Abonia, R., Schollmeyer, D. & Arteaga, D. (2011). *Acta Cryst. E* **67**, o2969.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Canturk, P., Kucukoglu, K., Topcu, Z., Gul, M. & Gul, H. I. (2008). *Arzneim. Forsch.* **58**, 686–691.
- Chen, H. T., Jing, Y. K., Ji, Z. Z. & Zhang, B. F. (1991). *YaoXueXueBao*, **26**, 183–192.
- Dimmock, J. R. & Kumar, P. (1997). *Curr. Med. Chem.* **4**, 1–22.
- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gul, M. (2005). PhD thesis, Kuopio University, Finland.
- Gul, M., Atalay, M., Gul, H. I., Nakao, C., Lappalainen, J. & Hanninen, O. (2005a). *Toxicol. Vitro*, **19**, 573–580.
- Gul, H. I., Calis, U. & Vepsalainen, J. (2004). *Arzneim. Forsch.* **54**, 359–364.
- Gul, H. I., Sahin, F., Gul, M., Ozturk, S. & Yerdelen, K. O. (2005b). *Arch. Pharm.* **338**, 335–338.

- Gul, H. I., Suleyman, H. & Gul, M. (2009). *Pharm. Biol.* **47**, 968–972.
- Kucukoglu, K., Gul, M., Atalay, M., Mete, E., Kazaz, C., Hanninen, O. & Gul, H. I. (2011). *Arzneim. Forsch. Drug Res.* **61**, 366–371.
- Mete, E., Gul, H. I., Bilginer, S., Algul, O., Topaloglu, M. E., Gulluce, M. & Kazaz, C. (2011a). *Molecules*, **16**, 4660–4671.
- Mete, E., Gul, H. I., Cetin-Atalay, R., Das, U., Sahin, E., Gul, M., Kazaz, C. & Dimmock, J. R. (2011b). *Arch. Pharm. Chem. Life Sci.* **11**, 333–339.
- Plati, J. T., Schmidt, R. A. & Wenner, W. (1949). *J. Org. Chem.* **14**, 873–878..
- Rigaku/MSC (2005). *CrystalClear*. Rigaku/MSC, The Woodlands, Texas, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Suleyman, H., Gul, H. I., Gul, M., Alkan, M. & Gocer, F. (2007). *Biol. Pharm. Bull.* **30**, 63–67.
- Tuzina, P., Fischer, A. & Somfai, P. (2006). *Acta Cryst. E* **62**, o2971–o2972.

# supplementary materials

*Acta Cryst.* (2012). E68, o2702–o2703 [doi:10.1107/S1600536812035106]

## N-(2-Benzoylethyl)propan-2-aminium chloride

**Abdullah Aydin, Mehmet Akkurt, Halise Inci Gul, Ebru Mete and Ertan Sahin**

### Comment

Mannich bases are generally formed by the reaction between formaldehyde, a secondary amine and a compound containing reactive hydrogen atoms. On occasion, aldehydes other than formaldehyde may be employed and the secondary amine may be replaced by ammonia and primary amines. This process is known as the Mannich reaction (Dimmock & Kumar, 1997).

Mannich bases display varied biological activities such as antimicrobial (Gul *et al.*, 2005; Mete *et al.*, 2011a), cytotoxic (Gul *et al.*, 2005; Mete *et al.*, 2011b; Kucukoglu *et al.*, 2011; Canturk *et al.*, 2008), anticancer (Dimmock & Kumar, 1997; Chen *et al.*, 1991; Gul, 2005), antiinflammatory (Suleyman *et al.*, 2007; Gul *et al.*, 2009), anticonvulsant (Gul *et al.*, 2004) and DNA topoisomeraseI inhibiting properties (Canturk *et al.*, 2008).

The geometric parameters of the title salt (I) in Fig. 1 are within the range of expected values for this type of compound (Allen *et al.*, 1987; Abonia *et al.*, 2011; Tuzina *et al.*, 2006).

The N—H···Cl hydrogen-bonding interactions between the free chloride anion and the organic cation link the molecules into hydrogen-bond dimers, forming an  $R^2_2(8)$  motif (Bernstein *et al.*, 1995; Etter, 1990). The dimers are linked by C—H···O hydrogen bonds, into chains extended along the [301] direction (Table 1, Fig. 2).

### Experimental

A mixture of the appropriate ketone (50 mmol), paraformaldehyde (50 mmol), and isopropylamine hydrochloride (27 mmol) was heated in an oil bath at 403 K. The reaction vessel was then removed from the oil bath and when the temperature of the mixture dropped to 338 K, ethyl acetate (40–80 ml) was added. The mixture was stirred at room temperature for 24 h and the resultant precipitate was collected and recrystallized from ether/methanol. The melting point was 445–447 K (lit. Plati *et al.*, 1949 m.p. 447–449 K) and the yield was 55% (Mete *et al.*, 2011b).


$^1\text{H-NMR}$   $\delta$  1.49 (d,  $J = 6.6$  Hz, 6H,  $\text{CH}(\text{CH}_3)_2$ ), 3.36–3.42 (m, 3H,  $\text{CH}(\text{CH}_3)_2$  and  $2 \times \text{H}-2$ ), 3.77 (t,  $J = 7.5$  Hz, 2H,  $2 \times \text{H}-3$ ), 7.36 (t,  $J = 7.3$  Hz, 2H,  $\text{H}-3'/5'$ ), 7.51 (t,  $J = 7.3$  Hz, 1H,  $\text{H}-4'$ ), 7.90 (d,  $J = 7.3$  Hz, 2H,  $\text{H}-2'/6'$ ), 9.56 (brs, 2H,  $\text{NH}_2^+$ );  $^{13}\text{C-NMR}$   $\delta$  19.4 ( $\text{CH}(\text{CH}_3)_2$ ), 35.2, 40.3, 51.2, 128.3, 128.9, 134.0, 136.0, 196.9; MS (EI)  $m/z$  (%): 176.2 ( $M-\text{CH}_3$ ) $^+$ , 192.1 ( $M+\text{H}$ ) $^+$ . IR (KBr, cm $^{-1}$ ): 2453 ( $\text{NH}_2^+$ ), 1678 (CO). Calcd for  $\text{C}_{12}\text{H}_{18}\text{ClNO}$  (227.73): C, 63.29; H, 7.97; N, 6.15. Found: C, 63.26; H, 8.18; N, 6.23.

### Refinement


H atoms of the  $\text{NH}_2$  group were located in a difference Fourier map. Their positions refined with restraints on the N—H bond lengths of 0.86 (2) Å, while their thermal parameters were refined as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ . H atoms bound to C atoms were positioned geometrically, with C—H = 0.93(aromatic), 0.97(methylene) and 0.98 Å (methine), and refined as riding with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$  for methyl H and  $1.2U_{\text{eq}}(\text{C})$  for the others.

**Computing details**

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear* (Rigaku/MSC, 2005); data reduction: *CrystalClear* (Rigaku/MSC, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

**Figure 1**

View of the title molecule with the atom numbering scheme. Displacement ellipsoids for non-H atoms are drawn at the 30% probability level.

**Figure 2**

The packing and hydrogen bonding of (I) viewed down the *b* axis. H atoms not involved in hydrogen bondings are omitted for the sake of clarity.

***N-(2-Benzoylethyl)propan-2-aminium chloride****Crystal data* $M_r = 227.72$ Monoclinic,  $P2_1/n$ 

Hall symbol: -P 2yn

 $a = 8.036 (5) \text{ \AA}$  $b = 8.656 (5) \text{ \AA}$  $c = 18.403 (5) \text{ \AA}$  $\beta = 97.174 (5)^\circ$  $V = 1270.1 (11) \text{ \AA}^3$  $Z = 4$  $F(000) = 488$  $D_x = 1.191 \text{ Mg m}^{-3}$ Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$ 

Cell parameters from 3090 reflections

 $\theta = 2.2\text{--}26.4^\circ$  $\mu = 0.28 \text{ mm}^{-1}$  $T = 294 \text{ K}$ 

Block, white

 $0.16 \times 0.13 \times 0.12 \text{ mm}$ *Data collection*Rigaku R-AXIS RAPID-S  
diffractometer

Radiation source: Sealed Tube

Graphite Monochromator monochromator

Detector resolution: 10.0000 pixels  $\text{mm}^{-1}$ 

dtpprofit.ref scans

Absorption correction: multi-scan  
(Blessing, 1995) $T_{\min} = 0.958, T_{\max} = 0.967$ 

15799 measured reflections

2334 independent reflections

1204 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.116$  $\theta_{\max} = 25.5^\circ, \theta_{\min} = 2.2^\circ$  $h = -9 \rightarrow 9$  $k = -9 \rightarrow 10$  $l = -22 \rightarrow 22$ *Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.075$  $wR(F^2) = 0.224$  $S = 1.05$ 

2334 reflections

145 parameters

2 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sitesH atoms treated by a mixture of independent  
and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0971P)^2 + 0.2008P]$   
where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.49 \text{ e \AA}^{-3}$  $\Delta\rho_{\min} = -0.19 \text{ e \AA}^{-3}$ *Special details*

**Geometry.** Bond distances, angles etc. have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement on  $F^2$  for ALL reflections except those flagged by the user for potential systematic errors. Weighted  $R$ -factors  $wR$  and all goodnesses of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The observed criterion of  $F^2 > \sigma(F^2)$  is used only for calculating - $R$ -factor-obs etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^* / U_{\text{eq}}$
O1	0.2324 (5)	0.4012 (5)	1.02649 (18)	0.1262 (18)
N1	0.3982 (4)	0.4236 (4)	0.8222 (2)	0.0689 (14)
C1	-0.1184 (6)	0.1831 (6)	0.9398 (3)	0.090 (2)

C2	-0.2586 (7)	0.1067 (7)	0.9575 (4)	0.114 (3)
C3	-0.2887 (7)	0.0999 (7)	1.0303 (4)	0.110 (3)
C4	-0.1824 (7)	0.1686 (8)	1.0833 (3)	0.103 (3)
C5	-0.0433 (7)	0.2458 (6)	1.0658 (3)	0.088 (2)
C6	-0.0099 (6)	0.2528 (5)	0.9940 (2)	0.0722 (17)
C7	0.1440 (6)	0.3359 (6)	0.9778 (3)	0.0828 (19)
C8	0.1897 (5)	0.3369 (5)	0.9007 (2)	0.0749 (17)
C9	0.3548 (5)	0.4217 (5)	0.8982 (2)	0.0772 (17)
C10	0.5624 (6)	0.5009 (5)	0.8146 (3)	0.0850 (19)
C11	0.6181 (8)	0.4545 (7)	0.7460 (4)	0.133 (3)
C12	0.5461 (6)	0.6727 (6)	0.8219 (3)	0.109 (3)
C11	0.40593 (14)	0.07597 (13)	0.77578 (6)	0.0828 (5)
H1	-0.09700	0.18770	0.89130	0.1080*
H1N	0.321 (5)	0.453 (5)	0.788 (2)	0.0990*
H2	-0.33240	0.06020	0.92100	0.1370*
H2N	0.402 (6)	0.332 (3)	0.803 (2)	0.0990*
H3	-0.38260	0.04780	1.04250	0.1320*
H4	-0.20350	0.16370	1.13180	0.1240*
H5	0.02880	0.29350	1.10240	0.1060*
H8A	0.10170	0.38750	0.86830	0.0900*
H8B	0.19980	0.23160	0.88380	0.0900*
H9A	0.34510	0.52680	0.91550	0.0930*
H9B	0.44310	0.37050	0.93010	0.0930*
H10	0.64530	0.46460	0.85460	0.1020*
H11A	0.53580	0.48440	0.70610	0.1990*
H11B	0.63260	0.34440	0.74560	0.1990*
H11C	0.72290	0.50400	0.74080	0.1990*
H12A	0.65050	0.72110	0.81500	0.1630*
H12B	0.51800	0.69720	0.86970	0.1630*
H12C	0.45930	0.70980	0.78540	0.1630*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.151 (3)	0.161 (4)	0.065 (2)	-0.063 (3)	0.007 (2)	-0.016 (2)
N1	0.065 (2)	0.066 (2)	0.076 (3)	-0.0063 (19)	0.0105 (16)	0.0006 (18)
C1	0.093 (4)	0.099 (4)	0.083 (3)	-0.006 (3)	0.027 (3)	-0.010 (3)
C2	0.082 (4)	0.143 (6)	0.119 (5)	-0.017 (3)	0.022 (3)	-0.019 (4)
C3	0.087 (4)	0.138 (5)	0.113 (5)	0.003 (3)	0.039 (4)	0.015 (4)
C4	0.091 (4)	0.136 (5)	0.086 (4)	0.025 (4)	0.028 (3)	0.020 (3)
C5	0.094 (4)	0.102 (4)	0.069 (3)	0.013 (3)	0.012 (3)	0.003 (3)
C6	0.078 (3)	0.072 (3)	0.068 (3)	0.009 (2)	0.015 (2)	-0.002 (2)
C7	0.097 (4)	0.086 (3)	0.065 (3)	0.002 (3)	0.009 (2)	-0.003 (2)
C8	0.084 (3)	0.075 (3)	0.067 (3)	-0.007 (2)	0.015 (2)	-0.001 (2)
C9	0.081 (3)	0.078 (3)	0.072 (3)	-0.004 (2)	0.007 (2)	0.002 (2)
C10	0.069 (3)	0.069 (3)	0.118 (4)	-0.005 (2)	0.016 (3)	0.003 (3)
C11	0.127 (5)	0.131 (5)	0.151 (6)	-0.037 (4)	0.061 (4)	-0.027 (4)
C12	0.090 (4)	0.069 (3)	0.168 (6)	-0.015 (3)	0.021 (3)	0.002 (3)
C11	0.0792 (8)	0.0703 (8)	0.0974 (9)	0.0004 (6)	0.0050 (6)	-0.0080 (6)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

O1—C7	1.212 (6)	C1—H1	0.9300
N1—C9	1.483 (5)	C2—H2	0.9300
N1—C10	1.502 (6)	C3—H3	0.9300
N1—H1N	0.87 (4)	C4—H4	0.9300
N1—H2N	0.87 (3)	C5—H5	0.9300
C1—C2	1.380 (8)	C8—H8A	0.9700
C1—C6	1.379 (7)	C8—H8B	0.9700
C2—C3	1.392 (10)	C9—H9A	0.9700
C3—C4	1.352 (9)	C9—H9B	0.9700
C4—C5	1.375 (8)	C10—H10	0.9800
C5—C6	1.382 (7)	C11—H11A	0.9600
C6—C7	1.493 (7)	C11—H11B	0.9600
C7—C8	1.509 (7)	C11—H11C	0.9600
C8—C9	1.522 (6)	C12—H12A	0.9600
C10—C12	1.500 (7)	C12—H12B	0.9600
C10—C11	1.448 (9)	C12—H12C	0.9600
C9—N1—C10	113.9 (3)	C5—C4—H4	120.00
C10—N1—H1N	111 (3)	C4—C5—H5	120.00
C9—N1—H1N	117 (3)	C6—C5—H5	120.00
C9—N1—H2N	113 (2)	C7—C8—H8A	110.00
C10—N1—H2N	107 (3)	C7—C8—H8B	110.00
H1N—N1—H2N	92 (4)	C9—C8—H8A	110.00
C2—C1—C6	120.0 (5)	C9—C8—H8B	110.00
C1—C2—C3	119.5 (6)	H8A—C8—H8B	108.00
C2—C3—C4	120.4 (5)	N1—C9—H9A	110.00
C3—C4—C5	120.3 (5)	N1—C9—H9B	110.00
C4—C5—C6	120.4 (5)	C8—C9—H9A	110.00
C5—C6—C7	118.5 (4)	C8—C9—H9B	110.00
C1—C6—C7	122.1 (4)	H9A—C9—H9B	108.00
C1—C6—C5	119.4 (5)	N1—C10—H10	108.00
O1—C7—C6	120.1 (5)	C11—C10—H10	108.00
C6—C7—C8	119.7 (4)	C12—C10—H10	108.00
O1—C7—C8	120.2 (4)	C10—C11—H11A	109.00
C7—C8—C9	110.3 (3)	C10—C11—H11B	109.00
N1—C9—C8	110.0 (3)	C10—C11—H11C	109.00
N1—C10—C12	110.2 (4)	H11A—C11—H11B	110.00
C11—C10—C12	113.1 (5)	H11A—C11—H11C	109.00
N1—C10—C11	109.2 (4)	H11B—C11—H11C	109.00
C2—C1—H1	120.00	C10—C12—H12A	109.00
C6—C1—H1	120.00	C10—C12—H12B	110.00
C1—C2—H2	120.00	C10—C12—H12C	109.00
C3—C2—H2	120.00	H12A—C12—H12B	110.00
C2—C3—H3	120.00	H12A—C12—H12C	109.00
C4—C3—H3	120.00	H12B—C12—H12C	109.00
C3—C4—H4	120.00	 	
C10—N1—C9—C8	-178.2 (3)	C4—C5—C6—C1	0.7 (8)

C9—N1—C10—C11	161.9 (4)	C4—C5—C6—C7	−179.1 (5)
C9—N1—C10—C12	−73.3 (5)	C5—C6—C7—C8	176.1 (4)
C2—C1—C6—C7	179.6 (5)	C1—C6—C7—O1	176.9 (5)
C6—C1—C2—C3	−0.4 (8)	C1—C6—C7—C8	−3.7 (7)
C2—C1—C6—C5	−0.2 (8)	C5—C6—C7—O1	−3.3 (7)
C1—C2—C3—C4	0.6 (9)	O1—C7—C8—C9	2.1 (6)
C2—C3—C4—C5	−0.1 (10)	C6—C7—C8—C9	−177.3 (4)
C3—C4—C5—C6	−0.6 (9)	C7—C8—C9—N1	−179.5 (4)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1N···C11 <sup>i</sup>	0.87 (4)	2.30 (4)	3.141 (4)	163 (4)
N1—H2N···C11	0.87 (3)	2.27 (3)	3.131 (4)	169 (4)
C10—H10···O1 <sup>ii</sup>	0.98	2.56	3.284 (7)	130

Symmetry codes: (i)  $-x+1/2, y+1/2, -z+3/2$ ; (ii)  $-x+1, -y+1, -z+2$ .