

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

- 26 Physicians for Human Rights. Obstruction and denial: health system disparities and COVID-19 in Daraa, Syria. December, 2020. https://phr.org/ wp-content/uploads/2020/12/PHR_Obstruction-and-Denial_Health-System-Disparities-and-COVID-19-in-Daraa-Syria_Dec-2020.pdf (accessed March 10, 2021).
- 27 Syrians for Truth and Justice. HTS stormed the free Hama Health Directorate and arrested doctors. Dec 19, 2018. https://stj-sy.org/en/1104/ (accessed March 10, 2021).
- 28 Khattab A. "It's like Judgment Day": Syrians recount horror of an underreported COVID-19 outbreak. Newlines Magazine, Oct 11, 2020. https://newlinesmag.com/reportage/its-like-judgment-day-syriansrecount-horror-of-an-underreported-covid-19-outbreak/ (accessed March 10, 2021).
- 29 Reuters. Syria says it has received COVID-19 vaccinations from "friendly country". Feb 25, 2021. https://www.reuters.com/article/us-healthcoronavirus-syria-vaccine-idUSKBN2AP1LA (accessed March 11, 2021).
- 30 Al-Khalidi S. WHO preparing to deliver vaccines across Syria from April despite conflict. Feb 4, 2021. https://www.reuters.com/article/us-healthcoronvirus-vaccine-syria/who-preparing-to-deliver-vaccines-across-syriafrom-april-despite-conflict-idINKBN2A32V7 (accessed March 11, 2021).
- 31 OCHA Services. Syria Humanitarian Response Plan 2020. Response plan/ appeal snapshot for 2020. https://fts.unocha.org/appeals/924/summary (accessed March 11, 2021).
- 32 Human Rights Watch. Rigging the system: government policies co-opt aid and reconstruction funding in Syria. June 28, 2019. https://www.hrw.org/ sites/default/files/report_pdf/syria0619_web4.pdf (accessed March 10, 2021).

- 33 Duclos D, Ekzayez A, Ghaddar F, et al. Localisation and cross-border assistance to deliver humanitarian health services in North-West Syria: a qualitative inquiry for The Lancet-AUB Commission on Syria. Confl Health 2019; published online May 27. https://doi.org/10.1186/s13031-019-0207-z.
- 34 Hill E, Triebert C. 12 hours. 4 Syrian hospitals bombed. One culprit: Russia. The New York Times, Oct 13, 2019.
- 35 DW News. German court hands down historic Syrian torture verdict. Feb 24, 2021. https://www.dw.com/en/german-court-hands-down-historicsyrian-torture-verdict/a-56670243 (accessed March 10, 2021).
- 36 Schuetze CF, Hubbard B. Syrian doctor accused of torture is arrested in Germany. The New York Times, June 22, 2020.
- 37 Syrian Network for Human Rights. At least 3,364 health care personnel still arrested/forcibly disappeared, 98% by the Syrian Regime. Feb 27, 2021. https:// sn4hr.org/wp-content/pdf/english/At_Least_3364_Health_Care_Personnel_ Still_Arrested_Forcibly_Disappeared_en.pdf(accessed March 10, 2021).
- 38 ACAPS, CrisisInSight. Humanitarian access overview. December, 2020. https://www.acaps.org/sites/acaps/files/products/files/20201214_acaps_ humanitarian_access_overview_december_2020_0.pdf (accessed March 10, 2021).
- 39 MedGlobal. Responding to the COVID-19 crisis in Syria: a five part approach through "Operation Breathe". March 5, 2021. https://medglobal.org/wpcontent/uploads/2021/03/Operation-Breathe-PDF-Report.pdf (accessed March 10, 2021).
- 40 Center for Operational Analysis and Research. Syrian public health after COVID-19: entry points and lessons learned from the pandemic response. March, 2021. https://coar-global.org/syrian-public-health-after-covid-19/ (accessed March 11, 2021).

The need to prioritise childhood tuberculosis case detection



Published Online March 22, 2021 https://doi.org/10.1016/ S0140-6736(21)00672-3

With 10 years left to the WHO End TB Strategy's interim milestones of 80% reduction in new tuberculosis cases and 90% reduction in tuberculosis deaths by 2030 compared with 2015,¹ little progress has been made. The COVID-19 pandemic has worsened the situation because of its negative impact on tuberculosis case detection and reduced access to tuberculosis treatment and prevention services globally.² In a worst-case scenario, COVID-19 might have resulted in up to 400000 excess tuberculosis deaths in 2020, which would mean the worldwide number was similar to that in 2012.3 Estimates suggest that the COVID-19 pandemic could cause an additional 6.3 million tuberculosis cases globally between 2020 and 2025, with the most vulnerable populations, especially children, at risk.⁴ These extreme outcomes would slow or reverse any progress made towards the tuberculosis treatment and prevention milestones and targets.5.6

WHO estimated that children younger than 15 years constituted 12% of the 10 million people who became ill with tuberculosis in 2019.⁴ The global annual numbers of tuberculosis cases accounted for by children have been on the rise, increasing from about 1 million in 2017 to 1.2 million in 2019,⁴⁷ with the highest burden of childhood tuberculosis in China, the Democratic Republic of the Congo, India, Indonesia, and Nigeria.⁸ Despite the global efforts to combat tuberculosis, an

estimated 192000 children died from tuberculosis in 2019.⁴ The high burden of childhood tuberculosis and poor outcomes are largely because of the difficulties in confirming the diagnosis of tuberculosis in children, arising from the non-specific nature of symptoms and signs of tuberculosis in children, the paucibacillary nature of childhood tuberculosis, the challenge of obtaining good-quality sputum samples, especially in young children, and difficulties in accessing health services.⁹ As a result, more than 90% of these children encounter delays in receiving treatment since they are either never diagnosed or are misdiagnosed.⁷ An estimated 568 000 missing childhood tuberculosis cases in 2019 were neither diagnosed nor adequately treated.⁴

A fundamental problem is the absence of large-scale contact screening and provision of tuberculosis preventive therapy to eligible children, which is largely due to insufficient health-care resources, workforce, and services in low-income and middle-income countries (LMICs).⁴ In 2019, more than 40% of eligible household child contacts were not screened for tuberculosis disease or infection.⁴ Therefore, access to and provision of tuberculosis contact screening and preventive treatment needs to be substantially expanded.

Alongside efforts to address multidrug-resistant tuberculosis in children and deliver less toxic and more

child-friendly tuberculosis treatments,⁷ improving childhood tuberculosis case detection is crucial. Efforts to prioritise case detection will also benefit from innovative community-based approaches to case finding, such as the need for enhanced screening for tuberculosis in children presenting at general child health services in LMICs. Given that many children with presumptive tuberculosis present to maternal and child health clinics and private care providers in LMICs,¹⁰ a close interaction between tuberculosis programmes and these services is needed. This approach is essential to reach the over 60% of children who acquire the infection outside the household and who would not be reached by household contact-based approaches, even with the most optimistic assumptions for coverage of household contact tracing.¹¹

Tuberculosis testing is reliant on sputum, a sample that is not easy to collect, especially in young children. Additionally, WHO-recommended rapid tuberculosis diagnostic tests such as Xpert-Ultra are not easily accessible at the primary care level in LMICs where they are most needed.⁷ The development of non-sputum biomarker-based tests for diagnosis of tuberculosis, especially in children who are typically paucibacillary, is now a recognised research priority.¹²

To achieve the WHO End TB Strategy targets and narrow the childhood tuberculosis case detection gap, tuberculosis vaccines are needed that are effective for both pre-exposure and post-exposure prevention and treatment.13 COVID-19 has shown how several efficacious vaccines can be developed within a short timeframe, given the right motivation and available funding. The BCG vaccine, which was developed 100 years ago, effectively prevents severe forms of the disease but has only moderate benefits for prevention of pulmonary tuberculosis.14 Having new tuberculosis vaccines would help reduce new tuberculosis cases and deaths. Given the demonstrated ability to develop effective COVID-19 vaccines rapidly, it is no longer justifiable to not equally invest in the development of tuberculosis vaccines.

To accelerate progress toward ending tuberculosis, there is the need for sustained increase in investments in tuberculosis control services that are delivered as part of comprehensive primary health-care services, and for strengthening health workforce capacity and health systems as part of overall efforts toward achieving universal health coverage in LMICs. Therefore, as we mark World Tuberculosis Day on March 24, 2021, the theme— The Clock is Ticking—is apt to convey a sense of urgency. As countries move towards recovery from the setbacks caused by COVID-19, it is a pivotal moment to make tuberculosis case detection in children a priority. The next few years will be crucial in the tuberculosis response, and it will take sustained efforts by many stakeholders to achieve the internationally agreed targets by 2030. Tuberculosis is preventable and treatable and no child with tuberculosis should miss out on the care they need.

TT reports grants from the UK Research and Innovation Global Challenges Research Fund (UKRI-GCRF) on childhood tuberculosis research and has a patent (Medical Research Council [MRC] Technology ref: A813/3133; TB Biomarkers Script IP Ref: P10280GB) on a childhood tuberculosis diagnostic biomarker pending. BK reports grants from UKRI-GCRF and a MRC programme grant for childhood tuberculosis research, and has a patent (MRC Technology ref: A813/3133; TB Biomarkers Script IP Ref: P10280GB) on a childhood tuberculosis diagnostic biomarker pending. EN declares no competing interests.

Esin Nkereuwem, Beate Kampmann, *Toyin Togun toyin.togun@lshtm.ac.uk

Vaccines and Immunity Theme, Medical Research Council Unit The Gambia at the London School of Hygiene & Tropical Medicine, Fajara, The Gambia (EN, BK, TT); The TB Centre and Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London WC1E 7HT, UK (EN, TT); The Vaccine Centre, London School of Hygiene and Tropical Medicine, London, UK (BK)

- 1 WHO. Resolution WHA67.1. Global strategy and targets for tuberculosis prevention, care and control after 2015. Geneva: World Health Organization, 2014.
- 2 Reid MJA, Silva S, Arinaminpathy N, Goosby E. Building a tuberculosis-free world while responding to the COVID-19 pandemic. *Lancet* 2020; 396: 1312–13.
- 3 Glaziou P. Predicted impact of the COVID-19 pandemic on global tuberculosis deaths in 2020. medRxiv 2020; published online May 4. https://doi.org/10.1101/2020.04.28.20079582 (preprint).
- 4 WHO. Global tuberculosis report 2020. Geneva, Switzerland: World Health Organization, 2020.
- 5 Bhatia V, Mandal PP, Satyanarayana S, Aditama TY, Sharma M. Mitigating the impact of the COVID-19 pandemic on progress towards ending tuberculosis in the WHO South-East Asia Region. WHO South East Asia J Public Health 2020; **9:** 95–99.
- 6 Togun T, Kampmann B, Stoker NG, Lipman M. Anticipating the impact of the COVID-19 pandemic on TB patients and TB control programmes. Ann Clin Microbiol Antimicrobs 2020; 19: 21.
- 7 WHO. Roadmap towards ending TB in children and adolescents. Geneva, Switzerland: World Health Organization, 2018.
- 8 Dodd PJ, Yuen CM, Sismanidis C, Seddon JA, Jenkins HE. The global burden of tuberculosis mortality in children: a mathematical modelling study. *Lancet Glob Health* 2017; 5: e898–06.
- 9 Dodd PJ, Gardiner E, Coghlan R, Seddon JA. Burden of childhood tuberculosis in 22 high-burden countries: a mathematical modelling study. Lancet Glob Health 2014; 2: e453-59.
- 10 Reuter A, Hughes J, Furin J. Challenges and controversies in childhood tuberculosis. *Lancet* 2019; **394:** 967–78.
- 11 Martinez L, Lo NC, Cords O, et al. Paediatric tuberculosis transmission outside the household: challenging historical paradigms to inform future public health strategies. *Lancet Respir Med* 2019; **7**: 544–52.
- 12 Denkinger CM, Kik SV, Cirillo DM, et al. Defining the needs for next generation assays for tuberculosis. J Infect Dis 2015; 211 (suppl 2): 529–38.
- 13 Treatment Action Group. Tuberculosis vaccines: pipeline report 2020. New York: Treatment Action Group, 2020. https://www. treatmentactiongroup.org/wp-content/uploads/2020/10/pipeline_TB_ vaccines_2020_final.pdf (accessed March 15, 2021).
- 14 Kaufmann SH, Hussey G, Lambert PH. New vaccines for tuberculosis. Lancet 2010; 375: 2110–19.