

Research



Cite this article: Smith MA-Y, Honegger KS, Turner G, de Bivort B. 2022 Idiosyncratic learning performance in flies. *Biol. Lett.* **18**: 20210424.
<https://doi.org/10.1098/rsbl.2021.0424>

Received: 10 August 2021
Accepted: 21 December 2021

Subject Areas:
behaviour, neuroscience

Keywords:
generalized learning, individuality, personality, *Drosophila*, Pavlovian conditioning

Author for correspondence:
Benjamin de Bivort
e-mail: debivort@oeb.harvard.edu

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5769072>.

Idiosyncratic learning performance in flies

Matthew A.-Y. Smith¹, Kyle S. Honegger², Glenn Turner³ and Benjamin de Bivort¹

¹Department of Organismic and Evolutionary Biology and Center for Brain Science, Harvard University, Cambridge, MA 02138, USA

²Ann and Robert H. Lurie Children's Hospital of Chicago, Chicago, IL 60611, USA

³Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA 20147, USA

BdB, 0000-0001-6165-7696

Individuals vary in their innate behaviours, even when they have the same genome and have been reared in the same environment. The extent of individuality in plastic behaviours, like learning, is less well characterized. Also unknown is the extent to which intragenotypic differences in learning generalize: if an individual performs well in one assay, will it perform well in other assays? We investigated this using the fruit fly *Drosophila melanogaster*, an organism long-used to study the mechanistic basis of learning and memory. We found that isogenic flies, reared in identical laboratory conditions, and subject to classical conditioning that associated odour with electric shock, exhibit clear individuality in their learning responses. Flies that performed well when an odour was paired with shock tended to perform well when the odour was paired with bitter taste or when other odours were paired with shock. Thus, individuality in learning performance appears to be prominent in isogenic animals reared identically, and individual differences in learning performance generalize across some aversive sensory modalities. Establishing these results in flies opens up the possibility of studying the genetic and neural circuit basis of individual differences in learning in a highly suitable model organism.

1. Introduction

Genetically identical *Drosophila melanogaster*, raised in identical environments, display individuality in numerous innate behaviours [1–8], including light preference [2], left–right turning [3], temperature preference [4], postural behaviours identified by unsupervised analyses [5] and object-tracking [6]. Work to date has focused exclusively on innate or spontaneous behaviours. But plastic behaviours, such as learning, also have the potential to exhibit individuality, as each animal may have an idiosyncratic propensity to respond to training stimuli [9]. Individual variation in learning within insect populations has been described as early as 1907, by Charles Turner [10,11] in ants and honeybees. To our knowledge, individual variation in learning among genetically identical flies has not been characterized.

Here, we present evidence that genetically identical flies exhibit individuality in their ability to learn odour associations. Drawing inspiration from a classical Pavlovian conditioning assay [12–14], animals are exposed to two stimuli simultaneously, a so-called ‘conditioned stimulus’ (CS+), to which their behavioural response will change across the conditioning, and a so-called ‘unconditioned stimulus’ (US), to which their response will remain invariant [15]. In addition, flies were exposed to a second odour, the ‘CS–’, without a US. The learned response to this training is likely to be avoidance of the CS+, as our US is aversive. Our experimental instrument, inspired by [9], allowed (i) measurement of

individual learning performance, (ii) the automated selection of different CS odorants and (3) the use of electric shock or optogenetic activation [16] of negative valence neural circuit elements as US [17]. With this instrument, we can test a fly's generalized learning performance through reversal learning trials (i.e. swapping the CS+ and CS− odours). This paradigm represents a more cognitively demanding form of learning compared to classical conditioning because it requires modification of the previous association [18–23]. We also examined the generality of learning differences by training the same flies across two aversive US modalities: shock and optogenetic stimulation of bitter taste receptor neurons. We found positive correlations in learning performance when varying either the CS odorant or US. Thus, individual learning performance in flies appears to generalize across some stimuli.

2. Material and methods

All flies were grown on cornmeal/dextrose food in incubators (25°C, 40% relative humidity, 12 : 12 h light : dark cycle). Behaviour experiments were conducted on females 7–8 days post eclosion. For optogenetic experiments, *Gr66a-LexA* and *LexAop-CsChrimson* flies were crossed to produce experimental F1 s. *Gr66a-LexAp65* was constructed using Sequence and Ligation Independent Cloning [24]. The *Gr66a* promoter fragment was the same 1798 bp segment used previously [25] and extended from the translation start site of the *Gr66a* open reading frame to the next upstream gene. This was joined to the start codon of the *LexA::p65* transcriptional activator from pBPLexA::p65Uw [26] in a vector backbone derived from pUASTattB [27] by removing the UAS sites. The construct was integrated into the attP18 site. In experimental groups receiving the optogenetic US, 10 µl of 100 mM all-trans-retinal was applied to the surface of fly food, and flies were housed on this food for at least 48 h. Flies were aspirated directly into the behavioural arenas without anaesthetization.

The assay instrument consisted of 15 linear tunnels with inlets at either end and a vent at the centre (figure 1*a,b*). In each trial, a single fly was placed into each tunnel and allowed to walk freely. Laminar airflow carrying odour stimuli enters the tunnels from either end and meets at the centre, forming a sharp boundary. From there, the odorized air is vented to the room (figure 1*b*). Odorants were generated by flowing clean air over liquid odorants in a series of vials, under the control of solenoids and mass-flow controllers, as described in [8]. Within the arena, flies were presented one pair of odours (e.g. methylcyclohexanol [MCH] versus octanol [OCT] or 1-pentanol versus 2-heptanone [HEPT]). Shock US was delivered via laser-cut grids of indium tin oxide installed on each tunnel floor. Eighty volt direct current pulses from a Grass SD9 Pulse Stimulator (20 Hz for 5 s) were delivered at 10-s intervals. For optogenetic experiments, 626 nm red LEDs were used to activate *CsChrimson* and pulsed at 20 Hz for 3 s with a 5 s interstimulus interval. See assay timelines in figures 1 and 2. Behaviour was recorded using a CMOS camera (Point Grey Firefly MV) with a longpass filter (Kodak Wratten Filter 87C) at 60 Hz. Tracking was performed using custom MATLAB scripts that used two-dimensional cross-correlation for tunnel and initial fly identification, and background subtraction to locate fly centroids.

With this instrument, we implemented three associative conditioning assays. Flies were subject to exactly one of these assays, all of which were conducted in a temperature-controlled environmental chamber in darkness at 25°C and 40% relative humidity. The start of each assay consisted of a 2 min pre-training period in which the CS+ and CS− odorants were present in the tunnels, allowing us to measure flies' untrained odour preference, which was variable as expected [8]. In the 'reversal assay', flies

were first subject to classical pairing of odour with shock and then a reversal pairing in which the CS+ and CS− odorants were swapped. Learned responses were assessed after each pairing in a choice between the CS+ and CS− odours without shock (i.e. a single extinction trial). The 'odour generalization assay' took the form of two reversal assays (using four odorants in total) conducted on successive days, between which the individual identity of flies in this assay was maintained by housing flies in modified 96-well plates (flyPlates, FlySorter, LLC; [28]). The 'US generalization assay' took the form of the reversal assay but replaced electric shock with optogenetic bitter US in the initial classical pairing. See schematic of assay phases in figures 1 and 2.

Individual learning responses were measured by the normalized magnitude of change in occupancy towards the CS− from pre-training to post-training. This metric has a value of 0 if flies exhibit no learning, 1 if they spend all their time post-training in the CS− compartment and −1 if they spend all their time post-training in the CS+ compartment. Normalizing by the pre-training preference response accounts for individual variation in baseline preference [8].

$$\text{learning response} = \begin{cases} \frac{\text{post} - \text{pre}}{1 - \text{pre}} & \text{if, post} - \text{pre} > 0 \\ \frac{\text{post} - \text{pre}}{\text{pre}} & \text{if, post} - \text{pre} \leq 0 \end{cases}$$

The correlation of learning responses across trials was calculated as the Pearson correlation coefficient, and all *p*-values are nominal. Data analysis was performed using custom MATLAB scripts. Raw data and analysis scripts [29] are available at <http://lab.debvort.org/individuality-in-learning> and <https://zenodo.org/record/4458572>.

3. Results

As expected, both training sessions in the reversal assay resulted in significant changes in mean OCT preference across flies (figure 1*d*). This mean change was not observed in control experiments (pairing the US with both CS+ and CS−, backward conditioning or presenting the CS alone; electronic supplementary material, figure S1). However, we also observed individual flies that appeared to not learn on a given trial, with similar preference for OCT pre- and post-training or increased OCT preference even when OCT was the CS+. These observations could reflect statistical noise, rather than individual variation in learning response. To test this, we examined the correlation between the learning response during the classical and reversal phases of the reversal assay. This correlation was positive and significant across individual flies ($r = 0.31$, $p = 0.02$; figure 1*e*), suggesting that individual animals have idiosyncratic learning responses that generalize across the identity of the CS+ odorant. (A few individuals exhibited the same odour preference after both classical and reversal phases, appearing to respond to one association but not the other, a result that could also be explained by these flies having a strong naive preferences that do not change across the assay.) Consistent individual differences in learning response were not correlated with a fly's activity (distance travelled) during the assay or initial odour preference (electronic supplementary material, figure S2), and we found no evidence that variation in learning could be explained by variation in prediction errors (electronic supplementary material, figure S1D).

The observation that individual performance following classical and reversal conditioning is correlated suggests that




Figure 1. Individuality in associative learning. (a) Schematic of the reversal assay. (b) Zoom-in view of the linear behavioural arenas, with odourant flowing into each half. (c) Diagram of training protocol (top). Grey numbers indicate the length in seconds of each stimulus phase. Note that the timing of US delivery differs in the classical and reversal phases. Position in the arena versus time kymographs of three specific flies undergoing conditioning. Magenta and green shading indicate the portions of each arena that are filled with OCT and MCH, respectively. (d) Octanol preference of flies before and after training with MCH as the CS+ (left) and with OCT as the CS+ (right). Points are individual flies. Coloured examples correspond to the individual flies highlighted in (c). p -values reflect paired t -tests. Thick black line represents the mean. (e) Scatterplot of individuals' learning responses for reversal versus classical conditioning trials ($r = 0.31$; $p = 0.02$; $n = 53$). Points are individual flies. Line is the best linear fit and shaded region is the 95% CI of the best-fit line.

learning ability may generalize across sensory channels in flies. To explore this possibility, we implemented the odour generalization assay in which flies were subject to classical and reversal conditioning with 1-pentanol and 2-heptanone

as CS odours, stored for 24 h and subject to classical and reversal conditioning with MCH and OCT (figure 2a). In addition, we substituted optogenetic stimulation of bitter taste neurons as the US instead of electric shock (electronic




Figure 2. Individual learning across odours and US modalities. (a) Schematic of the odour generalization assay. Grey numbers indicate the length in seconds of each stimulus phase. (b) Odorant preference of flies before and after training for all the learning trials of (a). Odorant labels indicate the CS+ in each trial. Points are individual flies. Thick black line represents the mean. p -Values reflect paired t -tests. (c) Correlation matrix for individual fly learning responses for all pairs of learning trials in (a) and (b). x - and y -Axes of scatter subplots correspond respectively to the learning responses of the CS+ condition indicated by the column and row of the matrix. Points are individual flies. Line is the best linear fit, and shaded region is the 95% CI of the best-fit line. (d) Schematic of the US generalization assay. Stimulus phases have the same durations as in figure 1c. (e) Octanol preference of flies before and after training with shock as the US (left) or optogenetic activation of bitter taste neurons (right). Points are individual flies. Thick black line represents the mean. p -values reflect paired t -tests. (f) Scatterplot of learning responses to the shock US trial versus the bitter taste US trial ($r = 0.45$; $p = 0.01$; $n = 47$). Points are individual flies. Line is the best linear fit, and shaded region is the 95% CI of the best-fit line, suggests learning responses to HEPT may not be idiosyncratic.

supplementary material, figure S3). This was done by expressing CsChrimson [16] in bitter taste neurons using a *Gr66a-LexA* driver and exposing flies to 626 nm LED illumination in place of the electric shocks. Replacing shock with bitter taste also let us assess whether individuality and correlation in learned responses to classical and reversal conditioning is US specific. In addition, by looking at learning performance after 24 h, we could assess whether individual variation in learning performance is stable over time. As we saw with shock-odour conditioning, flies subject to optogenetic

bitter-odour conditioning exhibited mean learned avoidance of the CS+ odour (figure 2b; electronic supplementary material, figure S4). We observed significant correlations in individual learning responses among almost all four conditioning variants in this experiment ($0.36 < r < 0.59$; $1.1 \times 10^{-7} < p < 2.8 \times 10^{-3}$; figure 2c). Two exceptions were MCH+ and 2-heptanone+ ($r = 0.06$; $p = 0.61$) and 2-heptanone+ and 1-pentanol+ in the odour generalization assay replicate ($r = 0.16$; $p = 0.17$), for which we have no confident explanation. These results suggest that individuality in learning

performance is largely odour CS- and US-independent and stable over at least 24 h.

A possible explanation of these results is individual variation in US encoding. Flies that receive stronger shocks show stronger learning responses [23], so spontaneous variation in the perception of a US (either shock or bitter taste) may affect the learning responses for many CS. We tested this in the US generalization assay by performing classical and reversal conditioning with OCT and MCH but switching between US within the same animals (figure 2*d*). Both classical and reversal sessions showed significant mean differences in odour preference (figure 2*e*). Comparing across these two aversive US modalities, we observed a positive correlation in learning responses ($r = 0.59$; $p = 8 \times 10^{-6}$; figure 2*d*). This suggests that in addition to generalizing across CS odour identity, individual differences in fly learning performance may generalize across aversive US modalities.

4. Discussion

Using a training instrument that (i) has versatile control over CS and US and (ii) tracks individual learned behaviour, we observed that flies are idiosyncratic in their learning performance in classical conditioning paradigms. Flies that perform well for one CS/US pair tend to perform well for other CS and US, suggesting that individual differences in learning performance generalize across CS odourants and aversive US modalities. We attempted learning experiments in other modalities (colour as a CS and optogenetic activation of sweet-sensitive neurons as a US) but did not see learning responses, likely a technical failure of our assay. Bees were recently shown to be similarly idiosyncratic, but without generalization between visual and olfactory CS modalities [30]. Our results, in a genetic model organism (see also [31]), provide a basis to probe the mechanistic basis of individuality in learning. Specifically, our results hint that the biological basis for such idiosyncrasy in olfactory learning originates more centrally in the brain than sensory circuit elements dedicated to encoding either CS or US. Stochastic physiological variation [1] in neurons mediating aversive US signals in general could account for individual variation that generalizes across aversive US. Such sites would be 'loci of

individuality' [32] for learning performance. Mushroom body dopaminergic neurons [33–37], particularly those in the protocerebral posterior lateral cluster [9], have been shown to mediate multiple aversive US signals including shock [38,39], bitter taste [40,41] and temperature [42]. Mushroom body output [43–46] and intrinsic neurons [47] are also promising candidates. But valence might also be encoded broadly across multiple populations of neurons [33,48,49], including elements in the sensory periphery [48,49]. Circuit elements known to exhibit high developmental stochasticity [8,50,51] may also be loci of individuality. Our results suggest that flies are a promising model for characterizing the circuit basis of individual variation in generalized learning ability, which is evident even among genetically identical individuals reared in the same environment.

Data accessibility. All raw data and analysis scripts are available at <http://lab.debivort.org/individuality-in-learning> and <https://zenodo.org/record/4458572>. Data were collected by the methods described in the manuscript. Raw values are stored in MATLAB files. Analysis was conducted by MATLAB scripts. Data files and analysis scripts are described individually in html read-me files at <http://lab.debivort.org/individuality-in-learning> and <https://zenodo.org/record/4458572> [29].

Authors' contributions. M.S.: conceptualization, data curation, formal analysis, funding acquisition, investigation, methodology, software, validation, visualization, writing—original draft, writing—review and editing; K.S.H.: conceptualization, investigation, methodology, writing—original draft, writing—review and editing; G.T.: resources, writing—original draft, writing—review and editing; B.d.B.: conceptualization, formal analysis, funding acquisition, investigation, methodology, project administration, supervision, visualization, writing—original draft, writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Competing interests. We declare we have no competing interests.

Funding. B.d.B. was supported by a Sloan Research Fellowship, a Klingenstein-Simons Fellowship Award, a Smith Family Odyssey Award, a Harvard/MIT Basic Neuroscience Grant, National Science Foundation grant no. IOS-1557913 and NIH/NINDS grant no. 1R01NS121874-01. G.T. was supported by the Howard Hughes Medical Institute. M.S. was supported by Harvard's Quantitative Biology Initiative.

Acknowledgements. We thank Matt Churgin for help with the manuscript and Ed Soucy, Brett Graham and Joel Greenwood of Harvard's Center for Brain Science Neuroengineering core for help with the stimulator.

References

- Honegger K, de Bivort B. 2018 Stochasticity, individuality and behavior. *Curr. Biol.* **28**, R8–R12. (doi:10.1016/j.cub.2017.11.058)
- Kain JS, Stokes C, de Bivort BL. 2012 Phototactic personality in fruit flies and its suppression by serotonin and white. *Proc. Natl. Acad. Sci. USA* **109**, 19 834–19 839. (doi:10.1073/pnas.1211988109)
- Buchanan SM, Kain JS, de Bivort BL. 2015 Neuronal control of locomotor handedness in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **112**, 6700–6705. (doi:10.1073/pnas.1500804112)
- Kain JS, Zhang S, Akhund-Zade J, Samuel ADT, Klein M, de Bivort BL. 2015 Variability in thermal and phototactic preferences in *Drosophila* may reflect an adaptive bet-hedging strategy. *Evolution* **69**, 3171–3185. (doi:10.1111/evo.12813)
- Todd JG, Kain JS, de Bivort BL. 2017 Systematic exploration of unsupervised methods for mapping behavior. *Phys. Biol.* **14**, 015002. (doi:10.1088/1478-3975/14/1/015002)
- Linneweber GA *et al.* 2020 A neurodevelopmental origin of behavioral individuality in the *Drosophila* visual system. *Science* **367**, 1112–1119. (doi:10.1126/science.aaw7182)
- Ayroles JF, Buchanan SM, O'Leary C, Kutt-Kakaria K, Grenier JK, Clark AG, Hartl DL, de Bivort BL. 2015 Behavioral idiosyncrasy reveals genetic control of phenotypic variability. *Proc. Natl. Acad. Sci. USA* **112**, 6706–6711. (doi:10.1073/pnas.1503830112)
- Honegger KS, Smith MA-Y, Churgin MA, Turner GC, de Bivort BL. 2019 Idiosyncratic neural coding and neuromodulation of olfactory individuality in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **22**, 23 292–23 297. (doi:10.1073/pnas.1901623116)
- Claridge-Chang A, Roorda RD, Vrontou E, Sjulson L, Li H, Hirsh J, Miesenböck G. 2009 Writing memories with light-addressable reinforcement circuitry. *Cell* **139**, 405–415. (doi:10.1016/j.cell.2009.08.034)
- Turner CH. 1907 The homing of ants: an experimental study of ant behavior. *J. Comp. Neurol. Psychol.* **17**, 367–434. (doi:10.1002/cne.920170502)
- Turner CH. 1911 Experiments on pattern-vision of the honey bee. *Biol. Bull.* **21**, 249–264. (doi:10.2307/1536017)

12. Quinn WG, Harris WA, Benzer S. 1974 Conditioned behavior in *Drosophila melanogaster*. *Proc. Natl Acad. Sci. USA* **71**, 708–712. (doi:10.1073/pnas.71.3.708)
13. Tully T, Quinn WG. 1985 Classical conditioning and retention in normal and mutant *Drosophila melanogaster*. *J. Comp. Physiol. A* **157**, 263–277. (doi:10.1007/BF01350033)
14. Davis RL. 1993 Mushroom bodies and *Drosophila* learning. *Neuron* **11**, 1–14. (doi:10.1016/0896-6273(93)90266-T)
15. Rescorla RA. 1967 Pavlovian conditioning and its proper control procedures. *Psychol. Rev.* **74**, 71–80. (doi:10.1037/h0024109)
16. Klapoetke NC *et al.* 2014 Independent optical excitation of distinct neural populations. *Nat. Methods* **11**, 338–346. (doi:10.1038/nmeth.2836)
17. Keene AC, Masek P. 2012 Optogenetic induction of aversive taste memory. *Neuroscience* **222**, 173–180. (doi:10.1016/j.neuroscience.2012.07.028)
18. Izquierdo A, Jentsch JD. 2012 Reversal learning as a measure of impulsive and compulsive behavior in addictions. *Psychopharmacol. (Berl.)* **219**, 607–620. (doi:10.1007/s00213-011-2579-7)
19. Foley BR, Marjoram P, Nuzhdin SV. 2017 Basic reversal-learning capacity in flies suggests rudiments of complex cognition. *PLoS ONE* **12**, e0181749. (doi:10.1371/journal.pone.0181749)
20. Mancini N, Hranova S, Weber J, Weiglein A, Schleyer M, Weber D, Thum AS, Gerber B. 2019 Reversal learning in *Drosophila* larvae. *Learn. Mem.* **26**, 424–435. (doi:10.1101/lm.049510.119)
21. Ren Q, Li H, Wu Y, Ren J, Guo A. 2012 A GABAergic inhibitory neural circuit regulates visual reversal learning in *Drosophila*. *J. Neurosci.* **32**, 11 524–11 538. (doi:10.1523/JNEUROSCI.0827-12.2012)
22. Wu Y, Ren Q, Li H, Guo A. 2012 The GABAergic anterior paired lateral neurons facilitate olfactory reversal learning in *Drosophila*. *Learn. Mem.* **19**, 478–486. (doi:10.1101/lm.025726.112)
23. Beck CDO, Schroeder B, Davis RL. 2000 Learning performance of normal and mutant *Drosophila* after repeated conditioning trials with discrete stimuli. *J. Neurosci.* **20**, 2944–2953. (doi:10.1523/JNEUROSCI.20-08-02944.2000)
24. Li MZ, Elledge SJ. 2007 Harnessing homologous recombination *in vitro* to generate recombinant DNA via SLIC. *Nat. Methods* **4**, 251–256. (doi:10.1038/nmeth1010)
25. Dunipace L, Meister S, McNealy C, Amrein H. 2001 Spatially restricted expression of candidate taste receptors in the *Drosophila* gustatory system. *Curr. Biol.* **11**, 822–835. (doi:10.1016/S0960-9822(01)00258-5)
26. Pfeiffer BD, Ngo T-TB, Hibbard KL, Murphy C, Jenett A, Truman JW, Rubin GM. 2010 Refinement of tools for targeted gene expression in *Drosophila*. *Genetics* **186**, 735–755. (doi:10.1534/genetics.110.119917)
27. Bischof J, Maeda RK, Hediger M, Karch F, Basler K. 2007 An optimized transgenesis system for *Drosophila* using germ-line-specific phiC31 integrases. *Proc. Natl Acad. Sci. USA* **104**, 3312–3317. (doi:10.1073/pnas.0611511104)
28. Alisch T, Crall JD, Kao AB, Zucker D, de Bivort BL. 2018 MAPLE (modular automated platform for large-scale experiments), a robot for integrated organism-handling and phenotyping. *Elife* **7**, e37166. (doi:10.7554/eLife.37166)
29. Smith MA-Y, Honegger K, Turner G, de Bivort BL. 2021 Data from: Idiosyncratic learning performance in flies generalizes across modalities. Zenodo Digital Repository. (doi:10.5281/zenodo.4458572)
30. Finke V, Baracchi D, Giurfa M, Scheiner R, Avarguès-Weber A. 2021 Evidence of cognitive specialization in an insect: proficiency is maintained across elemental and higher-order visual learning but not between sensory modalities in honey bees. *J. Exp. Biol. jeb.* **224**, jeb242470. (doi:10.1242/jeb.242470)
31. Lesar A, Tahir J, Wolk J, Gershov M. 2021 Switch-like and persistent memory formation in individual *Drosophila* larvae. *Elife* **10**, e70317. (doi:10.7554/eLife.70317)
32. Skutt-Kakaria K, Reimers P, Currier TA, Werkhoven Z, de Bivort BL. 2020 A neural circuit basis for context-modulation of individual locomotor behavior. *BioRxiv.* (doi:10.1101/797126)
33. Kim Y-C, Lee H-G, Han K-A. 2007 D1 dopamine receptor dDA1 is required in the mushroom body neurons for aversive and appetitive learning in *Drosophila*. *J. Neurosci.* **27**, 7640–7647. (doi:10.1523/JNEUROSCI.1167-07.2007)
34. Siju KP, Stih V, Aimon S, Gjorgjieva J, Portugues R, Grunwald Kadow I. 2020 Valence and state-dependent population coding in dopaminergic neurons in the fly mushroom body. *Curr. Biol.* **30**, 2104–2115. (doi:10.1016/j.cub.2020.04.037)
35. Hige T, Aso Y, Modi MN, Rubin GM, Turner GC. 2015 Heterosynaptic plasticity underlies aversive olfactory learning in *Drosophila*. *Neuron* **88**, 985–998. (doi:10.1016/j.neuron.2015.11.003)
36. Liu C *et al.* 2012 A subset of dopamine neurons signals reward for odour memory in *Drosophila*. *Nature* **488**, 512–516. (doi:10.1038/nature11304)
37. Burke CJ *et al.* 2012 Layered reward signalling through octopamine and dopamine in *Drosophila*. *Nature* **492**, 433–437. (doi:10.1038/nature11614)
38. Mao Z, Davis RL. 2009 Eight different types of dopaminergic neurons innervate the *Drosophila* mushroom body neuropil: anatomical and physiological heterogeneity. *Front. Neural. Circuits* **3**, 5. (doi:10.3389/neuro.04.005.2009)
39. Aso Y, Herb A, Ogueta M, Siwanowicz I, Templier T, Friedrich AB, Ito K, Scholz H, Tanimoto H. 2012 Three dopamine pathways induce aversive odor memories with different stability. *PLoS Genet.* **8**, e1002768. (doi:10.1371/journal.pgen.1002768)
40. Das G, Klappenbach M, Vrontou E, Perisse E, Clark C, Burke C, Waddell S. 2014 *Drosophila* learn opposing components of a compound food stimulus. *Curr. Biol.* **24**, 1723–1730. (doi:10.1016/j.cub.2014.05.078)
41. Masek P, Worden K, Aso Y, Rubin GM, Keene AC. 2015 A dopamine-modulated neural circuit regulating aversive taste memory in *Drosophila*. *Curr. Biol.* **25**, 1535–1541. (doi:10.1016/j.cub.2015.04.027)
42. Galili DS, Dilla KV, Lüdke A, Friedrich AB, Yamagata N, Wong JYH, Ho CH, Szyszka P, Tanimoto H. 2014 Converging circuits mediate temperature and shock aversive olfactory conditioning in *Drosophila*. *Curr. Biol.* **24**, 1712–1722. (doi:10.1016/j.cub.2014.06.062)
43. Aso Y *et al.* 2014 Mushroom body output neurons encode valence and guide memory-based action selection in *Drosophila*. *Elife* **3**, e04580. (doi:10.7554/eLife.04580)
44. Séjourné J *et al.* 2011 Mushroom body efferent neurons responsible for aversive olfactory memory retrieval in *Drosophila*. *Nat. Neurosci.* **14**, 903–910. (doi:10.1038/nn.2846)
45. Oswald D, Felsenberg J, Talbot CB, Das G, Perisse E, Huetteroth W, Waddell S. 2015 Activity of defined mushroom body output neurons underlies learned olfactory behavior in *Drosophila*. *Neuron* **86**, 417–427. (doi:10.1016/j.neuron.2015.03.025)
46. Bouzaiane E, Trannoy S, Scheunemann L, Plaçais P-Y, Preat T. 2015 Two independent mushroom body output circuits retrieve the six discrete components of *Drosophila* aversive memory. *Cell Rep.* **11**, 1280–1292. (doi:10.1016/j.celrep.2015.04.044)
47. Perisse E, Oswald D, Barnstedt O, Talbot CB, Huetteroth W, Waddell S. 2016 Aversive learning and appetitive motivation toggle feed-forward inhibition in the *Drosophila* mushroom body. *Neuron* **90**, 1086–1099. (doi:10.1016/j.neuron.2016.04.034)
48. Yu D, Ponomarev A, Davis R. 2004 Altered representation of the spatial code for odors after olfactory classical conditioning. *Neuron* **42**, 437–449. (doi:10.1016/S0896-6273(04)00217-X)
49. Rath L, Giovanni Galizia C, Szyszka P. 2011 Multiple memory traces after associative learning in the honey bee antennal lobe. *Eur. J. Neurosci.* **34**, 352–360. (doi:10.1111/j.1460-9568.2011.07753.x)
50. Caron SJ, Ruta V, Abbott LF, Axel R. 2013 Random convergence of olfactory inputs in the *Drosophila* mushroom body. *Nature* **497**, 113–117. (doi:10.1038/nature12063)
51. Chou Y, Spletter M, Yaksi E, Leong J, Wilson R, Luo L. 2010 Diversity and wiring variability of olfactory local interneurons in the *Drosophila* antennal lobe. *Nat. Neurosci.* **13**, 439–449. (doi:10.1038/nn.2489)
52. Vogt K, Yarali A, Tanimoto H. 2015 Reversing stimulus timing in visual conditioning leads to memories with opposite valence in *Drosophila*. *PLoS ONE* **10**, e0139797. (doi:10.1371/journal.pone.0139797)