

BMJ Open Postdischarge mortality in children with acute infectious diseases: derivation of postdischarge mortality prediction models

M O Wiens,¹ E Kumbakumba,² C P Larson,³ J M Ansermino,⁴ J Singer,^{1,5} N Kisson,⁶ H Wong,¹ A Ndamira,² J Kabakyenga,⁷ J Kiwanuka,² G Zhou⁸

To cite: Wiens MO, Kumbakumba E, Larson CP, *et al.* Postdischarge mortality in children with acute infectious diseases: derivation of postdischarge mortality prediction models. *BMJ Open* 2015;**5**:e009449. doi:10.1136/bmjopen-2015-009449

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2015-009449>).

Received 21 July 2015
Accepted 16 October 2015



CrossMark

For numbered affiliations see end of article.

Correspondence to

Dr Matthew Wiens;
mowiens@outlook.com

ABSTRACT

Objectives: To derive a model of paediatric postdischarge mortality following acute infectious illness.

Design: Prospective cohort study.

Setting: 2 hospitals in South-western Uganda.

Participants: 1307 children of 6 months to 5 years of age were admitted with a proven or suspected infection. 1242 children were discharged alive and followed up 6 months following discharge. The 6-month follow-up rate was 98.3%.

Interventions: None.

Primary and secondary outcome measures: The primary outcome was postdischarge mortality within 6 months following the initial hospital discharge.

Results: 64 children died during admission (5.0%) and 61 died within 6 months of discharge (4.9%). Of those who died following discharge, 31 (51%) occurred within the first 30 days. The final adjusted model for the prediction of postdischarge mortality included the variables mid-upper arm circumference (OR 0.95, 95% CI 0.94 to 0.97, per 1 mm increase), time since last hospitalisation (OR 0.76, 95% CI 0.61 to 0.93, for each increased period of no hospitalisation), oxygen saturation (OR 0.96, 95% CI 0.93 to 0.99, per 1% increase), abnormal Blantyre Coma Scale score (OR 2.39, 95% CI 1.18 to 4.83), and HIV-positive status (OR 2.98, 95% CI 1.36 to 6.53). This model produced a receiver operating characteristic curve with an area under the curve of 0.82. With sensitivity of 80%, our model had a specificity of 66%. Approximately 35% of children would be identified as high risk (11.1% mortality risk) and the remaining would be classified as low risk (1.4% mortality risk), in a similar cohort.

Conclusions: Mortality following discharge is a poorly recognised contributor to child mortality. Identification of at-risk children is critical in developing postdischarge interventions. A simple prediction tool that uses 5 easily collected variables can be used to identify children at high risk of death after discharge. Improved discharge planning and care could be provided for high-risk children.

Strengths and limitations of this study

- The primary strengths of this study are prospective and rigorous data collection, and near-complete follow-up.
- Further strengths include the derivation of multiple similar models to allow prediction in circumstances where not all variables may be available.
- Regression models can easily be incorporated into a mobile health-based tool for simple and rapid prediction by health workers.
- The primary limitations of this study are relatively few outcomes, and lack of external validity. Despite few outcomes, our models performed quite well.
- These limitations highlight the need for further research on this important but neglected topic.
- The identification of high risk does not imply that risk can be reduced. Further work is needed on the development of postdischarge interventions to reduce this burden.

BACKGROUND

Acute infectious diseases continue to be the most important contributor to the 6 million children younger than 5 years who die every year, particularly in Africa.¹ It is widely accepted that as a global community we have fallen short in reducing under-5 mortality, as demonstrated by the fact that most developing countries, especially those in sub-Saharan Africa will not achieve the fourth millennium development goal of a two-thirds reduction in child mortality.² An important but neglected contributor to infectious disease-related mortality is the vulnerable period following hospital discharge.

A recent systematic review of paediatric studies assessing postdischarge mortality in resource-poor countries found that post-discharge mortality often exceeds in-hospital

mortality.³ Thus, attention to at-risk populations post discharge is sorely needed. However, while several factors were consistently found to be associated with mortality following discharge, including malnutrition, HIV and severe pneumonia, easy identification is essential in order to develop targeted postdischarge interventions. Ideally, the unacceptably high risk of morbidity and mortality following discharge suggests that all children should be afforded follow-up care. However, significant resource constraints in the countries most affected by this issue preclude any significant intervention for all discharged children. Therefore, the ability to quickly and effectively identify at-risk children would be an invaluable step towards the implementation of life-saving postdischarge interventions. An important and easily identified dichotomy among hospital admissions are infectious diseases and non-infectious disease-related admissions such as trauma, cancer and congenital diseases. Although further divisions based on aetiology of infection or an underlying risk factor, such as malnutrition or HIV status, may be an attractive approach in risk stratification, significant difficulties in disease definitions and often overlapping risks makes this approach very difficult. The development of a robust yet simple risk-scoring algorithm could significantly advance a systematic and evidence-based approach in postdischarge care.

The purpose of this study was to derive simple prediction models that could efficiently stratify children according to postdischarge mortality risk.

METHODS

Population

Mbarara, a city of approximately 195 000, is the largest city in the South-western region of Uganda. This study was conducted at two hospitals in Mbarara. The Mbarara Regional Referral Hospital (MRRH) is the main referral hospital in South-western Uganda. It is a public hospital funded by the Uganda Ministry of Health. MRRH is associated with the Mbarara University of Science and Technology, and is a primary training site for its healthcare graduates. The paediatric ward admits approximately 5000 patients per year. The Holy Innocents Children's Hospital (HICH) is a faith-based children's hospital offering subsidised fee-for-service for outpatient and in-patient care in Mbarara. The HICH admits approximately 2500 patients per year.

This was a prospective observational study conducted between March 2012 and December 2013. This study was approved by the institutional review boards at the University of British Columbia (Canada) and the Mbarara University of Science and Technology (Uganda), as well as the Uganda National Council for Science and Technology and Office of the President. This study was voluntary and written informed consent was provided by a parent or guardian of all children who were enrolled.

Eligibility

All children aged 6 months to 5 years who were admitted with a proven or suspected infection were eligible for enrolment. The upper age limit was chosen to coincide with the under-5 target group of the millennium development goals. The lower age limit was chosen for logistic (census enrolment with limited research staff) and statistical considerations (group homogeneity). Participants already enrolled in the study were not eligible to be enrolled during subsequent admissions.

Study procedure

Following enrolment, a research nurse obtained and recorded clinical signs including a 1 min respiratory rate, blood pressure (automated), axillary temperature, Blantyre Coma Scale (BCS) score, and by using the Phone Oximeter,⁴ the 1 min photoplethysmogram (PPG), blood oxygen saturation (SpO₂) and heart rate. Anthropometric data (height, weight, mid-upper arm circumference (MUAC)) were also measured and recorded. Age-dependent demographic variables collected at enrolment were converted to age-corrected z-scores according to the WHO Child Growth Standards.⁵ The age-corrected heart rate and respiratory rate z-scores were obtained by standardising the raw measurements using the median and SD values provided by Fleming *et al.*⁶ The age-corrected z-scores for systolic blood pressure were calculated using participants' height, according to the procedures previously described.⁷

A blood sample was taken for measurement of haemoglobin, HIV and a malaria blood smear (microscopy). HIV status was determined using the national rapid diagnostic test serial algorithm.⁸ All positive tests on the Determine Antibody Test were confirmed by a separate test (UniGold). Children under 12 months of age with a positive test were confirmed using PCR. Haemoglobin was measured on a Beckman Coulter Ac.T Hematology Analyzer.

An interview was conducted with the participant's parent/guardian and information about previous admissions, distance from health facility, transportation costs, bed-net use, maternal education, maternal age, maternal HIV status, history of sibling deaths and drinking water safety were elicited. Participants received routine care during their hospital stay and were discharged at the discretion of the treating medical team. The discharge status of all enrolled participants was recorded as death, referral, discharged alive, and discharged against medical advice. The diagnoses made by the medical team were also recorded. On discharge, families with active telephone lines were contacted at months 2 and 4 to determine the vital status of the child. Families with no telephone access received in-person follow-up by a field officer. At approximately 6 months following discharge, all participants received in-person follow-up. In addition to postdischarge vital status, health seeking and

rehospitalisations since the initial discharge were also recorded.

Study data were collected and managed using REDCap electronic data capture tools hosted at the Child and Family Research Institute, Vancouver, Canada.⁹ REDCap (Research Electronic Data Capture) is a secure, web-based application designed to support data capture for research studies and provides: (1) an intuitive interface for validated data entry; (2) audit trails for tracking data manipulation and export procedures; (3) automated export procedures for seamless data downloads to common statistical packages and (4) procedures for importing data from external sources.

Candidate predictor variables were derived using a two-round modified Delphi approach. Briefly, 23 experts in relevant disciplines were solicited to complete an online survey and provide feedback on an initial list of proposed predictors. Predictors were evaluated on considerations of utility as predictors, availability, cost and resource-related applicability. Experts were asked to provide additional potential variables which were then evaluated during a second round of surveys. Data was evaluated by the research team and a final list of candidate predictor variables for modelling was then determined.¹⁰

Outcomes

The primary outcome was postdischarge mortality at any time during the 6-month postdischarge period.

Sample size

For the derivation of prediction models, standard calculations of sample size do not apply since these calculations do not account for the model development process (ie, selection of variables and the optimisation to achieve specified sensitivity and specificity cut-offs). For this study, we determined the sample size needed to validate the derived model and plan to use an equal number of patients for the derivation phase. For the validation study, assuming that the derived model achieves a sensitivity of 85% with at least 50% specificity, 100 events, corresponding to a total sample of approximately 1000 live-discharges (assuming a postdischarge mortality rate of 10%), would be needed to obtain 80% power for ensuring that the lower 95% confidence limit on sensitivity will be at least 75%. Since resources are scarce, a higher sensitivity at the expense of specificity would further limit practical application of such a model. An interim analysis of the study showed that the postdischarge mortality rate would likely not exceed 5% and enrolment was stopped when 1307 participants were enrolled.

Statistical analysis

All variables were assessed using univariate logistic regression to determine their level of association with the primary outcome. Continuous variables were assessed for model fit using the Hosmer-Lemeshow

test.¹¹ Missing data was imputed by the method of multivariate imputation using chained equations.¹² Following univariate analysis, candidate models were generated using a stepwise selection procedure minimising Akaike's Information Criterion (AIC). This method is considered asymptotically equivalent to cross-validation and bootstrapping.^{13 14} All models generated in this sequence having AIC values within 10% of the lowest value were considered as reasonable candidates. The final selection of a model was judged on model parsimony (the simpler the better), availability of the predictors (with respect to minimal resources and cost), and the attained sensitivity (with at least 50% specificity). All analyses were conducted using SAS V.9.3 (Carey, North Carolina, USA) and R 3.1.3 (Vienna, Austria; <http://www.R-project.org>). Additional models were created using the above process but with the absence of key variables used in deriving the primary model, including a model not including any variables likely to change over the course of admission. This was done to increase application in a variety of settings where certain variables may not be available.


RESULTS

During the period of study, 1822 participants were screened for eligibility, of which 516 (28%) were excluded. Reasons for exclusion included isolated malnutrition (n=192), readmission of previously enrolled participant (n=51), refusal of consent (n=22), cardiac disease (n=19), poisoning/drug reaction (n=19) and cancer (n=12) as well as a plethora of other non-infectious admissions (n=165). In total, 1307 participants admitted with a presumed or proven infection were enrolled at the time of their admission. During the course of admission, 64 (5.1%) participants died, and 1242 (94.9%) were discharged alive ([figure 1](#)). Among the children discharged, 54% were male, and the median age was 18.1 months (IQR 10.8–34.6). Pneumonia, malaria and gastroenteritis were the most common clinical discharge diagnoses and were present in 31%, 50%, and 8% of discharged participants, respectively. According to anthropometric variables collected at admission, 30% of participants were considered underweight (weight for age z-score <−2), 35% were considered wasted (weight for height/length z-score <−2), and 29% were considered stunted (height/length for age z-score <−2) ([table 1](#)). Missing observations were minimal ([table 2](#)).

Postdischarge mortality

The rate of successful follow-up during the postdischarge period was 98.3%, with only six participants receiving no follow-up during this period. Overall, 61 (4.9%) children died following discharge. Of those who died, the median time to death was 30 days (IQR 7–81). Of the 61 deaths, 41 (67%) occurred outside of a hospital and 20 (33%) occurred during a hospital

Figure 1 Consort diagram of study flow.



readmission. Thirty variables were tested for univariate associations with postdischarge mortality (table 2). MUAC was the variable with the highest area under the receiver operating characteristic (ROC) curve, 0.76 (95% CI 0.70 to 0.83), and was highly significant ($p < 0.0001$). Other anthropometric variables, including weight for age z-score, length/height for age z-score, and weight for length/height z-score were also highly associated with postdischarge mortality but had much lower areas under the ROC curve. Oxygen saturation was the most predictive of the non-anthropometric variables, with an area under the ROC curve of 0.65 (95% CI 0.57 to 0.73), followed by age and parasitaemia with areas under the ROC curve of 0.64 (95% CI 0.56 to 0.70) and 0.60 (95% CI 0.55 to 0.65), respectively. Other variables achieving statistical significance, but showing lower areas under the ROC curve, included systolic blood pressure, axillary temperature, HIV status, abnormal BCS (yes vs no), duration of illness prior to admission >7 days and time since last hospitalisation (analysed as continuous variable and ordered as <7 days, 7 to 30 days, 30 days to 1 year, >1 year, and never). Haemoglobin level, history of sibling deaths, maternal HIV status, maternal education and distance from admitting health facility were not associated with post-discharge mortality in the univariate analysis.

Multivariate prediction models

One primary model and three alternate models of equal sensitivity were developed for the prediction of 6-month postdischarge mortality (table 3). Two alternate models were developed while systematically excluding oxygen saturation, and HIV status, respectively, since these may not be routinely available in all clinical settings. A fourth model was developed excluding variables most likely to change over the course of admission (ie, clinical variables), giving the model utility for variables collected at any time throughout the hospital stay. The primary model included MUAC in mm, oxygen saturation (SpO_2) at admission (per cent), time since previous hospitalisation, the presence of abnormal BCS score at admission, and HIV status. The area under the ROC curve was 0.82 (95% CI 0.76 to 0.87) (figure 2). The model, at a cut-off of >80% sensitivity, had a final sensitivity of 82% (95% CI 0.75% to 0.87%) and a specificity of 66% (95% CI 64% to 69%). In a population similar to this model derivation cohort, we would expect the positive predictive value to be 11.1% and the negative predictive value to be 98.6% (table 4). The final model equation for the primary model was: $\text{logit}(p) = 7.71 + (-0.041; \text{MUAC}) + (-0.041; \text{SpO}_2) + (-0.28; \text{time period since last hospitalisation}) + (1.09; \text{HIV positive}) + (0.87; \text{BCS} < 5)$.

Table 1 General characteristics of discharged participants (N=1242)

Characteristic	Frequency (%)
Age, months	
<12	378 (30)
12–24 m	379 (30)
24–36 m	198 (16)
36–48 m	150 (12)
>48 m	138 (11)
Male sex	682 (55)
Length of stay, days	
<3	487 (39)
3–5	487 (39)
6–10	173 (14)
>10	96 (8)
Discharge AMA	120 (10)
Diagnoses	
Pneumonia	390 (31)
Clinical malaria	621 (50)
Parasitaemia	418 (34)
Gastroenteritis	96 (8)
SSTI	7 (0.5)
Meningitis	32 (2.5)
Tuberculosis	17 (1.4)
Measles	15 (1.2)
Comorbidities	
HIV	58 (4.7)
Sickle cell	7 (0.5)
Tuberculosis	21 (1.7)
Admission anthropometric characteristics	
Underweight (WAZ<−2)	347 (30)
Severe underweight (WAZ<−3)	188 (15)
Wasting (WHZ<−2)	436 (35)
Severe wasting (WHZ<−3)	232 (17)
Stunting (HAZ<−2)	357 (29)
Severe stunting (HAZ<−3)	187 (15)
MUAC<125 mm	183 (15)
MUAC<115 mm	96 (7.7)

AMA, against medical advice; HAZ, height/length for age z-score; MUAC, mid-upper arm circumference; SSTI, skin and soft tissue infection; WAZ, weight for age z-score; WHZ, weight for height/length z-score.

Model 2 excluded oxygen saturation (table 3). The final model included MUAC, time since last hospitalisation, HIV status and the presence of an abnormal BCS score. The area under the ROC curve was 0.81 (95% CI 0.75 to 0.87). This model had a sensitivity of 80% (95% CI 70% to 90%) and specificity of 68% (95% CI 65% to 70%), and would generate a positive and negative predictive value of 11.3% and 98.5%, respectively, in a population similar to the derivation cohort.

The third model excluded HIV status (table 3). This model had a final area under the ROC curve of 0.80 (95% CI 0.74 to 0.86), and a sensitivity of 80% (95% CI 70% to 90%) and specificity of 63% (95% CI 60% to 66%). The positive and negative predictive values were 10.2% and 98.4%, respectively.

The final model excluded all time-changing clinical parameters (eg, vital signs, SpO₂, coma score, etc) so as

to be applicable to data collected at any time during admission, including discharge. This model contained only three variables, MUAC, HIV status and the most recent hospitalisation. This model achieved good performance characteristics including an area under the curve (AUC) of 0.80 (95% CI 0.73 to 0.86). The sensitivity was 82% (95% CI 72% to 92%) and the specificity was 61% (95% CI 59% to 64%), and the positive and negative predictive values were 9.9% and 98.5%, respectively.

DISCUSSION

This study represents the first systematic approach to the development of a simple risk-scoring algorithm for post-discharge mortality following admission for an acute infectious illness using prospectively collected data. The variables used in these models are easy to collect and include MUAC, oxygen saturation, BCS score, time since last hospitalisation, and HIV status. Four prediction models were developed to ensure its effective application in a variety of clinical circumstances. All four models had very similar performance characteristics with the most parsimonious model including only MUAC, HIV status and time since last hospitalisation, with only marginally lower AUC than the full model with five variables. The models which were developed use only variables collected at admission and can, therefore, easily be incorporated into the discharge planning process during the hospital stay. Using these models, the identification of at-risk children would ensure that most children likely to die in the postdischarge period (about 80%) would be identified. These children have an average mortality risk of approximately over 10%, justifying the exploration of potentially life-saving interventions. Interventions found to be effective could likely be brought to scale without inordinately burdening the already stressed healthcare systems.

The development and implementation of predictive models into routine clinical care is not common in resource-poor countries. The high prevalence of overlapping diseases (such as pneumonia, malaria and malnutrition), and the difficulty in creating reliable diagnostic algorithms to identify eligible populations cause significant difficulty in the application of disease-specific models. To create models with uptake potential, these would need to be linked with existing clinical practices and resources, and would also require a shift in how infectious illness is viewed—not as an episodic disease but as a continuum beyond the acute episode. The Integrated Management of Childhood Illness (IMCI), while not a predictive tool per se, is an algorithm-based approach for the diagnosis and management of acute infectious illnesses.¹⁵ IMCI has seen significant uptake in many countries throughout sub-Saharan Africa, and has provided a systematic approach to the care of children within health facilities. More importantly, it has been shown to improve care in the regions where it has

Table 2 Univariate analysis of potential predictor variables

Variable	Missing observation	OR (95% CI)	AUC (95% CI)	p Value
Male sex	0	0.90 (0.54 to 1.51)	0.51 (0.45 to 0.58)	0.700
Age (months)	3	0.97 (0.97 to 0.97)	0.64 (0.56 to 0.70)	0.003
MUAC (mm)	14	0.97 (0.96 to 0.98)	0.76 (0.70 to 0.83)	<0.001
Weight for age z-score	5	0.66 (0.57 to 0.76)	0.68 (0.60 to 0.76)	<0.001
Weight for length/height z-score	15	0.81 (0.72 to 0.91)	0.62 (0.55 to 0.70)	<0.001
Length/height for age z-score	16	0.79 (0.70 to 0.89)	0.63 (0.56 to 0.71)	<0.001
HR for age z-score	3	0.86 (0.74 to 0.99)	0.61 (0.53 to 0.69)	0.036
HR (raw)	0	1.00 (0.99 to 1.01)	0.53 (0.47 to 0.62)	0.728
RR for age z-score	3	0.99 (0.92 to 1.06)	0.53 (0.45 to 0.60)	0.747
RR (raw)	0	1.01 (1.00 to 1.03)	0.57 (0.50 to 0.63)	0.100
SBP z-score	21	0.94 (0.79 to 1.12)	0.50 (0.45 to 0.61)	0.526
SBP (raw)	6	0.98 (0.96 to 1.00)	0.58 (0.50 to 0.66)	0.018
DBP (raw)	6	0.99 (0.97 to 1.01)	0.55 (0.50 to 0.65)	0.255
Temperature (transformed)	0	1.02 (0.90 to 1.16)	0.51 (0.45 to 0.57)	0.789
Temperature (raw)	0	0.76 (0.62 to 0.93)	0.58 (0.50 to 0.65)	0.007
SpO ₂ (raw)	13	0.94 (0.92 to 0.96)	0.65 (0.57 to 0.73)	<0.001
SpO ₂ (transformed)	13	1.04 (1.02 to 1.05)	0.65 (0.57 to 0.73)	<0.001
HIV positive (vs negative)	25	5.21 (2.55 to 10.65)	0.57 (0.52 to 0.62)	<0.001
Haemoglobin (g/dL)	10	0.95 (0.87 to 1.03)	0.56 (0.49 to 0.63)	0.227
Blantyre Coma Scale <5 (vs 5)	0	2.40 (1.27 to 4.57)	0.56 (0.50 to 0.61)	0.007
Positive blood smear (vs negative)	11	0.33 (0.16 to 0.68)	0.60 (0.55 to 0.65)	0.002
Illness >7 days prior to admission	1	0.50 (0.30 to 0.83)	0.58 (0.52 to 0.65)	0.008
Time since last hospitalisation*	3	0.75 (0.62 to 0.90)	0.59 (0.52 to 0.67)	0.003
Sibling deaths	0	1.54 (0.89 to 2.65)	0.55 (0.48 to 0.61)	0.121
Number of children in family	2	1.02 (0.92 to 1.13)	0.50 (0.43 to 0.58)	0.750
Boil all drinking water	0	0.82 (0.47 to 1.42)	0.52 (0.46 to 0.58)	0.471
Maternal age (years)	0	1.00 (0.97 to 1.04)	0.52 (0.41 to 0.57)	0.892
Maternal HIV (ref: negative)				
HIV positive, n=142	0	1.79 (0.87 to 3.67)	0.54 (0.48 to 0.61)	0.113
HIV status unknown, n=220	0	1.27 (0.64 to 2.52)		0.499
Maternal education (ref: <primary 3)				
Primary 3–7, n=630	0	1.18 (0.62 to 2.23)	0.54 (0.50 to 0.63)	0.619
Some secondary, n=269	0	0.72 (0.31 to 1.70)		0.457
Postsecondary, n=93	0	1.18 (0.41 to 3.36)		0.762
Bed-net use (ref: never)				
Sometimes	0	1.00 (0.48 to 2.09)	0.52 (0.45 to 0.59)	0.996
Always	0	0.85 (0.46 to 1.58)		0.612
Distance from hospital (ref: <30 min)				
30–60	0	0.71 (0.31 to 1.64)	0.56 (0.49 to 0.62)	0.421
>60	0	1.30 (0.70 to 2.41)		0.401

*Ordered as <7 days, 7 to 30 days, 30 days to 1 year, >1 year and never (analysed as continuous and coded and 1–5, respectively).

AUC, area under the curve; DBP, diastolic blood pressure; HR, heart rate; MUAC, mid-upper arm circumference; RR, respiratory rate; SBP, systolic blood pressure.

been implemented.¹⁶ However, the IMCI does not address the important issue of postdischarge vulnerability and therefore, fails to provide any guidance beyond the period of acute illness in the hospital, even though the postdischarge period will claim as many lives as the acute hospital period. The integration of a postdischarge risk score into IMCI could begin to address this need.

This study is participant to several limitations. A primary limitation of this study is the relatively low number of outcomes observed. Although our initial sample size estimates were to observe 100 outcomes, we only observed 61. Our comprehensive follow-up of participants ensured that missed outcomes are unlikely.

Further, the performance of our model was good, with the lower limits of the calculated 95% CIs for AUC, sensitivity and specificity remaining in an acceptable range. A further limitation is the lack of external validity. While our research sites represented the typical East African context, further research is required to ensure the validity of these models elsewhere, especially in areas with significant differences in the distribution of important diseases such as malaria, diarrhoea and pneumonia, and malnutrition. A limitation to application of the prediction models developed is that the risk score is based on a regression equation and cannot be easily computed without the assistance of a computer or similar device.

Table 3 Models developed for prediction of 6-month postdischarge mortality

Variable	Regression estimate	p Value	OR (95% CI)
Model 1—primary model, intercept=7.7172			
MUAC	−0.0462	<0.0001	0.95 (0.94 to 0.97)
SpO ₂	−0.0411	0.0029	0.96 (0.93 to 0.99)
Time since last hospitalisation	−0.2775	0.0085	0.76 (0.62 to 0.93)
HIV positive	1.0915	0.0064	2.98 (1.36 to 6.53)
Abnormal BCS score	0.8723	0.0150	2.39 (1.18 to 4.83)
Model 2—model without SpO ₂ , intercept=4.4538			
MUAC	−0.0505	<0.0001	0.95 (0.94 to 0.97)
Time since last hospitalisation	−0.2503	0.0153	0.78 (0.64 to 0.95)
HIV positive	1.0902	0.0061	2.98 (1.37 to 6.48)
Abnormal BCS score	1.0664	0.0022	2.91 (1.47 to 5.75)
Model 4—model without HIV, intercept=8.2813			
MUAC	−0.0492	<0.0001	0.95 (0.94 to 0.97)
SpO ₂	−0.0412	0.0027	0.96 (0.93 to 0.99)
Time since last hospitalisation	−0.2870	0.0058	0.75 (0.61 to 0.92)
Abnormal BCS score	0.8040	0.0248	2.23 (1.11 to 4.51)
Model 4—model without clinical variables, intercept=4.4511			
MUAC	−0.0492	<0.0001	0.95 (0.94 to 0.97)
HIV positive	1.0143	0.0108	2.76 (1.26 to 6.01)
Time since last hospitalisation	−0.2458	0.0164	0.78 (0.64 to 0.96)

BCS, Blantyre Coma Scale; MUAC, mid-upper arm circumference.

However, with the increasing prevalence of mobile phones in developing countries, health interventions are increasingly focused on utilising the computational power of mobile phones to implement life-saving technology. Several important health interventions use mobile technology to improve care.^{17–19}

It is clear that malnutrition plays a major role in post-discharge mortality. MUAC provided a significant proportion of the predictive power in our models by providing an AUC of 0.76, only 7% lower than the final

full model. No models meeting our prespecified criteria could be developed without the use of any anthropomorphic measure. The importance of malnutrition has also been clearly demonstrated in other studies of post-discharge mortality.^{20–22} Although first described over 50 years ago, environmental enteropathy (also called tropical enteropathy or environmental enteric dysfunction) has received significant attention in recent years. It has been suggested that changes in the gut microbiome and the small intestinal wall (flattened villi,

Figure 2 Performance of the primary prediction model derived with data from admission (AUC, area under the curve; NPV, negative predictive value; PPV, positive predictive value; ROC, receiver operating characteristic; Sens, sensitivity; Spec, specificity).




Table 4 Model characteristics at probability cut-offs ensuring model sensitivity of >80%

Model	AUC (95% CI)	Probability cut-off	Sensitivity (95% CI)	Specificity (95% CI)	PPV	NPV
1	0.82 (0.75 to 0.87)	0.035	82.0 (72.3 to 91.6)	66.2 (63.5 to 68.9)	11.1	98.6
2	0.81 (0.75 to 0.87)	0.040	80.3 (70.4 to 90.3)	67.5 (64.8 to 70.2)	11.3	98.5
3	0.80 (0.74 to 0.86)	0.031	80.3 (70.4 to 90.3)	63.4 (60.7 to 66.2)	10.2	98.4
4	0.80 (0.73 to 0.86)	0.035	82.0 (72.3 to 91.6)	61.4 (58.6 to 64.2)	9.9	98.5

AUC, area under the curve; NPV, negative predictive value; PPV, positive predictive value.

inflammation and increased permeability) soon after birth can lead to early and irreversible stunting, frequent diarrhoeal illness and persistent systemic subclinical inflammation.^{23–26} These appear to promote a vicious cycle of infection and malnutrition. While difficult to address, a focus on nutrition (micronutrient and macro-nutrient) before, during and following the acute phase of illness may reduce the exacerbation of this cycle. Half of the children who died during the course of this study did so after >30 days following discharge. Therefore, emphasis must also be placed on preventing reinfection in vulnerable children. Promotion of good health behaviour (including hygiene) during the postdischarge period is, therefore, likely to play an important role.

One further area for intervention is education on timely health seeking. Sixty-seven per cent of the deaths in this study occurred outside of a hospital context, but 28% of the out-of-hospital deaths occurred on the way to hospital. The education of mothers on the early warning signs of recurrent illness should also be emphasised during discharge since the common perception may be that recovery from infection brings a child back to a baseline level of risk, which is clearly not true. Since all children were enrolled during a hospital admission, physical inaccessibility was generally not an initial barrier. A previous study on the hospital burden of paediatric acute lower respiratory infections found that although 62% of children are treated in the hospital, 80% of deaths occur outside of the hospital.²⁷ While this study did not address the timing of the out-of-hospital deaths in relation to the hospital visit, it is possible that many of these deaths occurred in the vulnerable months following discharge.

CONCLUSION

This study has derived a parsimonious risk-scoring tool for paediatric postdischarge mortality. Further work is required in external validation of this tool and for the development of effective postdischarge interventions.

Author affiliations

¹School of Population and Public Health, University of British Columbia, Vancouver, British Columbia, Canada

²Department of Pediatrics, Mbarara University of Science and Technology, Mbarara, Uganda

³Center for International Child Health, BC Children's Hospital, Child and Family Research Institute, Vancouver, Canada

⁴Department of Pediatric Anesthesiology, BC Children's Hospital and University of British Columbia, Vancouver, Canada

⁵Canadian HIV Trials Network, St. Paul's Hospital and University of British Columbia, Vancouver, Canada

⁶Department of Pediatrics, BC Children's Hospital and University of British Columbia, Vancouver, Canada

⁷Maternal, Newborn and Child Health Institute, Mbarara University of Science and Technology, Mbarara, Uganda

⁸Department of Statistics, University of British Columbia, Vancouver, Canada

Twitter Follow Matthew Wiens at @matthew_wiens

Acknowledgements The authors gratefully acknowledge the contributions of Annet Twinomuguni, Justine Kamazima, Agaba Collins, Clare Komugisha, Solome Kobugenyi, Alexander Mutungi and Hassan Bariahikwa. Without whose dedication, this study could not have been completed.

Contributors MOW, EK, CPL, JMA, JS, HW, AN, JKa, NK and JKl contributed to the study design. MOW, EK and JMA were responsible for the data collection. MOW, JS, HW and GZ carried out the data analysis. MOW contributed to the drafting of the initial manuscript. MOW, EK, CPL, JMA, JS, NK, HW, AN, JKa, GZ and JKl participated in the reviewing and revising of the manuscript. All the authors gave their approval for the final version of the manuscript.

Funding This study was internally funded. MOW was supported by a fellowship from the Canadian Institutes of Health Research, the Michael Smith Foundation for Health Research and the Canadian Child Health Clinician Scientist Program.

Competing interests None declared.

Ethics approval University of British Columbia (Canada) and Mbarara University of Science and Technology (Uganda).

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Liu L, Oza S, Hogan D, *et al*. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet* 2015;385: 430–40.
2. Wang H, Liddell CA, Coates MM, *et al*. Global, regional, and national levels of neonatal, infant, and under-5 mortality during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384:957–79.
3. Wiens MO, Pawluk S, Kiskoos N. Pediatric post-discharge mortality in resource poor countries: a systematic review. *PLoS ONE* 2013;8: e66698.
4. Karlen W, Dumont G, Petersen C, *et al*. Human-centered phone oximeter interface design for the operating room: Pulse Oximeter interfaced to a mobile device for anesthesia monitoring in the developing world. *Proceedings of the International Conference on Health Informatics*. Rome, Italy: SciTePress, 2011:433–8.
5. The WHO Child Growth Standards [Internet]. 2006 (cited 1 Mar 2015). <http://www.who.int/childgrowth/standards/en/>

6. Fleming S, Thompson M, Stevens R, *et al*. Normal ranges of heart rate and respiratory rate in children from birth to 18 years of age: a systematic review of observational studies. *Lancet* 2011;377:1011–18.
7. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004;114:555–76.
8. National Implementation Guidelines for HIV Counselling and Testing in Uganda [Internet]. Kampala, Uganda, 2010. <http://library.health.go.ug/publications>
9. Harris PA, Taylor R, Thielke R, *et al*. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
10. Wiens MO, Kissoon N, Kumbakumba E, *et al*. Selecting candidate predictor variables for the modelling of post-discharge mortality: a protocol development project. Africa Health Sciences. In press.
11. Hosmer DW, Lemeshow S. Goodness of fit tests for the multiple logistic regression model. *Commun Stat* 1980;9:1043–69.
12. Van Buuren S, Groothuis-Oudshoorn K. Multivariate imputation by chained equations. *J Stat Softw* 2011;45:1–67.
13. Stone M. An asymptotic equivalence of choice of model by cross-validation and Akaike's Criterion. *J R Statist Soc* 1977;39:44–7.
14. Shibata R. Bootstrap estimate of Kullback-Leibler information for model selection. *Stat Sin* 1997;7:375–94.
15. World Health Organization. *Integrated Management of Childhood Illness: distance learning course*. Geneva: World Health Organization, 2014.
16. Nguyen DT, Leung KK, McIntyre L, *et al*. Does Integrated Management of Childhood Illness (IMCI) training improve the skills of health workers? A systematic review and meta-analysis. *PLoS ONE* 2013;8:e66030.
17. Lester RT, Ritvo P, Mills EJ, *et al*. Effects of a mobile phone short message service on antiretroviral treatment adherence in Kenya (WeTel Kenya1): a randomised trial. *Lancet* 2010;376:1838–45.
18. Obasola OI, Mabawonku I, Lagunju I. A review of E-health interventions for maternal and child health in sub-Saharan Africa. *Matern Child Health J* 2015;19:1813–24.
19. Mitchell M, Hedt-Gauthier BL, Msellemu D, *et al*. Using electronic technology to improve clinical care—results from a before-after cluster trial to evaluate assessment and classification of sick children according to Integrated Management of Childhood Illness (IMCI) protocol in Tanzania. *BMC Med Inform Decis Mak* 2013;13:95.
20. Moisi JC, Gatakaa H, Berkley JA, *et al*. Excess child mortality after discharge from hospital in Kilifi, Kenya: a retrospective cohort analysis. *Bull World Health Organ* 2011;89:725–32.
21. Islam MA, Rahman MM, Mahalanabis D, *et al*. Death in a diarrhoeal cohort of infants and young children soon after discharge from hospital: risk factors and causes by verbal autopsy. *J Trop Pediatr* 1996;42:342–7.
22. West TE, Goetghebuer T, Milligan P, *et al*. Long-term morbidity and mortality following hypoxaemic lower respiratory tract infection in Gambian children. *Bull World Health Organ* 1999;77:144–8.
23. Prendergast A, Kelly P. Enteropathies in the developing world: neglected effects on global health. *Am J Trop Med Hyg* 2012;86:756–63.
24. Keusch GT, Denno DM, Black RE, *et al*. Environmental enteric dysfunction: pathogenesis, diagnosis, and clinical consequences. *Clin Infect Dis* 2014;59(Suppl 4):207–12.
25. Kosek M, Guerrant RL, Kang G, *et al*. Assessment of environmental enteropathy in the MAL-ED cohort study: theoretical and analytic framework. *Clin Infect Dis* 2014;59(Suppl 4):239–47.
26. Jones KD, Thitiri J, Ngari M, *et al*. Childhood malnutrition: toward an understanding of infections, inflammation, and antimicrobials. *Food Nutr Bull* 2014;35:64–70.
27. Nair H, Simões EA, Rudan I, *et al*. Global and regional burden of hospital admissions for severe acute lower respiratory infections in young children in 2010: a systematic analysis. *Lancet* 2013;381:1380–90.