

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5-Nitro-1-nonyl-1*H*-benzimidazol-2(3*H*)-oneYounes Ouzidan,^a Youssef Kandri Rodi,^{a*} El Mokhtar Essassi,^b Lahcen El Ammari,^c Frank R. Fronczek^d and Ramaier Venkatraman^e

^aLaboratoire de Chimie Organique Appliquée, Université Sidi Mohamed, Ben Abdallah, Faculté des Sciences et Techniques, Route d'Immouzer, BP 2202 Fès, Morocco, ^bLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences Pharmacochimie, Av Ibn Battouta, BP 1014, Faculté des Sciences, Université Mohammed V-Agdal, Rabat, Morocco, ^cLaboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed V-Agdal, Avenue Ibn Battouta, BP 1014, Rabat, Morocco, ^dDepartment of Chemistry, Louisiana State University, Baton Rouge, LA 70803, USA, and ^eDepartment of Chemistry and Biochemistry, Jackson State University, Jackson, MS 39217, USA

Correspondence e-mail: kandri_rodii@yahoo.fr


Received 17 January 2011; accepted 15 February 2011

Key indicators: single-crystal X-ray study; $T = 90$ K; mean $\sigma(\text{C}-\text{C}) = 0.001$ Å; R factor = 0.040; wR factor = 0.113; data-to-parameter ratio = 31.6.

In the title molecule, $\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}_3$, the dihedral angle between the benzimidazole and nitro group planes is 5.34 (9°) and the dihedral angle between the benzimidazole and aliphatic chain mean planes is 73.23 (5°). The $\text{C}-\text{C}-\text{C}-\text{C}$ torsion angles (about $\pm 176^\circ$) of the nonyl group indicate an all-anti-periplanar conformation. In the crystal, adjacent molecules are linked by pairs of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds into inversion dimers. These molecules are further connected through $\text{C}-\text{H}\cdots\text{O}$ interactions, building tapes parallel to ($\bar{1}22$).

Related literature

For background to the pharmacological and biochemical properties of benzimidazolones, see: Gbadamassi *et al.* (1988); Singh *et al.* (2000); Derand *et al.* (2003); Badarau *et al.* (2009). For similar structures, see: Saber *et al.* (2010); Ouzidan *et al.* (2011).



Experimental

Crystal data

 $\text{C}_{16}\text{H}_{23}\text{N}_3\text{O}_3$ $M_r = 305.37$

Triclinic, $P\bar{1}$
 $a = 5.483$ (1) Å
 $b = 10.2092$ (15) Å
 $c = 14.746$ (3) Å
 $\alpha = 74.275$ (9°)
 $\beta = 79.727$ (6°)
 $\gamma = 83.410$ (8°)

$V = 779.9$ (2) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 90$ K
 $0.35 \times 0.27 \times 0.22$ mm

Data collection

Nonius KappaCCD diffractometer
 21087 measured reflections
 6349 independent reflections

5183 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.113$
 $S = 1.03$
 6349 reflections

201 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.45$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.28$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.88	1.89	2.7651 (9)	170
$\text{C6}-\text{H6}\cdots\text{O3}^{ii}$	0.95	2.58	3.3139 (11)	134

Symmetry codes: (i) $-x + 2, -y + 2, -z + 1$; (ii) $-x, -y + 1, -z + 1$.

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* (Otwinowski & Minor, 1997) and *SCALEPACK*; program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2342).

References

- Altomare, A., Burla, M. C., Camalli, M., Carrozzini, B., Casciarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Rizzi, R. (1999). *J. Appl. Cryst.* **32**, 339–340.
- Badarau, E., Suzenet, F., Bojarski, A. J., Adriana-Luminița Finaru, A. L. & Guillaumet, G. (2009). *Bioorg. Med. Chem. Lett.* **19**, 1600–1603.
- Derand, R., Bulteau-Pignoux, L. & Becq, F. (2003). *J. Membr. Biol.* **194**, 109–117.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Gbadamassi, M., Barascut, J. L., Imbach, J. L. & Gayral, P. (1988). *Eur. J. Med. Chem.* **23**, 225–232.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Ouzidan, Y., Kandri Rodi, Y., Butcher, R. J., Essassi, E. M. & El Ammari, L. (2011). *Acta Cryst.* **E67**, o283.
- Saber, A., Zouihri, H., Essassi, E. M. & Ng, S. W. (2010). *Acta Cryst.* **E66**, o1409.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Singh, S., Syme, C. A., Singh, A. K., Devor, D. C. & Bridges, R. J. (2000). *J. Pharmacol. Exp. Ther.* **296**, 600–611.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supplementary materials

Acta Cryst. (2011). E67, o669 [doi:10.1107/S1600536811005654]

5-Nitro-1-nonyl-1*H*-benzimidazol-2(3*H*)-one

Y. Ouzidan, Y. Kandri Rodi, E. M. Essassi, L. El Ammari, F. R. Fronczek and R. Venkatraman

Comment

Benzimidazoles are useful intermediates/subunits for the development of molecules of pharmaceutical or biological interest (Gbadamassi *et al.*, 1988). Benzimidazolone and its derivatives are also an important class of bioactive molecules in the field of drugs and pharmaceuticals (Derand *et al.*, 2003). They found potential applications in diverse therapeutic areas including, anti-hypertensives and anti-virals (Badarau *et al.*, 2009; Singh *et al.*, 2000). The structural studies of benzimidazolone, linked to an isopropenyl and nonyl group respectively, have been published by Saber *et al.* (2010) and Ouzidan *et al.* (2011).

The 5-nitro-1-nonyl-1*H*-benzimidazol-2(3*H*)-one molecule structure is built up from two fused six- and five-membered rings linked to C₉H₁₉ chain as shown in Fig.1. The aliphatic chain has all-antiperiplanar (all-*trans*) conformation. Furthermore, the fused-ring system and the nitro group are almost planar, with a maximum deviation of 0.0414 (8) Å and 0.0250 (7) Å for C4 and N1 respectively. The dihedral angle between the two rings and nitro group planes is 5.34 (9)°. The torsion angles C1 N2 C8 C9 and C13 C14 C15 C16 are 113.66 (8)° and 177.26 (7)° respectively.

In the crystal, adjacent molecules are linked by pairs of N—H···O hydrogen bonds into inversion dimers. These molecules are further connected through C—H···O hydrogen bonds into a tape parallel to the (-1 2 2) plane, as shown in Fig. 2 and Table 1.

Experimental

To 5-nitro-1*H*-benzimidazol-2(3*H*)-one (0.2 g, 1.1 mmol), potassium carbonate (0.30 g, 2.2 mmol) and tetra-*n*-butylammonium bromide (0.07 g, 0.2 mmol) in DMF (15 ml) was added 1-bromononane (0.43 ml, 2.2 mmol). Stirring was continued at room temperature for 6 h. The salt was removed by filtration and the filtrate concentrated under reduced pressure. The residue was separated by chromatography on a column of silica gel with ethyl acetate/hexane (1/2) as eluent. Colorless needle-shaped crystals were isolated when the solvent was allowed to evaporate [(m.p. 392–394 K (ethanol))].

Refinement

H atoms were located in a difference map and treated as riding with C—H = 0.99 Å, 0.98 Å, 0.95 Å, and 0.88 Å for —CH₂—, —CH₃, aromatic CH and NH respectively. All H atoms with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}$ (aromatic, methylene, N) and $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}$ (methyl).

Figures




Fig. 1. : Molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level. H atoms are represented as small circles.




Fig. 2. Partial packing view of the title compound, showing tapes in the (-1 2 2) plane, built up from molecules linked through N—H...O hydrogen bonds and intermolecular C—H...O contacts (dashed lines).

5-Nitro-1-nonyl-1*H*-benzimidazol-2(3*H*)-one

Crystal data

$C_{16}H_{23}N_3O_3$	$Z = 2$
$M_r = 305.37$	$F(000) = 328$
Triclinic, $P\bar{1}$	$D_x = 1.300 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 5.483 (1) \text{ \AA}$	Cell parameters from 5537 reflections
$b = 10.2092 (15) \text{ \AA}$	$\theta = 2.5\text{--}34.9^\circ$
$c = 14.746 (3) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\alpha = 74.275 (9)^\circ$	$T = 90 \text{ K}$
$\beta = 79.727 (6)^\circ$	Needle, colourless
$\gamma = 83.410 (8)^\circ$	$0.35 \times 0.27 \times 0.22 \text{ mm}$
$V = 779.9 (2) \text{ \AA}^3$	

Data collection

Nonius KappaCCD diffractometer	5183 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.023$
graphite	$\theta_{\text{max}} = 34.9^\circ$, $\theta_{\text{min}} = 2.8^\circ$
ω and φ scans	$h = -8 \rightarrow 8$
21087 measured reflections	$k = -15 \rightarrow 16$
6349 independent reflections	$l = -23 \rightarrow 22$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.040$	H-atom parameters constrained
$wR(F^2) = 0.113$	$w = 1/[\sigma^2(F_o^2) + (0.0578P)^2 + 0.1462P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
6349 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
201 parameters	$\Delta\rho_{\text{max}} = 0.45 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$
	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008),
	$F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Primary atom site location: structure-invariant direct methods Extinction coefficient: 0.023 (5)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.92795 (11)	1.07035 (6)	0.37556 (4)	0.01618 (12)
O2	0.40529 (12)	0.45052 (6)	0.71989 (4)	0.02109 (13)
O3	0.12666 (12)	0.41865 (6)	0.64166 (5)	0.02143 (13)
N1	0.78463 (12)	0.88673 (7)	0.50035 (5)	0.01369 (12)
H1	0.8751	0.8904	0.5431	0.016*
N2	0.61923 (12)	0.93607 (7)	0.36748 (4)	0.01292 (12)
N3	0.29292 (13)	0.48382 (7)	0.65097 (5)	0.01591 (13)
C1	0.79248 (13)	0.97452 (8)	0.41127 (5)	0.01294 (13)
C2	0.50448 (13)	0.82444 (7)	0.42883 (5)	0.01255 (13)
C3	0.61268 (13)	0.79137 (8)	0.51286 (5)	0.01240 (13)
C4	0.54726 (13)	0.68061 (8)	0.58770 (5)	0.01379 (13)
H4	0.6229	0.6565	0.6437	0.017*
C5	0.36219 (13)	0.60623 (7)	0.57568 (5)	0.01384 (13)
C6	0.24677 (14)	0.63920 (8)	0.49481 (6)	0.01500 (13)
H6	0.1193	0.5860	0.4912	0.018*
C7	0.31799 (14)	0.75013 (8)	0.41917 (5)	0.01431 (13)
H7	0.2421	0.7740	0.3632	0.017*
C8	0.56136 (14)	1.00911 (8)	0.27295 (5)	0.01453 (13)
H8A	0.6475	1.0948	0.2508	0.017*
H8B	0.3804	1.0338	0.2778	0.017*
C9	0.63787 (14)	0.92606 (8)	0.19926 (5)	0.01496 (13)
H9A	0.5714	0.9758	0.1402	0.018*
H9B	0.5589	0.8382	0.2237	0.018*
C10	0.91760 (14)	0.89612 (8)	0.17380 (5)	0.01536 (14)
H10A	0.9848	0.8397	0.2311	0.018*
H10B	1.0005	0.9829	0.1522	0.018*
C11	0.97314 (15)	0.82089 (8)	0.09493 (6)	0.01648 (14)
H11A	0.8959	0.7325	0.1186	0.020*
H11B	0.8936	0.8751	0.0400	0.020*
C12	1.24889 (15)	0.79376 (8)	0.06003 (6)	0.01714 (14)
H12A	1.3281	0.7349	0.1137	0.021*

supplementary materials

H12B	1.3293	0.8813	0.0386	0.021*
C13	1.29159 (14)	0.72462 (8)	-0.02204 (6)	0.01650 (14)
H13A	1.2188	0.6349	0.0010	0.020*
H13B	1.2018	0.7809	-0.0735	0.020*
C14	1.56428 (15)	0.70296 (8)	-0.06374 (6)	0.01707 (14)
H14A	1.6346	0.7928	-0.0918	0.020*
H14B	1.6575	0.6522	-0.0118	0.020*
C15	1.59873 (15)	0.62433 (9)	-0.14015 (6)	0.01812 (15)
H15A	1.5350	0.5330	-0.1111	0.022*
H15B	1.4981	0.6728	-0.1903	0.022*
C16	1.86819 (17)	0.60708 (10)	-0.18635 (7)	0.02506 (18)
H16A	1.9271	0.6967	-0.2215	0.038*
H16B	1.8795	0.5492	-0.2304	0.038*
H16C	1.9712	0.5643	-0.1368	0.038*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0171 (2)	0.0163 (3)	0.0153 (2)	-0.00631 (19)	-0.00140 (19)	-0.0029 (2)
O2	0.0265 (3)	0.0195 (3)	0.0162 (3)	-0.0041 (2)	-0.0049 (2)	-0.0008 (2)
O3	0.0224 (3)	0.0188 (3)	0.0235 (3)	-0.0096 (2)	-0.0006 (2)	-0.0048 (2)
N1	0.0147 (3)	0.0147 (3)	0.0125 (3)	-0.0040 (2)	-0.0026 (2)	-0.0035 (2)
N2	0.0138 (3)	0.0137 (3)	0.0117 (3)	-0.0029 (2)	-0.0018 (2)	-0.0032 (2)
N3	0.0177 (3)	0.0143 (3)	0.0153 (3)	-0.0031 (2)	0.0006 (2)	-0.0044 (2)
C1	0.0130 (3)	0.0138 (3)	0.0126 (3)	-0.0014 (2)	-0.0010 (2)	-0.0048 (2)
C2	0.0120 (3)	0.0132 (3)	0.0126 (3)	-0.0014 (2)	-0.0005 (2)	-0.0043 (2)
C3	0.0119 (3)	0.0133 (3)	0.0130 (3)	-0.0020 (2)	-0.0009 (2)	-0.0052 (2)
C4	0.0143 (3)	0.0145 (3)	0.0128 (3)	-0.0020 (2)	-0.0011 (2)	-0.0041 (2)
C5	0.0147 (3)	0.0123 (3)	0.0141 (3)	-0.0024 (2)	0.0000 (2)	-0.0034 (2)
C6	0.0140 (3)	0.0150 (3)	0.0168 (3)	-0.0028 (2)	-0.0017 (2)	-0.0051 (3)
C7	0.0132 (3)	0.0159 (3)	0.0148 (3)	-0.0020 (2)	-0.0030 (2)	-0.0046 (2)
C8	0.0158 (3)	0.0147 (3)	0.0125 (3)	0.0003 (2)	-0.0026 (2)	-0.0028 (2)
C9	0.0157 (3)	0.0172 (3)	0.0125 (3)	-0.0017 (2)	-0.0026 (2)	-0.0043 (2)
C10	0.0163 (3)	0.0171 (3)	0.0135 (3)	-0.0011 (2)	-0.0024 (2)	-0.0053 (3)
C11	0.0173 (3)	0.0189 (3)	0.0142 (3)	-0.0006 (3)	-0.0017 (2)	-0.0065 (3)
C12	0.0179 (3)	0.0196 (4)	0.0153 (3)	-0.0003 (3)	-0.0029 (3)	-0.0071 (3)
C13	0.0168 (3)	0.0184 (3)	0.0150 (3)	-0.0001 (3)	-0.0020 (2)	-0.0063 (3)
C14	0.0173 (3)	0.0181 (3)	0.0157 (3)	-0.0004 (3)	-0.0015 (3)	-0.0053 (3)
C15	0.0200 (3)	0.0183 (4)	0.0157 (3)	0.0012 (3)	-0.0015 (3)	-0.0056 (3)
C16	0.0217 (4)	0.0302 (5)	0.0221 (4)	0.0022 (3)	0.0013 (3)	-0.0094 (3)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.2387 (9)	C9—H9A	0.9900
O2—N3	1.2313 (9)	C9—H9B	0.9900
O3—N3	1.2352 (9)	C10—C11	1.5297 (11)
N1—C1	1.3715 (10)	C10—H10A	0.9900
N1—C3	1.3864 (9)	C10—H10B	0.9900
N1—H1	0.8800	C11—C12	1.5268 (11)

N2—C1	1.3838 (10)	C11—H11A	0.9900
N2—C2	1.3842 (10)	C11—H11B	0.9900
N2—C8	1.4628 (10)	C12—C13	1.5303 (11)
N3—C5	1.4646 (10)	C12—H12A	0.9900
C2—C7	1.3893 (10)	C12—H12B	0.9900
C2—C3	1.4109 (10)	C13—C14	1.5272 (11)
C3—C4	1.3787 (11)	C13—H13A	0.9900
C4—C5	1.3970 (11)	C13—H13B	0.9900
C4—H4	0.9500	C14—C15	1.5263 (11)
C5—C6	1.3924 (11)	C14—H14A	0.9900
C6—C7	1.3930 (11)	C14—H14B	0.9900
C6—H6	0.9500	C15—C16	1.5246 (12)
C7—H7	0.9500	C15—H15A	0.9900
C8—C9	1.5266 (11)	C15—H15B	0.9900
C8—H8A	0.9900	C16—H16A	0.9800
C8—H8B	0.9900	C16—H16B	0.9800
C9—C10	1.5284 (11)	C16—H16C	0.9800
C1—N1—C3	109.78 (6)	C9—C10—C11	110.76 (6)
C1—N1—H1	125.1	C9—C10—H10A	109.5
C3—N1—H1	125.1	C11—C10—H10A	109.5
C1—N2—C2	109.39 (6)	C9—C10—H10B	109.5
C1—N2—C8	124.01 (6)	C11—C10—H10B	109.5
C2—N2—C8	126.52 (6)	H10A—C10—H10B	108.1
O2—N3—O3	123.41 (7)	C12—C11—C10	114.85 (6)
O2—N3—C5	118.20 (7)	C12—C11—H11A	108.6
O3—N3—C5	118.39 (7)	C10—C11—H11A	108.6
O1—C1—N1	127.21 (7)	C12—C11—H11B	108.6
O1—C1—N2	125.74 (7)	C10—C11—H11B	108.6
N1—C1—N2	107.04 (6)	H11A—C11—H11B	107.5
N2—C2—C7	131.59 (7)	C11—C12—C13	112.22 (6)
N2—C2—C3	106.99 (6)	C11—C12—H12A	109.2
C7—C2—C3	121.42 (7)	C13—C12—H12A	109.2
C4—C3—N1	131.24 (7)	C11—C12—H12B	109.2
C4—C3—C2	121.98 (7)	C13—C12—H12B	109.2
N1—C3—C2	106.77 (6)	H12A—C12—H12B	107.9
C3—C4—C5	115.49 (7)	C14—C13—C12	114.38 (7)
C3—C4—H4	122.3	C14—C13—H13A	108.7
C5—C4—H4	122.3	C12—C13—H13A	108.7
C6—C5—C4	123.69 (7)	C14—C13—H13B	108.7
C6—C5—N3	118.32 (7)	C12—C13—H13B	108.7
C4—C5—N3	117.95 (7)	H13A—C13—H13B	107.6
C5—C6—C7	120.08 (7)	C15—C14—C13	112.45 (7)
C5—C6—H6	120.0	C15—C14—H14A	109.1
C7—C6—H6	120.0	C13—C14—H14A	109.1
C2—C7—C6	117.29 (7)	C15—C14—H14B	109.1
C2—C7—H7	121.4	C13—C14—H14B	109.1
C6—C7—H7	121.4	H14A—C14—H14B	107.8
N2—C8—C9	113.04 (6)	C16—C15—C14	113.56 (7)
N2—C8—H8A	109.0	C16—C15—H15A	108.9

supplementary materials

C9—C8—H8A	109.0	C14—C15—H15A	108.9
N2—C8—H8B	109.0	C16—C15—H15B	108.9
C9—C8—H8B	109.0	C14—C15—H15B	108.9
H8A—C8—H8B	107.8	H15A—C15—H15B	107.7
C8—C9—C10	115.31 (6)	C15—C16—H16A	109.5
C8—C9—H9A	108.4	C15—C16—H16B	109.5
C10—C9—H9A	108.4	H16A—C16—H16B	109.5
C8—C9—H9B	108.4	C15—C16—H16C	109.5
C10—C9—H9B	108.4	H16A—C16—H16C	109.5
H9A—C9—H9B	107.5	H16B—C16—H16C	109.5
C3—N1—C1—O1	-178.59 (7)	C3—C4—C5—N3	177.28 (6)
C3—N1—C1—N2	1.29 (8)	O2—N3—C5—C6	175.91 (7)
C2—N2—C1—O1	179.64 (7)	O3—N3—C5—C6	-3.41 (11)
C8—N2—C1—O1	-3.45 (12)	O2—N3—C5—C4	-1.82 (10)
C2—N2—C1—N1	-0.24 (8)	O3—N3—C5—C4	178.86 (7)
C8—N2—C1—N1	176.67 (6)	C4—C5—C6—C7	1.40 (12)
C1—N2—C2—C7	178.96 (8)	N3—C5—C6—C7	-176.19 (7)
C8—N2—C2—C7	2.15 (13)	N2—C2—C7—C6	178.47 (7)
C1—N2—C2—C3	-0.87 (8)	C3—C2—C7—C6	-1.72 (11)
C8—N2—C2—C3	-177.68 (6)	C5—C6—C7—C2	-0.33 (11)
C1—N1—C3—C4	176.91 (8)	C1—N2—C8—C9	113.65 (8)
C1—N1—C3—C2	-1.82 (8)	C2—N2—C8—C9	-69.97 (9)
N2—C2—C3—C4	-177.25 (7)	N2—C8—C9—C10	-66.44 (9)
C7—C2—C3—C4	2.90 (11)	C8—C9—C10—C11	-176.26 (6)
N2—C2—C3—N1	1.62 (8)	C9—C10—C11—C12	176.47 (7)
C7—C2—C3—N1	-178.23 (7)	C10—C11—C12—C13	-177.19 (7)
N1—C3—C4—C5	179.65 (7)	C11—C12—C13—C14	176.46 (7)
C2—C3—C4—C5	-1.79 (11)	C12—C13—C14—C15	175.71 (7)
C3—C4—C5—C6	-0.32 (11)	C13—C14—C15—C16	177.26 (7)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 \cdots O1 ⁱ	0.88	1.89	2.7651 (9)	170
C6—H6 \cdots O3 ⁱⁱ	0.95	2.58	3.3139 (11)	134

Symmetry codes: (i) $-x+2, -y+2, -z+1$; (ii) $-x, -y+1, -z+1$.

Fig. 1




Fig. 2

