

The associations of cardiorespiratory fitness, adiposity and sports participation with arterial stiffness in youth with chronic diseases or physical disabilities

Eero A Haapala^{1,2,3,*}, Kristel Lankhorst^{4,5,*}, Janke de Groot^{3,4,5},
 Maremka Zwinkels^{5,6}, Olaf Verschuren^{5,6}, Harriet Wittink⁴,
 Frank JG Backx⁷, Anne Visser-Meily^{6,7} and Tim Takken^{3,5}; on
 behalf of the HAYS study group

Abstract

Background: The evidence on the associations of cardiorespiratory fitness, body adiposity and sports participation with arterial stiffness in children and adolescents with chronic diseases or physical disabilities is limited.

Methods: Altogether 140 children and adolescents with chronic diseases or physical disabilities participated in this cross-sectional study. Cardiorespiratory fitness was assessed using maximal exercise test with respiratory gas analyses either using shuttle run, shuttle ride, or cycle ergometer test. Cardiorespiratory fitness was defined as peak oxygen uptake by body weight or fat-free mass. Body adiposity was assessed using waist circumference, body mass index standard deviation score and body fat percentage. Sports participation was assessed by a questionnaire. Aortic pulse wave velocity and augmentation index were assessed by a non-invasive oscillometric tonometry device.

Results: Peak oxygen uptake/body weight (standardised regression coefficient β -0.222 , 95% confidence interval (CI) -0.386 to -0.059 , $P=0.002$) and peak oxygen uptake/fat-free mass (β -0.173 , 95% CI -0.329 to -0.017 , $P=0.030$) were inversely and waist circumference directly (β 0.245 , 95% CI 0.093 to 0.414 , $P=0.002$) associated with aortic pulse wave velocity. However, the associations of the measures of cardiorespiratory fitness with aortic pulse wave velocity were attenuated after further adjustment for waist circumference. A higher waist circumference (β -0.215 , 95% CI -0.381 to -0.049 , $P=0.012$) and a higher body mass index standard deviation score (β 0.218 , 95% CI -0.382 to -0.054 , $P=0.010$) were related to lower augmentation index.

Conclusions: Poor cardiorespiratory fitness and higher waist circumference were associated with increased arterial stiffness in children and adolescents with chronic diseases and physical disabilities. The association between cardiorespiratory fitness and arterial stiffness was partly explained by waist circumference.

Keywords

Youth, arterial stiffness, exercise, cardiorespiratory fitness, obesity, chronic disease

Received 16 January 2017; accepted 14 March 2017

Introduction

Arteriosclerotic cardiovascular diseases are one of the leading causes of morbidity and mortality and the costs

⁴Research Group Lifestyle and Health, University of Applied Sciences, the Netherlands

⁵Partner of Shared Utrecht Pediatric Exercise Research (SUPER) Lab, the Netherlands

⁶Center of Excellence in Rehabilitation Medicine, University Medical Center Utrecht and De Hoogstraat Rehabilitation, the Netherlands

⁷Department of Rehabilitation, University Medical Center Utrecht, the Netherlands

*The first two authors shared first authorship.

Corresponding author:

Tim Takken, Child Development and Exercise Center, Wilhelmina Children's Hospital, P.O. Box 85090, 3508AB Utrecht, The Netherlands.
 Email: T.Takken@umcutrecht.nl

¹Faculty of Sport and Health Sciences, University of Jyväskylä, Finland

²Institute of Biomedicine, University of Eastern Finland, Finland

³Child Development and Exercise Center, Wilhelmina Children's Hospital, the Netherlands

related to arteriosclerosis demonstrate a considerable economic burden.¹ Arterial stiffness and endothelial dysfunction are the first signs of arteriosclerosis and they have already been established in children and adolescents.^{2,3} In adults, increased arterial stiffness has been found to predict subsequent clinical cardiovascular events.⁴ Therefore, early identification of children and adolescents with increased arterial stiffness is of importance in order to prevent arteriosclerotic cardiovascular diseases in later years.¹

Poor cardiorespiratory fitness has been associated with higher carotid and femoral artery stiffness and aortic intima media thickness (IMT) and elasticity in adolescents.^{5,6} Better cardiorespiratory fitness in adolescence also has been linked to lower femoral artery stiffness and carotid IMT at the age of 36 years.⁷ Furthermore, obesity has been consistently related to stiffer carotid and aortic arteries among youth.⁸ Higher levels of habitual physical activity have been associated with lower arterial stiffness, lower aortic IMT and improved endothelial function in children and adolescents in some,^{9–11} but not all studies.^{12–15} Furthermore, constantly high levels of vigorous physical activity between the ages of 13 and 36 years have been linked to lower arterial stiffness at the age of 36 years.¹⁶ Finally, the results of some previous studies suggest that exercise training has favourable effects on flow-mediated dilation as a marker of endothelial function in overweight and obese children and adolescents.¹⁷

Children and adolescents with chronic diseases or physical disabilities may have an increased risk of arteriosclerosis.¹⁸ The evidence suggests that children and adolescents with chronic diseases or physical disabilities have lower cardiorespiratory fitness,¹⁹ a higher prevalence of overweight and obesity,^{19,20} lower levels of physical activity,^{20,21} and they participate less often in organised sports,²⁰ than their apparently healthy or normally developing peers. Children and adolescents with chronic diseases or disabilities may also have increased arterial stiffness.²² Studies among adults also suggest that arterial stiffness is a particularly important marker of subsequent cardiovascular morbidity and mortality among those with chronic cardiovascular or metabolic diseases.⁴ However, one small study found no differences in arterial structure and function between children with cerebral palsy and normally developing children with similar levels of physical activity and body adiposity.²³ Nevertheless, there are no studies on the associations of cardiorespiratory fitness, body adiposity and sports participation with arterial stiffness in a large sample of children and adolescents with chronic diseases or physical disabilities.

The aim of the present study was to investigate the associations of cardiorespiratory fitness, body adiposity and sports participation with arterial stiffness in

children and adolescents with chronic diseases or physical disabilities. We also studied if there are thresholds for cardiorespiratory fitness and the measures of body adiposity that are associated with increased arterial stiffness.

Methods

Participants

The present analyses are based on the data from the Health in Adapted Youth Sports (HAYS) study and the Sport-2-Stay-Fit (S2SF) study. These studies are designed to investigate the associations of sports participation and exercise training with physical fitness, physical activity, body adiposity, cognition, cardiovascular health and quality of life in children and adolescents with chronic diseases or physical disabilities. The study designs are published in detail elsewhere.^{24,25}

The children and adolescents for the HAYS study and S2SF study were recruited in the Netherlands among different patient associations, paediatric physical therapy practices, Wilhelmina Children's Hospital in Utrecht, De Hoogstraat Rehabilitation Center in Utrecht, Fitkids practices, schools for special education and adapted sports organisations. The inclusion criteria for the HAYS study and the S2SF study were that participants had to understand the Dutch language, understand simple instructions and be able to perform physical fitness tests. All ambulatory children and adolescents and wheelchair users were eligible to participate. Children and adolescents using an electric wheelchair, having a progressive disease, who participated in other research projects, or who had contraindications for performing a maximal exercise test were excluded.

Ethical approval

The study protocols were approved by the medical ethics committee of the University Medical Center Utrecht, the Netherlands (METC number: 14-332/c and 14-118/m). All participants and the parents of participants under 18 years of age provided their informed written consent. Studies were conducted in accordance with the Helsinki Declaration.

Assessment of cardiorespiratory fitness

Peak oxygen uptake ($\dot{V}O_{2peak}$) as a measure of cardiorespiratory fitness was assessed either by an adapted 10-m incremental shuttle run test, by a 10-m incremental shuttle ride test, or by an incremental cardiopulmonary exercise test on an electronically braked Ergoline cycle ergometer (Ergoselect 200 K; Ergoline, Bitz,

Germany).²⁴ The test modality was based on the evaluation of children's ambulatory ability, mode of daily locomotion, type of sports and safety issues. All children with a congenital cardiopulmonary disease underwent an incremental cardiopulmonary exercise test with continuous electrocardiography monitoring on a cycle ergometer. Participants who were able to walk or run performed an incremental shuttle run test and wheelchair users performed a shuttle ride test. Altogether 64% of the participants performed a shuttle run test, 11% a shuttle ride test and 24% a cycle ergometer test. All the test procedures have been described in detail previously.²⁴

Respiratory gases were collected using paediatric masks (Hans-Rudolph, Shawnee, Kansas, USA) during the test by a calibrated metabolic cart (Cortex Metamax, Samcon bvba, Melle, Belgium). Regardless of testing modality, respiratory gases were measured directly by the breath-by-breath method from the 3-minute resting steady-state period to the post-exercise rest and were averaged over consecutive 10-s periods.

Children and adolescents were verbally encouraged to exercise until voluntary exhaustion. The exercise test was considered maximal if the subjective and objective criteria indicated maximal effort and maximal cardiorespiratory capacity (i.e. flushing, sweating, heart rate >180, respiratory exchange ratio >1.0, or plateau of $\dot{V}O_2$). Cardiorespiratory fitness was defined as $\dot{V}O_{2peak}$ per body weight and $\dot{V}O_{2peak}$ per kg fat-free mass (FFM).

Assessment of body size and body adiposity

Body size and composition were measured after emptying the bladder. Body height was measured by stadiometer (Seca, Hamburg, Germany) in ambulant participants. Among participants using a wheelchair the body height was measured in a supine position using a measuring tape. In the case of spasticity of the lower limbs, arm span width was measured to the nearest centimetre from middle fingertip to fingertip. Body weight was measured by standard scale (Seca, Hamburg, Germany). Participants who used a wheelchair and who were not able to stand on a scale were measured using a wheelchair scale (Stimag BV, Lisse, the Netherlands). Body mass index (BMI) was calculated as body weight (kg)/body height (m)² for ambulant participants and as body weight (kg)/arm span length. We used arm span adjustment of $\times 0.95$ and $\times 0.90$ for participants with central neurological disorders in mid-lumbar lesions and high lumbar/thoracic lesions, respectively.²⁶ Waist circumference was measured by an unstretchable measuring tape at the level of the navel. BMI standard deviation score (SDS) and waist circumference SDS were computed

using national reference values.²⁷ Body fat percentage and FFM were measured by bioelectrical impedance analyses using a Bodystat Quadscan 4000 device (EuroMedix, Leuven, Belgium).

Assessment of sports participation

To allow a comparison between children and adolescents with chronic diseases or physical disabilities who participated in organised sports and those without regular sports participation we recruited sports participants from a broad range of participation in sports including recreational level as well as high level competitive sports.²⁴ The study participants reported how often they participated in organised sports on a questionnaire. In the present study, we defined regular sports participation as any involvement in organised sports at least two times per week.

Assessment of arterial stiffness

Aortic pulse wave velocity (PWV_{ao}), as a measure of arterial stiffness and augmentation index (AIX%), as a measure of peripheral arterial tone, were assessed by a non-invasive oscillometric tonometry device (Arteriograph; TensioMed Ltd., Budapest, Hungary) from the right arm. PWV_{ao} and AIX% derived from the arteriograph analyses have been validated against an invasive method in adults. The correlation of invasively assessed PWV_{ao} and AIX% to PWV_{ao} and AIX% measured by the arteriograph has been found to be strong ($r > 0.90$) with a reasonable agreement between oscillometric tonometry and invasive methods.²⁸ Before the measurement, participants were asked to rest in a supine position for 10 minutes. Age and sex-specific SDS norm values for PWV_{ao} and AIX% were calculated based on the data of over 4500 Caucasian children and adolescents.^{29,30} A higher PWV_{ao} indicates higher aortic stiffness and a higher AIX% indicates higher peripheral arterial tone.

Systolic blood pressure was also assessed by the arteriograph device (TensioMed Ltd., Budapest, Hungary) in a supine position after 10 minutes rest.

Statistical methods

Basic characteristics between boys and girls were compared using the Student's *t*-test, the Mann-Whitney U-test, or the chi square test. The associations of cardiorespiratory fitness and body adiposity as independent variables with PWV_{ao} and AIX% as dependent variables were studied using linear regression analyses adjusted for age and sex. The differences in PWV_{ao} and AIX% between sport participants and non-sport participants were investigated using general

linear models (GLMs) adjusted for age and sex. The data on the associations of cardiorespiratory fitness, body adiposity and sports participation with PWVao and AIX% were also mutually adjusted.

All data were additionally controlled for systolic blood pressure, the mode of exercise testing and diagnosis (cardiovascular disease vs. other).

Receiver operating characteristics (ROC) curves were used to investigate the optimal cut-off for $\dot{V}O_{2\text{peak}}$, waist circumference, BMI SDS and body fat percentage to identify children and adolescents at or over +1 SD of PWVao and AIX% reference values.^{29,30} The area under the curve (AUC) is considered a measure of the effectiveness of the predictor variable to identify correctly children and adolescents at or above +1 SD of PWVao and AIX% (sensitivity) and to identify correctly participants (specificity) below +1 SD of PWVao and AIX%. An AUC of 1 represents the ability to identify perfectly children and adolescents at or above +1 SD of PWVao and AIX% from other participants, whereas an AUC of 0.5 indicates no greater predictive ability than chance alone.

The optimal cut-off was determined by the Youden index,³¹ which is the maximum value of J that is computed as: sensitivity + specificity - 1.

The effect modification of sex was investigated by GLMs. Because we found no statistically significant sex interactions between the measures of cardiorespiratory fitness and body adiposity and the outcome variables, we performed all analyses with data of boys and girls combined.

Student's t -test, the Mann-Whitney U-test, the chi square test, GLM analyses and the linear regression analyses were performed using the SPSS Statistics, Version 23.0 (IBM Corp., Armonk, NY, USA). The ROC curve analyses were performed using MedCalc Statistical Software, version 16.1 (MedCalc Software bvba, Ostend, Belgium). A P value of less than 0.05 was considered statistically significant.

Results

Descriptive characteristics

Complete data on variables used in the present analyses were available for 140 children and adolescents with chronic diseases or physical disabilities (86 boys, 54 girls). Children and adolescents who were excluded ($N=37$, 20 boys, 17 girls) from the present analyses because of the missing data did not differ from those who were included in waist circumference or BMI SDS. Children and adolescents who were included in the present analyses were slightly older than those who were excluded from the present analyses ($P=0.029$).

Among the included children and adolescents, boys were taller, had a lower body fat percentage, higher $\dot{V}O_{2\text{peak}}$ and a lower AIX% than girls (Table 1). Seventy-six (56%) of the included children and adolescents participated in sports. Altogether 17 children and adolescents had a cardiovascular disease, four had a pulmonary disease, 10 had a metabolic disease, 11 had a musculoskeletal/orthopaedic disability, 82 had a neuromuscular disease/disability, six had an immunological/haematological disease, three had cancer and seven had epilepsy (Table 1). There were no age differences among the children and adolescents in different diagnosis groups ($P=0.682$).

Associations of cardiorespiratory fitness body adiposity and sports participation with arterial stiffness

$\dot{V}O_{2\text{peak}}$ per body weight and $\dot{V}O_{2\text{peak}}$ per FFM were inversely and waist circumference was directly associated with PWVao after adjustment for age and sex (Table 2). However, the relationship of $\dot{V}O_{2\text{peak}}$ per body weight ($\beta -0.133$, 95% confidence interval (CI) -0.315 to 0.050 , $P=0.152$) and $\dot{V}O_{2\text{peak}}$ per FFM ($\beta -0.137$, 95% CI -0.291 to 0.017 , $P=0.082$) to PWVao were no longer statistically significant after further adjustment for waist circumference. Additional adjustment for systolic blood pressure, the mode of exercise testing, or diagnosis had no effect on these associations (data not shown).

Waist circumference and BMI SDS were inversely associated with AIX% after adjustment for age and sex. Further adjustment for the measures of cardiorespiratory fitness or sports participation had no effect on these associations (data not shown). However, further adjustment for systolic blood pressure slightly attenuated the association between waist circumference and AIX% ($\beta -0.166$, 95% CI -0.343 to 0.010 , $P=0.064$). Additional adjustment for the mode of exercise testing, or diagnosis had no effect on these associations (data not shown).

Sports participation was not associated with PWVao or AIX% after adjustment for age and sex ($P > 0.40$). The associations of cardiorespiratory fitness, body adiposity, sports participation with PWVao and AIX% remained similar when PWVao SDS and AIX% SDS were used as outcome measures or waist circumference SDS was used as an independent variable (data not shown).

Ability of cardiorespiratory fitness and body adiposity to identify children and adolescents with increased arterial stiffness

The ROC curve analyses revealed that the optimal cut-off for waist circumference to identify children and

Table 1. Descriptive characteristics.

	All	Boys (N = 86)	Girls (N = 54)	P value
Age (years)	14.3 (2.7)	14.1 (2.7)	14.7 (2.9)	0.078
Diagnosis group				0.320
Cardiovascular disease (%)	12.1	15.1	7.4	
Pulmonary disease (%)	2.9	4.7	0.0	
Metabolic disease (%)	7.1	4.7	11.1	
Musculoskeletal/orthopaedic disability (%)	7.9	9.3	5.6	
Neuromuscular disease/disability (%)	58.6	57.0	61.1	
Immunological/haematological disease (%)	4.3	3.5	5.6	
Cancer (%)	2.1	1.2	3.7	
Epilepsy (%)	5.0	4.7	5.6	
Body height (cm)	160 (14.3)	162.7 (15.9)	157.9 (10.9)	0.033
Body weight (kg)	54.8 (16.6)	56.7 (18.4)	51.7 (12.7)	0.059
Body mass index ^a (kg/m ²)	20.0 (5.2)	20.2 (5.6)	19.7 (5.1)	0.681
Body mass index standard deviation score	0.68 (1.3)	0.83 (1.3)	0.43 (1.3)	0.078
Prevalence of overweight (%)	39.3	43	33.3	0.253
Waist circumference (cm)	75.7 (13.3)	75.6 (14.2)	75.9 (12.1)	0.888
Waist circumference standard deviation score	0.8 (1.3)	0.7 (1.5)	1.1 (1.1)	0.062
Body fat percentage (%)	23.9 (9.7)	21.2 (9.4)	28.1 (8.7)	<0.001
Peak oxygen uptake (L/min)	2.1 (1.0)	2.5 (0.9)	1.8 (0.4)	<0.001
Peak oxygen uptake (ml/kg/min) ^a	40.0 (15.0)	44.0 (15.0)	36.0 (11.0)	<0.001
Peak oxygen uptake (ml/FFM/min) ^a	52.9 (13.7)	56.5 (14.8)	51.1 (8.1)	<0.001
Aortic pulse wave velocity (m/s) ^a	5.8 (1.3)	5.8 (1.2)	5.8 (1.1)	0.098
Aortic pulse wave velocity standard deviation score ^a	-0.17 (1.71)	-0.21 (1.7)	-0.04 (1.7)	0.189
Aortic augmentation index (%) ^a	9.0 (10.4)	7.7 (11.4)	10.1 (10.3)	0.005
Aortic augmentation index (%) standard deviation score ^a	0.19 (1.23)	0.17 (1.2)	0.25 (1.2)	0.245

FFM: fat-free mass.

The data are mean (SD), median (interquartile range),^a or percentages and the P values from the t-test for independent samples for continuous variables with normal distribution and Mann-Whitney U-test for continuous variables with skewed distribution, or chi-square for prevalence of diseases/disabilities and overweight.

Table 2. Associations of cardiorespiratory fitness and body adiposity with arterial stiffness in 140 children and adolescents with chronic diseases or physical disabilities.

	Aortic pulse wave velocity (m/s)			Aortic augmentation index (%)		
	B	95% CI	P value	B	95% CI	P value
Peak oxygen uptake (ml/kg/min)	-0.222	-0.386 to -0.059	0.008	-0.100	-0.271 to 0.070	0.247
Peak oxygen uptake (ml/fat-free mass/min)	-0.173	-0.329 to -0.017	0.030	-0.120	-0.281 to 0.041	0.142
Waist circumference (cm)	0.254	0.093 to 0.414	0.002	-0.215	-0.381 to -0.049	0.012
Body mass index standard deviation score	0.141	-0.026 to 0.307	0.097	-0.218	-0.382 to -0.054	0.010
Body fat percentage (%)	0.036	-0.127 to 0.198	0.664	0.016	-0.150 to 0.183	0.847

The data are standardised regression coefficients and their 95% confidence intervals (CI) adjusted for age and sex.

adolescents with 1 SD or greater of PWV SDS was more than 73 cm (95% CI 66 to 78) with a sensitivity of 81%, a specificity of 53% and the Youden index of 0.3386 (Figure 1). The corresponding cut-off for waist circumference SDS was greater than -0.05 units (95%

CI -0.09 to -0.02) with a sensitivity of 81%, a specificity 54% and the Youden index of 0.3477. The optimal cut-off for $\dot{V}O_{2peak}$ per body weight was less than 35 ml/kg/min (95% CI 27 to 40) with a sensitivity of 48%, a specificity of 76% and the Youden index of 0.2453.




Figure 1. Receiver operating characteristics curves of the efficacy of the measures of cardiorespiratory fitness and body adiposity to identify children and adolescents with chronic diseases or physical disabilities with increased aortic pulse wave velocity (≥ 1 SD from reference values.²⁸ AUC indicates the area under the curve (95% confidence interval; CI).




Figure 2. Receiver operating characteristics curve of the efficacy of cardiorespiratory fitness to identify children and adolescents with chronic diseases or physical disabilities with increased peripheral arterial tone (augmentation index (AIX%) ≥ 1 SD from reference values.²⁹ AUC indicates the area under the curve (95% confidence interval; CI). FFM: fat-free mass.

These analyses indicated that children and adolescents with a higher waist circumference and a lower $\dot{V}O_{2\text{peak}}$ per body weight were more likely to have increased arterial stiffness.

The optimal cut-off for $\dot{V}O_{2\text{peak}}$ per FFM to identify children and adolescents with 1 SD or greater of AIX% SDS was less than 51 ml/kg FFM/min (95% CI 46 to 66) with a sensitivity of 64%, a specificity of 66% and the Youden index of 0.3036, suggesting that children and adolescents with poorer cardiorespiratory fitness had increased AIX% compared to those with better cardiorespiratory fitness (Figure 2).

Discussion

In the present study, we found inverse associations of the measures of cardiorespiratory fitness with aortic stiffness among children and adolescents with chronic disease or physical disabilities. However, the relationships between cardiorespiratory fitness and aortic stiffness were partly explained by waist circumference. We also found that a higher waist circumference had strong and consistent relationship to a higher aortic stiffness. Furthermore, higher waist circumference and BMI SDS were associated with a lower peripheral arterial tone as indicated by a lower AIX%. Finally, we observed no relationship between sports participation and arterial stiffness.

To the best of our knowledge, this is the first study on the associations of directly assessed $\dot{V}O_{2\text{peak}}$ with

arterial stiffness among children and adolescents with chronic diseases or physical disabilities. In conjunction with some other studies, we found that better cardiorespiratory fitness was linked to lower aortic stiffness.^{5,6} However, the associations of $\dot{V}O_{2\text{peak}}$ per body weight and $\dot{V}O_{2\text{peak}}$ per FFM with arterial stiffness were partly explained by waist circumference. Similarly, one study found that an inverse association between cardiorespiratory fitness and arterial stiffness was explained by body adiposity among children aged 10 years.³² These results are in line with the observations that the associations between $\dot{V}O_{2\text{peak}}$ relative to body weight and cardiometabolic risk factors are confounded by body adiposity.^{33,34}

In our previous study, maximal work load by FFM as a measure of cardiorespiratory fitness was inversely associated with arterial stiffness in children aged 6–8 years independent of body fat percentage.³⁵ We found that $\dot{V}O_{2\text{peak}}$ relative to FFM was attenuated after further adjustment for waist circumference, but remained at borderline significance. Furthermore, controlling for BMI SDS or body fat percentage had no effect on the relationship between $\dot{V}O_{2\text{peak}}$ relative to FFM and arterial stiffness. These results together indicate that better cardiorespiratory fitness is related to more compliant arteries and that this relationship is not completely caused by lower levels of body adiposity in individuals with higher cardiorespiratory fitness. However, more research on the associations of cardiorespiratory fitness with arterial fitness with an appropriate scaling of $\dot{V}O_{2\text{peak}}$ is warranted to clarify further the independent role of higher cardiorespiratory fitness of reduced arterial stiffness in children and adolescents. Nevertheless, our results suggest that in addition to weight management improvements in cardiorespiratory fitness may improve arterial stiffness in children and adolescents with chronic diseases or physical disabilities.

Previous studies have suggested a threshold of 35–46 mL/kg/min for $\dot{V}O_{2\text{peak}}$ to identify apparently healthy children with increased cardiometabolic risk.³⁶ We found that a $\dot{V}O_{2\text{peak}}$ per body weight lower than 35 mL/kg/min was the optimal threshold for identifying those with increased PWVao and the threshold for identifying children and adolescents with increased AIX% was 51 mL/FFM/min. These slight differences in the cut-offs are probably due to the differences in the outcome measures, methods used to assess cardiorespiratory fitness and to measure or estimate $\dot{V}O_{2\text{peak}}$ as well as the populations studied. We also analysed boys and girls together to achieve a better statistical power, whereas other studies have provided separate cut-offs for boys and girls. However, our results along with the results from previous studies suggest that $\dot{V}O_{2\text{peak}}$ below 35–40 mL/kg/min is related to increased cardiometabolic risk in children and adolescents.

Obesity has been consistently linked to increased arterial stiffness in children and adolescents.⁸ Similarly, we found that waist circumference in particular was directly associated with arterial stiffness in children and adolescents with chronic diseases and physical disabilities independent of cardiorespiratory fitness. However, we found relatively weak associations of BMI SDS and body fat percentage with arterial stiffness. An explanation for these inconsistent findings may be that abdominal adiposity has a pronounced negative impact on arterial health in children and adolescents.³⁷ Furthermore, we found that higher levels of body adiposity were related to lower arterial tone at rest. We³⁵ and others³⁸ have previously observed inverse associations between body adiposity and peripheral arterial tone in children and adolescents. One reason for these observations may be that overweight may increase arterial diameter and thereby compensate the negative effects of body adiposity on arterial stiffness.³⁸

We observed that waist circumference exceeding 73 cm and waist circumference SDS above –0.05 units were the optimal cut-offs for identifying children and adolescents with increased arterial stiffness. These cut-offs corresponding to an average waist circumference in the Dutch population suggest that waist circumference already at normal range at the population level is associated with increased arterial stiffness in children and adolescents with chronic diseases or physical disabilities. These findings are supported by the study showing an increased cardiovascular mortality over a 40-year follow-up already among those who had a BMI at the mid-normal range during adolescence.³⁹

Previous studies have found either inverse or no association between habitual physical activity and the measures of arterial stiffness in children and adolescents.^{13,40,41} We observed no differences in arterial stiffness between sports participants and non-sports participants. One reason for our observations may be that sports participation among children and adolescents with chronic diseases or physical disabilities is insufficient in frequency, duration, or intensity to elicit favourable effects on arterial stiffness. Accordingly, there is some evidence that despite the high prevalence of sports participation, the majority of children and adolescents with chronic diseases and physical disabilities fail to meet the physical activity recommendations.⁴² Another explanation may be that there may not be a large difference in total habitual physical activity among sports participants and non-participants.²³

The strengths of the present study include valid and reproducible methods used to assess cardiorespiratory fitness, body composition and arterial stiffness in a relatively large sample of children and adolescents with chronic diseases or physical disabilities. This study

had few limitations. We were not able to assess the pubertal status of the participants, nor were dietary factors recorded. We also used sports participation as a proxy for physical activity instead of objectively measured physical activity. We defined chronic disease using a non-categorical approach that considers chronically ill young people as one population rather than specific disease classes.⁴³ The current sample size prohibited a sub-group analysis in specifying the disease classes. Furthermore, we analysed boys and girls combined because of the limited statistical power to analyse boys and girls separately. We found no evidence on the effect modification of sex, but the results of some previous studies suggest that the association between cardiorespiratory fitness and arterial health is stronger in boys than in girls.⁷ Furthermore, longitudinal studies are needed to study whether the effects of arterial stiffness during childhood and adolescence on cardiovascular diseases in adulthood are different in boys and girls.

In conclusion, lower cardiorespiratory fitness and a higher waist circumference were associated with a higher arterial stiffness in children and adolescents with chronic diseases or physical disabilities in the present study. However, the associations of cardiorespiratory fitness were partly explained by waist circumference. Furthermore, we found no association between sports participation and arterial stiffness. We also found that thresholds of less than 35 mL/kg/min for $\dot{V}O_{2peak}$, more than 73 cm for waist circumference and over -0.05 units for waist circumference SDS were associated with increased arterial stiffness. Taken together, these results suggest that interventions aiming to decrease body adiposity and to improve cardiorespiratory fitness may improve arterial stiffness in children and adolescents with chronic diseases or physical disabilities. However, intervention studies are warranted to confirm the findings of the present study.

The HAYS study group

FJG Backx (Department of Rehabilitation, Physical Therapy Science and Sports, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, the Netherlands), KWJE van der Ende-Kastelijn (University of Applied Sciences, Utrecht, the Netherlands), JF de Groot (University of Applied Sciences, Utrecht, the Netherlands), KM Lankhorst (University of Applied Sciences, Utrecht, the Netherlands), TCW Nijboer (Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, the Netherlands), T Takken (Child Development and Exercise Center, University Medical Center Utrecht, Utrecht, the Netherlands), DW Smits (Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, the Netherlands), OW Verschuren (Brain Center Rudolf Magnus, University Medical Center Utrecht and De Hoogstraat Rehabilitation, Utrecht, the Netherlands), JMA Visser-Meily (Department of Rehabilitation, Physical

Therapy Science and Sports, Brain Center Rudolf Magnus, University Medical Center Utrecht, and De Hoogstraat Rehabilitation, Utrecht, the Netherlands), MJ Volman (Faculty of Social Sciences, Department of General and Special Education, Utrecht University, Utrecht, the Netherlands), HW Wittink (University of Applied Sciences, Utrecht, the Netherlands).

Author contribution

EAH, KL, AVM, OV, JG, FJGB, HW, MZ and TT contributed to conception or design of the study and analysis or interpretation of the data of the present study. EAH and KL drafted the manuscript and AVM, OV, JG, FJGB, HW, MZ and TT critically revised the manuscript. All authors gave final approval and agree to be accountable for all aspects of work ensuring integrity and accuracy. The paper is original and it or parts of it have not been published elsewhere.

Acknowledgements

The authors would like to thank all children and adolescents who participated the HAYS and the F2SF studies and the study groups for the skillful contribution in performing these studies.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: the study was funded by an unconditional grant of the Dutch Organization of Health Research (ZONMW) grant number 525001005. Eero Haapala was financially supported by Jenny and Antti Wihuri Foundation, Päivikki and Sakari Sohlberg Foundation, Eemil Aaltonen Foundation and the University of Jyväskylä.

References

- Piepoli M, Hoes A, Agewall S, et al. 2016 European Guidelines on cardiovascular disease prevention in clinical practice: the Sixth Joint Task Force of the European Society of Cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of 10 societies and by invited experts) developed with the special contribution of the European Association for Cardiovascular Prevention and Rehabilitation (EACPR). *Eur Heart J* 2016; 37: 2315–2381.
- McGill HC, McMahan CA, Herderick EE, et al. Origin of atherosclerosis in childhood and adolescence. *Am J Clin Nutr* 2000; 72: 1307–1315.
- Fernhall B, Agiovlasitis S, Rowland T, et al. Arterial function in youth: window into cardiovascular risk. *J Appl Physiol* 2008; 105: 325–333.

4. Vlachopoulos C, Aznaouridis K and Stefanadis C. Prediction of cardiovascular events and all-cause mortality with arterial stiffness. A systematic review and meta-analysis. *J Am Coll Cardiol* 2010; 55: 1318–1327.
5. Pahkala K, Laitinen TT, Heinonen OJ, et al. Association of fitness with vascular intima-media thickness and elasticity in adolescence. *Pediatrics* 2013; 132: 77–84.
6. Ferreira I, Twisk JWR, Stehouwer CDA, et al. Longitudinal changes in $\dot{V}O_{2\max}$: Associations with carotid IMT and arterial stiffness. *Med Sci Sport Exerc* 2003; 35: 1670–1678.
7. Ferreira I, Twisk JWR, Van Mechelen W, et al. Current and adolescent levels of cardiopulmonary fitness are related to large artery properties at age 36: the Amsterdam Growth and Health Longitudinal Study. *Eur J Clin Invest* 2002; 32: 723–731.
8. Cote AT, Phillips AA, Harris KC, et al. Obesity and arterial stiffness in children: systematic review and meta-analysis. *Arterioscler Thromb Vasc Biol* 2015; 35: 1038–1044.
9. Pahkala K, Heinonen OJ, Lagström H, et al. Vascular endothelial function and leisure-time physical activity in adolescents. *Circulation* 2008; 118: 2353–2359.
10. Pahkala K, Heinonen OJ, Simell O, et al. Association of physical activity with vascular endothelial function and intima-media thickness. *Circulation* 2011; 124: 1956–1963.
11. Ried-Larsen M, Grøntved A, Kristensen PL, et al. Moderate-and-vigorous physical activity from adolescence to adulthood and subclinical atherosclerosis in adulthood: prospective observations from the European Youth Heart Study. *Br J Sports Med* 2015; 49: 107–112.
12. Walker DJ, MacIntosh A, Kozyrskyj A, et al. The associations between cardiovascular risk factors, physical activity, and arterial stiffness in youth. *J Phys Act Health* 2013; 10: 198–204.
13. Melo X, Santa-Clara H, Pimenta NM, et al. Intima-media thickness in 11–13 years-old children: variation attributed to sedentary behavior, physical activity, cardiorespiratory fitness and waist circumference. *J Phys Act Health* 2014; 12: 610–617.
14. Reed KE, Warburton DER, Lewanczuk RZ, et al. Arterial compliance in young children: the role of aerobic fitness. *Eur J Cardiovasc Prev Rehabil* 2005; 12: 492–497.
15. Idris NS, Evelein AMV, Geerts CC, et al. Effect of physical activity on vascular characteristics in young children. *Eur J Prev Cardiol* 2015; 22: 656–664.
16. Van De Laar RJ, Ferreira I, Van Mechelen W, et al. Lifetime vigorous but not light-to-moderate habitual physical activity impacts favorably on carotid stiffness in young adults: the Amsterdam Growth and Health Longitudinal Study. *Hypertension* 2010; 55: 33–39.
17. Dias KA, Green DJ, Ingul CB, et al. Exercise and vascular function in child obesity: a meta-analysis. *Pediatrics* 2015; 136: 648–659.
18. Rimmer JA, Rowland JL. Physical activity for youth with disabilities: a critical need in an underserved population. *Dev Neurorehabil* 2015; 11: 141–148.
19. Kotte EMW, Winkler AMF and Takken T. Fitkids exercise therapy program in the Netherlands. *Pediatr Phys Ther* 2013; 25: 7–13.
20. Neter JE, Schokker DF, de Jong E, et al. The prevalence of overweight and obesity and its determinants in children with and without disabilities. *J Pediatr* 2011; 158: 735–739.
21. Carlon SL, Taylor NF, Dodd KJ, et al. Differences in habitual physical activity levels of young people with cerebral palsy and their typically developing peers: A systematic review. *Disabil Rehabil* 2013; 35: 647–655.
22. Cheung YF. Arterial stiffness in the young: assessment, determinants, and implications. *Korean Circ J* 2010; 40: 153–162.
23. Martin AA, Cotie LM, Timmons BW, et al. Arterial structure and function in ambulatory adolescents with cerebral palsy are not different from healthy controls. *Int J Pediatr* 2012; 2012: 168209.
24. Lankhorst K, van der Ende-Kastelijns K, de Groot J, et al. Health in Adapted Youth Sports Study (HAYS): health effects of sports participation in children and adolescents with a chronic disease or physical disability. *Springer Plus* 2015; 4: 796.
25. Zwinkels M, Verschuren O, Lankhorst K, et al. Sport-2-Stay-Fit study: health effects of after-school sport participation in children and adolescents with a chronic disease or physical disability. *BMC Sports Sci Med Rehabil* 2015; 7: 22.
26. Dosa NP, Foley JT, Eckrich M, et al. Obesity across the lifespan among persons with spina bifida. *Disabil Rehabil* 2009; 31: 914–920.
27. Schönbeck Y, Talma H, von Dommelen P, et al. Increase in prevalence of overweight in dutch children and adolescents: a comparison of nationwide growth studies in 1980, 1997 and 2009. *PLoS One* 2011; 6: 27608.
28. Horváth IG, Németh A, Lenkey Z, et al. Invasive validation of a new oscillometric device (Arteriograph) for measuring augmentation index, central blood pressure and aortic pulse wave velocity. *J Hypertens* 2010; 28: 2068–2075.
29. Hidvégi EV, Illyés M, Benczúr B, et al. Reference values of aortic pulse wave velocity in a large healthy population aged between 3 and 18 years. *J Hypertens* 2012; 30: 1.
30. Hidvégi EV, Illyés M, Molnár FT, et al. Influence of body height on aortic systolic pressure augmentation and wave reflection in childhood. *J Hum Hypertens* 2015; 29: 495–501.
31. Perkins NJ and Schisterman EF. The inconsistency of ‘optimal’ cutpoints obtained using two criteria based on the receiver operating characteristic curve. *Am J Epidemiol* 2006; 163: 670–675.
32. Sakuragi S, Abhayaratna K, Gravenmaker KJ, et al. Influence of adiposity and physical activity on arterial stiffness in healthy children: the Lifestyle of our Kids Study. *Hypertension* 2009; 53: 611–616.
33. Shaibi G, Cruz M, Ball G, et al. Cardiovascular fitness and the metabolic syndrome in overweight Latino youths. *Med Sci Sport Exerc* 2005; 37: 922–928.
34. Tompuri T, Lintu N, Savonen K, et al. Measures of cardiorespiratory fitness in relation to measures of body size

- and composition among children. *Clin Physiol Funct Imaging* 2015; 35: 469–477.
35. Veijalainen A, Tompuri T, Haapala EA, et al. Associations of cardiorespiratory fitness, physical activity, and adiposity with arterial stiffness in children. *Scand J Med Sci Sports* 2016; 26: 943–950.
 36. Adegboye ARA, Anderssen SA, Froberg K, et al. Recommended aerobic fitness level for metabolic health in children and adolescents: a study of diagnostic accuracy. *Br J Sports Med* 2011; 45: 722–728.
 37. Arnberg K and Larnekjær A. Central adiposity and protein intake are associated with arterial stiffness in overweight children. *J Nutr* 2012; 142: 878–885.
 38. Cote AT, Harris KC, Panagiotopoulos C, et al. Childhood obesity and cardiovascular dysfunction. *J Am Coll Cardiol* 2013; 62: 1309–1319.
 39. Twig G, Yaniv G, Levine H, et al. Body-mass index in 2.3 million adolescents and cardiovascular death in adulthood. *N Engl J Med* 2016; 374: 2430–2440.
 40. Palve KS, Pahkala K, Magnussen CG, et al. Association of physical activity in childhood and early adulthood with carotid artery elasticity 21 years later: the Cardiovascular Risk in Young Finns Study. *J Am Heart Assoc* 2014; 3: e000594.
 41. Haapala EA, Väistö J, Veijalainen A, et al. Associations of objectively measured physical activity and sedentary time with arterial stiffness in pre-pubertal children. *Ped Exerc Sci*. Epub ahead of print 2017. DOI: 10.1123/pes.2016-0168.
 42. Ng K, Rintala P, Tynjälä J, et al. Physical activity trends of Finnish adolescents with long-term illnesses or disabilities from 2002 to 2014. *J Phys Act Health* 2016; 13: 816–821.
 43. Stein RE and Jessop DJ. A noncategorical approach to chronic childhood illness. *Public Health Rep* 1982; 97: 354–362.