





# Evidence for Quinol Oxidation Activity of ImoA, a Novel NapC/NirT Family Protein from the Neutrophilic Fe(II)-Oxidizing Bacterium *Sideroxydans lithotrophicus* ES-1

Abhiney Jain,<sup>a</sup> Anaísa Coelho,<sup>b</sup> Joana Madjarov,<sup>b</sup>  Catarina M. Paquete,<sup>b</sup>  Jeffrey A. Gralnick<sup>a</sup>

<sup>a</sup>BioTechnology Institute and Department of Plant and Microbial Biology, University of Minnesota — Twin Cities, St. Paul, Minnesota, USA

<sup>b</sup>Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Lisbon, Oeiras, Portugal

Abhiney Jain and Anaísa Coelho contributed equally to this article. The order was determined based on who generated the first partial draft of the manuscript.

**ABSTRACT** *Sideroxydans* species are important chemolithoautotrophic Fe(II)-oxidizing bacteria in freshwater environments and play a role in biogeochemical cycling of multiple elements. Due to difficulties in laboratory cultivation and genetic intractability, the electron transport proteins required for the growth and survival of this organism remain understudied. In *Sideroxydans lithotrophicus* ES-1, it is proposed that the Mto pathway transfers electrons from extracellular Fe(II) oxidation across the periplasm to an inner membrane NapC/NirT family protein encoded by *Slit\_2495* to reduce the quinone pool. Based on sequence similarity, *Slit\_2495* has been putatively called CymA, a NapC/NirT family protein which in *Shewanella oneidensis* MR-1 oxidizes the quinol pool during anaerobic respiration of a wide range of substrates. However, our phylogenetic analysis using the alignment of different NapC/NirT family proteins shows that *Slit\_2495* clusters closer to NirT sequences than to CymA. We propose the name ImoA (inner membrane oxidoreductase) for *Slit\_2495*. Our data demonstrate that ImoA can oxidize quinol pools in the inner membrane and is able to functionally replace CymA in *S. oneidensis*. The ability of ImoA to oxidize quinol *in vivo* as opposed to its proposed function of reducing quinone raises questions about the directionality and/or reversibility of electron flow through the Mto pathway in *S. lithotrophicus*.

**IMPORTANCE** Fe(II)-oxidizing bacteria play an important role in biogeochemical cycles. At circumneutral pH, these organisms perform extracellular electron transfer, taking up electrons from Fe(II) outside the cell, potentially through a porin-cytochrome complex in the outer membrane encoded by the Mto pathway. Electrons from Fe(II) oxidation would then be transported to the quinone pool in the inner membrane via periplasmic and inner membrane electron transfer proteins. Directly demonstrating the functionality of genes in neutrophilic iron oxidizers is challenging due to the absence of robust genetic methods. Here, we heterologously expressed a NapC/NirT family tetraheme cytochrome ImoA, encoded by *Slit\_2495*, an inner membrane protein from the Gram-negative Fe(II)-oxidizing bacterium *Sideroxydans lithotrophicus* ES-1, proposed to be involved in extracellular electron transfer to reduce the quinone pool. ImoA functionally replaced the inner membrane *c*-type cytochrome CymA in the Fe(III)-reducing bacterium *Shewanella oneidensis*. We suggest that ImoA may function primarily to oxidize quinol in *S. lithotrophicus*.

**KEYWORDS** Fe(II)-oxidizing bacteria, quinol oxidoreductase, extracellular electron transfer, Mto, extracellular matrix electron transfer, iron oxidizing bacteria

The contribution of chemolithoautotrophic Fe(II)-oxidizing bacteria to iron cycling in the environment has been widely recognized in recent years (1). These bacteria rely on Fe(II) as the electron donor while reducing oxygen to fix carbon dioxide (1–4). Despite the

**Editor** Dianne K. Newman, California Institute of Technology

**Copyright** © 2022 Jain et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Catarina M. Paquete, [cpaquete@itqb.unl.pt](mailto:cpaquete@itqb.unl.pt), or Jeffrey A. Gralnick, [gralnick@umn.edu](mailto:gralnick@umn.edu).

The authors declare no conflict of interest.

**Received** 28 July 2022

**Accepted** 30 August 2022

**Published** 15 September 2022


relevance of these bacteria, the understanding of electron transfer proteins required for their growth and survival in the environment is primarily based on genomic analysis, heterologous expression, and biochemical analysis (5–9) due to their genetic intractability. Although a genetic system has been reported for the obligate Fe(II)-oxidizing bacterium *Mariprofundus ferrooxydans* (10), methods to generate gene deletions in this strain are still lacking, which is a critical tool for validating predictions of functionality.

In Gram-negative bacteria, Fe(II) oxidation would likely require either an outer membrane protein or a periplasmic protein with access to extracellular environment, e.g., through an outer membrane porin, to oxidize Fe(II) extracellularly (5). The electrons released from Fe(II) oxidation would then be transferred to quinone pools in the inner membrane via periplasmic and inner membrane electron transferring proteins (5). Based on genomic analysis, the *bc<sub>1</sub>* complex (Complex III) has been proposed to be the inner membrane quinone reductase in most chemolithoautotrophic Fe(II)-oxidizing bacteria used to generate a proton gradient to drive NADH production (5). A similar function has also been proposed for a NapC/NirT family tetraheme containing protein encoded by *Slit\_2495* during Fe(II) oxidation in *Sideroxydans lithotrophicus* ES-1 (9), a freshwater chemolithoautotroph (5–7). The proposed function of *Slit\_2495* is based on its sequence similarity with CymA, another NapC/NirT protein known to oxidize quinol during reduction of multiple electron acceptors in *Shewanella oneidensis* MR-1 (11). The *S. lithotrophicus* ES-1 genome also encodes a putative *bc<sub>1</sub>* complex (7), making the possible quinone reductase role of *Slit\_2495* functionally redundant (8). Since *S. lithotrophicus* ES-1 is not yet genetically tractable, we heterologously expressed *Slit\_2495* and performed *in vivo* assays in a *S. oneidensis* mutant where CymA was replaced by *Slit\_2495*. Our results revealed that *Slit\_2495* can functionally replace CymA in several modes of anaerobic respiration. *Slit\_2495* has been putatively called CymA in the literature (6, 8). However, phylogenetic analysis shows that *Slit\_2495* is more closely related to NirT from *Pseudomonas* than to CymA from *Shewanella*. Based on our analysis, we propose that *Slit\_2495* be named *imoA* for inner membrane oxidoreductase.

Phylogenetic analysis of ImoA was performed with respect to related NapC/NirT family proteins. We included sequences of CymA from *Shewanella* species, NirT from *Pseudomonas* species, NrfH from *Desulfovibrio* species, the membrane-bound cytochrome *c<sub>552</sub>* sequences from *Nitrosomonas* species along with NapC/NirT family protein sequences from 2 *Sideroxydans* species. Sequences from TorC or DorC were not included in the analysis since both proteins contain 5 heme binding sites compared to 4 heme binding sites in ImoA. Our phylogenetic analysis showed that ImoA clustered close to NapC sequences from *Escherichia coli* and NirT sequences from *Pseudomonas* (Fig. 1). Furthermore, the protein sequence of ImoA is 64% identical to NirT from *Pseudomonas stutzeri* and 54% identical to NapC from *E. coli*, whereas it is only 33% identical to CymA from *S. oneidensis* MR-1.

We tested the ability of ImoA to oxidize quinol pools in the inner membrane by expressing it in a *S. oneidensis* mutant lacking *cymA*. Expression of *imoA* complemented the Fe(III) citrate and anode reduction activities in *S. oneidensis* mutant lacking *cymA* (Fig. 2A and B), consistent with ImoA oxidizing quinols *in vivo*. Furthermore, expression of *imoA* partially complemented the growth defects of the *cymA* mutant strain with dimethyl sulfoxide (DMSO) (Fig. 2C), fumarate (Fig. 2D), and nitrate (Fig. 2E) as electron acceptors, consistent with the ability of ImoA to transfer electrons to multiple periplasmic redox proteins in *S. oneidensis*.


**IMPLICATIONS** In *S. lithotrophicus* ES-1 ImoA is a *c*-type tetraheme cytochrome of the NapC/NirT family proposed to act as a quinone reductase, receiving electrons from iron oxidizing extracellular electron uptake pathway to reduce the quinone pool (5–7). While ImoA may indeed mediate inward electron flow in *S. lithotrophicus*, another quinone reductase (a putative *bc<sub>1</sub>* complex) is also encoded in the genome, and it is unclear what, if any, advantages would be gained by having 2



**FIG 1** Phylogenetic analysis of ImoA with respect to other NapC/NirT proteins. Maximum likelihood tree was built using 2,000 bootstrap replications. Clusters of different NapC/NirT protein families used in the analysis are identified by the brackets labeled with specific proteins and black arrow points to ImoA sequence. The tree is drawn to scale, and the scale bar represents substitutions per site.

different quinone reductase systems. While ImoA was previously proposed to function as a quinone reductase during Fe(II) oxidation (5), the results presented in this work (Fig. 2) suggest the possibility that ImoA may oxidize quinol pools. However, *S. lithotrophicus* has been reported to be metabolically limited, and known to grow only using either Fe(II) or thiosulfate oxidation (7). The potential function of ImoA to oxidize quinol in *S. lithotrophicus* is intriguing from both physiological and ecological perspectives. Fe(II) oxidation pathways are dependent on electron transport chains that require quinone reduction rather than a quinol oxidation (5). If ImoA oxidizes quinol, as indicated by our data, what role could it be playing in *S. lithotrophicus*? By analyzing the *S. lithotrophicus* ES-1 genome, we identified genes that encode putative periplasmic redox proteins that are known to either directly or indirectly accept electrons from NapC/NirT family quinol oxidizing proteins, including *nirS* (Slit\_1129) and several homologs of *nirM*. The protein sequence encoded by *Slit\_1129* is 68% identical to NirS from *P. aeruginosa*, while NirM is known to transfer electrons from NirT to NirS during nitrite reduction (12). Interestingly, MtoD is homologous to NirM (28% identity and 43% similarity), suggesting that *S. lithotrophicus* may be able to reduce nitrite, though this capability has not been demonstrated (7). Our work suggests that *S. lithotrophicus*, and other ImoA-containing Fe(II)-oxidizing bacteria may be more metabolically versatile than initially observed. Importantly, heterologous protein expression can provide insight into the function of novel electron transfer pathways from challenging environmental microbes, complementing biochemical approaches and motivating development of native genetic methods.

All methods are described in the supplementary information (TEXT S1).



**FIG 2** Functionality of ImoA to oxidize quinols in CymA-dependent respiratory pathways. (A) Resting cell Fe(III) citrate reduction: *S. oneidensis* containing pBBR1MCS-2 ( $\Delta$ ),  $\Delta cymA$  containing pBBR1MCS-2 ( $\diamond$ ),  $\Delta cymA$  containing pBBR1MCS-2::cymA ( $\blacksquare$ ) and  $\Delta cymA$  containing pBBR1MCS-2::imoA ( $\bullet$ ). Error bars represent standard deviation of the mean from experiments performed in triplicate. (B) Current density produced by the different  $\Delta cymA$  strains containing pBBR1MCS-2::imoA (black line), pBBR1MCS-2::cymA (gray line), or empty pBBR1MCS-2 (gray dotted line) in a bioelectrochemical reactor. Error bars shown in gray represent standard deviation of the mean from experiments performed in two triplicates. ImoA can functionally replace CymA on growth with different electron acceptors including fumarate (C), DMSO (D) and nitrate (E) while using lactate as the electron donor. *S. oneidensis* containing pBBR1MCS-2 ( $\Delta$ ),  $\Delta cymA$  containing the empty pBBR1MCS-2 vector ( $\diamond$ ),  $\Delta cymA$  containing pBBR1MCS-2::imoA ( $\bullet$ ) and  $\Delta cymA$  containing pBBR1MCS-2::cymA ( $\blacksquare$ ). Error bars represent standard deviation of the mean from experiments performed in triplicate.

### SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

**TEXT S1**, PDF file, 0.1 MB.

## ACKNOWLEDGMENTS

This work was funded by national funds through FCT–Fundação para a Ciência e a Tecnologia, I.P. (FCT), Project MOSTMICRO-ITQB with refs UIDB/04612/2020 and UIDP/04612/2020, and projects PTDC/BIA-BQM/4143/2021 and PD/BD/135153/2017 integrated in the PhD Program in NMR applied to chemistry, materials, and biosciences (PD/00065/2013). This project also received funding from the European Union's Horizon 2020 Research and Innovation Program under the grant agreement number BioCat 867360.

This work was also supported by the Office of Naval Research (N0014-21-1-2166) and the National Science Foundation (MCB-1815584).

We declare that there are no conflicts of interest.

## REFERENCES

- Emerson D, Fleming EJ, McBeth JM. 2010. Iron-oxidizing bacteria: an environmental and genomic perspective. *Annu Rev Microbiol* 64:561–583. <https://doi.org/10.1146/annurev.micro.112408.134208>.
- Krepeski ST, Hanson TE, Chan CS. 2012. Isolation and characterization of a novel biomineral stalk-forming iron-oxidizing bacterium from a circum-neutral groundwater seep. *Environ Microbiol* 14:1671–1680. <https://doi.org/10.1111/j.1462-2920.2011.02652.x>.
- Emerson D, Rentz JA, Lilburn TG, Davis RE, Aldrich H, Chan CS, Moyer CL. 2007. A novel lineage of proteobacteria involved in formation of marine Fe-oxidizing microbial mat communities. *PLoS One* 2:e667. <https://doi.org/10.1371/journal.pone.0000667>.
- Emerson D, Moyer C. 1997. Isolation and characterization of novel iron-oxidizing bacteria that grow at circumneutral pH. *Appl Environ Microbiol* 63:4784–4792. <https://doi.org/10.1128/aem.63.12.4784-4792.1997>.
- He S, Barco RA, Emerson D, Roden EE. 2017. Comparative genomic analysis of neutrophilic iron (II) oxidizer genomes for candidate genes in extracellular electron transfer. *Front Microbiol* 8:1584. <https://doi.org/10.3389/fmicb.2017.01584>.
- Liu J, Wang Z, Belchik SM, Edwards MJ, Liu C, Kennedy DW, Merkley ED, Lipton MS, Butt JN, Richardson DJ, Zachara JM, Fredrickson JK, Rosso KM, Shi L. 2012. Identification and characterization of MtoA: a decaheme c-type cytochrome of the neutrophilic Fe(II)-oxidizing bacterium *Sideroxydans lithotrophicus* ES-1. *Front Microbiol* 3:37. <https://doi.org/10.3389/fmicb.2012.00037>.
- Emerson D, Field EK, Chertkov O, Davenport KW, Goodwin L, Munk C, Nolan M, Woyke T. 2013. Comparative genomics of freshwater Fe-oxidizing bacteria: implications for physiology, ecology, and systematics. *Front Microbiol* 4:254. <https://doi.org/10.3389/fmicb.2013.00254>.
- Beckwith CR, Edwards MJ, Lawes M, Shi L, Butt JN, Richardson DJ, Clarke TA. 2015. Characterization of MtoD from *Sideroxydans lithotrophicus*: a cytochrome c electron shuttle used in lithoautotrophic growth. *Front Microbiol* 6:332. <https://doi.org/10.3389/fmicb.2015.00332>.
- Jain A, Gralnick JA. 2020. Evidence for auxiliary anaerobic metabolism in obligately aerobic Zetaproteobacteria. *ISME J* 14:1057–1062. <https://doi.org/10.1038/s41396-020-0586-6>.
- Jain A, Gralnick JA. 2021. Engineering lithoheterotrophy in an obligate chemolithoautotrophic Fe(II) oxidizing bacterium. *Sci Rep* 11:1–6. <https://doi.org/10.1038/s41598-021-81412-3>.
- Marritt SJ, Lowe TG, Bye J, McMillan DG, Shi L, Fredrickson J, Zachara J, Richardson DJ, Cheesman MR, Jeuken LJ, Butt JN. 2012. A functional description of CymA, an electron-transfer hub supporting anaerobic respiratory flexibility in *Shewanella*. *Biochem J* 444:465–474. <https://doi.org/10.1042/BJ20120197>.
- Hasegawa N, Arai H, Igarashi Y. 2001. Two c-type cytochromes, NirM and NirC, encoded in the *nir* gene cluster of *Pseudomonas aeruginosa* act as electron donors for nitrite reductase. *Biochem Biophys Res Commun* 288:1223–1230. <https://doi.org/10.1006/bbrc.2001.5919>.