

**STUDY PROTOCOL**

**Open Access**

# The value of Innovative ICT guided disease management combined with Telemonitoring in Outpatient clinics for Chronic Heart failure patients. Design and methodology of the IN TOUCH study: a multicenter randomised trial

Arjen E de Vries<sup>1\*</sup>, Richard M de Jong<sup>1</sup>, Martje HL van der Wal<sup>1</sup>, Tiny Jaarsma<sup>3</sup>, Rene B van Dijk<sup>2</sup> and Hans L Hillege<sup>1</sup>

## Abstract

**Background:** Although the value of telemonitoring in heart failure patients is increasingly studied, little is known about the value of the separate components of telehealth: ICT guided disease management and telemonitoring. The aim of this study is to investigate the value of telemonitoring added to ICT guided disease management (DM) on the quality and efficiency of care in patients with chronic heart failure (CHF) after a hospitalisation.

**Methods/Design:** The study is divided in two arms; a control arm (DM) and an intervention arm (DM+TM) in 10 hospitals in the Netherlands. In total 220 patients will be included after worsening of CHF (DM: N = 90, DM+TM: N = 130). Total follow-up will be 9 months. Data will be collected at inclusion and then after 2 weeks, 4.5 and 9 months. The primary endpoint of this study is a composite score of: 1: death from any cause during the follow-up of the study, 2: first readmission for HF and 3: change in quality of life compared to baseline, assessed by the Minnesota Living with Heart failure Questionnaire. The study has started in December 2009 and results are expected in 2012.

**Conclusions:** The IN TOUCH study is the first to investigate the effect of telemonitoring on top of ICT guided DM on the quality and efficiency of care in patients with worsening HF and will use a composite score as its primary endpoint.

**Trial registration:** Netherlands Trial Register (NTR): NTR1898

## Background

Heart failure (HF) is the most common hospital discharge diagnosis in elderly patients [1]. Between the age of 70 and 80 years the incidence of HF is 10 to 20%. HF is associated with high mortality and morbidity, readmission rates and costs [1]. The readmission rates vary between 25% and 50% within 6 months after the first hospitalisation for HF, with a higher readmission rate within the first month after discharge[2,3]. The costs related to HF contribute to 1-2% of all healthcare

expenditures and are mainly the result of hospital stay [4-6]. Because of an increasing shortage of resources, HF is a major public health problem and therefore, a more effective and efficient organisation of care for HF patients needs to be reconsidered. A first step in organising treatment and care for patients with chronic HF more efficiently, was the implementation of specialised outpatient HF clinics. In the recent European Society of Cardiology (ESC) guidelines, HF management programmes are strongly recommended for all patients with HF [1] and HF clinics are considered as 'usual care' in several European countries [7]. A widely used way to implement HF management is the use of specific disease management (DM) programs.

\* Correspondence: a.e.de.vries@thorax.umcg.nl

<sup>1</sup>Thoraxcenter, Department of Cardiology, University Medical Centre Groningen, Groningen, The Netherlands

Full list of author information is available at the end of the article

DM can be defined as an intervention, designed to manage a chronic disease and to reduce hospital readmissions, using a systematic approach to care and potentially employing multiple treatment modalities [8]. Control and cost effectiveness are substantial components of a DM program. Randomised studies suggest that DM programs can reduce readmissions for HF or cardiovascular disease with 30% [7,9,10] and significantly decrease mortality rates [11]. Yu et al [12] described that DM for HF patients, as recommended by the ESC guidelines, [1] are effective in reducing hospital readmissions and mortality rate [13]. However, inconsistent findings for readmission and mortality rates have been found, probably due to the variety of components and practical applications of the DM programs.

We recently reported results of the COACH study, a study on the effect of a nurse led DM program on clinical outcome [14], in which the positive effects of a DM program on readmission were not confirmed, although there was a trend to a reduction of mortality in the intervention groups. The INH study [15] on the effect of DM in HF, showed that a DM program compared to usual care could reduce mortality but not hospitalisation rates. Important components of this program were patient education, optimisation of medical therapy, psychosocial support and an easy access to healthcare. An important aspect for the treatment of HF patients is the prescription of HF related medication at an optimal dose i.e. ACE-inhibitors, beta-blockers, and aldosteronantagonists. The up titration to optimal dosage is an aspect that often takes place at a HF outpatient clinic. However, data from the Euro Heart Failure Survey showed us that guideline adherence for HF medication although improving still is not optimal [16]. In the IMPROVE study, dedicated HF clinics were associated with greater use of cardiac resynchronisation therapy and a better HF education, but not with better guideline adherence to medication [17]. Health information technology, integrated into a DM program might facilitate adherence to guidelines of health professionals [18]. With new information and communication technology (ICT), healthcare providers can be supported in the diagnosis, treatment and follow up of HF patients by expert computerised systems, based on guidelines and protocols [19]. These systems can be used to optimise medication according to guidelines and provide structural support and education [20]. We were the first to report promising findings on ICT guided DM in terms of higher doses of recommended HF medication and lower readmissions [21,22]. Another promising ICT tool is telemonitoring. Telemonitoring is often used to monitor patients at home and guide patients to take action in case of deterioration, but it also can be used to up-titrate medication according to guidelines at distance [23]. There is support that remote monitoring of patients with HF can reduce hospitalisation and mortality rates, [24,25] however results on clinical outcome and efficacy are inconclusive

and limited [26-28]. There are also recent study's that where not successful in their primary endpoints [29,30]. Furthermore, cost-effectiveness of these systems has not been thoroughly evaluated. It can be concluded that the overall effects of telemonitoring are inconclusive. To summarise, due to a growing population of patients with HF and an expected shortage of healthcare providers in the near future, there is a need to seek to more cost effective and efficient ways of providing optimal care for HF patients, including a better adherence to guidelines. ICT guided DM tools in combination with telemonitoring could be of important value [31,32]. At the same time there is substantial data that the adaptation and implementation of those systems is lacking [33]. The experiences with such a system however are fragmented. User resistance is described as a major obstacle in the adoption of these computerised tools. More insight in user resistance and experienced barriers in using ICT guided DM tools is needed to successfully implementing such tools [34].

The IN TOUCH study will investigate the effect of telemonitoring in addition to an ICT guided DM system on the quality and efficiency of care for patients after worsening HF. This is the first study investigating a combination of two newly developed ICT interventions in a group of chronic HF patients on clinical outcome, adherence to guidelines, cost effectiveness and quality of life.

This study will add important information to other telemonitoring studies because of its strong commitment to ICT guided DM, the chosen composite endpoint, a strong focus on cost-effectiveness and the investigation of the influence of user aspects as resistance and barriers that accompany the use of modern healthcare related ICT tools.

## Methods/Design

### Study hypothesis

Telemonitoring added to ICT guided DM improves prognosis and quality of life in patients with HF compared to ICT guided DM alone.

### Aim of the study

The primary aim of this study is to assess the effect of telemonitoring on top of an ICT guided DM system in patients after worsening HF on the combined endpoint of death, readmission and quality of life, compared to patients treated with ICT guided DM alone.

Secondary aims of the study are;

- To assess the effect of telemonitoring in addition to an ICT guided DM system compared to an ICT guided DM alone on the separate components of the combined endpoint (death, readmission and quality of life).

- To determine the cost benefit ratio of ICT guided DM with telemonitoring compared to ICT guided DM alone.

### Study design

A multicentre, randomised study in which in total 220 HF patients (NYHA II-IV) will be included. Patients will be randomised to the ICT guided DM (control) arm (N = 90) or into the ICT guided DM with telemonitoring (intervention) arm (N = 130) (Figure 1).

### Study population

All patients admitted to the intensive care/coronary care unit or cardiology ward for HF or visiting the outpatient clinic with worsening HF who need treatment or adjustment of oral or intravenous diuretics, can be included in the study. Other inclusion criteria are; evidence of structural underlying heart disease, left ventricular ejection fraction  $\leq 45\%$  and age of at least 18 years. Reasons for exclusion are myocardial infarction in the past month, cardiac invasive intervention (percutaneous coronary intervention, coronary arterial bypass, valve replacement, heart transplantation, or cardiac resynchronisation therapy) in the past 6 months or planned in the next 3 months, weight  $> 200$  kg, actual haemodialysis and the use of other telemonitoring systems (Table 1).

### Primary endpoint

The primary endpoint of the study is a composite, weighted score consisting of values for mortality, HF

### Table 1 In and exclusion criteria IN TOUCH

#### Inclusion criteria

Worsening HF defined as signs of fluid retention (peripheral oedema/ congestion)  
 needing an increase of the dose of diuretics (i.v. or oral)  
 Evidence for structural underlying heart disease  
 Documented reduced left ventricular ejection fraction (LVEF)  $\leq 45\%$   
 18 years of age  
 Patients have to be able to understand content of and willing to provide informed consent.

#### Exclusion criteria

History of myocardial infarction in the previous month  
 Life expectation less than 1 year  
 Undergone cardiac invasive intervention within the last 6 months (PCI, CABG, HTX, valve replacement, CRT implantation)  
 A planned procedure for PCI, CABG, HTX, CRT implantation or valve replacement in the following 3 months  
 Evaluation for heart transplantation prior to or during the study  
 Weight more than 200 kilogram  
 The use of telemonitoring devices at home  
 Haemodialysis


readmission and change in quality of life between end of the study and baseline measured with the Minnesota Living with HF Questionnaire (MLHFQ), adapted from the A-HeFT study [35] (Table 2). A readmission for HF is defined as an overnight hospital stay for HF or directly related to HF. The readmissions for HF will be blinded adjudicated by an endpoint committee. When data on quality of life are missing, the worst-case score for that component of the composite endpoint will be used in the analysis.

### Secondary endpoints

Secondary endpoints of the study are the separate components of the primary endpoint. Other secondary endpoints are the total number and duration of all hospital

**Table 2 score system for primary endpoint IN TOUCH**

| End point                                    | Score    |
|--|----------|
| Death (at any time during study)             | -3       |
| Survival to end of study                     | 0        |
| First readmission for heart failure          | -1       |
| No readmission for heart failure             | 0        |
| <i>Change in quality of life at 9 months</i> |          |
| Improvement $\geq 20$ units                  | +2       |
| Improvement by 10 until 19 units             | +1       |
| No improvement by -9 until +4 units          | 0        |
| Worsening by +5 until +9 units               | -1       |
| Worsening by $\geq 10$ units                 | -2       |
| <i>Possible score</i>                        | -6 to +2 |



**Figure 1** Flowchart of the inclusion.

admissions, treatment according to the guidelines using the criteria of the Guidelines Adherence Indicator-3 [36], number of visits to the outpatient HF clinic, patient and carer satisfaction, and cost-benefit ratio.

#### **Ethics statement**

On March 2009 this study design has become ethical approval (M09.070323) given by the medical ethical commission (MEC) of the medical university of Groningen (UMCG).

The study has started (first patient) in December 2009.

#### **Incremental cost-effectiveness ratio (ICER)**

Costs A distinction will be made between intervention costs and resource utilization costs. The intervention costs consist of the costs of the DM system and the costs of the telemonitoring devices and will be calculated as a lump-sum over the study's follow-up period. Resource utilisation costs will be estimated by preparing a structured data collection form to collect detailed information regarding scheduled and non-scheduled outpatient clinic visits and hospital admissions (both HF and non-HF related), ward type (e.g. intensive/coronary care unit, cardiology, general internal medicine), and cardiovascular procedures/operations. In addition, a patient questionnaire will be administered at 4.5 months and 9 months of follow-up to collect complementary data on general practitioner visits, home care utilization, and nursing home admissions. Unit costs will be estimated by using the Dutch guidelines for cost calculations and inflated to current price levels using a general consumer price index. Indirect costs, such as productivity losses, will not be taken into account.

#### **Quality Adjusted Life Years (QALYs)**

Preference-based quality of life scores will be obtained by administering the EQ-5D to all patients in both the control group and the two intervention groups at baseline, 4.5 months of follow-up, and 9 months of follow-up. QALYs will subsequently be estimated by calculating the area of the two trapezoids that result from linear extrapolation of the three quality of life scores.

#### **Cost-effectiveness**

The balance between costs and effects will be assessed by estimating the incremental cost per QALY gained (ICER) for the intervention group compared with the control group.

The time horizon over which the costs and effects of the different treatment strategies will be compared is equivalent to that observed during the period of the study (i.e. no future projections will be made). Uncertainty surrounding the ICER will be represented through the use of cost-effectiveness acceptability curves, which show, for each possible value of  $\lambda$  (i.e. the societal

willingness-to-pay for one additional QALY), the probability that the intervention will be cost-effective.

#### **Randomisation and data collection**

Patients can be randomised during admission for HF, and in case of worsening of HF at the outpatient clinic. After confirmation of eligibility and written informed consent, patients will be included into the study. Patients will be randomised into the control or intervention group. Patient characteristics and clinical variables will be collected at baseline, and 2 weeks and 9 months after discharge. Echocardiography, ECG, and laboratory analysis will be performed at baseline during hospitalisation and at the end of the study. Quality of life as part of the primary endpoint, measured with the Minnesota Living with HF Questionnaire, will be collected at baseline and at 9 months. Data about utilisation of resources will be collected prospectively and comprise components of direct costs, i.e. scheduled and non-scheduled outpatient visits and hospital admissions.

Questionnaires about self-care behaviour, anxiety, depression, compliance and medical technology assessment (MTA) will be completed before discharge, and at 4.5 and 9 months (Table 3). Patients should be included preferably as soon as possible but at least within a period of time of 14 days after discharge or after the first visit at the HF clinic with worsening HF.

#### **Control group**

##### ***ICT guided disease management system without telemonitoring***

Patients in this group receive care guided by an ICT DM system. This system supports DM in a fully automatic way, assisting the HF nurse to optimise pharmacological and non-pharmacological treatment, evaluate treatment and adjust therapy to optimal levels, according to current HF guidelines. The patient will receive tailored education and counselling on the HF regimen, symptom management and improvement of the pharmacological and non pharmacological regimen.

The system mainly works as a computer decision support system. Based on the input of data from physical examination, medical history, questionnaires and nursing assessment, the system provides an advice to healthcare providers according to the actual guidelines, including up titration of HF medication to optimal doses.

#### **Intervention group**

##### ***ICT guided disease management with telemonitoring***

Patients in the DM with telemonitoring group will be treated with the above described ICT guided DM system in combination with the following integrated telemonitoring devices that will be installed at the patients' home;

**Table 3 Questionnaires and medical assessment used in the IN TOUCH**

| Questionnaires  | Baseline | 2 weeks | 4,5 month | 9 month |
|---|----------|---------|-----------|---------|
| Minnesota Living with Heart failure Questionnaire (MLHFQ) | •        |         |           | •       |
| Revised Heart failure Compliance                          | •        |         |           | •       |
| Disability Rating Index (VAS)                             | •        |         |           | •       |
| Hospital Anxiety and Depression scale (HADS)              | •        |         |           | •       |
| Heart failure self-care behaviour scale (EHFScB)          | •        |         |           | •       |
| Medical technology assessment (MTA)                       |          |         | •         | •       |
| EQ-5D   | •        |         | •         | •       |
| Satisfaction questionnaire for patients and providers     |          |         |           | •       |
| <b>Medical assessment</b>                                 |          |         |           |         |
| NYHA score  | •        | •       |           | •       |
| Echocardiography  | •        |         |           |         |
| ECG   | •        | •       |           | •       |
| Physical examination                                      | •        | •       |           | •       |
| Laboratory tests  | •        | •       |           | •       |

- weighing scale; patients will be instructed to weigh daily.

- blood pressure meter; patients will be instructed to measure their blood pressure daily during up titration of medication.

- ECG; patients have to perform an ECG twice a week during up titration of beta-blockers.

- Health monitor; an interactive monitor collects data from the weighing scale, blood pressure meter and ECG device and will respond to the patients' collected data. Data will be directly transmitted to the DM system in the hospital.

In case of a deviation of individualised predefined ranges of weight, blood pressure or heart rhythm, the health monitor will automatically generate supplementary questions directly to the patient to evaluate the actual health situation. This data, measurements and subjective patient information (predefined multiple choice questions about HF symptoms) will be transferred to the computerised DM in the hospital. Furthermore, the health monitor will generate an advice about the non-pharmacological treatment, for example regarding compliance with fluid and sodium restriction. When the data collected by the system deviates from predefined ranges, the HF nurse will be informed automatically by mobile phone and email. In that case, the HF nurse will contact the patient by phone within two hours and further discusses symptoms and treatment. If data are outside any reference range in combination with symptoms of deterioration, patients will receive a message from the health monitor that they will be contacted by the HF nurse.

#### Sample size calculation

Group sample sizes of 130 patients for the group treated with telemonitoring and ICT guided DM and 90 patients treated with ICT guided DM achieve 80% power to

detect superiority for telemonitoring using a one-sided, two-sample t-test. The margin of equivalence is 0.0. The true difference between the means is assumed to be 0.8. The significance level (alpha) of the test is 0.025. The data are drawn from populations with standard deviations of - 2.0 and + 2.0. These figures are based on a publication [37] in which the effect of the combination of isosorbidedinitrate and hydralazin was investigated in patient with HF. In this study a difference of 0.4 was demonstrated regarding the composite endpoint. Using telemonitoring, we expect to find a larger difference especially in the QoL domain. It has to be addressed that no information is currently available regarding the sensitivity of the composite endpoint in a setting of new onset or worsening HF patients.

#### Statistical analyses

The primary analysis will consist of a comparison of the composite endpoint scores (Table 2) between ICT guided DM in combination with telemonitoring, compared to ICT guided DM without telemonitoring. The test will be performed using a two-sample t-test, and two-sided 95% confidence intervals will be constructed to describe the treatment differences. An analysis of covariance will be used to test for the treatment effect controlling for different baseline characteristics. The results will be analysed using an intention-to-treat analysis including the full set of all randomised patients (primary efficacy population). The primary efficacy population will be analysed at endpoint for composite score. Secondary endpoints involving e.g. individual evaluation of deaths, hospitalisations, and QOL after 9 months will be analysed over the entire course of the study using appropriate methods. Two types of economic analyses will be performed: these include a cost-consequence analysis (CCA) for a disaggregated examination of resource costs and health

outcomes associated with the alternative intervention; and cost-effectiveness analysis (CEA) in which the alternative intervention is examined in light of total cost per unit of health outcome. Thus CCA will be performed using the primary outcome of the study as the measure of effectiveness. For this, the annual cost per patient treated to postpone or prevent one patient experiencing a cardiovascular death or hospital admission for worsening HF within the trial will be calculated [38]. For CEA, the incremental cost-effectiveness ratios (ICERs), in terms of cost per life year gained (LYG), will be estimated given that there was a significant increase in survival with DM and telemonitoring. CEA will not be performed in case no reduction will be observed in cardiovascular or all cause mortality. In case of missing data for the composite end point score, the worst-case scenario will be assumed for the primary analyses. Patients lost to follow-up will be assumed to have died (-3) and those without a quality of life measurement will be assigned the worst score (-2).

### Study organisation

To include 220 patients in this study, 10 hospitals in the Netherlands are participating in the IN TOUCH study. The first patients are recruited in December 2009; the end of the study is expected to be in September 2012.

### Support and monitoring

The study is supported and monitored by the Trial Coordination Centre (TCC), a contract research organisation for clinical trials. Both the quality of the research data and of the intervention will be structurally monitored according to the GCP guidelines and TCC standards (ISO 9000:2001).

### Discussion

In the last decades, the introduction of ICT in healthcare promised an improved quality of care while reducing work load, and resulting in a more cost effective system. This might be realised by the use of computer guided decision support systems and telemonitoring [20,31,32,39,40]. The IN TOUCH study is the first to investigate the effect of an ICT guided DM system in combination with telemonitoring in patients with HF. In recent years there has been much research in the field of HF and DM. This has resulted in the implementation of DM programmes in the ESC guidelines [1]. The development of these guidelines has enabled the creation of computer aided decision support programmes [39]. In the recent guidelines the application of ICT by DM programs is recommended. Although there has been progression in the development of ICT guided tools and these tools have become practical equipment for enhanced decision making, healthcare providers still experience great barriers in using and implementing them for several, often unclear reasons [33,34].

Mortality and morbidity are important outcomes in studies in HF patients. Quality of life becomes a more important issue for patients and may be even more important for them than survival or readmissions. Therefore, a composite primary endpoint, including quality of life, mortality and morbidity was chosen for the IN TOUCH study. Another reason for this composite endpoint is that for HF a high standard of care has been established which makes it more difficult to investigate the added effect of new therapeutic options. Beside the statistical advantage of a composite endpoint, the study becomes less costly and results of promising new treatment may occur earlier with this selected design. The strength of this study is the important role of ICT guided DM, telemonitoring, the composite endpoint, including quality of life, a strong focus on cost effectiveness and the emphasis on experienced user resistance and barriers regarding ICT tools in healthcare.

### Abbreviations

ICT: Information Communication Technology; DM: Disease Management; HF: Heart Failure; ESC: European Society of Cardiology; NYHA: New York Heart Association; MLHFQ: Minnesota Living with Heart Failure Questionnaire; QALYs: Quality Adjusted Life Years; ICER: Incremental Cost Effectiveness Ratio; MTA: Medical Technology Assessment; ECG: Electrocardiography; CCA: Cost-Consequence Analysis; CEA: Cost Effectiveness Analysis; TCC: Trial Coordination Centre

### Acknowledgements

This project was funded by the Dutch Government of health, section pharmaceuticals and medical technology (GMT)

### Author details

<sup>1</sup>Thoraxcenter, Department of Cardiology, University Medical Centre Groningen, Groningen, The Netherlands. <sup>2</sup>Martini hospital Groningen, Department of Cardiology, Groningen, The Netherlands. <sup>3</sup>Linköpings universitet, Department of Social- and Welfare Studies, Norrköping, Sweden.

### Authors' contributions

RD, RJ, TJ and HH designed the study. AV drafted the article and all authors contributed to the final concept. All authors have read and approved the final manuscript.

### Competing interests

The authors declare that they have no competing interests.

Received: 25 March 2011 Accepted: 13 July 2011

Published: 13 July 2011

### References

1. Dickstein K, Cohen-Solal A, Filippatos G, et al: ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the diagnosis and treatment of acute and chronic heart failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur J Heart Fail* 2008, **10**(10):933-989.
2. Krumholz HM, Parent EM, Tu N, et al: Readmission after hospitalization for congestive heart failure among Medicare beneficiaries. *Arch Intern Med* 1997, **157**(1):99-104.
3. Aranda JM, Johnson JW, Conti JB: Current trends in heart failure readmission rates: analysis of Medicare data. *Clin Cardiol* 2009, **32**(1):47-52.

4. Berry C, Murdoch DR, McMurray JJ: **Economics of chronic heart failure.** *Eur J Heart Fail* 2001, **3**(3):283-291.
5. Lee WC, Chavez YE, Baker T, Luce BR: **Economic burden of heart failure: a summary of recent literature.** *Heart Lung* 2004, **33**(6):362-371.
6. Stewart S: **Financial aspects of heart failure programs of care.** *Eur J Heart Fail* 2005, **7**(3):423-428.
7. McAlister FA, Lawson FM, Teo KK, Armstrong PW: **A systematic review of randomized trials of disease management programs in heart failure.** *Am J Med* 2001, **110**(5):378-384.
8. Ellrodt G, Cook DJ, Lee J, Cho M, Hunt D, Weingarten S: **Evidence-based disease management.** *JAMA* 1997, **278**(20):1687-1692.
9. Phillips CO, Wright SM, Kern DE, Singa RM, Shepperd S, Rubin HR: **Comprehensive discharge planning with postdischarge support for older patients with congestive heart failure: a meta-analysis.** *JAMA* 2004, **291**(11):1358-1367.
10. Windham BG, Bennett RG, Gottlieb S: **Care management interventions for older patients with congestive heart failure.** *Am J Manag Care* 2003, **9**(6):447-459.
11. Clark RA, Inglis SC, McAlister FA, Cleland JG, Stewart S: **Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis.** *BMJ* 2007, **334**(7600):942.
12. Yu DS, Thompson DR, Lee DT: **Disease management programmes for older people with heart failure: crucial characteristics which improve post-discharge outcomes.** *Eur Heart J* 2006, **27**(5):596-612.
13. van der Wal MH, Jaarsma T, van Veldhuisen DJ: **Non-compliance in patients with heart failure; how can we manage it?** *Eur J Heart Fail* 2005, **7**(1):5-17.
14. Jaarsma T, van der Wal MH, Lesman-Leegte I, et al: **Effect of moderate or intensive disease management program on outcome in patients with heart failure: Coordinating Study Evaluating Outcomes of Advising and Counseling in Heart Failure (COACH).** *Arch Intern Med* 2008, **168**(3):316-324.
15. Coletta AP, Cleland JG, Cullington D, Clark AL: **Clinical trials update from Heart Rhythm 2008 and Heart Failure 2008: ATHENA, URGENT, INH study, HEART and CK-1827452.** *Eur J Heart Fail* 2008, **10**(9):917-920.
16. Komajda M, Follath F, Swedberg K, et al: **The EuroHeart Failure Survey programme—a survey on the quality of care among patients with heart failure in Europe. Part 2: treatment.** *Eur Heart J* 2003, **24**(5):464-474.
17. Albert NM, Fonarow GC, Yancy CW, et al: **Influence of dedicated heart failure clinics on delivery of recommended therapies in outpatient cardiology practices: findings from the Registry to Improve the Use of Evidence-Based Heart Failure Therapies in the Outpatient Setting (IMPROVE HF).** *Am Heart J* 2010, **159**(2):238-244.
18. Chaudhry B, Wang J, Wu S, et al: **Systematic review: impact of health information technology on quality, efficiency, and costs of medical care.** *Ann Intern Med* 2006, **144**(10):742-752.
19. Garcia-Lizana F, Sarría-Santamera A: **New technologies for chronic disease management and control: a systematic review.** *J Telemed Telecare* 2007, **13**(2):62-68.
20. Tierney WM, Overhage JM, Takesue BY, et al: **Computerizing guidelines to improve care and patient outcomes: the example of heart failure.** *J Am Med Inform Assoc* 1995, **2**(5):316-322.
21. Vries de AE, Dijk van RB, Hendriks M: **Medical nurses and the use of expert software in the treatment of patients with congestive heart failure: first year experience.** *Eur J Heart Fail Supplement I* 2002.
22. de Vries AE, van der Wal MH, Bedijn W, et al: **Follow-up and treatment of an instable patient with heart failure using telemonitoring and a computerised disease management system: A case report.** *Eur J Cardiovasc Nurs* 2011.
23. Antonicelli R, Mazzanti I, Abbatecola AM, Parati G: **Impact of home patient telemonitoring on use of beta-blockers in congestive heart failure.** *Drugs Aging* 2010, **27**(10):801-805.
24. Klersy C, De Silvestri A, Gabutti G, Regoli F, Auricchio A: **A meta-analysis of remote monitoring of heart failure patients.** *J Am Coll Cardiol* 2009, **54**(18):1683-1694.
25. Riley JP, Cowie MR: **Telemonitoring in heart failure.** *Heart* 2009, **95**(23):1964-1968.
26. Wootton R, Gramotnev H, Hailey D: **A randomized controlled trial of telephone-supported care coordination in patients with congestive heart failure.** *J Telemed Telecare* 2009, **15**(4):182-186.
27. Clarke M, Shah A, Sharma U: **Systematic review of studies on telemonitoring of patients with congestive heart failure: a meta-analysis.** *J Telemed Telecare* 2011, **17**(1):7-14.
28. Polisena J, Tran K, Cimon K, et al: **Home telemonitoring for congestive heart failure: a systematic review and meta-analysis.** *J Telemed Telecare* 2010, **16**(2):68-76.
29. Chaudhry SI, Matterna JA, Curtis JP, et al: **Telemonitoring in Patients with Heart Failure.** *N Engl J Med* 2010.
30. Koehler F, Winkler S, Schieber M, et al: **Telemedical Interventional Monitoring in Heart Failure (TIM-HF), a randomized, controlled intervention trial investigating the impact of telemedicine on mortality in ambulatory patients with heart failure: study design.** *Eur J Heart Fail* 2010, **12**(12):1354-1362.
31. Garg AX, Adhikari NK, McDonald H, et al: **Effects of computerized clinical decision support systems on practitioner performance and patient outcomes: a systematic review.** *JAMA* 2005, **293**(10):1223-1238.
32. Craig JC, Irwig LM, Stockler MR: **Evidence-based medicine: useful tools for decision making.** *Med J Aust* 2001, **174**(5):248-253.
33. Varonen H, Kortteisto T, Kaila M: **What may help or hinder the implementation of computerized decision support systems (CDSSs): a focus group study with physicians.** *Fam Pract* 2008, **25**(3):162-167.
34. Kawamoto K, Houlihan CA, Balas EA, Lobach DF: **Improving clinical practice using clinical decision support systems: a systematic review of trials to identify features critical to success.** *BMJ* 2005, **330**(7494):765.
35. Taylor AL, Lindenfeld J, Ziesche S, et al: **Outcomes by gender in the African-American Heart Failure Trial.** *J Am Coll Cardiol* 2006, **48**(11):2263-2267.
36. Komajda M, Lapuerta P, Hermans N, et al: **Adherence to guidelines is a predictor of outcome in chronic heart failure: the MAHLER survey.** *Eur Heart J* 2005, **26**(16):1653-1659.
37. Taylor AL, Ziesche S, Yancy C, et al: **Combination of isosorbide dinitrate and hydralazine in blacks with heart failure.** *N Engl J Med* 2004, **351**(20):2049-2057.
38. Mauskopf JA, Paul JE, Grant DM, Stergachis A: **The role of cost-consequence analysis in healthcare decision-making.** *Pharmacoeconomics* 1998, **13**(3):277-288.
39. Walsh MN, Yancy CW, Albert NM, et al: **Electronic health records and quality of care for heart failure.** *Am Heart J* 2010, **159**(4):635-642.
40. Inglis SC, Clark RA, McAlister FA, et al: **Structured telephone support or telemonitoring programmes for patients with chronic heart failure.** *Cochrane Database Syst Rev* 2010, **8**:CD007228.

#### Pre-publication history

The pre-publication history for this paper can be accessed here:  
<http://www.biomedcentral.com/1472-6963/11/167/prepub>

doi:10.1186/1472-6963-11-167

**Cite this article as:** de Vries et al.: The value of Innovative ICT guided disease management combined with Telemonitoring in Outpatient clinics for Chronic Heart failure patients. Design and methodology of the IN TOUCH study: a multicenter randomised trial. *BMC Health Services Research* 2011 **11**:167.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

