



# Complete Genome Sequence of *Acinetobacter schindleri* SGAir0122 Isolated from Singapore Air

Carmon Kee,<sup>a</sup>  Ana Carolina M. Junqueira,<sup>a\*</sup> Akira Uchida,<sup>a</sup> Rikky W. Purbojati,<sup>a</sup> James N. I. Houghton,<sup>a</sup> Caroline Chénard,<sup>b</sup> Anthony Wong,<sup>a</sup> Megan E. Clare,<sup>a</sup> Kavita K. Kushwaha,<sup>a</sup> Deepa Panicker,<sup>a</sup> Alexander Putra,<sup>a</sup> Nicolas E. Gaultier,<sup>a</sup> Balakrishnan N. V. Premkrishnan,<sup>a</sup> Cassie E. Heinle,<sup>a</sup> Vineeth Kodengil Vettath,<sup>a</sup> Daniela I. Drautz-Moses,<sup>a</sup> Stephan C. Schuster<sup>a</sup>

<sup>a</sup>Singapore Centre for Environmental Life Sciences Engineering, Nanyang Technological University, Singapore

<sup>b</sup>Asian School of the Environment, Nanyang Technological University, Singapore

**ABSTRACT** *Acinetobacter schindleri* strain SGAir0122 was isolated from tropical air samples collected in Singapore. The prevalence of nosocomial infection caused by this Gram-negative bacterium indicates its clinical significance as an opportunistic human pathogen. Its complete genome consists of one chromosome of 3.105 Mb and a plasmid of 181 kb.

*Acinetobacter schindleri* is a nonmotile, aerobic, Gram-negative bacterium (1). *A. schindleri* is an opportunistic human pathogen affecting patients in intensive care units and immunocompromised patients (2), contributing to nosocomial infections and outbreaks worldwide (3). This species can be found in a wide range of natural environments, such as in soil and water, as well as in hospitals (1), due to its ability to survive on inert surfaces for a long duration (4). Phenotypic identification of the isolates remains indiscriminative and may lead to underestimation of infections caused by *A. schindleri*; thus, the complete genome sequences of *A. schindleri* strains may be of great help in providing molecular markers for identification.

Air samples were collected outdoor in Singapore (1.345°N, 103.679°E) using an Andersen single-stage impactor (SKC, Inc., USA). Strain SGAir0122 was isolated from trypticase soy agar (Becton, Dickinson, USA) incubated at 30°C after airborne microorganisms were impacted onto it. The isolated strain was cultured in Luria-Bertani broth overnight at room temperature. Prior to DNA extraction, an axenic culture was obtained by streaking. Genomic DNA was extracted using the Wizard genomic DNA purification kit (Promega, USA) according to the manufacturer's recommended protocol. Whole-genome shotgun libraries were constructed using a SMRTbell template prep kit version 1.0 (Pacific Biosciences, USA), followed by single-molecule real-time (SMRT) sequencing conducted on a PacBio RS II (Pacific Biosciences) sequencer.

The PacBio RS II platform provided 68,481 long reads (208-fold coverage) for the isolate. The complete processing of the assembly included preassembly, *de novo* assembly with the Hierarchical Genome Assembly Process (HGAP) version 3 implemented in the PacBio SMRT Analysis algorithm version 2.3.0 (5), polishing with Quiver (5), and error correction using Pilon version 1.16 (6) to improve accuracy. The consensus assembly generated two contigs, one chromosome with 3,104,554 bp (202-fold coverage), and one plasmid with 180,768 bp (321-fold coverage). The chromosomal contig showed an average G+C content of 42.7%.

The complete genome presented 99.3% marker identity with that of the *Acinetobacter* genus, as predicted by the phylum-specific automated phylogenomic inference pipeline Phyla\_AMPHORA using a minimum confidence of 1.00 (7). Further taxonomical identification was conducted using the average nucleotide identity method with the Microbial Species Identifier (MiSI) (8), which revealed a 97.5% similarity with the

Received 18 May 2018 Accepted 22 May 2018 Published 28 June 2018

**Citation** Kee C, Junqueira ACM, Uchida A, Purbojati RW, Houghton JN, Chénard C, Wong A, Clare ME, Kushwaha KK, Panicker D, Putra A, Gaultier NE, Premkrishnan BN, Heinle CE, Vettath VK, Drautz-Moses DI, Schuster SC. 2018. Complete genome sequence of *Acinetobacter schindleri* SGAir0122 isolated from Singapore Air. Genome Announc 6:e00567-18. <https://doi.org/10.1128/genomeA.00567-18>.

**Copyright** © 2018 Kee et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Stephan C. Schuster, [scschuster@ntu.edu.sg](mailto:scschuster@ntu.edu.sg).

\* Present address: Ana Carolina M. Junqueira, Departamento de Genética, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.

C.K. and A.C.M.J. contributed equally to this work.

available reference genome of *A. schindleri* CIP 107287 (GenBank accession number KB849571).

Annotation of the genome was performed with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) version 4.2 (9). The complete genome was predicted to contain 3,206 genes, including 2,850 protein-coding genes, 21 rRNA subunits (5S, 16S, and 23S), 86 tRNAs, and 4 noncoding RNAs, as well as 245 pseudogenes. Functional annotation using the Rapid Annotations using Subsystems Technology (RAST) server (10–12) classified 96 genes as being associated with virulence and defense, indicating a moderate virulence. There were 120 genes found to have a role in stress response and 20 genes found to be involved in secondary metabolism. This could confer the organism's advantages to outcompete other bacteria with secondary metabolites and contribute to its long-term survival on inert surfaces.

**Accession number(s).** The complete genome sequence of *A. schindleri* strain SGAir0122, including the plasmid pSGAir0122, has been deposited in DDBJ/EMBL/GenBank under the accession numbers [CP025618](#) (chromosome) and [CP025619](#) (plasmid pSGAir0122).

## ACKNOWLEDGMENT

This work was supported by a Singapore Ministry of Education Academic Research Fund tier 3 grant (MOE2013-T3-1-013).

## REFERENCES

- Bergogne-Berezin E, Towner KJ. 1996. *Acinetobacter* spp. as nosocomial pathogens: microbiological, clinical, and epidemiological features. Clin Microbiol Rev 9:148–165.
- Humphreys H, Towner KJ. 1997. Impact of *Acinetobacter* spp. in intensive care units in Great Britain and Ireland. J Hosp Infect 37:281–286. [https://doi.org/10.1016/S0195-6701\(97\)90144-4](https://doi.org/10.1016/S0195-6701(97)90144-4).
- Forster DH, Daschner FD. 1998. *Acinetobacter* species as nosocomial pathogens. Eur J Clin Microbiol Infect Dis 17:73–77. <https://doi.org/10.1007/BF01682159>.
- Dortet L, Legrand P, Soussy C-J, Cattoir V. 2006. Bacterial identification, clinical significance, and antimicrobial susceptibilities of *Acinetobacter ursingii* and *Acinetobacter schindleri*, two frequently misidentified opportunistic pathogens. J Clin Microbiol 44:4471–4478. <https://doi.org/10.1128/JCM.01535-06>.
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. Nat Methods 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated for comprehensive microbial variant detection and genome assembly improvement. PLoS One 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
- Wang Z, Wu M. 2013. A phylum-level bacterial phylogenetic marker database. Mol Biol Evol 30:1258–1262. <https://doi.org/10.1093/molbev/mst059>.
- Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC, Pati A. 2015. Microbial species delineation using whole genome sequences. Nucleic Acids Res 43:6761–6771. <https://doi.org/10.1093/nar/gkv657>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. Nucleic Acids Res 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Arrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. BMC Genomics 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). Nucleic Acids Res 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, Olson R, Overbeek R, Parrello B, Pusch GD, Shukla M, Thomason JA, Stevens R, Vonstein V, Wattam AR, Xia F. 2015. RASTtk: a modular and extensible implementation of the RAST algorithm for building custom annotation pipelines and annotating batches of genomes. Sci Rep 5:8365. <https://doi.org/10.1038/srep08365>.