

RESEARCH ARTICLE

Open Access



# Antimicrobial activities of saponins from *Melanthera elliptica* and their synergistic effects with antibiotics against pathogenic phenotypes

Cyrille Ngoufack Tagousop<sup>1</sup>, Jean-de-Dieu Tamokou<sup>2\*</sup> , Irene Chinda Kengne<sup>2</sup>, David Ngnokam<sup>1\*</sup> and Laurence Voutquenne-Nazabadioko<sup>3</sup>

## Abstract

**Background:** Resistance of bacteria and fungi to antibiotics is one of the biggest problems that faces public health. The present work was designated to evaluate the antimicrobial activities of saponins from *Melanthera elliptica* and their synergistic effects with standard antibiotics against pathogenic phenotypes. The plant extract was prepared by maceration in methanol. The methanol extract was partitioned into ethyl acetate and *n*-butanol extracts. Column chromatography of the *n*-butanol extract followed by purification of different fractions led to the isolation of four saponins. Their structures were elucidated on the basis of spectra analysis, and by comparison with those from the literature. The antimicrobial activities of the extracts/compounds alone and their combinations with tetracycline and fluconazole were evaluated using the broth microdilution method through the determination of minimum inhibitory concentration (MIC) and minimum microbicidal concentration.

**Results:** Four compounds: 3-*O*- $\beta$ -*D*-glucuronopyranosyl-oleanolic acid (**1**), 3-*O*- $\beta$ -*D*-glucuronopyranosyloleanolic acid 28-*O*- $\beta$ -*D*-glucopyranosyl ester (**2**), 3-*O*- $\beta$ -*D*-glucopyranosyl(1  $\rightarrow$  2)- $\beta$ -*D*-glucuronopyranosyl oleanolic acid (**3**) and 3-*O*- $\beta$ -*D*-glucopyranosyl(1  $\rightarrow$  2)- $\beta$ -*D*-glucuronopyranosyl oleanolic acid 28-*O*- $\beta$ -*D*-glucopyranosyl ester (**4**) were isolated. Compounds **1**, **2** and **3** showed the largest antibacterial activities (MIC = 8–128  $\mu$ g/mL) whereas compound **4** displayed the highest antifungal activities (MIC = 8–16  $\mu$ g/mL). The antibacterial activities of compounds **1** and **2** (MIC = 16–32  $\mu$ g/mL) against multi-drug-resistant *Escherichia coli* S2 (1) and *Shigella flexneri* SDINT are equal to those of vancomycin (MIC = 16–32  $\mu$ g/mL) used as reference antibiotic.

**Conclusions:** The present study showed significant antimicrobial activity of compounds **1**, **2**, **3** and **4** against the tested microorganisms. The saponins act in synergy with the tested standard antibiotics. This synergy could lead to new options for the treatment of infectious diseases and emerging drug resistance.

**Keywords:** *Melanthera elliptica*, Asteraceae, Saponins, Antibacterial, Antifungal, Synergy, Additive

\*Correspondence: jtamokou@yahoo.fr; jean.tamokou@univ-dschang.org;  
dngnokam@yahoo.fr; ngnokam@univ-dschang.org

<sup>1</sup> Research Unit of Environmental and Applied Chemistry, Department of Chemistry, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon

<sup>2</sup> Research Unit of Microbiology and Antimicrobial Substances, Department of Biochemistry, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon

Full list of author information is available at the end of the article



## Introduction

The development and spread of resistance to currently available antibiotics is a global concern [1]. Many investigations have reported multidrug-resistant pathogens, including *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexneri*, *Staphylococcus aureus*, *Candida*, *Cryptococcus*, *Aspergillus* and *Fusarium* [2, 3]. With the increase in bacterial and fungal resistance to antibiotics, plant-based antimicrobials have attracted attention in the scientific research. The use of natural antimicrobial compounds is important not only in food preservation, but also in the control of human and plant infectious diseases [4]. The use of natural products with therapeutic properties, whether vegetable, animal, and mineral, for a long time were the main sources of important therapeutic agents as well as important raw materials for the manufacture of traditional and modern medicines [5]. Many bioassay reports have indicated the presence of antimicrobial compounds including saponins from several medicinal plants in Cameroon [6–8]. The combination of saponins with antibiotics therapeutic approach may lead to new ways to treat infectious diseases. Therefore, many researchers have studied experimentally the synergistic effect resulting from the association of antibiotics with various plant extracts [9–12].

Saponins constitute a large group of glycosides having in their breasts a hydrophilic part consisting of monosaccharides and a lipophilic part commonly called genin. They are characterized by their surfactant properties because they dissolve in water forming foaming solutions. They are abundant in many plants and divided into two groups namely steroidal saponins and triterpenoid saponins. Many biological activities of saponins have been reported such as antibacterial, antifungal, antiviral, anti-inflammatory, anti-ulcer, haemolytic and hepatoprotective properties [13–18]. In addition, the saponins demonstrated the potential to act synergistically with antibiotics, contributing to the recycling of old antibiotics that were once considered ineffective due to resistance problems [19–22]. Hence, the synergistic effect of combinations of saponins with standard antibiotics will open a complete new field of possible applications in medicine to overcome resistance in multidrug resistant microorganisms.

*Melanthera elliptica* O. Hoffm., belonging to Asteraceae family, is a perennial bushy herb to 1.3 m high, commonly found in tropical West and Central Africa [23]. The leaves are used in folk medicine to treat cough, malaria and stomach ache [24]. In Cameroon, the herbalists use the leaf decoction to treat gonorrhoea, feverish headache and pulmonary haemorrhage. The aqueous and ethanol whole plant extract of *Melanthera scandens* showed antibacterial activities against a group of

microorganisms that are implicated in either typhoid fever and/or other gastrointestinal disorders such as diarrhea and dysentery [25, 26]. The phytochemical screening of the ethanolic extract of the leaves of *M. scandens* revealed the presence of cardiac glycosides, tannins, saponins, terpenes and flavonoids [27]. To the best of our knowledge no study has been carried on the phytochemical analysis and antimicrobial properties of *M. elliptica*. This work was carried out in order to evaluate the antimicrobial activities of saponins from *M. elliptica* and their synergistic effects with standard antibiotics against pathogenic phenotypes namely *E. coli*, *S. flexneri*, *S. aureus*, *Candida albicans*, *Candida parapsilosis* and *Cryptococcus neoformans*.

## Methods

### General experimental procedures

The isolation of secondary metabolites was done using column chromatography on Merck silica gel 60 (70–230 mesh) and gel permeation on Sephadex LH-20. TLC was carried out on silica gel GF254 pre-coated plates developed with ternary systems consisting of AcOEt–MeOH–H<sub>2</sub>O in the proportion (8-1-1 or 8-2-1). Detection was accomplished by spraying with 50% H<sub>2</sub>SO<sub>4</sub> followed by heating at 100 °C, or by visual inspection under UV lamp at 254 and 365 nm. In order to determine the structures of the isolated compounds, the 1D and 2D NMR spectra were made. In fact, <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker Avance III 600 spectrometer equipped with a cryoprobe (<sup>1</sup>H at 600 MHz and <sup>13</sup>C at 150 MHz). The 2D NMR experiments were recorded using Bruker microgram firmware (Xwin-NMR software version 2.1 TopSpin 3.2). Chemical shifts ( $\delta$ ) were reported in parts per million (ppm) using the residual solvent signals as a secondary reference to the TMS ( $\delta=0$ ), while the coupling constants (*J* values) are given in Hertz (Hz).

### Plant material

The aerial parts of *M. elliptica* were collected at FOTO Village (Menoua Division, Western region of Cameroon) in January 2014. Authentication was done by Mr. Fulbert Tadjouteu, a Botanist of the Cameroon National Herbarium, Yaoundé, where a voucher specimen (N°8002/HNC) was deposited.

### Extraction and isolation

The air-dried plant material (4 kg) was powdered and extracted at room temperature with methanol (3 × 20 L, 72 h). The solvent was evaporated under reduced pressure, leaving an extract (220 g). Part of this extract (210 g) was suspended in water (300 mL) and successively extracted with equal volumes (500 mL) of ethyl acetate

(EtOAc) and *n*-BuOH (saturated with water) yielding respectively 65 g and 30 g extracts after evaporation to dryness.

A part of the *n*-BuOH extract (25 g) was subjected to silica gel column chromatography using EtOAc–MeOH (100:0 → 40:60) gradient elution. Forty-seven fractions of 250 mL each were collected and combined on the basis of their TLC profiles to give five fractions: A (1–7), B (8–12), C (13–15), D (16–47) and E (24–47). Fraction B (3 g) was also purified on silica gel column chromatography eluted with EtOAc to give compound **1** (75 mg). Fraction C (5 g) mainly yielded compounds **2** (30 mg) and **3** (30 mg) after multiple chromatographic steps over silica gel using EtOAc–MeOH–H<sub>2</sub>O (95:5:2) and (90:5:5) as eluents. Fraction D (6 g) eluted by EtOAc–MeOH (80:20) on column chromatography separation over silica gel yielded three sub-fractions (D<sub>1</sub>–D<sub>3</sub>). Compounds **4** (35 mg) was obtained from fraction D<sub>2</sub> after multiple chromatography separation over silica gel using EtOAc–MeOH–H<sub>2</sub>O (80:10:10) and (80:20:10) as eluents.

## Antimicrobial assay

### Microorganisms

The studied microorganisms were three strains of bacteria (*S. aureus* ATCC 25923, *E. coli* S2 (1) and *S. flexneri* SDINT) and three strains of yeasts (*C. parapsilosis* ATCC 22019, *C. albicans* ATCC 9002 and *C. neoformans* IP95026) taken from our laboratory collection. The bacterial and fungal species were grown at 37 °C and maintained on nutrient agar (NA, Conda, Madrid, Spain) and Sabouraud Dextrose Agar (SDA, Conda) slants respectively.

### Determination of minimum inhibitory concentration (MIC) and minimum microbicidal concentration (MMC)

Minimum inhibitory concentration values were determined by a broth micro-dilution method as described earlier [28] with slight modifications. Each test sample was dissolved in dimethylsulfoxide (DMSO) and the solution was then added to Mueller Hinton Broth (MHB) for bacteria or Sabouraud Dextrose Broth (SDB) for yeasts to give a final concentration of 8192 µg/mL. This was serially diluted twofold to obtain a concentration range of 0.125–4096 µg/mL. Then, 100 µL of each concentration were added in each well (96-well microplate) containing 95 µL of MHB or SDB and 5 µL of inoculum for final concentrations varying from 0.0625 to 2048 µg/mL. The inoculum was standardized at  $2.5 \times 10^5$  cells/mL for yeasts and  $10^6$  CFU/mL for bacteria using a JENWAY 6105 UV/Vis spectrophotometer. The final concentration of DMSO in each well was < 1% [preliminary analyses with 1% (v/v) DMSO did not inhibit the growth of the test organisms]. The negative control well consisted

of 195 µL of MHB or SDB and 5 µL of the standard inoculum. The cultured micro plates were covered; then, the contents of each well were mixed thoroughly using a plate shaker (Flow Laboratory, Germany) and incubated at 35 °C for 24 h (bacteria) and 48 h (yeasts) under shaking. The assay was repeated three times. The MIC values of samples were determined by adding 50 µL of a 0.2 mg/mL *p*-iodonitrotetrazolium violet solution followed by incubation at 35 °C for 30 min. Viable microorganisms reduced the yellow dye to a pink color. MIC values were defined as the lowest sample concentrations that prevented this change in color indicating a complete inhibition of microbial growth. For the determination of MMC values, a portion of liquid (5 µL) from each well that showed no growth of microorganism was plated on Mueller Hinton Agar or SDA and incubated at 35 °C for 24 h (for bacteria) or 35 °C for 48 h (for yeasts). The lowest concentrations that yielded no growth after this subculturing were taken as the MMC values [29]. Vancomycin (Sigma-Aldrich, Steinheim, Germany) and fluconazole (Merck, Darmstadt, Germany) were used as positive controls for bacteria and yeasts, respectively.


### Combined effect of antibiotics and isolated compounds

The interaction between combination of reference antibiotics and isolated compounds (**1**, **3** and **4**) was performed by using the broth microdilution method. The antimicrobial activity of the isolated compounds in the presence of reference antibiotics ( $1/4 \times \text{MIC}$ ) and that of reference antibiotics in the presence of the isolated compounds ( $1/4 \times \text{MIC}$ ) were evaluated as described above. The preliminary tests allowed the selection of MIC/4 as the sub-inhibitory concentrations of the samples. The fractional inhibitory concentration (FIC) index for combinations of two antibacterial agents was calculated according to the following equation: FIC index = FIC A + FIC C; where FIC A = MIC of antibiotic in combination/MIC of antibiotic alone; FIC C = MIC of the isolated compound in combination/MIC of the isolated compound alone. The FIC indices were interpreted as follows:  $\leq 0.5$ , synergy;  $> 0.5-1$ , addition;  $> 1$  and  $\leq 4$ , indifference and  $> 4$ , antagonism [30]. All the experiments were performed in triplicate.

## Results

### Chemical analysis

The purification of *n*-BuOH extract of aerial parts of *M. elliptica* led to the isolation of four saponins (Fig. 1). Structures of these compounds were assigned on the basis of spectroscopic data (<sup>1</sup>H and <sup>13</sup>C NMR, <sup>1</sup>H-<sup>1</sup>H COSY, HSQC, HMBC, and ROESY) and by the comparison of their spectroscopic data with those reported in the literature. These compounds



**Fig. 1** Chemical structures of saponins (1–4) isolated from *n*-BuOH extract of areal parts of *M. elliptica*. **1:** 3-O- $\beta$ -D-glucuronopyranosyl-oleanolic acid; **2:** 3-O- $\beta$ -D-glucuronopyranosyl-oleanolic acid 28-O- $\beta$ -D-glucopyranosyl ester; **3:** 3-O- $\beta$ -D-glucopyranosyl(1  $\rightarrow$  2)- $\beta$ -D-glucuronopyranosyl-oleanolic acid; **4:** 3-O- $\beta$ -D-glucopyranosyl(1  $\rightarrow$  2)- $\beta$ -D-glucuronopyranosyl-oleanolic acid 28-O- $\beta$ -D-glucopyranosyl ester

are: 3-*O*- $\beta$ -D-glucuronopyranosyl-oleanolic acid (**1**) [31], 3-*O*- $\beta$ -D-glucuronopyranosyloleanolic acid 28-*O*- $\beta$ -D-glucopyranosyl ester (**2**) [31], 3-*O*- $\beta$ -D-glucopyranosyl(1  $\rightarrow$  2)- $\beta$ -D-glucuronopyranosyl oleanolic acid (**3**) [32] and 3-*O*- $\beta$ -D-glucopyranosyl(1  $\rightarrow$  2)- $\beta$ -D-glucuronopyranosyl oleanolic acid 28-*O*- $\beta$ -D-glucopyranosyl ester (**4**) [33].

#### Antimicrobial activity of the isolated saponins

The antimicrobial activities of MeOH, *n*-BuOH and EtOAc extracts as well as their isolated compounds were examined by microdilution susceptibility assay against three bacterial and three fungal strains selected on the basis of their relevance as human pathogens. The results showed that the extracts and isolated compounds exhibited variable MICs, depending on the microbial strains (Table 1). The MeOH and *n*-BuOH extracts were active only against bacterial species whereas the EtOAc extract displayed antimicrobial activity against both bacterial and fungal species. The EtOAc extract was the most

active extract followed in decreasing order by *n*-BuOH and MeOH extracts.

The lowest MIC value was recorded on *S. aureus* with compound **1** and on *C. neoformans* with compound **4**, whereas the lowest MBC value was obtained on *S. aureus* with compounds **1** and **2** and on *C. neoformans* with compounds **1** and **4**. However, the highest MIC value was recorded on *C. parapsilosis* and *C. albicans* with compound **1** and the highest MBC value was obtained on *C. parapsilosis*, *C. albicans* and *C. neoformans* with compound **1**. The antibacterial activities of compounds **1** and **2** against *E. coli* S2 (1) and *S. flexneri* SDINT are equal to those of vancomycin used as reference antibiotic.

#### Antimicrobial activity of the combination between antibiotics and saponins

The effect of the interaction between the isolated saponins (**1**, **3** and **4**) and antibiotics (vancomycin and fluconazole) was studied and the results are depicted in Tables 2, 3 and 4. The MIC values of the isolated saponins in combination with antibiotics at the

**Table 1** Antimicrobial activity (MIC and MMC in  $\mu\text{g/mL}$ ) of extracts, isolated compounds and reference antimicrobial drugs

Extracts/compounds	Inhibition parameters	<i>E. coli</i>	<i>S. flexneri</i>	<i>S. aureus</i>	<i>C. parapsilosis</i>	<i>C. albicans</i>	<i>C. neoformans</i>
MeOH extract	MIC	512	256	128	> 2048	> 2048	> 2048
	MMC	1024	512	128	/	/	/
	MMC/MIC	2	2	1	/	/	/
<i>n</i> -BuOH extract	MIC	256	128	128	> 2048	> 2048	> 2048
	MMC	512	256	128	/	/	/
	MMC/MIC	2	2	1	/	/	/
EtOAc extract	MIC	128	128	64	1024	1024	512
	MMC	256	128	128	1024	1024	1024
	MMC/MIC	2	1	2	1	1	2
<b>1</b>	MIC	32	16	8	32	64	16
	MMC	32	32	16	64	64	16
	MMC/MIC	1	2	2	2	1	1
<b>2</b>	MIC	32	16	16	> 256	> 256	32
	MMC	32	32	16	/	/	64
	MMC/MIC	1	2	1	/	/	2
<b>3</b>	MIC	128	64	32	128	128	32
	MMC	256	128	64	256	256	64
	MMC/MIC	2	2	2	2	2	2
<b>4</b>	MIC	256	128	64	16	16	8
	MMC	> 256	256	128	32	64	16
	MMC/MIC	/	2	2	2	4	1
Ref <sup>a</sup>	MIC	32	16	0.5	0.5	1	2
	MMC	32	16	0.5	0.5	1	2
	MMC/MIC	1	1	1	1	1	1

/: not determined; MIC minimum inhibitory concentration, MMC minimum microbicidal concentration

<sup>a</sup> Fluconazole for yeasts and vancomycin for bacteria

**Table 2 Antimicrobial activities of the isolated compounds in the presence of reference antibiotics<sup>a</sup> at 1/4 of MIC as a function of microorganisms**

Microorganisms	Compound 1 alone		Compound 1 with ATB <sup>a</sup> at 1/4 of MIC		Compound 3 alone		Compound 3 with ATB <sup>a</sup> at 1/4 of MIC		Compound 4 alone		Compound 4 with ATB <sup>a</sup> at 1/4 of MIC	
	MIC	MIC	FIC	MIC	MIC	FIC	MIC	FIC	MIC	MIC	FIC	
<i>E. coli</i>	32	8	0.25	128	16	0.125	256	16	0.0625	16	0.0625	
<i>S. flexneri</i>	16	8	0.5	64	16	0.25	128	4	0.0312	4	0.0312	
<i>S. aureus</i>	8	1	0.125	32	16	0.50	64	2	0.0312	2	0.0312	
<i>C. parapsilosis</i>	32	4	0.125	128	4	0.0312	16	2	0.125	2	0.125	
<i>C. albicans</i>	64	4	0.0625	128	4	0.0312	16	1	0.0625	1	0.0625	
<i>C. neoformans</i>	16	2	0.125	32	2	0.0625	8	0.5	0.0625	0.5	0.0625	

MIC minimal inhibitory concentration, FIC fractional inhibitory concentration, ATB antibiotics

<sup>a</sup> Fluconazole for yeasts and tetracycline for bacteria were tested at the concentration of MIC/4 on the corresponding microorganism

**Table 3 Antimicrobial activities of the reference antibiotics in the presence of compounds 1, 3 and 4 at 1/4 of MIC as a function of microorganisms**

Microorganisms	ATB <sup>a</sup> alone		ATB <sup>a</sup> with compound 1 at 1/4 of MIC		ATB <sup>a</sup> with compound 3 at 1/4 of MIC		ATB <sup>a</sup> with compound 4 at 1/4 of MIC	
	MIC	MIC	FIC	MIC	FIC	MIC	FIC	
<i>E. coli</i> S2 (1)	32	1	0.0312	0.25	0.00781	0.125	0.00390	
<i>S. flexneri</i>	16	0.25	0.0156	0.125	0.00781	0.0312	0.00195	
<i>S. aureus</i>	2	0.125	0.0625	0.0625	0.0312	0.0312	0.0156	
<i>C. parapsilosis</i>	0.5	0.25	0.50	0.0625	0.125	0.0312	0.0625	
<i>C. albicans</i>	1	0.125	0.125	0.0625	0.0625	0.0625	0.0625	
<i>C. neoformans</i>	2	0.125	0.0625	0.0625	0.0312	0.0312	0.0156	

MIC minimal inhibitory concentration, FIC fractional inhibitory concentration, ATB antibiotics

<sup>a</sup> Fluconazole for yeasts and tetracycline for bacteria; compounds 1, 3 and 4 were tested at the concentration of MIC/4 on the corresponding microorganism

**Table 4 Fractional inhibitory concentrations (FIC) calculated for the combination of reference antibiotics and compounds 1, 3 and 4 as a function of microorganisms**

Microorganisms	Compound 1 + ATB <sup>a</sup>		Compound 3 + ATB <sup>a</sup>		Compound 4 + ATB <sup>a</sup>	
	∑ FIC	Interpretation	∑ FIC	Interpretation	∑ FIC	Interpretation
<i>E. coli</i>	0.2812	Synergy	0.132	Synergy	0.0664	Synergy
<i>S. flexneri</i>	0.515	Additive	0.257	Synergy	0.0331	Synergy
<i>S. aureus</i>	0.25	Synergy	0.531	Additive	0.0468	Synergy
<i>C. parapsilosis</i>	0.625	Additive	0.156	Synergy	0.187	Synergy
<i>C. albicans</i>	0.187	Synergy	0.0937	Synergy	0.125	Synergy
<i>C. neoformans</i>	0.187	Synergy	0.0937	Synergy	0.0781	Synergy

∑ FIC sum of fractional inhibitory concentrations, ATB antibiotics

<sup>a</sup> Fluconazole for yeasts and tetracycline for bacteria

concentration of MIC/4 are smaller than that of the isolated saponins used alone, suggesting an increase in the activity of saponins in combination with antibiotics (Table 2). The MIC values of antibiotics (vancomycin and fluconazole) in combination with the isolated

saponins at the concentration of MIC/4 are also smaller than those of vancomycin or fluconazole alone, indicating an increase in the activity of vancomycin or fluconazole in combination with the isolated saponins at the concentration of MIC/4 (Table 3).

According to the obtained results, the combination of compound **1** with tetracycline showed a synergistic effect against *E. coli* and *S. aureus* while an additive effect was observed in *S. flexneri* (Table 4). The application of fluconazole with compound **1** led to a synergistic effect on *C. albicans* and *C. neoformans* and to an additive effect on *C. parapsilosis*. Combination of compound **3** with tetracycline had a synergistic effect against *E. coli* and *S. shigella*, and an additive effect against *S. aureus*. A synergistic effect was recorded on *C. parapsilosis*, *C. albicans* and *C. neoformans* with the combination of compound **3** and fluconazole. The interaction between compound **4** and tetracycline/fluconazole led to a synergistic effect against the tested bacterial and fungal strains. The combinatory effect of saponins and antibiotics was tested against six pathogenic microorganisms including two multi-drug-resistant (MDR) pathogenic bacteria [*E. coli* S2 (1) and *S. flexneri* SDINT]. The MIC of tetracycline changed to the susceptible range (MIC:  $\leq 1$  mg/mL) when saponins at the concentration of MIC/4 was added. This result provides strong evidence for the therapeutic potential of saponins in combination with defined antibiotic drugs to fight MDR pathogenic bacteria.

## Discussion

The findings of the present study indicated that the MeOH, *n*-BuOH and EtOAc extracts as well as their isolated saponins showed antimicrobial activity against different bacterial and fungal strains, but at different levels. According to the literature, saponins and extracts of *Melanthera scandens* are known for their antibacterial activities [15, 18, 25, 26]. The results of this study also revealed that all the tested microorganisms were sensitive to EtOAc extract, suggesting that the fractionation enhanced the antimicrobial activity of this extract and diluted that of *n*-BuOH extract. According to Tamokou et al. [34], a plant extract is considered to be highly active if the MIC < 100  $\mu$ g/mL; significantly active when  $100 \leq \text{MIC} \leq 512$   $\mu$ g/mL; moderately active when  $512 < \text{MIC} \leq 2048$   $\mu$ g/mL; weakly active if MIC > 2048  $\mu$ g/mL and not active when MIC > 10 mg/mL. Hence, the *n*-BuOH and MeOH extracts were significantly active ( $100 \leq \text{MIC} \leq 512$   $\mu$ g/mL) against the bacterial strains and weakly active (MIC > 2048  $\mu$ g/mL) against the tested yeasts whereas the EtOAc extract was highly active (MIC < 100  $\mu$ g/mL) against *S. aureus*, significantly active on *E. coli*, *S. flexneri* and *C. neoformans* and moderately active ( $512 < \text{MIC} \leq 2048$   $\mu$ g/mL) vis-à-vis *C. parapsilosis* and *C. albicans*. These findings highlight the traditional use of the *M. elliptica* in the treatment of infectious diseases, especially those caused by the tested microorganisms.

Antimicrobial cut-off points of pure compounds have been defined as follows: highly active: MIC below 1  $\mu$ g/mL (or 2.5  $\mu$ M), significantly active:  $1 \leq \text{MIC} \leq 10$   $\mu$ g/mL (or  $2.5 \leq \text{MIC} < 25$   $\mu$ M), moderately active:  $10 < \text{MIC} \leq 100$   $\mu$ g/mL (or  $25 < \text{MIC} \leq 250$   $\mu$ M), weakly active:  $100 < \text{MIC} \leq 1000$   $\mu$ g/mL (or  $250 < \text{MIC} \leq 2500$   $\mu$ M) and not active: MIC > 1000  $\mu$ g/mL (or > 2500  $\mu$ M) [34]. Based on this, the antimicrobial activity of compound **1** on *S. aureus* as well as that of compound **4** on *C. neoformans* can be considered significant. The strain of *E. coli* S2 (1) and *S. flexneri* [12, 35, 36] included in the present study were MDR clinical isolates and these were resistant to commonly used drugs such as tetracycline, streptomycin, ampicillin, nalidixic acid, *co*-trimoxazole, furazolidone, etc. However, most of the tested saponins displayed moderated antibacterial activities against these bacterial strains, suggesting that their administration may represent an alternative treatment against multidrug resistant *E. coli* and *S. flexneri*.

With regard to the structure–activity relationship analysis, the four saponins with the same basic skeleton, showed different degrees of antimicrobial activity. Compounds **1**, **2** and **3** with less than 3 sugar moieties showed the largest antibacterial activities with the best MIC value recorded with compound **1** on *S. aureus* whereas compound **4** with 3 sugar moieties displayed the highest antifungal activities with the best MIC value recorded with this compound on *C. neoformans*. In general, compound **1** was the most active followed in decreasing order by compound **4**, compound **3** and compound **2**. These observations show that the sugar moieties should be responsible for the difference in the observed activity.

The Gram-positive bacterium, *S. aureus* was found to be more sensitive compared to Gram-negative bacteria, *E. coli* and *S. flexneri*. The findings of the present study showed that the antimicrobial activities varied with the bacterial and fungal strains. These variations may be due to genetic differences between the microorganisms.

With regard to the MIC and MMC values, a lower MMC/MIC ( $\leq 4$ ) value signifies that a minimum amount of plant extracts/pure compounds is used to kill the microbial species, whereas, a higher value signifies the use of comparatively more amount of sample for the control of any microorganism [37].

Combined antibiotic therapy has been shown to delay the emergency of microbial resistance and may also produce favorable synergistic effects in the treatment of infectious diseases. Drug synergism between known antibiotics and bioactive plant extracts is one of the novel ways to overcome the resistance mechanisms of microorganisms. Since lipophilic saponins targeting the biomembrane usually accompany polar polyphenols in phytomedicine preparations, we decided to investigate

their effect as single substances and in combination with standard antibiotics to gain further insight into potential synergistic effects of herbal medicine. In this study, the application of saponins with tetracycline or fluconazole showed synergistic effect on most of the tested microbial strains. This is very promising since the observed synergistic effect could lead to new options for the treatment of infectious diseases and emerging drug resistance. This result corroborates that of Brahim et al. [38] who showed that the combination of saponins extracts and fluconazole exhibited a total synergism against *C. albicans*, *C. parapsilosis*, *Candida krusei* and *Candida glabrata*. Synergistic activities were also verified when combining saponins with tetracycline, erythromycin and ciprofloxacin against *S. aureus* XU212, *S. aureus* RN4220 and *S. aureus* SA1199B [39]. The saponin of *Quillaja saponaria* enhanced the sensitivity of the Gram-negative bacteria *Proteus mirabilis* towards colistin [21]. In a murine wound model induced by *S. aureus* or *E. faecalis* infected BALB/c mice, the coapplication of local saponin (HSM-1) and intraperitoneal treatment with vancomycin or daptomycin resulted in a higher reduction of bacteria in infected BALB/c mice as compared to the respective antibiotic treatment alone [22]. These observations reinforce the advantageous antimicrobial effect of phytochemical-antibiotic combinations.

## Conclusions

The overall results of the present work provide baseline information for the possible use of *M. elliptica* extracts and mostly the isolated saponins in the treatment of infectious diseases caused by the tested microorganisms. In addition, the isolated saponins of this plant could be used in association with tetracycline and fluconazole to combat multidrug resistant pathogens subject to further toxicological and pre-clinical studies.

## Abbreviations

<sup>13</sup>C-NMR: carbon thirteen nuclear magnetic resonance; <sup>1</sup>H NMR: proton nuclear magnetic resonance; 2D NMR: two-dimension nuclear magnetic resonance; ATCC: American Type Culture Collection; CC: column chromatography; COSY: correlation spectroscopy; DMSO: dimethylsulfoxide; EtOAc: ethyl acetate; HMBC: heteronuclear multiple bond connectivities; HNC: *Herbier National du Cameroun*; HR-ESI-MS: high resolution electron impact mass spectrometry; HR-TOFESIMS: high-resolution time of flight electrospray ionization mass spectrometry; HSQC: the heteronuclear single quantum coherence; IR: infra-red; MBC: minimum bactericidal concentration; MDR: multi-drug-resistant; MeOH: methanol; MHA: Mueller–Hinton agar; MHB: Mueller–Hinton broth; MIC: minimum inhibitory concentration; MMC: minimum microbicidal concentration; NA: nutrient agar; *n*-BuOH: *n*-butanol; NMR: nuclear magnetic resonance; Rf: retention factor; TLC: thin layer chromatography; TMS: tetramethylsilane; TOF-ESIMS: time of flight electrospray ionization mass spectrometry; UV: ultra-violet.

## Authors' contributions

CNT and ICK contributed to the data collection and analysis. JDT designated the study, did the biological assays and helped in manuscript writing and

editing. JDT, DN and LVN supervised and revised the manuscript critically for important intellectual content. All authors read and approved the final manuscript.

## Author details

<sup>1</sup> Research Unit of Environmental and Applied Chemistry, Department of Chemistry, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon. <sup>2</sup> Research Unit of Microbiology and Antimicrobial Substances, Department of Biochemistry, Faculty of Science, University of Dschang, P.O. Box 67, Dschang, Cameroon. <sup>3</sup> Groupe Isolement et Structure, Institut de Chimie Moléculaire de Reims (ICMR), CNRS, UMR 7312, Bat. 18, BP.1039, 51687 Reims cedex 2, France.

## Acknowledgements

The study was supported in part by the University of Dschang and the Cameroonian Ministry of Higher Education.

## Competing interests

The authors declare that they have no competing interests.

## Availability of data and materials

The datasets supporting the conclusions of this article are presented in this paper. Plant materials used in this study have been identified at the Cameroon National Herbarium where voucher specimens are deposited.

## Consent to participate

Not applicable.

## Ethics approval and consent to participate

Not applicable.

## Funding

No funding.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 23 March 2018 Accepted: 15 September 2018

Published online: 20 September 2018

## References

- Chanda SK, Rakholiya K (2011) Combination therapy: synergism between natural plant extracts and antibiotics against infectious diseases. In: Mendez-Vilas A (ed) Science against microbial pathogens: communicating current research and technological advances. Formatex Research Center, Badajoz, pp 520–529
- Grundmann H, de Kraker M, Davey P (2011) Clinical impact of antimicrobial resistance: design matters. *Lancet Infect Dis* 11(5):344–345
- Lai CC, Wang CY, Chu CC, Tan CK, Lu CL, Lee YC, Huang YT, Lee PI, Hsueh PR (2011) Correlation between antibiotic consumption and resistance of Gram-negative bacteria causing healthcare-associated infections at a University Hospital in Taiwan from 2000 to 2009. *J Antimicrob Chemother* 66:1374–1382
- Choi HS (2010) Antioxidative activity. In: Sawamura M (ed) *Citrus essential oils: flavor and fragrance*. Wiley, New Jersey, pp 231–243
- Ishrat JB, Laizuman N, Farhana AR, Obaydul H (2011) Antibacterial, cytotoxic and antioxidant activity of chloroform, *n*-hexane and ethyl acetate extract of plant *Amaranthus spinosus*. *Int J PharmTech Res* 3(3):1675–1680
- Lunga PK, Tamokou JD, Fodouop SPC, Kuiate JR, Tchoumboue J, Gatsing D (2014) Antityphoid and radical scavenging properties of the methanol extracts and compounds from the aerial part of *Paullinia pinnata*. *Springerplus* 3:302
- Lunga PK, Qin X-J, Yang XW, Kuiate JR, Du ZZ, Gatsing D (2014) Antimicrobial steroidal saponin and oleanane-type triterpenoid saponins from *Paullinia pinnata*. *BMC Complement Altern Med* 14:369
- Fankam AG, Kuete V, Voukeng IK, Kuiate JR, Pages JM (2011) Antibacterial activities of selected Cameroonian spices and their synergistic effects



- with antibiotics against multidrug-resistant phenotypes. *BMC Complement Altern Med* 11:104
9. Fouedjou TR, Teponno RB, Quassinti L, Bramucci M, Petrelli D, Vitali LA, Fiorini D, Taponjoud AL, Barboni L (2014) Steroidal saponins from the leaves of *Cordyline fruticosa* (L.) A. Chev. and their cytotoxic and antimicrobial activity. *Phytochem Lett* 7:62–68
  10. Olajuyigbe OO, Afolayan AJ (2012) Synergistic interactions of methanolic extract of *Acacia meansii* De Wild. with antibiotics against bacteria of clinical relevance. *Int J Mol Sci* 13:8915–8932
  11. Stefanovic O, Comic L (2012) Synergistic antibacterial interaction between *Melissa officinalis* extracts and antibiotics. *J Appl Pharm Sci* 2:1–5
  12. Tamokou JD, Fosso MJA, Njouendou AJ, Ekom SE, Kengne IC (2017) Antibacterial activity of three Cameroonian medicinal plants traditionally used for the treatment of urinary tract infections and their synergistic effects with amoxicillin and serum. *J Adv Med Pharm Sci* 15(2):1–14
  13. Zhang S, Hu Z (1985) Anti-ulcerogenic effects of Ginseng flower saponins in the rat *Zhongyao Tongbao* 10: 331. *Chem Abstr* 104:512
  14. Okubo K, Kudou S, Uchida T, Yoshiki Y, Yoshikoshi M, Tomomura M (1994) Soyabean saponin and isoflavonoids: structure and antiviral activity against Human Immunodeficiency virus in vitro. In: *ACS Symp Ser*, pp 546
  15. Sen S, Makkar HPS, Muetzel S, Becker K (1998) Effect of *Quillaja saponaria* saponins and *Yucca schidigera* plant extract on growth of *Escherichia coli*. *Letts Appl Microbiol* 27:35–38
  16. Chao AC, Nguyen JV, Broughall M, Recchia J, Kensil CR, Daddona PE, Fix JA (1998) Enhancement of intestinal model compound transport by DS-1, a modified Quillaia saponin. *J Pharm Sci* 87(11):1395–1399
  17. Just MJ, Recsio MG, Gner RM, Cuellar MJ, Marez S, Bilia AR, Rios J (1998) Anti-inflammatory activity of unusual lupane saponins from *Buleurum frutescens*. *Planta Med* 64(5):404–407
  18. Woldemichael GM, Wink M (2001) Identification and biological activities of triterpenoid saponins from *Chenopodium quinoa*. *J Agric Food Chem* 49(5):2327–2332
  19. Kim HS, Han SS, Oh KW, Jeong TS, Nam KY (1987) Effect of ginseng saponin on the antimicrobial activities of some antibiotics. *Korean J Mycol* 15(2):87–91
  20. Sung WS, Lee DG (2008) The combination effect of Korean red ginseng saponins with kanamycin and cefotaxime against methicillin-resistant *Staphylococcus aureus*. *Biol Pharm Bull* 31(8):1614–1617
  21. Arabski M, Wasik S, Dworecki K, Kaca W (2009) Laser interferometric and cultivation methods for measurement of colistin/ampicillin and saponin interactions with smooth and rough of *Proteus mirabilis* lipopolysaccharides and cells. *J Microbiol Methods* 77(2):178–183
  22. Cirioni O, Myszka H, Dawgul M, Ghiselli R, Orlando F, Silvestri C, Brescini L, Kamysz W, Guerrieri M, Giacometti A (2011) In vitro activity and in vivo efficacy of the saponin diosgenyl 2-amino-2-deoxy- $\beta$ -D-glucopyranoside hydrochloride (HSM1) alone and in combination with daptomycin and vancomycin against Gram-positive cocci. *J Med Microbiol* 60(9):1337–1343
  23. Burkill HM (1985) The useful plants of west tropical Africa. vol 1, 2nd ed. Royal Botanic Gardens, Kew, UK, p 960
  24. Yapi AB, Kassi NJ, Fofie NBY, Zirihi GN (2015) Etude ethnobotanique des Asteraceae médicinales vendues sur les marchés du district autonome d'Abidjan (Côte d'Ivoire). *Int J Biol Chem Sci* 9(6):2633–2647
  25. Bolou GEK, Bagré I, Ouattara K, Djaman AJ (2011) Evaluation of the antibacterial activity of 14 medicinal plants in Côte d'Ivoire. *Trop J Pharm Res* 10(3):335–340
  26. Kokora PA, Ackah JAAB, Nanga ZY, Kra MAK, Loukou GY, Coulibaly A, Djaman J (2013) Antibacterial activity of ethanolic and aqueous extracts of four medicinal plants on the in vitro growth of *Escherichia coli* and *Staphylococcus aureus*. *J Drug Deliv Ther* 3(3):113–116
  27. Okokon JE, Udoh AE, Frank SG, Amazu LU (2012) Anti-inflammatory and analgesic activities of *Melanthera scandens*. *Asian Pac J Trop Biomed* 2(2):144–148
  28. Nyaa TBL, Taponjoud AL, Barboni L, Tamokou JD, Kuate JR, Tane P, Park H (2009) NMR assignment and antimicrobial/antioxidant activities of 1-hydroxyeuscaphic acid from the seeds of *Butyrospermum parkii*. *Nat Prod Sci* 15:76–82
  29. Tamokou JD, Kuate JR, Tane M, Kenla Nwemeguela TJ, Tane P (2011) The antimicrobial activities of extract and compounds isolated from *Brillantaisia lamium*. *Iran J Med Sci* 36:24–31
  30. Bone RC (1994) Gram-positive organisms and sepsis. *Arch Intern Med* 154(1):26–34
  31. Vidal-Ollivier E, Balansard G, Faure R, Babadjamian A (1989) Revised structures of triterpenoid saponins from the flowers of *Calendula officinalis*. *J Nat Prod* 52:1156–1159
  32. Paphassarang S, Raynaud J, Lussignol M (1989) Triterpenoid saponins from *Polyscias scutellarza*. *J Nat Prod* 52:239–242
  33. Tan N, Zhou J, Zhao S (1999) Advances in structural elucidation of glucuronide oleanane-type triterpene carboxylic acid 3, 28-O-bisdesmosides (1962–1997). *Phytochemistry* 52(2):153–192
  34. Tamokou JD, Mbaveng TA, Kuete V (2017) Antimicrobial activities of African medicinal spices and vegetables. In: Kuete V (ed) *Medicinal spices and vegetables from Africa: therapeutic potential against metabolic, inflammatory, infectious and systemic diseases*. 1st edn, Chap 8. Academic Press, Cambridge, Elsevier, pp 207–237
  35. Acharyya S, Sarkar P, Saha DR, Patra A, Ramamurthy T, Bag PK (2015) Intracellular and membrane damaging activities of methyl gallate isolated from *Terminalia chebula* against multi-drug resistant *Shigella species*. *J Med Microbiol* 64(8):901–909
  36. Jouda JB, Tamokou JD, Djama Mbazono C, Douala-Meli C, Sarkar P, Bag PK, Wandji J (2016) Antibacterial and cytotoxic cytochalasins from the endophytic fungus *Phomopsis* sp. harbored in *Garcinia kola* (Heckel) nut. *BMC Complement Altern Med* 16:462
  37. Djouossi MG, Tamokou JD, Ngnokam D, Kuate JR, Taponjoud AL, Harakat D, Nazabadioko LV (2015) Antimicrobial and antioxidant flavonoids from the leaves of *Oncoba spinosa* Forssk. (Salicaceae). *BMC Complement Altern Med* 15:134
  38. Brahim MAS, Fadli M, Markouk M, Hassani L, Larhsini M (2015) Synergistic antimicrobial and antioxidant activity of saponins-rich extracts from *Paronychia argentea*. *Eur J Med Plants* 7(4):193–204
  39. Monte J, Abreu AC, Borges A, Simões LC, Simões M (2014) Antimicrobial activity of selected phytochemicals against *Escherichia coli* and *Staphylococcus aureus* and their biofilms. *Pathogens* 3:473–498

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

