

## REVIEW ARTICLE

# Global colistin use: a review of the emergence of resistant *Enterobacterales* and the impact on their genetic basis

Ulrike Binsker<sup>1,\*†</sup>, Annemarie Käsbohrer<sup>1,2</sup> and Jens A. Hammerl<sup>1,‡</sup>

<sup>1</sup>Unit Epidemiology, Zoonoses and Antimicrobial Resistance, Department Biological Safety, German Federal Institute for Risk Assessment, Diedersdorfer Weg 1, 12277 Berlin, Germany and <sup>2</sup>Department for Farm Animals and Veterinary Public Health, Institute of Veterinary Public Health, University of Veterinary Medicine Vienna, Veterinärplatz 1, 1210 Vienna, Austria

\*Corresponding author: Unit Epidemiology, Zoonoses and Antimicrobial Resistance, Department of Biological Safety, German Federal Institute for Risk Assessment, Diedersdorfer Weg 1, 12277 Berlin, Germany. Tel: +49 30 18412 24340; E-mail: [Ulrike.Binsker@bfr.bund.de](mailto:Ulrike.Binsker@bfr.bund.de)

One sentence summary: Colistin resistance in *Enterobacterales* is highly complex and mutations in chromosomally encoded genes are of great concern due to the use of colistin as last-line drug.

Editor: Ehud Banin

†Ulrike Binsker, <https://orcid.org/0000-0002-1963-1916>

‡Jens A. Hammerl, <https://orcid.org/0000-0002-6930-4358>

## ABSTRACT

The dramatic global rise of MDR and XDR *Enterobacterales* in human medicine forced clinicians to the reintroduction of colistin as last-resort drug. Meanwhile, colistin is used in the veterinary medicine since its discovery, leading to a steadily increasing prevalence of resistant isolates in the livestock and meat-based food sector. Consequently, transmission of resistant isolates from animals to humans, acquisition via food and exposure to colistin in the clinic are reasons for the increased prevalence of colistin-resistant *Enterobacterales* in humans in the last decades. Initially, resistance mechanisms were caused by mutations in chromosomal genes. However, since the discovery in 2015, the focus has shifted exclusively to mobile colistin resistances (*mcr*). This review will advance the understanding of chromosomal-mediated resistance mechanisms in *Enterobacterales*. We provide an overview about genes involved in colistin resistance and the current global situation of colistin-resistant *Enterobacterales*. A comparison of the global colistin use in veterinary and human medicine highlights the effort to reduce colistin sales in veterinary medicine under the One Health approach. In contrast, it uncovers the alarming rise in colistin consumption in human medicine due to the emergence of MDR *Enterobacterales*, which might be an important driver for the increasing emergence of chromosome-mediated colistin resistance.

**Keywords:** polymyxin; antimicrobial use; chromosome; One Health; *mcr*; lipid A

## INTRODUCTION

Antimicrobial resistance (AMR) is recognized as one of the greatest challenges for human health worldwide. Over-prescribing and over-using of antibiotics in human and veterinary medicine

has led to the development of multidrug-resistant (MDR; at least one antimicrobial agent in three or more antibiotic classes), extensively drug-resistant (XDR; species are only susceptible to two antimicrobial drug classes) and pandrug-resistant

Received: 29 March 2021; Accepted: 4 October 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of FEMS. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited

(PDR; resistant to almost all commercially available antimicrobials) Gram-negative bacteria (Magiorakos et al. 2012). Rising AMR causes difficult-to-treat infections, therapeutic complications, longer hospital stays and increased mortality. Especially, extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriales*, such as *Escherichia coli* (*E. coli*) and *Enterobacter* spp., as well as carbapenem-resistant *Enterobacteriales* (CRE), particularly *Klebsiella* spp., have been increasingly associated with high morbidity rates due to limited treatment options. According to the European Centre for Disease Prevention and Control, more than 670 000 bacterial infections can be attributed to MDR bacteria, which causes 33 000 death annually in Europe (European Centre for Disease Prevention and Control 2019). As a result, it is estimated that MDR infections and complications cost the health-care system 1.1 billion annually in Europe. The enormous lack of novel antimicrobials active against these MDR Gram-negative bacteria, particularly those producing carbapenemases, requires the growing use of last-resort antibiotics, such as colistin (Falagas and Kasiakou, 2005; Grundmann et al. 2017). In contrast, colistin has been continuously used in the global livestock production for prophylactic, therapeutic and even for growth promotion purposes, which has already been banned in Europe since 2006 (European Commission 2005). The frequent application of antibiotics in food-producing animals is associated with selection of resistant zoonotic strains that can be transmitted directly from animal to human or indirectly via the food chain and eventually causing difficult-to-treat diseases in humans (Marshall and Levy, 2011). Therefore, concepts have been developed and implemented to reduce and limit the use of antibiotics in animals and humans. As such, the globally active One Health Commission focuses on the protection of human health by protecting animal and environmental health, biodiversity and food safety. The One Health approach emerged internationally to primarily address emerging and re-emerging zoonotic diseases (Min, Allen-Scott and Buntain 2013). However, when the concept grew, further interdependent areas were included. Nowadays, One Health also aims to address the urgent problem of AMR by reducing the use of antimicrobials in food-producing animals, which means human health and animal health are interconnected. Thus, controlling animal and zoonotic diseases is effective in controlling human diseases.

The present review outlines the current global colistin resistance situation in *Enterobacteriales* and summarizes the consumption of colistin in the veterinary sector and human medicine in different countries worldwide. The focus of the review was placed on the so far identified bacterial mechanisms mediating colistin resistance highlighting naturally-occurring mutations in chromosomally encoded genes. Since the discovery in 2015, the scientific interest has shifted almost exclusively to plasmid-mediated and transmissible colistin resistance (*mcr*). In contrast to chromosomal-mediated resistance, plasmid-mediated resistance can easily disseminate among different enterobacterial genera, which occurs predominantly in the livestock sector. It can be assumed that with the continuous effort to reduce colistin consumption in veterinary medicine under the One Health approach, the prevalence of *mcr*-bearing plasmids will decrease to a low but stable level in the future. In absence of colistin as selective pressure, bacteria will remove their redundant *mcr*-plasmids due to the energy-consuming replication mechanism of the plasmids. On the other hand, chromosomally mediated colistin resistance is predominantly described in human clinical *Enterobacteriales* isolates and their prevalence will increase especially in human medicine, where colistin is increasingly used as a last resort

treatment for carbapenemase-producing pathogens. Chromosomal alterations in the core genome are characterized by a high stability and irreversibility. This may become a significant health problem when chromosomal mutations accumulate in key human pathogenic lineages. Furthermore, this review stresses the urgent need in the routine monitoring of colistin resistance in *Enterobacteriales* isolated from human and veterinary niches, the food chain and the environment.

### Enterobacteriales – a major host for colistin resistance

The family of *Enterobacteriales* is a large group of Gram-negative bacteria and members, such as *Escherichia*, *Klebsiella*, *Salmonella*, *Enterobacter*, *Serratia*, *Citrobacter* and *Proteus*, are common inhabitants of the gastrointestinal tract of humans and animals (Guentzel, 1996; Marshall and Levy, 2011). However, due to the ability to acquire and disseminate a wide variety of antibiotic resistances, some members of the *Enterobacteriales* family are among the most significant public health problems worldwide (Smet et al. 2010; Navon-Venezia, Kondratyeva and Carattoli 2017). In early 2017, the World Health Organization (WHO) published a pathogen priority list, which included CRE as “critical” antibiotic-resistant bacteria that represent an enormous threat to public health (World Health Organization 2017). In addition, in humans, they account for about 80% of Gram-negative isolates with a variety of diseases including urinary tract infections, pneumonia, diarrhea, meningitis, sepsis, endotoxic shock and others (Oliveira and Reygaert, 2021). Comparable to humans, certain pathogenic lineages of, for example *E. coli* and *Salmonella*, can cause infections in animals such as diarrhea and colibacillosis (Catry et al. 2015). Colistin resistances have been increasingly described in *Enterobacteriales* of human and animal origin (Monaco et al. 2014; European Centre for Disease Prevention and Control 2020; Homeier-Bachmann et al. 2021). Those resistances were initially thought to be located on the chromosome, until 2015, when the first mobile colistin resistance (*mcr*) gene carried by a plasmid has been found (Liu et al. 2016). The occurrence of *mcr*-genes in *Enterobacteriales* has been described in detail elsewhere and is outside of the scope of the present review. A short overview is given in the supplementary material and Table S1(Supporting Information).

### Antibacterial effect of colistin

The polymyxins have bactericidal activity against most members of the *Enterobacteriales* family including *E. coli*, *Klebsiella*, *Salmonella*, *Shigella* and *Enterobacter*, as well as other clinically relevant Gram-negative pathogens such as *Acinetobacter baumannii* and *Pseudomonas aeruginosa*. On the other hand, the polymyxins demonstrated no activity towards Gram-negative and Gram-positive cocci and Gram-positive bacilli. In addition, polymyxins lack activity against intrinsically resistant species, including *Neisseria*, *Serratia*, *Stenotrophomonas*, *Providencia* and *Proteus* spp., *Burkholderia pseudomallei*, *Morganella morganii* and *Edwardsiella tarda* as well as anaerobic bacteria (Muyembe, Vandepitte and Desmyter 1973; Shimizu, Iyobe and Mitsuhashi 1977; Storm, Rosenthal and Swanson 1977; Pogue et al. 2011).

The polyanionic lipopolysaccharide (LPS) structure of Gram-negative bacteria, consisting of a lipid A moiety, a conserved oligosaccharide core (2-keto-3-deoxyoctonic acid, Kdo) and an O-antigen group, is the main target of colistin. The bactericidal effect of colistin is based on its amphipathicity and a multi-step mechanism. Lipid A produced by most species carries a negative charge due to the presence of free phosphate groups. Divalent

cations such as calcium ( $\text{Ca}^{2+}$ ) and magnesium ( $\text{Mg}^{2+}$ ) stabilize LPS by binding to the phosphate groups (Schnaitman, 1971). Initially, colistin establishes an electrostatic interaction with lipid A. The positively-charged diaminobutyric acid (Dab) residues of colistin bind the negatively charged phosphate groups of lipid A and replace  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  ions in a competitive manner thereby destabilizing the LPS and reducing the outer membrane integrity (Velkov et al. 2010). Thereafter, the N-terminal fatty acid side chain and the hydrophobic domain of colistin ( $\text{Leu}^6\text{-L-Leu}^7$ ) insert into the outer membrane, leading to expansion of the lipid monolayer. The increased membrane permeability causes release of periplasmic substances, uptake of colistin into the periplasm, probable integration of the molecule into the inner membrane and eventually cell death (Dixon and Chopra, 1986). Although colistin initially interacts with lipid A, the detailed mechanism of its activity has not yet been fully deciphered and is subject to current investigations (Moffatt, Harper and Boyce 2019).

## RESISTANCE MECHANISMS

### LPS modifying enzymes and their regulators

The composition of LPS is explained very comprehensively by Raetz and colleagues, which is highly heterogeneous and is adapted to environmental stimuli, e.g. pH changes or the presence of cationic antimicrobial peptides. LPS often carries non-stoichiometric substitutions in lipid A and in the inner core. Its heterogeneity is achieved at different position of the LPS structure by (i) modification of lipid A, (ii) changes in the inner core as well as (iii) truncation of the outer core. LPS changes underlie a tight regulation and mediate resistance to cationic antimicrobial peptides, such as colistin (Raetz et al. 2007). The interaction of positively charged Dab residues of colistin with the negatively charged phosphoryl-groups of lipid A is pivotal for its bactericidal activity. Consequently, polymyxin resistance is achieved by remodeling of LPS by incorporation of pEtN and L-Ara4N leading to a reduction of the LPS net negative charge by shielding phosphate and carboxyl groups, which in turn impedes binding of polymyxins.

The biosynthesis of L-Ara4N and addition to LPS requires the enzymes Ugd, ArnB, ArnC, ArnA, ArnD, ArnT, ArnE and ArnF. In *E. coli* and *Salmonella* spp., L-Ara4N is preferentially linked to the 4'phosphate group of lipid A by ArnT, but it can also be found at the 1'position (Raetz et al. 2007). Opposite to L-Ara4N, pEtN is primarily added to the 1'phosphate group, however, can also be linked to the 4'position, when L-Ara4N is absent (Raetz et al. 2007).

The proteins responsible for the regulation, biosynthesis and addition of L-Ara4N and pEtN are chromosomally encoded and respond to the presence of environmental signals or mutational changes. In the following, only those proteins are mentioned for which a genetic alteration could be detected in naturally occurring colistin-resistant animal or human isolates.

### *pmrCAB* operon

The operon of *Enterobacteriales* encodes the two-component signal transduction system (TCS) PmrAB (BasRS in *E. coli*) including the phosphoethanolamine transferase PmrC (EptA in *E. coli*), which modifies the 1'-phosphate group of lipid A. Interruption of the regulatory mechanisms due to critical genetic changes provokes constitutive activation of PmrA or PmrB, subsequent over-expression of LPS modifying enzymes and thus colistin resistance.

In total, two hotspots for missense mutations in PmrA (BasR) affecting the amino acids (aa) G53 and R81 within the phosphate receiver (REC) domain have been identified in *Enterobacteriales*, of which mutations at position G53 have been experimentally confirmed to mediate colistin resistance (Table 1; Sun et al. 2009; Diene et al. 2013; Olaitan et al. 2014; Quesada et al. 2015; Nordmann, Jayol and Poirel 2016; Bourrel et al. 2019; Janssen et al. 2020). Among different species, the histidine kinase gene *pmrB* seems to be the more common site for gain-of-function mutations compared to the response regulator gene *pmrA*. Hot spots for mutations were located in L14, P94, E121, T156, V161 and G206 affecting the N-terminus, the HAMP and the HisKA domain of the protein (Table 2; Sun et al. 2009; Quesada et al. 2015; Delannoy et al. 2017; Sato et al. 2018; Bourrel et al. 2019; Kathayat et al. 2020). However, only the missense mutation at the position G206 was experimentally confirmed (Sato et al. 2018). Notably, mutational studies of EnvZ in *E. coli*, a homolog of PmrB, showed that mutations in the TM1 domain, the HAMP domain and the HisKA domain increase the ratio of kinase activity to phosphatase activity (Park and Inouye, 1997; Hsing et al. 1998; Zhu and Inouye, 2002). Thus, it is possible that mutations in the same domains of PmrB would lead to an increased kinase/phosphatase activity and consequently an increased transfer of the phosphate to PmrA.

*Escherichia coli* and *Salmonella* spp. exhibit most of the mutations that lead to colistin resistance in the genes of the PmrAB TCS, predominantly in the histidine-kinase PmrB. In contrast, fewer genetic alterations within PmrAB have been reported for klebsiellae.

Mutations in the transferase PmrC have been found in colistin-resistant *E. coli* and *Klebsiella* strains by *in silico* analysis (Table 3; Mathur et al. 2018, Choi et al. 2020). Notably, *pmrC* contains multiple missense mutations, but additional mutations in other colistin resistance-related genes have been found in the same isolates. Therefore, the contribution of gene alterations in *pmrC* to colistin resistance has not yet been deciphered.

### PhoPQ two-component system

The PhoQ sensor kinase has been shown to respond to low environmental  $\text{Mg}^{2+}$  concentrations, changes in pH and the presence of antimicrobial peptides resulting in activation and phosphorylation of the PhoP response regulator (Fig. 3). PhoP controls the expression of genes involved in magnesium transport and modification of LPS. Interestingly, PhoPQ was identified to regulate the expression of the small RNA *mgrR*, which is a negative regulator of the phosphoethanolamine transferase EptB (Moon and Gottesman, 2009). Additionally, PhoPQ contribute to colistin resistance by indirectly activating the PmrAB TCS via PmrD (Kox, Wosten and Groisman 2000; Kato, Latifi and Groisman 2003; Rubin et al. 2015). Missense mutations and deletions in PhoP have been identified in the REC and Trans.reg.C domains as well as inter-domain regions (Table 4; Cheng et al. 2015; Jayol et al. 2015; Delannoy et al. 2017; Dagher et al. 2020). Furthermore, mutations in PhoQ occurred in several functional domains, but also in inter-domain regions (Table 5; Choi and Ko 2014; Cheng et al. 2015; Olaitan et al. 2015; Halaby et al. 2016; Nordmann, Jayol and Poirel 2016; Luo et al. 2017; Gentile et al. 2020).

### PmrD adaptor protein

PmrD is a small protein connecting the two TCS's PmrAB and PhoPQ (Fig. 3). PmrD binds to the phosphorylated form of PmrA

Table 1. Mutations in the response regulator PmrA of Enterobacterales.

Bacterial genera	ST	PmrA protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference	
<i>E. coli</i>	ns	222 (BasR)	REC (1–112)	G15R	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)	
	ns	222 (BasR)	REC (1–112)	S29G	ns	Not confirmed	Chicken feces	Younba et al. (2019)	
	ns	222 (BasR)	REC (1–112)	S39I	4	Not confirmed	Swine feces, also mutation in PmrA R81S	Quesada et al. (2015)	
	131	222 (BasR)	REC (1–112)	G53A	8	Confirmed	Human clinical blood isolate	Janssen et al. (2020)	
	ns	222 (BasR)	REC (1–112)	G53A or G53C or G53E or G53R or G53S or G53V or G53W	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)	
<i>Klebsiella</i> spp.	ns	222 (BasR)	REC (1–112)	A80V	ns	Not confirmed	Avian pathogenic <i>E. coli</i>	Kathayat et al. (2020)	
	ns	222 (BasR)	REC (1–112)	R81L or R81S	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)	
	131	222 (BasR)	REC (1–112)	L105P	16	Confirmed	Human clinical specimen	Sato et al. (2018)	
	ns	222 (BasR)	–	G144S	ns	Not confirmed	Diseased pig (also observed in sensitive strains)	Delannoy et al. (2017)	
	ns	223	REC (1–112)	S42N	ns	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)	
<i>Enterobacter</i> spp.	ns	223	REC (1–112)	G53C or G53S	32 / 128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	223	REC (1–112)	G53C or G53S	ns	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)	
	ns	223	REC (1–112)	E57G	ns	Not confirmed	Human clinical isolate, also mutation in PmrB T246A	Samuelsen et al. (2017)	
	ns	223	REC (1–112)	D86E	>8	Not confirmed	Human clinical isolate, also mutation in PmrA G35C	Samuelsen et al. (2017)	
	ns	222	REC (1–112)	G53S	> 16	Confirmed	Human clinical specimen	Diene et al. (2013)	
	ns	222	REC (1–112)	S64C	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)	
	ns	222	REC (1–112)	L216W	ns	Not confirmed	Rectal swab specimen, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)	
	ns	222	REC (1–112)	E217I	ns	Not confirmed	Rectal swab specimen, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)	

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis.

Table 2. Mutations in the histidine kinase PmrB of Enterobacteriales.

Bacterial genera	ST	PmrB protein, domain length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
E. coli	648	363 (BasS)	-	Δ6-11 RPISLR	16	Confirmed	Human rectal swab isolate	Janssen et al. (2020)
	ns	363 (BasS)	-	Δ7-12	ns	Not confirmed	Human clinical specimen	Poirel, Jayol and Nordmann (2017)
E. coli	59	363 (BasS)	-	L10P	4	Confirmed	Human urinary tract isolate	Cannatelli et al. (2017)
	131	363 (BasS)	-	L10R	16	Confirmed	Human clinical blood isolate	Janssen et al. (2020)
	ns	363 (BasS)	-	L14Q	ns	Not confirmed	Avian pathogenic E. coli	Kathayat et al. (2020)
	38	363 (BasS)	TM1 (15-34)	G19E	4	Confirmed	Human rectal swab isolate	Janssen et al. (2020)
	131	363 (BasS)	TM1 (15-34)	Δ27-45 LISVFWL-WHESFEQQLFE	16	Confirmed	Human clinical specimen	Sato et al. (2018)
	ns	363 (BasS)	TM2 (66-88)	C84R or C84Y	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)
	ns	363 (BasS)	HAMP	T92P	ns	Not confirmed	Avian pathogenic E. coli	Kathayat et al. (2020)
	ns	363 (BasS)	HAMP	R93P	ns	Not confirmed	Diseased pig	Kuang et al. (2020)
	ns	363 (BasS)	HAMP	P94A or P94L or P94Q or P94S	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)
	10	363 (BasS)	HAMP	A118T	>32	Not confirmed	Human clinical specimen	Luo et al. (2017)
E. coli	ns	363 (BasS)	HAMP	E121K or E121Q	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)
	14/131	363 (BasS)	HAMP	E123D	>32	Not confirmed	Human clinical specimen	Luo et al. (2017)
	ns	363 (BasS)	HAMP	E123D	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)
	ns	363 (BasS)	HAMP	T147A	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)
	ns	363 (BasS)	HisKA (142-202)	T156A	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)
	ns	363 (BasS)	HisKA (142-202)	T156K	ns	Not confirmed	Human clinical specimen	Poirel et al. (2017)
	ns	363 (BasS)	HisKA (142-202)	T156M	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)
	ns	363 (BasS)	HisKA (142-202)	A159P	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)
	ns	363 (BasS)	HisKA (142-202)	A159V	ns	Not confirmed	Human clinical specimen	Poirel et al. (2017)
	ns	363 (BasS)	HisKA (142-202)	G160E	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)
E. coli	ns	363 (BasS)	HisKA (142-202)	V161G	4	Not confirmed	Swine feces	Quesada et al. (2015)
	ns	363 (BasS)	HisKA (142-202)	E166K	ns	Not confirmed	Human clinical specimen	Bourrel et al. (2019)
	416/131	363 (BasS)	-	G206D	4/8	Confirmed	Human clinical specimen	Sato et al. (2018)
	641	363 (BasS)	HATPase.c (249-357)	D283G	8	Not confirmed	Calf caecum, also mutation in PmrB Y358N	Rebelo et al. (2018)
	ns	363 (BasS)	HATPase.c (249-357)	D283G	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)

Table 2. Continued

Bacterial genera	ST	PmrB protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>Klebsiella</i> spp.	38	363 (BasS)	HATPase.c (249–357)	Y315F	>32	Not confirmed	Human clinical specimen	Luo et al. (2017)
	ns	363 (BasS)	HATPase.c (249–357)	V351I	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)
	101/410	363 (BasS)	–	Y358N	>32	Not confirmed	Human clinical specimen	Luo et al. (2017)
	131	417 (BasS)	HAMP2	Extra HAMP domain	16	Confirmed	Human clinical blood isolate	Janssen et al. (2020)
	ns	365	TM1 (13–35)	L17Q	32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
	17	365	TM1 (13–35)	G20S	256	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)
	258	365	–	V46E	ns	Not confirmed	Patient rectal swab	Gentile et al. (2020)
	646	365	–	Q56S	4	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)
	512	365	TM2 (67–89)	L82R	4	Confirmed	Human clinical specimen	Cannatelli et al. (2014)
	ns	365	TM2 (67–89)	S85R	ns	Not confirmed	Human feces from healthy individual	Olaitan et al. (2014)
512	365	HAMP (90–142)	P95L	ns	Not confirmed	Human rectal swab isolate	Gentile et al. (2020)	
512	365	HAMP (90–142)	Δ129–134 ALNQLV	>8	Not confirmed	Human rectal swab isolate	Giordano et al. 2019	
ns	365	HAMP (90–142)	T140P	ns	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)	
ns	365	HAMP (90–142)	H156R	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)	
14,258,15,101	365	HisKA (143–203)	T157P	3–6	Confirmed	Human clinical specimen	Jayol et al. 2014	
15	365	HisKA (143–203)	T157P	32	Not confirmed	Human clinical specimen	Cheng et al. (2015)	
86	365	–	T246A	64	Not confirmed	Human clinical specimen	Cheng et al. (2020)	
11	365	HATPase.c (250–358)	R256G	>512	Not confirmed	Human blood isolate, further mutations in other genes mediating colistin resistance	Cheng et al. (2015)	
ns	365	HATPase.c (250–358)	H333Y	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)	
23	365	HATPase.c (250–358)	P344L	4	Not confirmed	Human blood isolate, further mutations in other genes mediating colistin resistance	Cheong et al. (2020)	
45	365	TM1 (13–35)	Δ11–14	16	Confirmed	Human rectal swab	Olaitan et al. (2015)	
32	365	HAMP (89–141)	R92P	4 – >16	Not confirmed	Poultry farm	Jovic et al. (2020)	
32	365	HisKA (142–202)	V164M or V164G	4 – >16	Not confirmed	Poultry farm	Jovic et al. (2020)	

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis; Δ: deletion.

Table 3. Mutations in the phosphoethanolamine transferase PmrC of *Enterobacteriales*.

Bacterial genera	ST	PmrC protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>E. coli</i>	1	547	Transferase domain,-	T148A, K233T	8–16	Not confirmed	Isolate from animal and plant quarantine agency, also mutations in other genes mediating colistin resistance	Choi et al. (2020)
<i>Klebsiella</i> spp.	11, 14 and 231	546	TM1	C27F	4–16	Not confirmed	Human clinical specimen, also mutations in other genes mediating colistin resistance	Mathur et al. (2018)
		546	–	V39L				
		546	–	V42L				
		546	Transferase domain	R152H				
		546	Sulfatase domain	S260L				
		546	Sulfatase domain	S257L				
		546	Sulfatase domain	A279G				
		546	Sulfatase domain	Q319R				
		546	Sulfatase domain	D477N				

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico*.

Table 4. Mutations in the response regulator PhoP of Enterobacteriales.

Bacterial genera	ST	PhoP protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>E. coli</i>	ns	223	REC (1–112)	V108M	ns	not confirmed	Diseased pig	Delannoy et al. (2017)
	ns	223	Trans_reg-C (145–220)	A182P	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)
<i>Klebsiella</i> spp.	29	223	REC (1–112)	V3F	>2048	Not confirmed	Human clinical specimen	Cheng et al. (2015)
	ns	223	REC (1–112)	L12Q	ns	Not confirmed	Human clinical specimen	Macetic et al. (2020)
	ns	223	REC (1–112)	L26Q	ns	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)
	646	223	REC (1–112)	A30S	4	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)
	17	223	REC (1–112)	L87P	256	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)
	11	223	REC (1–112)	S86L	128	Not confirmed	Human clinical specimen	Cheng et al. (2015)
	ns	223	Trans_reg-C (145–220)	D191Y	12	Confirmed	Human clinical specimen	Jayol et al. (2015)
<i>Enterobacter</i> spp.	ns	223	REC (1–112)	D46V	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	223	REC (1–112)	I47F	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	223	REC (1–112)	I49F	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	223	-	ΔE140	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	223	-	ΔF141	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	223	-	I143D	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	223	-	N144A	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	223	Trans_reg-C (145–220)	Δ148–163	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis; Δ: deletion.



Table 5. Mutations in the histidine kinase PhoQ of *Enterobacteriales*.

Bacterial genera	ST	PhoQ protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>E. coli</i>	354	486	-	N346K	>32	Not confirmed	Human clinical specimen	Luo et al. (2017)
	3997	486	HATPase-c (374-480)	E375K	4	Not confirmed	Human rectal swab isolate from healthy individuals	Olaitan et al. (2015)
<i>Klebsiella</i> spp.	ns	488	-	R16C	>128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
	43	488	TM1 (20-42)	A21S	16	Confirmed	Human clinical specimen	Halaby et al. (2016)
	ns	488	TM1 (20-42)	V24G	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	11	488	TM1 (20-42)	L26P	64	Not confirmed	Human clinical specimen	Cheng et al. (2015)
	ns	488	TM1 (20-42)	L30Q	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	ns	488	-	K46N	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	512	488	-	S56R	ns	Not confirmed	Human rectal swab isolate	Gentile et al. (2020)
	17	488	-	A70K, D90H	256	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)
	17	488	-	P72N, D90N	256	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)
	646	488	-	D74E, Q92S	4	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)
646	488	-	I75L	4	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)	
944	488	-	E77D, K94E	4	Not confirmed	Chicken meat, also mutation in other genes involved in colistin resistance	Chaalal et al. (2021)	

Table 5. Continued

Bacterial genera	ST	PhoQ protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
	258	488	-	L87P	ns	Not confirmed	Human rectal swab isolate	Gentile et al. (2020)
	ns	488	-	L96P	ns	Not confirmed	Human clinical specimen	Olaitan et al. (2014)
	ns	488	-	I109N	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	11	488	-	D150G	128	Not confirmed	Human clinical specimen, additional mutation in MgrB and PmrB R156G	Cheng et al. (2015)
	ns	488	-	S174N	4	Not confirmed	Human clinical specimen	Choi and Ko (2014)
	ns	488	TM2 (194-216)	P208H	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	15	488	HAMP (215-266)	V258F	64	Not confirmed	Human clinical specimen, additional mutation in MgrB	Cheng et al. (2015)
	ns	488	HisKA (274-482)	Q310L	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	ns	488	HisKA (274-482)	H339D	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	ns	488	HisKA (274-482)	L348Q	ns	Not confirmed	Human clinical specimen	Olaitan et al. (2014)
	ns	488	HisKA (274-482)	A351P	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	ns	488	HisKA (274-482)	G385S	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	ns	488	HisKA (274-482)	P420S	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	512	489 (insertion)		D266.267insD	ns	Not confirmed	Human rectal swab isolate, ins799/801(GAC)	Gentile et al. (2020)

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis

following activation by PhoP (Kato and Groisman, 2004; Mitrophanov et al. 2008; Rubin et al. 2015). Phosphorylated and therefore activated PmrA is protected from dephosphorylation resulting in binding to its targets such as the promoter of the *pmrHF1-JKLM* operon, which encodes genes responsible for the modification of LPS. Interestingly, gene multiplications of *pmrD* positively correlate with colistin resistance levels in *Salmonella*, but not in *E. coli* (Hjort, Nicoloff and Andersson 2016). Furthermore, *pmrD* is not present in all *Enterobacter* species (Guerin et al. 2016).

Missense mutations in PmrD in colistin-resistant *Enterobacteriales* have been found in human clinical *Klebsiella pneumoniae* (*K. pneumoniae*) isolates and *E. coli* strains of animal origin (Table 6; Kim et al. 2019; Cheong et al. 2020). However, the parallel occurrence of mutations in MgrB, PhoPQ or PmrAB does not clarify the contribution of mutations in PmrD to the colistin resistance phenotype. Notably, a missense mutation at position K82 has been identified in several *E. coli* strains and represents eventually a hot spot (Kim et al. 2019). Furthermore, this stand-alone mutation occurred in strains without other detectable genetic alterations eventually highlighting this aa substitution as critical for colistin resistance. Nevertheless, experimental confirmation using site-directed mutagenesis or complementation experiments is required.

### MgrB regulator

MgrB is a 47 aa regulatory transmembrane peptide, which is produced upon activation of the PhoPQ system (Lippa and Goulian, 2009). Through interactions with the periplasmic domain of PhoQ, MgrB acts as a feedback inhibitor of the PhoPQ system in different bacteria (Fig. 3). Within *mgrB* various groups of genetic changes have been identified including missense mutations, non-sense mutations, deletion of individual nucleotides or deletion of the entire *mgrB* locus as well as insertions of additional aas and IS elements (Table 7; Cannatelli et al. 2014; Olaitan et al. 2014; Cheng et al. 2015; Nordmann, Jayol and Poirel 2016; Haeili et al. 2017; Esposito et al. 2018). Almost any position of the *mgrB* gene can be affected, which predominantly leads to functional inactivation of the peptide. Thus, the PhoPQ system becomes upregulated, which, in turn, activates the Pmr system responsible for modification of the LPS. In colistin-resistant *K. pneumoniae* strains, the disruption of the *mgrB* gene plays a significant role, with a prevalence of up to 59% of human clinical isolates tested (Cannatelli et al. 2014). A study analyzing 973 clinical *K. pneumoniae* isolates showed that the insertions of IS elements (IS5-like, IS1F, ISKpn13, ISKpn14 and IS10R) are the most common genetic alteration of *mgrB*, followed by partial or complete deletion of the gene, missense mutations and finally nonsense mutations (Hamel et al. 2020). It seems that the high frequency of mutational changes and the inactivation of MgrB do not have major detectable consequences for the fitness and virulence of the *K. pneumoniae* strains. In contrast to *K. pneumoniae*, only two missense mutations in MgrB of *E. coli* have been reported, but their contribution to colistin resistance has not yet been confirmed, since no complementation or site-directed mutagenesis has been performed (Delannoy et al. 2017). Noteworthy, the *phoPQ* operon in *E. coli* is regulated not only by MgrB but also by the small RNA MicA, which eventually negates the contribution of mutational changes in MgrB to colistin resistance (Janssen et al. 2020). In *Enterobacter* a few mutations in MgrB have been identified by *in silico* analysis, however, concurrent mutations in PmrAB or PhoP of the same isolate were present, making it difficult to assess their significance for colistin resistance (Dagher et al. 2020).

### CrrAB two-component system

CrrA and CrrB belong to a third TCS, which has been investigated in the context of colistin resistance in *K. pneumoniae*. The physiologic role of the TCS is still unknown. The *crrAB* operon is variably expressed in *K. pneumoniae* and has also been found in *Enterobacter* spp., but is not encoded in the *E. coli* chromosome (Wright et al. 2015). CrrB mutations have been reported to increase CrrC expression, which positively regulates the PmrAB TCS, thereby resulting in elevated transcription of the *pmrC* gene and the *pmrHF1JKLM* operon (Fig. 3; Cheng et al. 2016). The *pmrHF1JKLM* operon can also be directly activated by the CrrAB TCS, which has been shown by using CrrB mutants in *K. pneumoniae* (McConville et al. 2020). A single missense mutation has been identified in the response regulator CrrA of *K. pneumoniae*, whereas several confirmed missense mutations in the histidine kinase CrrB were found to induce colistin resistance (Table 8; Wright et al. 2015; Cheng et al. 2016; Jayol et al. 2017; Pishnian, Haeili and Feizi 2019).

### The altered proteins YciM, LpxM, RamA and OmpW of *K. pneumoniae*

In *K. pneumoniae*, additional genes seem to be associated with the colistin-resistant phenotype. YciM (LapB in *E. coli*) has not been well-characterized in *K. pneumoniae*, however, its homolog in *E. coli* is involved in maintaining the cell wall integrity by regulation of LPS biosynthesis (Mahalakshmi et al. 2014; Nicolaes et al. 2014). A total of two missense mutations in YciM have been detected in human clinical isolates of which the mutation V43G has been experimentally confirmed to cause colistin resistance (Table 9; Halaby et al. 2016; Boszczowski et al. 2019). LpxM (MsbB) is responsible for the acylation of Lipid A in *Enterobacteriales* (Somerville et al. 1996; Khan et al. 1998). Interestingly, the loss of LpxM leads to increased colistin susceptibility (Clements et al. 2007). Four mutations have been found in *K. pneumoniae*, but only the mutation V30G was confirmed to contribute to colistin resistance (Halaby et al. 2016; Boszczowski et al. 2019). Very recently, a single missense mutation has been described in the global regulator RamA of a colistin-resistant human clinical *K. pneumoniae* isolate (Table 9; Macesic et al. 2020). RamA is responsible for the activation of gene expression necessary for the biosynthesis and modification of lipid A (De Majumdar et al. 2015). Furthermore, the same study detected an IS-insertion in the outer membrane protein OmpW, likely leading to its functional inactivation, in a clinical *K. pneumoniae* strain resistant to colistin (Macesic et al. 2020). This is the first indication of the involvement of OmpW in the colistin-resistant phenotype of *Enterobacteriales*. In contrast, OmpW of *A. baumannii*, which is a homolog to OmpW of *E. coli*, showed reduced expression levels in colistin-resistant strains and was involved in binding to colistin (Vila, Marti and Sanchez-Cespedes 2007; Catel-Ferreira et al. 2016).

There are excellent publications on the mechanisms of polymyxin resistance and the importance of chromosomal mutations, but these studies also include *in vitro* induced mutations or mention only a portion of all the so far identified genes. We provide here a summary of fourteen chromosomally encoded genes that have been analyzed in the context of colistin resistance in *Enterobacteriales* of animal and human origin. However, five genes (*prnC*, *pmrD*, *crrA*, *ramA* and *ompW*) were not experimentally confirmed to mediate colistin resistance and additional mutations in other colistin resistance genes were found in the same isolate. The cumulative appearance of genetic alterations in different colistin resistance genes seems to be a

Table 6. Mutations in the connector protein PmrD of *Enterobacteriales*.

Bacterial genera	ST	PmrD protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>E. coli</i>	ns	88	-	N11D	ns	Not confirmed	Animal samples, also mutations in other genes involved in colistin resistance	Kim et al. (2019)
	ns	88	-	M20	ns	Not confirmed	Animal samples, also mutations in other genes involved in colistin resistance	Kim et al. (2019)
	ns	88	-	A27T	ns	Not confirmed	Animal samples, also mutations in other genes involved in colistin resistance	Kim et al. (2019)
	ns	88	-	K35N	ns	Not confirmed	Animal samples, also mutations in other genes involved in colistin resistance	Kim et al. (2019)
	ns	88	-	A52V	ns	Not confirmed	Animal samples, also mutations in other genes involved in colistin resistance	Kim et al. (2019)
<i>Klebsiella</i> spp.	3054, 224, 6488, 2035, 278, 448, 906, 4038, 156 and 548	88	-	K82T	4-32	Not confirmed	Animal sample	Kim et al. (2019)
	ns	81	-	Q9R	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	A12S	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	S13M	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	A14T	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	L16S	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)

Table 6. Continued

Bacterial genera	ST	PmrD protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
	ns	81	-	R18C	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	A25T	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	E27A	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	R38H	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	R40Q	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	D50N	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	T60A	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	R66L	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	N67K	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	T77A	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	N78K	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	A79L	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	G80D	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)
	ns	81	-	K81G	ns	Not confirmed	Human blood isolate, also mutation in other genes involved in colistin resistance	Cheong et al. (2020)

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis

Table 7. Mutations in the regulator MgrB of Enterobacteriales.

Bacterial genera	ST	MgrB protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference	
<i>E. coli</i>	ns	47	-	V8A	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)	
	ns	47	-	Q33R	ns	Not confirmed	Diseased pig	Delannoy et al. (2017)	
	ns	47	-	K3*	64–128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	47	-	L9*	8–12	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)	
	ns	47	-	I13*	16–32	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)	
	491	47	-	A14S	12–4	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)	
	726	47	-	C16*	>128	Not confirmed	Healthy broiler, also mutation in PmrB R256G and CrrB T150R	Pishman et al. (2019)	
	101	47	-	L17R	32	Confirmed	Human blood isolate	Esposito et al. (2018)	
	ns	47	-	W20R	32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	258	47	-	L24H	32	Confirmed	Human clinical specimen	Cannatelli et al. (2014)	
507	47	-	V26*	32	Not confirmed	Human feces from healthy individuals	Olaitan et al. (2014)		
<i>Klebsiella</i> spp.	ns	47	-	M27K	32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	1309	47	-	C28F	4	Not confirmed	Human feces from healthy individual	Olaitan et al. (2014)	
	258	47	-	C28Y	32	Confirmed	Human clinical specimen	Cannatelli et al. (2014)	
	ns	47	-	C28*	32–>128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	47	-	Q30*	32–128	Confirmed	Human clinical specimen	Nordmann et al. (2016); Haeili et al. (2017)	
	ns	47	-	D31N	ns	Not confirmed	Human clinical specimen	Olaitan et al. (2014)	
	101	47	-	V32G	2–56	Confirmed	Human blood isolate	Esposito et al. (2018)	
	ns	47	-	Q33*	32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	47	-	F35I	ns	Not confirmed	Human feces from healthy individual	Olaitan et al. (2014)	
	258/512	47	-	G37S	16–32	Confirmed	Human clinical specimen	Cannatelli et al. (2014)	
	ns	47	-	C39Y	64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	47	-	C39*	128	Confirmed	Human clinical specimen	Haeili et al. (2017)	
	ns	47	-	N42Y, K43I	32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	47	-	I45T	64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	47	-	P46S	64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	47	-	W47R or W47*	4 or 32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	62	-	*48Y	ns	Confirmed	Human clinical specimen	Cheng et al. (2015)	
	ns	Additional insertions	-	V7::ISEcp1/blaCTX-M-15 or V7::ISIR	64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	
	ns	Additional insertions	Additional insertions	Additional insertions	V12::IS102-like	32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
					L14::IS102-like or L14::IS903b or L14::IS2 or L14::IS1R	>128 64 64 128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
W20::IS1R					64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)	

Table 7. Continued

Bacterial genera	ST	MgrB protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
	ns	Additional insertions	-	V23::IS903b-like or V23::IS903-like	64 or 128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
	ns	Additional insertions	-	F24::IS5-like or F24::ISKpn13 or F24::ISKpn26-like	16-128 128 64- > 128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
	ns	Additional insertions	-	N25::SKpn26-like or N25::IS903B or N25::ISKpn14 or N25::IS5-like element	32 32 128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
258/512		Additional insertions	-	N25::IS5-like element	16-32 / 16-32	Confirmed	Human clinical specimen	Cannatelli et al. (2014)
ns		Additional insertions	-	D31::IS903b	64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
512		Additional insertions	-	F35::IS1F-like element	8-32	Confirmed	Human clinical specimen	Cannatelli et al. (2014)
ns		Additional insertions	-	I38::IS1R-like or I38::IS903b-like	12.8 64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
ns		Additional insertions	-	C39::IS1R	8	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
ns		Additional insertions	-	I41::ISKpn26-like or I41::IS1R	12.8 32	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
147		Additional insertions	-	I41::ISKpn14	8-16	Confirmed	Human clinical specimen	Cannatelli et al. (2014)
ns		Additional insertions	-	N42::ISKpn14	64	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
512		Additional insertions	-	N42::IS5-like element	32	Confirmed	Human clinical specimen	Cannatelli et al. (2014)
ns		Additional insertions	-	Q43::IS1R	128	Not confirmed	Human clinical specimen	Nordmann et al. (2016)
ns		Insertions upstream of mgrB	-	nt -62 to -26	32-128	Not confirmed	Human clinical specimen	Nordmann, Jayol and Poirel (2016)
512		Deletions	-	Δnt18/27	32	Confirmed	Human clinical specimen, frameshift and premature termination	Cannatelli et al. (2014)
512		Deletions	-	Δnt19	128-256	Confirmed	Human blood isolate, frameshift and premature termination	Esposito et al. (2018)
258/512		Deletions	-	ΔmgrB locus	64/8	Confirmed	Human clinical specimen	Cannatelli et al. (2014)

Table 7. Continued

Bacterial genera	ST	MgrB protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>Enterobacter</i> spp.	258	Deletions	-	ΔmgrB locus	16	Confirmed	Human clinical specimen, deletion from -400 to +599	Cannatelli et al. (2014)
	512	Deletions	-	Δnt61/70	ns	Not confirmed	Human rectal swab isolate	Gentile et al. (2020)
	258/512	Deletions	-	Δnt47	32 / 8	Confirmed	Human clinical specimen, frameshift and premature termination	Cannatelli et al. (2014)
	512	Deletions	-	Δnt 109/119	32	Confirmed	Frameshift and premature termination	Cannatelli et al. (2014)
	258	47 (non-sense mutation)	-	c88t	64	Confirmed	Human blood isolate, non-sense mutation and premature termination	Esposito et al. (2018)
	ns	47	-	V38S or V38I	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	47	-	C39G	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	47	-	A40K	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	47	-	I41M	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	47	-	N42S	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
	ns	47	-	K43G	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)
ns	47	-	I45Y	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)	
ns	47	-	P46G	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)	
ns	47	-	W47V or W47S	ns	Not confirmed	Human rectal swab isolate, also mutation in other genes involved in colistin resistance	Dagher et al. (2020)	

ST, sequence type; ns, not specified; confirmed (grey background): experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis; \*, stop codon results in termination and truncated protein; Δ, deletion; ::, insertion



Table 8. Mutations in the CcrAB two-component system of Enterobacterales.

Bacterial genera	ST	Protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>Klebsiella</i> spp.	11	CrrA (234)		A83V	>128	Not confirmed	Dead broiler, also mutation in PmrB R256G	Pishnian, Haeili and Feizi (2019)
<i>Klebsiella</i> spp.	258a	CrrB (353)	–	Q10L	16	Confirmed	Human clinical isolate	Wright et al. (2015)
	ns	CrrB (353)	TM1 (12–34)	Y31H	512	Confirmed	Human clinical isolate	Cheng et al. (2016)
	ns	CrrB (353)	HAMP (81–135)	F84S	>128	Confirmed	Human clinical isolate	Jayol et al. (2017)
	ns	CrrB (353)	HAMP (81–135)	L87V	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	258a	CrrB (353)	HAMP (81–135)	L94M	16	Confirmed	Human clinical isolate	Wright et al. (2015)
	ns	CrrB (353)	HisKA (136–200)	W140R	2048	Confirmed	Human clinical isolate	Cheng et al. (2016)
	ns	CrrB (353)	HisKA (136–200)	N141I or N141Y	2048 / >128	Confirmed	Human clinical isolate	Cheng et al. (2016), Jayol et al. (2017)
	ns	CrrB (353)	HisKA (136–200)	P151S or P151L	1024 / >128	Confirmed	Human clinical isolate	Cheng et al. (2016); Jayol et al. (2017)
	ns	CrrB (353)	HisKA (136–200)	G183V	>128	Confirmed	Human clinical isolate	Jayol et al. (2017)
	ns	CrrB (353)	HisKA (136–200)	L191F	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
<i>C. freundii</i>	ns	CrrB (353)	HisKA (136–200)	S195N	2048	Confirmed	Human clinical isolate	Cheng et al. (2016)
	ns	CrrB (353)	–	S322W	ns	Not confirmed	Human clinical specimen	Macesic et al. (2020)
	117	CrrB (353)	HAMP (81–135)	A91T	256	Not confirmed	Human clinical specimen	Rocha et al. (2020)

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis

Table 9. Additional colistin resistance-associated genes in *Enterobacterales*

Bacterial genera	ST	Protein, length [aa]	Protein domain (residues)	Amino acid change	Resulting colistin MICs mg/L	Experimentally confirmed mutation	Bacterial source and comments	Reference
<i>Klebsiella</i> spp.	43	YciM (LapB) (389)	-	V43G	48	Confirmed	Human clinical isolate	Halaby et al. (2016)
	11	YciM (LapB) (389)	-	N212T	ns	Not confirmed	Human clinical isolate, also mutation in other genes involved in colistin resistance	Boszczowski et al. (2019)
<i>Klebsiella</i> spp.	11	LpxM (324)	-	N6K	ns	Not confirmed	Human clinical isolate, also mutation in other genes involved in colistin resistance	Boszczowski et al. (2019)
	43, 11, 23, 340	LpxM (324) LpxM (324)	- -	V30G S285G	8 ns	Confirmed Not confirmed	Human clinical isolate Human clinical isolate, also mutation in other genes involved in colistin resistance	Halaby et al. (2016) Boszczowski et al. (2019)
<i>Klebsiella</i> spp.	11	LpxM (324)	-	P321T	ns	Not confirmed	Human clinical isolate, also mutation in other genes involved in colistin resistance	Boszczowski et al. (2019)
	ns	RamA	-	V82A	ns	Not confirmed	Human clinical isolate	Macesic et al. (2020)

ST: sequence type; ns: not specified; confirmed (grey background); experimentally confirmed mutation mediating colistin resistance; not confirmed: mutation found by *in silico* analysis.

common phenomenon. Interestingly, Macesic *et al.* (2020) found a positive correlation between the number of mutations and the MIC value of a colistin-resistant isolate. Overall, the majority of suggested missense mutations in those fourteen genes were found by *in silico* analysis and not experimentally validated, which complicates the identification of genes and associated mutations crucial for the colistin resistance phenotype in *Enterobacterales*. Therefore, the summarized mutations (Tables 1–9) are likely to be an overestimation of the actual number of genetic changes that cause resistance to colistin. Furthermore, many studies analyze the sequence of only a limited number of genes, e.g. only *pmrAB*, and may not identify the critical mutation(s) responsible for colistin resistance. Vice versa, genetic changes are also found in colistin resistance determinants of susceptible isolates, which complicates the identification of causative mutations in resistant strains. Most likely, not all genes that play a role in the colistin-resistant phenotype have been identified so far. Interestingly, *Enterobacterales* isolates have also been found to have a dual resistance mechanism as they harbor a plasmid-borne *mcr*-gene and mutational alterations in colistin resistance genes (Garcia-Menino *et al.* 2019; Zakaria, Edward and Mohamed 2021). Those observations raise new questions that need to be addressed: Which mechanism was developed first and did it provide sufficient resistance? Is there an additive effect of both resistance mechanism? Did one mechanism even facilitate the acquisition of the second mechanism? Is the presence of the *mcr*-gene only accidental, since other beneficial genes are encoded on the plasmid?

### Species-specific colistin resistance mechanisms

Especially in *K. pneumoniae*, two more resistance mechanisms have been described which are not based on mutational changes in chromosomal genes. The overproduction and shedding of anionic capsular polysaccharide prevents cationic polymyxins to reach their target on the outer membrane (Llobet, Tomas and Bengoechea 2008). In addition, the overexpression of efflux pumps, such as AcrAB and KpnEF, has been suggested as an effective mechanism to exfiltrate the antibiotic from the bacterial cell (Padilla *et al.* 2010; Srinivasan and Rajamohan, 2013; Naha *et al.* 2020).

### Colistin and the polymyxin family

The family of polymyxins comprises five antimicrobial compounds (polymyxin A, B, C, D and E). Due to their reduced renal toxicity compared to the other polymyxins, only polymyxin B and E (colistin) are used as last-resort defense against severe infections with CRE in human medicine (Li, Nation and Kaye 2019). The polymyxins share a similar structure and are pentacationic polypeptides consisting of a cyclic heptapeptide linked to a linear tripeptide, whose N-terminus is acylated with a fatty acid moiety. Colistin is a secondary metabolite peptide, which is nonribosomal produced by the soil bacterium *Paenibacillus polymyxa* (formerly named *Bacillus polymyxa*). Since its introduction in the 1950s, colistin has been used continuously in the veterinary medicine to treat and prevent animal infectious diseases caused by Gram-negative bacteria. For the treatment of human infections, colistin was initially used therapeutically in Japan and in Europe during the 1950s and in the United States in 1959. However, the intravenous formulations of colistin and polymyxin B were gradually abandoned in most parts of the world in the early 1980s. Colistin was restricted

to ophthalmic and topical use owing to concerns about neurotoxicity and nephrotoxicity (Ryan *et al.* 1969; Brown, Dorman and Roy 1970; Koch-Weser *et al.* 1970). Thereafter, colistin was re-introduced for systemic treatment of lung infections due to MDR, Gram-negative bacteria in patients with cystic fibrosis (Conway *et al.* 1997; Ledson *et al.* 1998). Given the increased detection of colistin-resistant bacteria in livestock animals and animal-related food products as well as the need to retain the efficacy of antimicrobials to treat MDR infections in humans, the use of colistin in veterinary medicine is being re-evaluated. Throughout this review, and unless otherwise indicated, the term polymyxins is used to refer to the two clinically relevant compounds, polymyxin B and colistin.

## COLISTIN USAGE IN EUROPE

### Colistin resistance in the food, animal and livestock sector

Colistin is mainly administered orally in form of premix, powder and oral solutions in feed, drinking water or during milk replacer diets for the treatment of gastrointestinal tract infections caused by non-invasive *E. coli*. Colistin products are given to an enormous amount of different animal species including pigs, poultry, cattle, sheep, goats, laying hens and rabbits, but also to milk-producing species such as cattle, sheep and goats. Within the European Union (EU) and the European Economic Area (EEA), colistin and polymyxin B are authorized at national level and have been used since the 1950s (Table S2, Supporting Information). Colistin was widely used among the European countries in food-producing animals and their consumption extended beyond the treatment of infections to include pro- and meta-phylaxis purposes. At that time, primary indications were the treatment and prevention of diarrhoea in pigs caused by *E. coli* and *Salmonella* spp., the treatment of neonatal diarrhoea in piglets and veal calves caused by *E. coli*, as well as the treatment of mild colibacillosis in poultry (Timmerman *et al.* 2006; Pardon *et al.* 2012).

However, the extensive use of colistin has led to the emergence and spread of AMR pathogenic and commensal bacteria in the intestinal tract of food-producing animals. Resistant bacteria could colonize the human microbiota via the food chain through handling and/or consumption of contaminated food products. In light of this, initial restrictions on the use of colistin were implemented in the EU in 2006, prohibiting the supplementation of animal feed with antibiotics to promote animal growth (Regulation 1831/2003/EC; European Commission 2005). Importantly, in 2014 the EU implemented mandatory susceptibility testing to colistin for bacteria isolated from food-producing animals covered by the national monitoring programs (Regulation 2013/652/EU; European Commission 2013).

In 2011, the sales of antimicrobials were collected on EU level and published standardized and corrected for the total weight of treated animals. Following the reports of 25 EU/EEA members, polymyxin was the 5th most sold class of antimicrobials after tetracyclines (37%), penicillins (23%), sulphonamides (11%) and macrolides (8%) (European Medicines Agency (EMA), European Surveillance of Veterinary Antimicrobial Consumption 2014). In 2012, the overall consumption of polymyxins was approximately 600 times higher in food-producing animals compared to humans within the 19 member states of the EU/EEA, which reported complete data for both the animal sector and human medicine and after controlling for biomass (ECDC – European Centre for Disease Prevention and Control; EFSA –

European Food Safety Authority and EMA – European Medicines Agency 2015). The latest data comparing the consumption of polymyxins among 28 EU/EEA member states show that the population-weighted mean of consumption is 340 times higher in food-producing animals compared to human medicine (ECDC – European Centre for Disease Prevention and Control, EFSA – European Food Safety Authority and EMA – European Medicines Agency 2017). However, polymyxin usage among countries varied dramatically within the sector. In addition, there was no association of the consumption of polymyxin between food-producing animals and humans within a country. As example, in 2014, Italy, Spain and Portugal reported the highest polymyxin usage in food-producing animals, whereas Greece, Ireland and United Kingdom consumed most of the polymyxin in human medicine after correction for biomass. Following the implementation of mandatory monitoring of AMR in zoonotic and commensal bacteria, a statistically significant positive correlation between polymyxin usage in animals and the emergence of polymyxin resistance in *E. coli* could be shown in 2014–2015 (ECDC – European Centre for Disease Prevention and Control, EFSA – European Food Safety Authority and EMA – European Medicines Agency 2017). In order to secure colistin (polymyxin E) as last-resort antibiotic for the human medicine and in accordance with the One Health approach, effort has been made to limit the use of polymyxins in food-producing animals. Overall, the sales of polymyxin in the veterinary sector were reduced by 54% in the EU between 2011 and 2018 (Fig. 1). The European Medicine Agency (EMA) recommended in their advice on colistin use in 2016 that EU member states with high polymyxin use should reduce the consumption in livestock below 5 mg/PCU by 2020 (European Medicines Agency (EMA) 2016). The latest data on polymyxin consumption by the EMA demonstrate that still six countries exceed this threshold in 2018, including Cyprus, Germany, Hungary, Poland, Portugal and Romania (European Medicines Agency – EMA, European Surveillance of Veterinary Antimicrobial Consumption 2020). However, the use of polymyxin cannot be completely abolished at present. Polymyxin is of great importance for the treatment of intestinal infections in pigs, poultry and veal calves caused by *Salmonella* spp. or *E. coli* due to its narrow bactericidal spectrum against Gram-negative bacteria (European Medicines Agency – EMA 2016).

### Human medicine


Colistin is used as last-line antimicrobial for treating infections with CRE that belong to MDR isolates and their occurrence has already been reported worldwide (Grundmann et al. 2010). Especially immunocompromised patients, e.g. in intensive care units or oncology wards, are susceptible to CRE infections (Satlin, Jenkins and Walsh 2014; Satlin and Walsh, 2017). Such infections are difficult to treat and with limited therapeutic options thereby leading to high morbidity and mortality rates (Parisi et al. 2015). The control of CRE infections can be achieved by two commercially available forms of colistin, colistin sulphate for oral and topical use and the prodrug CMS (syn. colistin methanesulphate, colistin sulphonyl methate and pentasodium colistin methanesulphate) for parenteral use. Both drug forms can also be delivered by inhalation. CMS is microbiologically inactive and is less toxic than colistin sulphate. Following administration, CMS is hydrolyzed to colistin mediating the antibacterial effects. Besides colistin, also polymyxin B is licensed within the EU/EEA, but only for topical use. Colistin has been used for the treatment of infections at different body sites,

e.g. bacteremia and ventilator-associated pneumonia. Especially in combination with other antibiotics such as tigecycline or carbapenems, colistin has been the preferred treatment option for carbapenemase-producing *Enterobacterales* (Li, Nation and Kaye 2019).

As reported for other antibiotics, colistin resistances emerged rapidly following its re-introduction in human medicine (Meletis et al. 2011; Capone et al. 2013). Several studies reported increasing colistin resistance rates in carbapenemase-producing *K. pneumoniae* in individual hospitals in Greece of 0% in 2007, 8.13% in 2008, 24.3% in 2009, 21.7% in 2013 and worryingly, an average of 40.4% between the years 2014 and 2016 in 15 participating hospitals (Meletis et al. 2011, 2015; Galani et al. 2018). In 2013, colistin resistance rates for CRE isolates for Spain and Italy were 31% and 43%, respectively (Monaco et al. 2014, Pena et al. 2014). A recent study reported current and slightly decreased colistin resistance rates in an Italian hospital of 20.1% in 2017, 31.2% in 2018 and 26.9% in 2019 (Basso et al. 2020). During the years 2007–2014, Norway identified a prevalence of chromosomally mediated colistin resistance of 21% among the tested human clinical isolates (Samuelsen et al. 2017). Germany reported a prevalence of 13.3% of colistin resistance among carbapenemase-producing *K. pneumoniae* isolated between 2011 and 2016 (Koppe et al. 2018). Especially outbreaks with colistin-/CREs are of great concern due to dramatically limited treatment options (Antoniadou et al. 2007; Mezzatesta et al. 2011; Mammina et al. 2012; Weterings et al. 2015; Haller et al. 2019). Alarmingly, the overall use of colistin in human medicine increased steadily in the EU between 2005 and 2018, probably due to an increase in MDR-resistant isolates (Fig. 1). However, there is a strong geographical heterogeneity regarding the colistin consumption among the European countries, with increased use in Greece, Malta and United Kingdom and no consumption in Austria, Portugal, or Germany (ESAC-Net interactive database, accessed 22nd October 2020). The choice of antibiotics to treat colistin-resistant XDR isolates depends on the infection type and body site, the susceptibility of the isolate as well as the pharmacokinetic/pharmacodynamic properties and potential side effects of the antimicrobial (Petrosillo, Taglietti and Granata 2019).

Noteworthy, the United Kingdom, Sweden and Greece have implemented a mandatory notification system of bloodstream infections caused by colistin-resistant bacteria (Anderson, Cecchini and Mossialos 2020).

In 2016, the WHO classified polymyxins into the group of critically important antimicrobials (CIA) with highest priority (HPCIA) for human medicine (World Health Organization (WHO) 2019). Complementary to this, the World Organization for Animal Health (OIE) included polymyxins in their list of veterinary antimicrobial agents into the class of high importance (World Organisation for Animal Health – OIE 2018). In 2017, in response to the steadily increasing AMR, the EU has stressed the One Health approach to combat antibiotic resistance in the animal and human medicine and to prevent transmission of zoonotic diseases (European Commission 2017). In 2019, The EMA updated its 2014 advice on the categorization of antibiotics used in veterinary medicine, which could pose a risk for human public health. Polymyxins are classified into category B (“Restrict”), which includes antimicrobials from the WHO HPCIA list, and should only be used in food-producing and companion animals for the treatment of infections when there is no alternative antibiotic from category C or D (EMA/688114/2020; European Medicines Agency – EMA 2019).



**Figure 1.** Comparison of changes in sales and consumption of polymyxin in veterinary and human medicine within the European Union. Left: Averaged percentages of sales of polymyxin relative to total sales (in tons of active ingredient) in five European countries for the period 2005–2009 using the data from the first ESVAC report (European Medicines Agency 2011). The period 2010–2017 shows the averaged percentage of sales of polymyxin relative to the total sales (in mg of active ingredient by Population Correction Unit (PCU)) of the reporting countries for each year (European Medicines Agency 2012; European Medicines Agency 2013; European Medicines Agency 2014; European Medicines Agency 2015; European Medicines Agency 2016; European Medicines Agency 2017; European Medicines Agency 2018; European Medicines Agency 2019; European Medicines Agency 2020). Right: Annual average consumption of polymyxin [in Defined Daily Doses (DDD) per 1000 inhabitants per day] in the community and hospital sector in Europe, including Switzerland, using data provided by the ESAC-Net interactive database (<https://www.ecdc.europa.eu/en/antimicrobial-consumption/surveillance-and-disease-data/database>, accessed October 2020). For both sectors, polymyxins include colistin (polymyxin E) and polymyxin B.

Since the monitoring of colistin resistance in *Enterobacteriales* of livestock is mandatory, the surveillance of colistin-resistant isolates from human cases is still in its infancy. EU-wide data on colistin resistance in human isolates are collected only for the zoonotic agent *Salmonella*, but only a quarter of the EU/EEA member states reported to the ECDC (EFSA – European Food Safety Authority and ECDC – European Centre for Disease Prevention and Control 2019). The burden of AMR in Europe is assessed through the ECDC and the European Antimicrobial Resistance Surveillance Network (EARS-Net) in collaboration with national institutions. However, EARS-Net collects data from a limited number of bacterial species isolated from human blood and cerebrospinal fluid. As a result, infections affecting the urinary and respiratory tract caused by resistant bacteria, such as *E. coli* or *Klebsiella* spp., are not recorded. Moreover, colistin resistance in *Enterobacteriales* is not assessed by EARS-Net as it is not included in the initial routine antimicrobial susceptibility testing and may only be examined by national laboratories

(European Centre for Disease Prevention and Control – ECDC 2020). In 2019, the ECDC initiated a carbapenem-and/or colistin-resistant *Enterobacteriales* (CCRE) project, which is a European network aiming to complement the phenotypic data collected by EARS-Net with WGS-based data, to address the needs described above. It is desirable however, to expand this effort to the surveillance data collected from the veterinary sector. As Tacconelli *et al.* pointed out, national and European-wide surveillance systems of AMR in livestock, the food chain and humans are very heterogeneous and need enhancement as well as improved multisectorial collaboration (Tacconelli *et al.* 2018). However, the most important basis for comparable data is the methodology for reliable colistin susceptibility testing. Standard broth microdilution (BMD) is the only recommended method by the European Committee on Antimicrobial Susceptibility Testing and Clinical and Laboratory Standards Institute for determination of the MIC values of colistin (European Committee on Antimicrobial Susceptibility Testing – EUCAST

2014), (Performance Standards for Antimicrobial Susceptibility Testing – CLSI 2020). However, this method is not automated, therefore time-consuming and difficult to incorporate into the routine of diagnostic laboratories. Other applied methodologies are broth microdilution, disc diffusion, agar dilution, Etest, Rapid Polymyxin NP test and automated systems. For a reliable MIC determination, most likely two different systems should be combined such as an automated system and BMD, since so far all methods show specific weaknesses (Jayol et al. 2018; Garcia-Menino et al. 2020).

## GLOBAL USE OF COLISTIN: THE INCIDENCE AND MONITORING OF RESISTANCE

The global use of antibiotics in livestock animals is growing rapidly due to the growing global population and the increased production and consumption of animal protein as a result of increased incomes in fast-developing middle-income countries. The current OIE report states that still nine countries use colistin as growth promoter (World Organisation for Animal Health – OIE 2021). Contrary to the OIE report, Olaitan et al. (2021) pointed out that the majority of low- and middle-income countries still use colistin as feed additive. Discrepancies could be due to incomplete or inaccurate reporting to the OIE. In human medicine, the worldwide antibiotic consumption increased by 36% between 2000 and 2010, with Brazil, Russia, India, China and South Africa (BRICS) accounting for three-quarters of this increase, however, representing only 40% of the world's population. Thereof, the increase in colistin use in hospitals in BRICS countries for the same period corresponds to 13% (Laxminarayan et al. 2016).

The prevalence of the global colistin resistance among human clinical *Enterobacterales* isolated between 2012 and 2013 was 1.6% with a regional distribution of the resistance in Europe (1.8%), North America (1.3%), Latin America (1.5%), Middle East-Africa (1.4%) and the Asia-Pacific (1.3%) (Bradford et al. 2016). The most abundant resistant genera was *Enterobacter* spp., followed by *K. pneumoniae* and *E. coli*. By using the surveillance data from the ATLAS database, the global incidence of colistin-resistant human clinical *Enterobacterales* increased between 2014 and 2019 from 2.6 to 3.6%, showing a regional distribution of 2.4–3.4% in Europe, 1.2–2.6% in North America, 2.7–4.3% in Latin America, 3.3–6.7% in Asia, 2.1–2.6% in Africa and 0.6–2.7% in Oceania (Fig. 2).

### Asia

#### China

China is a leading consumer and producer of colistin worldwide. Between 2011 and 2015, about 2875 metric tons of colistin were used in food-producing animals annually (Shen et al. 2016). In November 2016, the Ministry of Agriculture announced the withdrawal of colistin as feed additive to promote animal growth, which took into effect in April 2017 (Walsh and Wu, 2016). Thereafter, the production of colistin sulfate premix decreased steadily, from 27 170 tons in 2015 to 2497 tons in 2018 with the most significant decrease between 2016 and 2017 (Wang et al. 2020). Polymyxin B and colistin became available for the use in humans in October 2017 and December 2018, respectively (Table S2, Supporting Information).

#### India

Data from 2016 on colistin shipments showed strong exports of the antimicrobial from China to India, South Korea and Vietnam. Antimicrobial use in chickens is estimated to increase 5-fold by


2030 compared with 2013 (Davies and Walsh, 2018). However, data on colistin-resistant *Enterobacterales* from livestock animals are lacking. The Indian Antimicrobial Resistance Surveillance & Research Network collects data from AMR profiles from clinically relevant pathogens. Between 2016 and 2018, the prevalence of colistin resistance in *E. coli* and *K. pneumoniae* causing hospital-acquired infections was 1.1% and 8%, respectively (Walia et al. 2019). Alarming, 70 and 83.3% of *E. coli* and *K. pneumoniae* isolated from hospital wastewater were resistant to colistin (Bardhan, Chakraborty and Bhattacharjee 2020). Another study analyzing the presence of colistin-resistant bacteria on raw food, including meat, fish and vegetables, found that 46.4% of the samples contained colistin-resistant bacteria, of which the most prevalent species were *E. coli* and *Klebsiella* species (Ghafur et al. 2019). Remarkably, since July 2019 the Indian Union Ministry of Health and Welfare has banned colistin for the use in food-producing animals, which goes beyond the use as growth promoter and includes the prohibition for pro- and metaphylaxis purposes (section 26A of Drugs and Cosmetics Act, 1940; Table S2, Supporting Information).

In 2019, the rate of colistin resistance in *Enterobacterales* isolated from clinical specimen in Asia was 25.4% for *Enterobacter* spp., 4.0% for *Klebsiella* spp. and 1.2% for *E. coli* (Fig. 2). Only few studies from Asia investigated chromosome-mediated colistin resistance in *Enterobacterales*, which mainly comprises the analysis of clinical *Klebsiella* spp. isolates from Lebanon, Taiwan, Turkey, Iran and United Arab Emirates (Cheng et al. 2015; Okdah et al. 2017; Can et al. 2018; Moubareck et al. 2018; Dagher et al. 2019; Jafari et al. 2019). Described resistance mechanisms were disruption of the *mgrB* gene, as well as missense mutations in *MgrB*, *PhoQ* and *PmrB*. A total of two studies from South Korea and Japan found missense mutations in *PhoPQ* and *PmrAB* two-component system in clinical *E. cloacae* strains, but their contribution to colistin resistance was not confirmed (Hong and Ko, 2019; Uechi et al. 2019).

A systematic review covering southeast Asia identified reports on chromosomal mutations in *mgrB* in *Klebsiella* spp. isolates (Malchione et al. 2019). Susceptibility to colistin is not routinely tested in these countries, resulting in limited data and underestimation of resistance levels. Colistin resistance has also been described for *Salmonella* spp. strains from chicken and turkey in South Korea and Taiwan, but the mechanism of resistance has not been specified (Yeh et al. 2018; Seo et al. 2019).

### Australia

Australia has approved polymyxin B but not colistin or its formulations for the use in livestock animals (Commonwealth of Australia 2018; Table S2, Supporting Information). Studies analyzing resistant bacteria from poultry and egg layer flocks found neither colistin-resistant *Enterobacterales* nor mobilizable genes conferring colistin resistance (Abraham et al. 2019; Bean et al. 2020; Veltman et al. 2021). Colistin and polymyxin B are authorized for the use in human medicine (Australian Commission on Safety and Quality in Health Care 2019). The Australian National Alert System for Critical Antimicrobial Resistances (CARAlert) monitors only transmissible colistin resistance in clinical *Enterobacterales* of human origin. Only a few studies reported infections caused by colistin-resistant *Enterobacterales* in humans. In accordance with the data provided by the ATLAS database, the percentage of resistant isolates collected between 2007 and 2016 was 2.1% (Fig. 2; Ellem et al. 2017). 98% of the isolates analyzed were negative for *mcr-1*, suggesting that they carry either a different *mcr*-gene or a chromosomal mutation (Ellem et al. 2017).




**Figure 2.** Global trends in colistin resistance in human clinical *Enterobacteriales*. Data were obtained from the ATLAS database (<https://atlas-surveillance.com>, accessed February 2021), which includes data from the TEST (Tigecycline Evaluation Surveillance Trial) surveillance program, the AWARE (Assessing Worldwide Antimicrobial Resistance Evaluation) as well as INFORM (International Network for Optimal Resistance Monitoring) program. Resistance is shown as percentage from *Enterobacter* spp., *Klebsiella* spp. and *E. coli* using the data from all surveillance programs (ATLAS data source) and MIC values >2 mg/L according to the EUCAST breakpoint. (A) Global trend of colistin resistance among clinical *Enterobacter* spp., *Klebsiella* spp. and *E. coli* as well as combined genera from 2014 to 2019. (B) Trends of colistin resistance in clinical *Enterobacteriales* in different continents from 2014 to 2019. (A and B) Data reporting countries were: Europe: Austria, Belgium, Croatia, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Netherlands, Poland, Portugal, Romania, Russia, Spain, Sweden, Switzerland, Turkey, Ukraine and United Kingdom; North America: Canada, United States; Latin America: Argentina, Brazil, Chile, Colombia, Costa Rica, Dominican Republic, Guatemala, Mexico, Panama and Venezuela; Asia: China (incl. Hong Kong and Taiwan), Japan, Korea, South, Malaysia, Philippines and Thailand; Africa: Israel, Jordan, Kenya, Kuwait, Morocco, Nigeria, Qatar, Saudi Arabia and South Africa; Oceania: Australia and New Zealand.

## North America

### USA

Colistin has been approved but never been marketed for the use in food-producing animals in the United States (U.S.; <https://www.center4research.org/8094-2/>, accessed June 2021). Since 2009, polymyxin B has been approved and used as antimicrobial (Table

S2, Supporting Information; U.S. Food and Drug Administration – FDA 2016; <https://animaldrugatfda.fda.gov/adafda/views/#/home/searchResult>, accessed June 2021). Therefore, the presence of colistin resistance in *Enterobacteriales* remains very low with a prevalence of 0.1% in animals at slaughter and 0.02% from animal meat (Meinersmann et al. 2017; Wang et al. 2020). Within the National Antimicrobial Resistance Monitoring System (NARMS),



**Figure 3.** Regulatory network of LPS-modifying proteins involved in colistin resistance in *Enterobacterales*. The PhoPQ TCS is activated by low  $Mg^{2+}$  concentrations, low pH and the presence of antimicrobials peptides, such as colistin, leading to the expression of the regulator MgrB, the adaptor protein PmrD and the sRNA *mgrR*. MgrB exerts negative feedback on PhoQ, while mutations in MgrB typically result in the constitutive activation of the PhoPQ TCS. The sRNA *mgrR* impedes the expression of EptB. The adaptor protein PmrD activates the PmrAB TCS leading to the expression of multiple target genes responsible for LPS biosynthesis and modification. PmrA also becomes activated by the CrrAB TCS via the adaptor protein CrrC. Gain-of function mutations in CrrB can also result in the activation of gene expression of the *pmrHFIJKLM* operon without involvement of the PmrAB TCS. In addition, mutations in the proteins YciM and LpxM have been found to confer colistin resistance. The plus symbol indicates positive regulation and the yellow star highlights alterations in proteins/genes, which may lead to colistin resistance.

which collects bacteria from humans, animals and food, isolates are not tested for susceptibility to colistin. The latest report states that NARMS sequenced the genomes of 55 000 *Salmonella* and *E. coli* isolates and found only *mcr-9.1* in 55 *Salmonella* strains from humans, animals and meat, as well as in five *E. coli* isolated from meat (National Antimicrobial Resistance Monitoring System – NARMS 2018, Wang et al. 2020). Macesic et al. (2020) found chromosomal alterations in colistin resistance genes of *K. pneumoniae* in 36% of patients. According to the United States Committee on Antimicrobial Susceptibility Testing, reliable detection of colistin resistance in human clinical isolates is lacking due to the use of test systems (e.g. Etest, disk diffusion) other than the recommended reference method, and the absence of colistin in most automated test systems, which accounts for over 90% of the U.S. test results (Pogue et al. 2020).

#### Canada

The Public Health Agency of Canada has announced that polymyxin B is approved for the use in livestock animals (Canadian Ministry of Agriculture 2019). An EMA report from 2016 states that colistin was not approved for the use in veterinary medicine. Unfortunately, we were not able to conclusively clarify the statement with the source of the Public Health Agency of Canada indicated in the EMA report or with other sources (European Medicines Agency – EMA 2016). However, a loophole in the regulation gave farmers the opportunity to import and use unlicensed, non-prescription antimicrobial combinations of third-generation cephalosporins and penicillin containing

colistin in their livestock, which were used very frequently on dairy farms (Saini et al. 2012; Webb et al. 2017). In 2016, the loophole of “importation for own use” has been recognized and regulatory changes have been proposed that would prohibit those practices (<https://canadagazette.gc.ca/rp-pr/p1/2016/2016-07-02/html/reg2-eng.html>, accessed August 2021). In addition, polymyxin B and colistin are used in the human medicine (Public Health Agency of Canada 2018). The Canadian Ward Surveillance Study monitors AMR in multiple pathogens in 15 hospitals across the country. Across samples collected from 2007 to 2016 the prevalence of colistin resistance in *E. coli*, *Klebsiella* spp. and *Enterobacter* spp. was 0.2%, 5.8% and 18.1%, respectively (Zhan et al. 2019). Unfortunately, the trend of colistin resistance development within the period was not calculated for the corresponding isolates (Lagace-Wiens et al. 2019).

In 2019, the overall prevalence of colistin resistance among clinical *Enterobacterales* in North America was 10.4% for *Enterobacter* spp., 0.8% for *Klebsiella* spp. and 0.3% for *E. coli* (Fig. 2).

#### Central and South America

The Pan American Center PANAFOTSA oversees AMR in zoonotic agents, while AMR in community- and hospital-acquired pathogens is monitored by the ReLAVRA (Latin American Network for Antimicrobial Resistance Surveillance), which covers 19 member states. However, uniform data on the overall colistin susceptibility in *Enterobacterales* of animal and human origin from member states are not available. Colistin is no



longer approved for the use in livestock in Nicaragua, Costa Rica, Peru, Paraguay and Argentina (Table S2, Supporting Information). Besides Guatemala, El Salvador, Panama, Venezuela, Guyana, Suriname and French Guiana, which did not provide information, colistin is approved as a therapeutic in the other countries of Latin America (unpublished, personal communication with PANAFTOSA, February 2021). A study analyzing colistin resistance in *Enterobacteriales* collected from hospitals in six South American countries between 2015 and 2017 found a prevalence of resistance of 5.6% in *E. cloacae*, 4.9% in *K. pneumoniae*, 1.4% in *K. aerogenes*, 0.9% in *K. oxytoca* and 0.8% in *E. coli*, which is similar to the data provided by the ATLAS database (Stone and Ponce-de-Leon, 2020; Fig. 2). In 2019, resistance to colistin among human clinical *Enterobacter* spp., *Klebsiella* spp. and *E. coli* was 12.8%, 3.5% and 0.3% respectively (Fig. 2).

### Brazil

Besides China, Brazil is one of the largest poultry producers and exporter globally (Food and Agriculture Organization of the United Nations – FAO 2020). In 2016, Brazil prohibited the use of colistin as feed-additive in food-producing animals (Brazil. Governmental Normative Instruction IN-45 2016). However, due to the lack of sufficient studies it is not possible to assess the colistin resistance levels in livestock animals. Morales et al. (2012) reported a prevalence of 6.3% of colistin-resistant *E. coli* in swine samples. The colistin resistance rate among human clinical *Enterobacteriales* increased from 6.6% in 2010 to 9.9% in 2014 using samples from nine hospitals of São Paulo, Brazil (Rossi et al. 2017). Another hospital reported a resistance rate of 5% in *Enterobacteriales* mediated by chromosomal mutations (2%) and *mcr*-genes (3%; Rocha et al. 2020).

### Colombia

A study from Colombia analyzing colistin-resistant clinical isolates found that only 2.3% of the *Enterobacteriales* carried *mcr*-1, suggesting that the majority of isolates have a chromosomal mutation that leads to colistin resistance (Saavedra et al. 2017). However, analyzed isolates contain the serovars *S. Enteritidis* and *S. Dublin*, which seem to exhibit increased intrinsic resistance to colistin, eventually leading to an overestimation of the overall number of resistant strains.

The majority of studies analyzing the prevalence of colistin resistance in *Enterobacteriales* in Latin American countries focus on the presence of *mcr*-genes rather than chromosomal mutations. In total, two studies described disruptions of the *mgrB*-gene leading to colistin resistance in clinical *K. pneumoniae* strains isolated in Argentina and Uruguay (Alvarez et al. 2018; Escalante et al. 2020). Reports from other Latin American countries, such as Mexico, Argentina, Chile, Peru, Ecuador, Venezuela and Uruguay focus exclusively on the identification of *mcr*-1 in *E. coli* obtained from livestock, companion animals and human clinical specimen (Delgado-Blas et al. 2016; Ortega-Paredes, Barba and Zurita 2016; Dominguez et al. 2017, 2019; Garza-Ramos et al. 2018; Gutierrez et al. 2019; Merida-Vieyra et al. 2019; Rumi et al. 2019; Coppola et al. 2020; Loayza-Villa et al. 2020; Papa-Ezdra et al. 2020).

### Africa


Livelihoods of 250–300 million people in Africa is financed by animal husbandry, and the use of colistin in livestock remained largely unregulated (Van et al. 2020). For example, in 2015 in Morocco, colistin was the most frequently (27.85% of treatments) used antimicrobial for treatments in the broiler sector and the

second most commonly used antimicrobial by active ingredient. Furthermore, colistin was overdosed in most of the administrations (Rahmatallah et al. 2018). Similarly, colistin resistance in *E. coli* from South African poultry increased steadily from 3.9% in 2009 to 12.08% in 2015 (Theobald et al. 2019). Following the prohibition of colistin as a feed additive in 2016, resistance levels in avian *E. coli* in South Africa decreased to 1.77% (Table S2, Supporting Information). End of 2017, the Africa CDC founded the Anti-Microbial Resistance Surveillance Network (AMRSNET) for the monitoring of resistant organism in the animal and human health sector. Due to the recent establishment of the network, data regarding the overall prevalence of colistin-resistant *Enterobacteriales* in Africa are not yet available. Figure 2 shows resistance levels of 8.6% for *Enterobacter* spp., 2.4% for *Klebsiella* spp. and 0.19% of *E. coli* clinical isolates in 2019. A systematic review by Olowo-okere and Yacouba identified studies regarding colistin-resistant bacteria from Algeria, Egypt, Tunisia, South Africa, Libya, São Tomé and Príncipe and Nigeria. Overall, colistin resistance was most frequently described for *E. coli* isolates obtained from human clinical samples. Furthermore, studies reported both chromosomal and plasmid-mediated resistance mechanisms, of which plasmid-mediated colistin resistance was the most prevalent, accounting for up to 72.2%. Mutational changes were found in *pmrA/B* of *E. coli* and *K. pneumoniae* and additionally in *mgrB* of *K. pneumoniae* (Olowo-okere and Yacouba, 2020).

A study performed in South Africa analyzing colistin resistance mechanisms in human clinical strains collected between 2016 and 2017 notified a prevalence of *mcr*-1 in 55% of *E. coli* and 71% of *Klebsiella* isolates. The analysis of the chromosomal-mediated colistin resistance showed that genetic alterations occurred predominantly in *pmrB* and *mgrB* of *E. coli* and *Klebsiella* isolates, respectively (Snyman et al. 2021). However, colistin-resistant *Enterobacteriales* were isolated not only from hospitalized patients but, more worryingly, also from healthy hotel employees in Zanzibar, Tanzania, with a prevalence of 59.3%. Overall, 55% of the colistin-resistant *E. coli* isolates carried *mcr*-1, whereas none of *K. pneumoniae* harbored *mcr*-1 to *mcr*-8 (Budel et al. 2019). In addition to animals and humans, two reports from South Africa and Tunisia demonstrated colistin resistance among cefotaxime-resistant *E. coli* (76.5%) as well as ESBL-producing *Enterobacteriales* (10.8%) that were *mcr*-negative, isolated from waste water (Adegoke et al. 2020; Hassen et al. 2020).

### Transmission of colistin-resistant *Enterobacteriales* in a One Health perspective

Globalization connects the different areas of life but also facilitates the spread of AMR. Humans and animals, including domestic animals and wildlife, continuously interact with each other and share often the same habitat. The excessive use of colistin in animals resulted in the selection for resistance affecting both human and animal health. Overall, three different pathways for the transmission of resistant bacteria can be recognized: (i) transmission between animals and humans, (ii) transmission to humans/animals via contaminated food and (iii) transmission via the environment (Fig. 4). The transmission of colistin-resistant bacteria occurs in direct contact among animals, humans and between them (Budel et al. 2020). Especially human individuals in constant contact with animals, e.g. farm workers and veterinarians, are at greater risk of acquiring resistant microorganisms (Marshall and Levy, 2011). Slaughterhouses and farms are main places of inter-species transmission, where colistin-resistant bacteria are transferred from animals



**Figure 4.** Possible transmission routes of colistin-resistant *Enterobacteriales*. Colistin-resistant *Enterobacteriales* emerge as a result of the use of colistin in the livestock sector, in animal clinics and the hospitals. Resistant isolates can disseminate between different areas of life, which is indicated by the red arrows.

to humans in the event of contamination during the slaughter process, but also may enter the food chain and the environment via sewage. Chromosome-mediated colistin-resistant *E. coli* were found in livestock in Europe, Asia and Africa (Rebello et al. 2018; Kim et al. 2019; Budel et al. 2020; Massella et al. 2020).

Backyard livestock and small family farms are common husbandry systems for food-producing animals in Asia where antimicrobials are often overused resulting in high resistance levels (Hallenberg et al. 2019; Kawahara et al. 2019). Furthermore, poor hygiene and close contact with farm staff facilitates the transmission of colistin-resistant *Enterobacteriales* from animals to humans (Trung et al. 2017). Noteworthy, transmission of colistin-resistant *E. coli* also occurs between animals on integrated poultry-fish farming systems where chickens are kept over fish ponds and feces is excreted into the ponds. In 2017, the most common antimicrobial used in poultry-fish farms in Myanmar was Octamix (amoxicillin and colistin sulfate) in over 40% of poultry flocks and 6% of *E. coli* isolated from feces were resistant to colistin (Gibson et al. 2020).

Besides farm animals, also companion animals are increasingly discussed to serve as vehicle for a potential bacterial transmission (Joosten et al. 2020; Marin et al. 2021). Notably, a study from Switzerland uncovered the contamination of surfaces in companion animal clinics with colistin-resistant *Enterobacteriales* and worryingly, the colonization of employees with resistant strains (Schmidt et al. 2020). Although farm workers, veterinarians and pet owners represent only a limited number of people having close contact with animals, they still provide an entry point for the transmission of resistant *Enterobacteriales* into the community and hospitals. Several studies demonstrated precisely the spread of colistin-resistant pathogens, in hospital outbreaks. *Klebsiella* spp. in particular, with mutations in chromosomally encoded genes, is a major cause of the clonal dissemination in clinical settings around the world (Mezzatesta et al. 2011; Mammìna et al. 2012; Goel et al. 2014; Giani et al. 2015; Weterings et al. 2015; Jayol et al. 2016; Kocsis et al. 2017; Avgoulea et al. 2018; Guducuoglu et al. 2018; Haller et al. 2019). But also clinical *Enterobacter* spp. isolates harboring colistin resistance have

been described (Hong, Lee and Ko 2018). Transmission of resistant bacteria during surgery, the acquisition from hospital surfaces, or solely the transfer between patients and health care workers via hand contact may increase the risk for the dissemination of resistant pathogens in hospital settings (as already shown for methicillin-resistant *Staphylococcus aureus* or *A. baumannii*; Blanco, O'Hara and Harris 2019).

Resistant bacteria can also reach the consumer through the consumption of contaminated food, which displays a far more complex route of transmission. Sources for foodborne transmission are the consumption of animal-related products, such as meat and fish. Colistin-resistant *K. pneumoniae* lacking *mcr* genes were isolated from fish and poultry meat from Europe, Africa and Asia (Ghafur et al. 2019; Diaz-Jimenez et al. 2020; Chaalal et al. 2021). Furthermore, colistin-resistant *S. Abony* strains with mutations in chromosomal genes were located in fish farms (Antunes et al. 2018). A less noticed vehicle for transmission of colistin-resistant bacteria are vegetables and seafood, where the consumption in the raw state may display a greater risk (Ghafur et al. 2019). In addition, international food trade may facilitate the introduction and spread of colistin-resistant *Enterobacteriales*. Finally, resistant bacteria can also disseminate via waste material, such as sewage, contaminating the environment (Savin et al. 2020). Especially water is an efficient route for bacterial transmission into nature and wild life. Colistin-resistant *Enterobacteriales* have been found in several species of wild animals, such as mice, deer and sea lions across several countries (Wasyl et al. 2018; Hernandez-Castro et al. 2020; Skarzynska et al. 2020; Zanardi et al. 2020).

Altogether, these findings highlight the importance of national monitoring programs and global routine surveillance of colistin resistance in zoonotic bacteria of animal, food and human origin, which provides scientific data for the assessment of AMR burden as well as for strategic interventions.

### Colistin-resistant *Enterobacteriales* and associated sequence types

Studies regarding AMR in livestock animals focused primarily on *E. coli* and *Salmonella*. As early as 1975, it was determined that the administration of sub-therapeutic quantities of antibiotics is sufficient to develop resistance in *E. coli* in the gastrointestinal tract of chicken. Strikingly, the farm workers also acquired resistant *E. coli* in their intestine (Levy, Fitzgerald and Maccone 1976). Due to the uninterrupted use, colistin resistance has been continuously reported in *E. coli* and *Salmonella* isolated from farm animals.

Especially in healthcare setting, CRE, predominantly *K. pneumoniae* and *Enterobacter* spp., emerged in recent years as a major threat in the group of antibiotic-resistant pathogens (Peleg and Hooper, 2010; Chavda et al. 2016). Mortality rates are high due to limited treatment options and successful dissemination of certain strains. CRE have been reported from several countries, but especially the Mediterranean countries, such as Greece, Italy, Malta and Israel, report the rapid spread of endemic clones in many hospitals (Leavitt et al. 2007; Samra et al. 2007; Capone et al. 2013; Glasner et al. 2013; Albiger et al. 2015). In Europe, 23 European countries report a worsened epidemiological situation of CRE between the years 2010 and 2018 (Brolund et al. 2019). Colistin, together with tigecycline and gentamicin, is among the few antimicrobial available to treat infections with CRE (Petrosillo et al. 2013). Therefore, the emergence of resistance to colistin and other last-option antimicrobials especially in, but not restricted to, CRE is important to monitor.

## Escherichia coli

*Escherichia coli* is a part of the normal intestinal microbiome in animals and humans. However, *E. coli* is also the most prevalent bacterial agent causing community- and hospital-acquired infections such as urinary tract and bloodstream infections. To date, no association could be established between colistin resistance in clinical isolates and consumption in human medicine due to lack of data on *E. coli* (ECDC – European Centre for Disease Prevention and Control, EFSA – European Food Safety Authority and EMA – European Medicines Agency 2017). In contrast, in food-producing animals, a strong positive correlation was revealed between the resistance to and the consumption of colistin. The European Food Safety Authority (EFSA) and the European Centre for Disease Prevention and Control (ECDC) identified in their report from 2019a prevalence of colistin resistance of 0.3% and 0.8% in fattening pigs and calves, respectively (EFSA – European Food Safety Authority and ECDC – European Centre for Disease Prevention and Control 2019). Data for poultry were only available for 2015, when colistin resistance was reported in 1.7% of broilers and 5.7% of turkeys (EFSA – European Food Safety Authority and ECDC – European Centre for Disease Prevention and Control 2018). Only Italy reported on a voluntary basis data regarding colistin resistance in meat products, which was 5.3% in pig meat and 3.1% in bovine meat (EFSA – European Food Safety Authority and ECDC – European Centre for Disease Prevention and Control 2019).

Certain sequence types occur repeatedly among *E. coli* strains isolated from humans and food-related animals and worryingly, belong to high-risk clones, which display a major health concern. Several studies from Europe and Asia identified the *E. coli* ST10 as a common isolate in livestock animals and humans with bloodstream or urinary tract infections, which is globally distributed and able to develop chromosomally encoded colistin resistance (Fig. 5 and Table S3, Supporting Information; Luo et al. 2017; Kim et al. 2019; Janssen et al. 2020). *Escherichia coli* ST131 is currently the most important human clone worldwide and is frequently associated with chromosomal colistin resistance (Sato et al. 2016, 2018; Luo et al. 2017; Dafopoulou et al. 2020; Dagher et al. 2020; Snyman et al. 2021). Human clinical *E. coli* of the pandemic clonal group ST131 with chromosomal mutations in colistin-resistance determinants were reported predominantly from Asia and Europe, but also from Africa (Fig. 5 and Table S3, Supporting Information).

The review of the scientific literature revealed a predominant number of studies analyzing the presence of *mcr*-genes in *E. coli* strains obtained from livestock animals and animal products, as opposed to only a few reports dealing with chromosome-mediated colistin resistance in those isolates.

## Klebsiella pneumoniae

*Klebsiella pneumoniae* colonizes the respiratory and gastrointestinal tract of humans (Bagley, 1985; Martin and Bachman, 2018). XDR and PDR *Klebsiella* spp. strains are major cause of healthcare-associated infections and outbreaks leading to difficult-to-treat diseases, e.g. lower respiratory tract infections, urinary tract infections and bloodstream infections (Martin and Bachman, 2018). An example of this are carbapenemase-producing *K. pneumoniae* strains belonging to pandemic clones, such as ST11, ST147, ST258 and ST525, which have been reported to acquire resistances to numerous unrelated antimicrobial agents. (Comandatore et al. 2013; Monaco et al. 2014; Pena et al. 2014; Giani et al. 2015; Oteo et al. 2016; Samuelsen et al. 2017;


Diaz-Jimenez et al. 2020; Gentile et al. 2020). Colistin resistance caused by chromosomal alterations in isolates with the above mentioned sequence types is reported from countries in Africa, Asia, Europe as well as North and South America from human clinical specimen (Fig. 5 and Table S3, Supporting Information; Cannatelli et al. 2014; Jaidane et al. 2018; Teo et al. 2019; Macesic et al. 2020; de la Cadena et al. 2021). Additionally, colistin resistant ST11 is also isolated from animal sources from Africa and Asia (Pishnian, Haeili and Feizi 2019; Budel et al. 2020). The prevalence of chromosome-mediated colistin resistance among carbapenemase-producing isolates ranges from 6 to 80% among EU member states (Pena et al. 2014; Bonura et al. 2015; Jayol et al. 2016; Oteo et al. 2016; Otter et al. 2017; Samuelsen et al. 2017; Hamel et al. 2020). High colistin resistance rates among carbapenem-resistant klebsiellae, ranging from 27 to 61%, were also reported from Asia, South Africa and South America (Sampaio and Gales, 2016; Jafari et al. 2019; Al-Zalabani et al. 2020; Kopotsa, Mbelle and Sekyere 2020; Shankar et al. 2021). Notably, a strong correlation was found between emerging polymyxin resistance in *K. pneumoniae* isolates and consumption of polymyxin in the hospital sector (ECDC – European Centre for Disease Prevention and Control, EFSA – European Food Safety Authority and EMA – European Medicines Agency 2017).

In contrast to studies regarding colistin-resistant *E. coli* isolates from the livestock sector, most of the publications about human-pathogenic colistin-resistant *Klebsiella* strains analyzed the contribution of mutational changes in chromosomal genes.

## Salmonella enterica

Non-typhoidal *Salmonella* is a major cause of food poisoning resulting in gastrointestinal infections that range from asymptomatic to clinically severe illness. In 2018, salmonellosis was the second most common gastrointestinal infection in the EU/EEA (European Food Safety Authority – EFSA) and European Centre for Disease Prevention and Control – ECDC 2019). The predominant risk factor for the acquisition of *Salmonella* is the consumption of contaminated food, such as meat, eggs, vegetables and dairy products. The largest proportion of colistin-resistant *Salmonella* strains derived from livestock animals in the EU were found in cattle with a prevalence of 14.5% (EFSA – European Food Safety Authority and ECDC – European Centre for Disease Prevention and Control 2019). In calf carcasses under 1 year of age, 3.7% of *Salmonella* isolates were resistant to colistin. However, all resistant strains derived from cattle and calf carcasses belong to serovar *S. Dublin*. The ECDC and EFSA joint report from 2019 refers to a study from 2012, which suggests that the serovars *S. Dublin* and *S. Enteritidis* exhibit increased intrinsic resistance levels to colistin. However, only two chromosomal genes were analyzed in the mentioned study and this review summarizes additional genes involved in colistin resistance (Agero et al. 2012). In addition, other yet unidentified mechanisms or mutations in these serovars could mediate colistin resistance.

Furthermore, EU member states reported colistin resistance in 1.9% of *Salmonella* strains recovered from fattening pigs and 0.6% of *Salmonella* spp. from fattening pig carcasses, which belonged to different serovars (EFSA – European Food Safety Authority and ECDC – European Centre for Disease Prevention and Control 2019). In 2017, resistances to colistin were reported in 4.7% of all human *Salmonella* isolates with 88.9% of the resistant isolates belonging to either *S. Enteritidis* or *S. Dublin* (EFSA – European Food Safety Authority and ECDC – European Centre for Disease Prevention and Control 2019). Only seven EU member states reported data regarding colistin-resistant *Salmonella*



**Figure 5.** Landscape of *Enterobacteriales* sequence types associated with chromosomal mutations leading to colistin resistance. Worldwide prevalence of chromosomal-mediated colistin resistance in different sequence types from *E. coli*, *K. pneumoniae*, *Enterobacter* spp. and *Salmonella* spp. isolated from human, animal, food and wildlife. Detailed information is given in Table S2 (Supporting Information) in addition with information regarding sequence types of each species associated with *mcr*-gene

strains, six of which detected colistin resistance, suggesting an underestimation of the actual resistance levels. ST32 is the most common sequence type among serovar Infantis, which is an increasingly important avian serovar. ST32 isolates from poultry farms in Serbia were found to harbor chromosomal colistin resistance (Jovicic et al. 2020). Additionally, *S. Newport* is an emerging serovar in human infections and chromosomal colistin resistance has been found in isolates from Europe and Asia (Olaitan et al. 2015; Jajere, 2019; Elbediwi et al. 2020). Chromosomal colistin resistance was also reported in further *S. enterica* isolated from human and animal samples from Asia and Europe (Fig. 5 and Table S3, Supporting Information; Luo et al. 2020).

### Enterobacter spp.

*Enterobacter* spp. are part of the animal and human gut microbiome but also emerged as opportunistic human pathogens causing bacteremia, respiratory, urinary and gastrointestinal infections (Sanders and Sanders, 1997). Several studies observed that the prevalence of colistin resistance among human clinical *Enterobacteriales* is higher in *Enterobacter* spp. than *E. coli* and *Klebsiella* spp. and was 0.7% in a global surveillance program, 1.5% in Tunisia, 4.2% in Spain, 4–20% in UK, 27.2% in Taiwan for *Enterobacter* species (Fig. 2; Maalej et al. 2012; Bradford et al. 2016; Prim et al. 2017; Jean et al. 2018; Mushtaq et al. 2020). Among CRE isolates, the prevalence of colistin resistance was as high as 54.1% for *Enterobacter* (Teo et al. 2019). The most prevalent colistin-resistant species detected were *E. asburiae* and *E. cloacae* but also found in *E. aerogenes*, *E. bugandensis* and other species (Bradford et al. 2016; Mushtaq et al. 2020; ATLAS database, <https://atlas-surveillance.com/>, accessed June 2021). A total of

two major sequence types of carbapenem-resistant *E. cloacae* complex ST171 and ST78 have been described as epidemic, but chromosomal colistin resistance has been observed only in an ST78 isolate from Asia (Gomez-Simmonds et al. 2018; Teo et al. 2019). So far, there is no accumulation of chromosomal mutations in a particular sequence type and different sequence types from human clinical samples have been described from Europe and Asia (Fig. 5 and Table S3, Supporting Information; Majewski et al. 2014; Teo et al. 2019; Wand and Sutton, 2020). Worryingly, *Enterobacter* spp. exhibit the phenomenon of heteroresistance, which poses a significant problem for antimicrobial susceptibility testing in clinical settings (Hong, Lee and Ko 2018). Colistin heteroresistance can cause the 'skip well' phenomenon when using the BMD method for susceptibility testing, which is characterized by no bacterial growth at a certain antibiotic concentration, but growth takes place at higher antibiotic concentration (Landman, Salamera and Quale 2013). As a result, colistin heteroresistance may lead to treatment failure in clinical settings and may explain the high prevalence of colistin resistance among *Enterobacter* spp. (Fig. 2; Band et al. 2016). Interestingly, heteroresistance is observed more frequently in isolates belonging to a particular species or *E. cloacae* complex (Guerin et al. 2016).

### CONCLUSION

The global prevalence of colistin resistance in *Enterobacteriales* and the significance of chromosomal mutations in mediating this resistance was assessed in the present review. To our knowledge, this study represents a critical comprehensive review on colistin-resistant *Enterobacteriales*, including a comparison of

their dissemination among European countries in the veterinary and human medicine, highlighting the role of mutational changes in chromosomal encoded genes. In the last 5 years, studies on colistin resistance comprised mainly the distribution of *mcr*-genes. However, resistance mechanisms seem to be more complex than previously thought. The genetic background of the bacterial species and the presence of supporting factors might play an important role and some enterobacterial genera (for example *Enterobacter* spp.) might contribute more than others to the development of colistin-resistance in *Enterobacterales*. Therefore, the impact of chromosomal mutations and their rate of emergence should not be overlooked. Our goal, with presenting these data, is to obtain a better understanding on the molecular basis of colistin resistance, which is necessary to be able to comprehend the development and spread of resistant isolates within the animal and human community.

Overall, the molecular basis of colistin resistance in *Enterobacterales* is very complex and not yet fully understood, whereby further clarification is urgently needed due to the increasing use of colistin as last-line antimicrobial in the clinic. Those findings highlight the need for routine WGS to define whether AMR is based on transfer of resistance determinants between different strains, or even species, or caused by spread of resistant strains. The genotypic results should be combined with experimental functional studies to understand the principles of colistin resistance. More importantly, the information should be linked with epidemiological data on AMR obtained from monitoring programs of the veterinary sector and human medicine. There is an urgent need for improved monitoring programs with real-time data reporting, especially for resistance towards those antimicrobials, which are used as last-line option for the treatment of serious infections in humans.

## SUPPLEMENTARY DATA

Supplementary data are available at [FEMSRE](#) online.

## ACKNOWLEDGMENTS

We want to thank all the scientists without their dedicated work this review would not have been possible. We also would like to thank the editor and the anonymous reviewers for their critical reading, thoughtful comments and constructive suggestions. Background world maps shown in Figure 2 and 5 were provided by Esri (<https://www.arcgis.com/apps/mapviewer/index.html?w=8b3d38c0819547faa83f7b7aca80bd76>).

## FUNDING

This work was supported by the German Federal Ministry of Health [GÜCCI, grant number 60-0103-08.P105] and the German Federal Institute for Risk Assessment [grant number 1322-648].

**Conflicts of interest.** None declared.

## REFERENCES

- Abraham S, O'Dea M, Sahibzada S et al. *Escherichia coli* and *Salmonella* spp. isolated from Australian meat chickens remain susceptible to critically important antimicrobial agents. *PLoS ONE* 2019;14:e0224281.
- Adegoke AA, Madu CE, Aiyegoro OA et al. Antibigram and beta-lactamase genes among cefotaxime resistant *E. coli* from wastewater treatment plant. *Antimicrob Resist Infect Cont* 2020;9:46.
- Agero Y, Torpdahl M, Zachariassen C et al. Tentative colistin epidemiological cut-off value for *Salmonella* spp. *Foodborne Pathog Dis* 2012;9:367–9.
- Al-Zalabani A, AlThobyane OA, Alshehri AH et al. Prevalence of *Klebsiella pneumoniae* antibiotic resistance in Medina, Saudi Arabia, 2014–2018. *Cureus* 2020;12:e9714.
- Albiger B, Glasner C, Struelens MJ et al. Carbapenemase-producing *Enterobacteriaceae* in Europe: assessment by national experts from 38 countries, May 2015. *Eurosurveillance* 2015;20:17–34.
- Alvarez VE, Campos J, Galiana A et al. Genomic analysis of the first isolate of KPC-2-producing *Klebsiella pneumoniae* from Uruguay. *J Glob Antimicrob Resist* 2018;15:109–10.
- Anderson M, Cecchini M, Mossialos E. *Challenges to Tackling Antimicrobial Resistance: Economic and Policy Responses*. European Observatory on Health Systems and Policies. Cambridge: Cambridge University Press, 2020.
- Antoniadou A, Kontopidou F, Poulakou G et al. Colistin-resistant isolates of *Klebsiella pneumoniae* emerging in intensive care unit patients: first report of a multiclonal cluster. *J Antimicrob Chemother* 2007;59:786–90.
- Antunes P, Campos J, Mourao J et al. Inflow water is a major source of trout farming contamination with *Salmonella* and multidrug resistant bacteria. *Sci Total Environ* 2018;642:1163–71.
- Australian Commission on Safety and Quality in Health Care. AURA 1019: *Third Australian Report on Antimicrobial Use and Resistance in Human Health*. 2019.
- Avgoulea K, Di Pilato V, Zarkotou O et al. Characterization of extensively drug-resistant or pandrug-resistant sequence Type 147 and 101 OXA-48-producing *Klebsiella pneumoniae* causing bloodstream infections in patients in an intensive care unit. *Antimicrob Agents Chemother* 2018;62:e02457–17.
- Bagley ST. Habitat association of *Klebsiella* species. *Infect Control Hosp Epidemiol* 1985;6:52–8.
- Band VI, Crispell EK, Napier BA et al. Antibiotic failure mediated by a resistant subpopulation in *Enterobacter cloacae*. *Nat Microbiol* 2016;1:16053.
- Bardhan T, Chakraborty M, Bhattacharjee B. Prevalence of colistin-resistant, carbapenem-hydrolyzing proteobacteria in hospital water bodies and out-falls of West Bengal, India. *Int J Environ Res Pub Health* 2020;17:1007.
- Basso M, Zago D, Pozzetto I et al. Intra-hospital acquisition of colonization and infection by *Klebsiella pneumoniae* strains producing carbapenemases and carriage evolution: A longitudinal analysis in an Italian teaching hospital from January 2017 to August 2019. *Int J Infect Dis* 2020;92:81–8.
- Bean DC, Wigmore SM, Momin MHFA et al. Polymyxin resistant bacteria in Australian poultry. *Front Sustain Food Syst* 2020;4:550318.
- Blanco N, O'Hara LM, Harris AD. Transmission pathways of multidrug-resistant organisms in the hospital setting: a scoping review. *Infect Control Hosp Epidemiol* 2019;40:447–56.
- Bonura C, Giuffre M, Aleo A et al. An update of the evolving Epidemic of bla<sub>KPC</sub> carrying *Klebsiella pneumoniae* in Sicily, Italy, 2014: Emergence of multiple non-ST258 clones. *PLoS ONE* 2015;10:e0132936.
- Boszczowski I, Salomao MC, Moura ML et al. Multidrug-resistant *Klebsiella pneumoniae*: genetic diversity, mechanisms of resistance to polymyxins and clinical outcomes in a tertiary teaching hospital in Brazil. *Rev Inst Med Trop Sao Paulo* 2019;61:e29.

- Bourel AS, Poirel L, Royer G et al. Colistin resistance in Parisian inpatient faecal *Escherichia coli* as the result of two distinct evolutionary pathways. *J Antimicrob Chemother* 2019;**74**:1521–30.
- Bradford PA, Kazmierczak KM, Biedenbach DJ et al. Correlation of beta-lactamase production and colistin resistance among *Enterobacteriaceae* isolates from a global surveillance program. *Antimicrob Agents Chemother* 2016;**60**:1385–92.
- Brazil. Governmental Normative Instruction IN-45. *Diario Oficial da Uniao*. 2016. Nov 11 [cited 2021 June 22]. 2016.
- Brolund A, Lagerqvist N, Byfors S et al. Worsening epidemiological situation of carbapenemase-producing *Enterobacteriaceae* in Europe, assessment by national experts from 37 countries, July 2018. *Eurosurveillance* 2019;**24**:17–24.
- Brown JM, Dorman DC, Roy LP. Acute renal failure due to overdosage of colistin. *Med J Aust* 1970;**2**:923.
- Budel T, Kuenzli E, Campos-Madueno EI et al. On the island of Zanzibar people in the community are frequently colonized with the same MDR *Enterobacteriales* found in poultry and retailed chicken meat. *J Antimicrob Chemother* 2020;**75**:2432–41.
- Budel T, Kuenzli E, Clement M et al. Polyclonal gut colonization with extended-spectrum cephalosporin- and/or colistin-resistant *Enterobacteriaceae*: a normal status for hotel employees on the island of Zanzibar, Tanzania. *J Antimicrob Chemother* 2019;**74**:2880–90.
- Can F, Menekse S, Ispir P et al. Impact of the ST101 clone on fatality among patients with colistin-resistant *Klebsiella pneumoniae* infection. *J Antimicrob Chemother* 2018;**73**:1235–41.
- Canadian Ministry of Agriculture. *Use of Over-the-Counter Antibiotics in BC Livestock and Poultry, 2002-2018*. 2019.
- Cannatelli A, Giani T, Aiezza N et al. An allelic variant of the PmrB sensor kinase responsible for colistin resistance in an *Escherichia coli* strain of clinical origin. *Sci Rep* 2017;**7**:5071.
- Cannatelli A, Giani T, D'Andrea MM et al. MgrB inactivation is a common mechanism of colistin resistance in KPC-producing *Klebsiella pneumoniae* of clinical origin. *Antimicrob Agents Chemother* 2014;**58**:5696–703.
- Capone A, Giannella M, Fortini D et al. High rate of colistin resistance among patients with carbapenem-resistant *Klebsiella pneumoniae* infection accounts for an excess of mortality. *Clin Microbiol Infect* 2013;**19**:E23–30.
- Catel-Ferreira M, Marti S, Guillon L et al. The outer membrane porin OmpW of *Acinetobacter baumannii* is involved in iron uptake and colistin binding. *FEBS Lett* 2016;**590**:224–31.
- Catry B, Cavaleri M, Baptiste K et al. Use of colistin-containing products within the European Union and European Economic Area (EU/EEA): development of resistance in animals and possible impact on human and animal health. *Int J Antimicrob Agents* 2015;**46**:297–306.
- Chaalal N, Touati A, Yahiaoui-Martinez A et al. Colistin-resistant *Enterobacteriales* isolated from chicken meat in Western Algeria. *Microb Drug Resist* 2021;**27**:991–1002.
- Chavda KD, Chen L, Fouts DE et al. Comprehensive genome analysis of carbapenemase-producing *Enterobacter* spp.: new insights into phylogeny, population structure, and resistance mechanisms. *Mbio* 2016;**7**:e02093–16.
- Cheng YH, Lin TL, Lin YT et al. Amino acid substitutions of CrrB responsible for resistance to colistin through CrrC in *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 2016;**60**:3709–16.
- Cheng YH, Lin TL, Pan YJ et al. Colistin resistance mechanisms in *Klebsiella pneumoniae* strains from Taiwan. *Antimicrob Agents Chemother* 2015;**59**:2909–13.
- Cheong HS, Kim SY, Seo J et al. Colistin resistance and extensive genetic variations in PmrAB and PhoPQ in *Klebsiella pneumoniae* isolates from South Korea. *Curr Microbiol* 2020;**77**:2307–11.
- Choi MJ, Ko KS. Mutant prevention concentrations of colistin for *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* clinical isolates. *J Antimicrob Chemother* 2014;**69**:275–7.
- Choi Y, Lee JY, Lee H et al. Comparison of fitness cost and virulence in chromosome- and plasmid-mediated colistin-resistant *Escherichia coli*. *Front Microbiol* 2020;**11**:798.
- Clements A, Tull D, Jenney AW et al. Secondary acylation of *Klebsiella pneumoniae* lipopolysaccharide contributes to sensitivity to antibacterial peptides. *J Biol Chem* 2007;**282**:15569–77.
- Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing (CLSI)*. 30th edn. Clinical and Laboratory Standards Institute. 2020. CLSI supplement M100.
- Comandatore F, Sasser D, Ambretti S et al. Draft genome sequences of two multidrug resistant *Klebsiella pneumoniae* ST258 isolates resistant to colistin. *Genome Announc* 2013;**1**:e00113–12.
- Commonwealth of Australia. *Importance Ratings and Summary of Antibacterial Uses in Human and Animal Health in Australia*. 2018.
- Conway SP, Pond HN, Watson A et al. Intravenous colistin sulphomethate in acute respiratory exacerbations in adult patients with cystic fibrosis. *Thorax* 1997;**52**:987–93.
- Coppola N, Freire B, Umpierrez A et al. Transferable resistance to highest priority critically important antibiotics for human health in *Escherichia coli* strains obtained from livestock feces in Uruguay. *Front Vet Sci* 2020;**7**:588919.
- Dafopoulou K, Xavier BB, Zarkotou O et al. Detection of colistin resistance in the highly virulent *Escherichia coli* ST131 H30Rx clone in Greece. *J Glob Antimicrob Resist* 2020;**20**:31–2.
- Dagher TN, Al-Bayssari C, Chabou S et al. Intestinal carriage of colistin-resistant *Enterobacteriaceae* at Saint Georges Hospital in Lebanon. *J Glob Antimicrob Resist* 2020;**21**:386–90.
- Dagher TN, Azar E, Al-Bayssari C et al. First detection of colistin-resistant *Klebsiella pneumoniae* in association with NDM-5 carbapenemase isolated from clinical Lebanese patients. *Microb Drug Resist* 2019;**25**:925–30.
- Davies M, Walsh TR. A colistin crisis in India. *Lancet Infect Dis* 2018;**18**:256–7.
- de la Cadena E, Mojica MF, Garcia-Betancur JC et al. Molecular analysis of polymyxin resistance among carbapenemase-producing *Klebsiella pneumoniae* in Colombia. *Antibiotics* 2021;**10**:284.
- De Majumdar S, Yu J, Fookes M et al. Elucidation of the RamA regulon in *Klebsiella pneumoniae* reveals a role in LPS regulation. *PLoS Pathog* 2015;**11**:e1004627.
- Delannoy S, Devendec LLe, Jouy E et al. Characterization of colistin-resistant *Escherichia coli* isolated from diseased pigs in France. *Front Microbiol* 2017;**8**:2278.
- Delgado-Blas JF, Ovejero CM, Abadia-Patino L et al. Coexistence of *mcr-1* and *bla<sub>NDM-1</sub>* in *Escherichia coli* from Venezuela. *Antimicrob Agents Chemother* 2016;**60**:6356–8.
- Diaz-Jimenez D, Garcia-Menino I, Fernandez J et al. Chicken and turkey meat: consumer exposure to multidrug-resistant *Enterobacteriaceae* including *mcr*-carriers, uropathogenic *E. coli* and high-risk lineages such as ST131. *Int J Food Microbiol* 2020;**331**:108750.
- Diene SM, Merhej V, Henry M et al. The rhizome of the multidrug-resistant *Enterobacter aerogenes* genome reveals how new

- “Killer Bugs” are created because of a sympatric lifestyle. *Mol Biol Evol* 2013;**30**:369–83.
- Dixon RA, Chopra I. Leakage of periplasmic proteins from *Escherichia coli*-mediated by polymyxin-B nonapeptide. *Antimicrob Agents Chemother* 1986;**29**:781–8.
- Dominguez JE, Faccione D, Tijet N et al. Characterization of *Escherichia coli* carrying *mcr-1*-plasmids recovered from food animals from Argentina. *Front Cell Infect Microbiol* 2019;**9**:41.
- Dominguez JE, Figueroa Espinosa RA, Redondo LM et al. Plasmid-mediated colistin resistance in *Escherichia coli* recovered from healthy poultry. *Rev Argent Microbiol* 2017;**49**:297–8.
- ECDC (European Centre for Disease Prevention and Control), EFSA (European Food Safety Authority), and EMA (European Medicines Agency). ECDC/EFSA/EMA second joint report on the integrated analysis of the consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals – Joint Inter-agency Antimicrobial Consumption and Resistance Analysis (JIACRA) Report. *EFSA J* 2017;**15**:1–135.
- ECDC (European Centre for Disease Prevention and Control), EFSA (European Food Safety Authority) and EMA (European Medicines Agency). ECDC/EFSA/EMA first joint report on the integrated analysis of the consumption of antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals. Stockholm/Parma/London: ECDC/EFSA/EMA. *EFSA J*, 2015;**13**: 1–114.
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control). The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2016. *EFSA J* 2018;**16**:270.
- EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control). The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2017. *EFSA Journal* 2019;**17**:278.
- Elbediwi M, Pan H, Biswas S et al. Emerging colistin resistance in *Salmonella enterica* serovar Newport isolates from human infections *Emerg Microb Infect* 2020;**9**:535–8.
- Ellem JA, Ginn AN, Chen SCA et al. Locally acquired *mcr-1* in *Escherichia coli*, Australia, 2011 and 2013. *Emerg Infect Dis* 2017;**23**:1160–3.
- Escalante EG, Condor KY, Di Conza JA et al. Phenotypic detection of plasmid-mediated colistin resistance in *Enterobacteriaceae*. *J Clin Microbiol* 2020;**58**:e01555–19.
- Esposito EP, Cervoni M, Bernardo M et al. Molecular epidemiology and virulence profiles of colistin-resistant *Klebsiella pneumoniae* blood isolates from the hospital agency “Ospedale dei Colli,” Naples, Italy. *Front Microbiol* 2018;**9**:1463.
- European Centre for Disease Prevention and Control (ECDC). *Antimicrobial resistance in the EU/EEA (EARS-Net) - Annual Epidemiological Report 2019*. Stockholm: ECDC, 2020.
- European Centre for Disease Prevention and Control (ECDC). *Antimicrobial resistance in the EU/EEA (EARS-Net) -Annual Epidemiological Report 2019*, 2020.
- European Centre for Disease Prevention and Control (ECDC). *Surveillance of antimicrobial resistance in Europe 2018: Annual report of the European Antimicrobial Resistance Surveillance Network (EARS-Net)*. Stockholm: ECDC. 2019.
- European Commission. *A European One Health Action Plan against Antimicrobial Resistance (AMR)*. 2017.
- European Commission. *Ban On Antibiotics As Growth Promoters In Animal Feed Enters Into Effect*. 2005 [cited 2020 8th December]; Available from: [https://ec.europa.eu/commission/presscorner/detail/en/IP\\_05\\_1687](https://ec.europa.eu/commission/presscorner/detail/en/IP_05_1687).
- European Commission. Commission implementing decision 2013/652/EU of 12 November 2013 on the monitoring and reporting of antimicrobial resistance in zoonotic and commensal bacteria. *Off J Eur Union* 2013;**303**:26.
- European Committee on Antimicrobial Susceptibility Testing (EUCAST). *Breakpoint Tables for Interpretation of MICs and Zone Diameters, Version 9.0*. 2019. 2014.
- European Food Safety Authority (EFSA) and European Centre for Disease Prevention and Control (ECDC). The European Union One Health 2018 zoonoses report. *EFSA J* 2019;**17**:276.
- European Medicines Agency (EMA), European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 26 EU/EEA countries in 2012'. Fourth ESVAC report. (EMA/333921/2014). 2014, 1–115.
- European Medicines Agency (EMA), European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 31 European countries in 2018'. Tenth ESVAC report (EMA/24309/2020). 2020, 104.
- European Medicines Agency (EMA). Categorisation of antibiotic-sin the European Union. 2019.
- European Medicines Agency (EMA). Sales of veterinary antimicrobial agents in 19 EU/EEA countries in 2010. 2012;(EMA/88728/2012):74.
- European Medicines Agency (EMA). Trends in the sales of veterinary antimicrobial agents in nine European countries (2005–2009). 2011;(EMA/238630/2011):77.
- European Medicines Agency (EMA). Updated advice on the use of colistin products in animals within the European Union: development of resistance and possible impact on human and animal health. 2016:56.
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 25 EU/EEA countries in 2011. 2013;(EMA/236501/2013):98.
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 26 EU/EEA countries in 2013. 2015;(EMA/387934/2015):162.
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 29 European countries in 2014. 2016;(EMA/61769/2016):176.
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 30 European countries in 2015. 2017;(EMA/184855/2017):184.
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 30 European countries in 2016. 2018;(EMA/275982/2018):184.
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 31 European countries in 2017. 2019;(EMA/294674/2019):109.
- European Medicines Agency, European Surveillance of Veterinary Antimicrobial Consumption. Sales of veterinary antimicrobial agents in 31 European countries in 2018. 2020;(EMA/24309/2020):104.

- Falagas ME, Kasiakou SK. Colistin: the revival of polymyxins for the management of multidrug-resistant Gram-negative bacterial infections. *Clin Infect Dis* 2005;**40**:1333–41.
- Food and Agriculture Organization of the United Nations (FAO). *Meat market review*. 2020.
- Galani I, Karaiskos I, Karantani I et al. Epidemiology and resistance phenotypes of carbapenemase-producing *Klebsiella pneumoniae* in Greece, 2014 to 2016. *Eurosurveillance* 2018;**23**:2–13.
- Garcia-Menino I, Diaz-Jimenez D, Garcia V et al. Genomic characterization of prevalent *mcr-1*, *mcr-4*, and *mcr-5* *Escherichia coli* within swine enteric colibacillosis in Spain. *Front Microbiol* 2019;**10**:2469.
- Garcia-Menino I, Lumbreras P, Valledor P et al. Workflow for an accurate detection of colistin-resistant and *mcr*-positive *E. coli*. *Antibiotics-Basel* 2020;**9**:861.
- Garza-Ramos U, Tamayo-Legorreta E, Arellano-Quintanilla DM et al. Draft genome sequence of a multidrug- and colistin-resistant *mcr-1*-producing *Escherichia coli* isolate from a swine farm in Mexico. *Microbiol Resour Announcements* 2018;**6**:e00102–18.
- Gentile B, Grottola A, Orlando G et al. A retrospective whole-genome sequencing analysis of carbapenem and colistin-resistant *Klebsiella pneumoniae* nosocomial strains isolated during an MDR surveillance program. *Antibiotics-Basel* 2020;**9**:246.
- Ghafur A, Shankar C, GnanaSoundari P et al. Detection of chromosomal and plasmid-mediated mechanisms of colistin resistance in *Escherichia coli* and *Klebsiella pneumoniae* from Indian food samples. *J Glob Antimicrob Resist* 2019;**16**:48–52.
- Giani T, Arena F, Vaggelli G et al. Large nosocomial outbreak of colistin-resistant, carbapenemase-producing *Klebsiella pneumoniae* traced to clonal expansion of an *mgrB* deletion mutant. *J Clin Microbiol* 2015;**53**:3341–4.
- Gibson JS, Wai H, Oo SSML et al. Antimicrobials use and resistance on integrated poultry-fish farming systems in the Ayeyarwady Delta of Myanmar. *Sci Rep* 2020;**10**:16149.
- Giordano C, Klak A, Barnini S et al. Reduced Fitness Costs of *mcr-1.2* Compared to Mutated *pmrB* in Isogenic Colistin-Resistant KPC-3-Producing *Klebsiella pneumoniae*. *mSphere* 2019;**4**:e00551–19.
- Glasner C, Albiger B, Buist G et al. Carbapenemase-producing *Enterobacteriaceae* in Europe: a survey among national experts from 39 countries, February 2013. *Eurosurveillance* 2013;**18**:9–15.
- Goel G, Hmar L, De MS et al. Colistin-resistant *Klebsiella pneumoniae*: report of a cluster of 24 cases from a new oncology center in eastern India. *Infect Control Hosp Epidemiol* 2014;**35**:1076–7.
- Gomez-Simmonds A, Annavajhala MK, Wang Z et al. Genomic and geographic context for the evolution of high-risk carbapenem-resistant *Enterobacter cloacae* complex clones ST171 and ST78. *Mbio* 2018;**9**:e00542–18.
- Grundmann H, Glasner C, Albiger B et al. Occurrence of carbapenemase-producing *Klebsiella pneumoniae* and *Escherichia coli* in the European survey of carbapenemase-producing *Enterobacteriaceae* (EuSCAPE): a prospective, multinational study. *Lancet Infect Dis* 2017;**17**:153–63.
- Grundmann H, Livermore DM, Giske CG et al. Carbapenem-nonsusceptible *Enterobacteriaceae* in Europe: conclusions from a meeting of national experts. *Eurosurveillance* 2010;**15**:22–34.
- Guducuoglu H, Gursoy NC, Yakupogullari Y et al. Hospital outbreak of a colistin-resistant, NDM-1-and OXA-48-producing *Klebsiella pneumoniae*: high mortality from pandrug resistance. *Microb Drug Resist* 2018;**24**:966–72.
- Guentzel MN. *Escherichia, Klebsiella, Enterobacter, Serratia, Citrobacter, and Proteus*. In: Baron S. (ed). *Medical Microbiology*. Galveston, TX: The University of Texas Medical Branch at Galveston, 1996.
- Guerin F, Isnard C, Sinel C et al. Cluster-dependent colistin hetero-resistance in *Enterobacter cloacae* complex. *J Antimicrob Chemother* 2016;**71**:3058–3061.
- Gutierrez C, Zenis J, Legarraga P et al. Genetic analysis of the first *mcr-1* positive *Escherichia coli* isolate collected from an outpatient in Chile. *Braz J Infect Dis* 2019;**23**:203–6.
- Haeili M, Javani A, Moradi J et al. MgrB alterations mediate colistin resistance in *Klebsiella pneumoniae* isolates from Iran. *Front Microbiol* 2017;**8**:2470.
- Halaby T, Kucukkose E, Janssen AB et al. Genomic characterization of colistin heteroresistance in *Klebsiella pneumoniae* during a nosocomial outbreak. *Antimicrob Agents Chemother* 2016;**60**:6837–43.
- Hallenberg GS, Borjesson S, Sokerya S et al. Detection of *mcr*-mediated colistin resistance in *Escherichia coli* isolates from pigs in small-scale farms in Cambodia. *Antimicrob Agents Chemother* 2019;**63**:e02241–18.
- Haller S, Kramer R, Becker K et al. Extensively drug-resistant *Klebsiella pneumoniae* ST307 outbreak, north-eastern Germany, June to October 2019. *Eurosurveillance* 2019;**24**:2–7.
- Hamel M, Chatzipanagiotou S, Hadjadj L et al. Inactivation of *mgrB* gene regulator and resistance to colistin is becoming endemic in carbapenem-resistant *Klebsiella pneumoniae* in Greece: a nationwide study from 2014 to 2017. *Int J Antimicrob Agents* 2020;**55**:105930.
- Hassen B, Abbassi MS, Benlabidi S et al. Genetic characterization of ESBL-producing *Escherichia coli* and *Klebsiella pneumoniae* isolated from wastewater and river water in Tunisia: predominance of CTX-M-15 and high genetic diversity. *Environ Sci Pollut Res Int* 2020;**27**:44368–77.
- Hernandez-Castro R, Rodriguez-Santiago J, Tellez-Sosa J et al. Molecular and genome characterization of colistin-resistant *Escherichia coli* isolates from wild sea lions (*Zalophus californianus*). *Braz J Microbiol* 2020;**51**:2009–14.
- Hjort K, Nicoloff H, Andersson DI. Unstable tandem gene amplification generates heteroresistance (variation in resistance within a population) to colistin in *Salmonella enterica*. *Mol Microbiol* 2016;**102**:274–89.
- Homeier-Bachmann T, Heiden SE, Lubcke PK et al. Antibiotic-resistant *Enterobacteriaceae* in wastewater of abattoirs. *Antibiotics-Basel* 2021;**10**:568.
- Hong YK, Ko KS. PmrAB and PhoPQ variants in colistin-resistant *Enterobacter* spp. isolates in Korea. *Curr Microbiol* 2019;**76**:644–9.
- Hong YK, Lee JY, Ko KS. Colistin resistance in *Enterobacter* spp. isolates in Korea. *J Microbiol* 2018;**56**:435–40.
- Hsing WH, Russo FD, Bernd KK et al. Mutations that alter the kinase and phosphatase activities of the two-component sensor EnvZ. *J Bacteriol* 1998;**180**:4538–46.
- Jafari Z, Harati AA, Haeili M et al. Molecular epidemiology and drug resistance pattern of carbapenem-resistant *Klebsiella pneumoniae* isolates from Iran. *Microb Drug Resist* 2019;**25**:336–43.
- Jaidane N, Bonnin RA, Mansour W et al. Genomic insights into colistin-resistant *Klebsiella pneumoniae* from a Tunisian teaching hospital. *Antimicrob Agents Chemother* 2018;**62**:e01601–17.



- Jajere SM. A review of *Salmonella enterica* with particular focus on the pathogenicity and virulence factors, host specificity and antimicrobial resistance including multidrug resistance. *Vet World* 2019;12:504–21.
- Janssen AB, Bartholomew TL, Marciszewska NP et al. Non-clonal emergence of colistin resistance associated with mutations in the BasRS two-component system in *Escherichia coli* bloodstream isolates. *Msphere* 2020;5. DOI: 10.1128/mSphere.00143-20.
- Janssen AB, van Hout D, Bonten MJM et al. Microevolution of acquired colistin resistance in Enterobacteriaceae from ICU patients receiving selective decontamination of the digestive tract. *J Antimicrob Chemother* 2020;75: 3135–43.
- Jayol A, Nordmann P, Brink A et al. Heteroresistance to colistin in *Klebsiella pneumoniae* associated with alterations in the PhoPQ regulatory system. *Antimicrob Agents Chemother* 2015;59:2780–4.
- Jayol A, Nordmann P, Brink A et al. High-level resistance to colistin mediated by various mutations in the *ctrB* gene among carbapenemase-producing *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* 2017;61:e01423–17.
- Jayol A, Nordmann P, Lehours P et al. Comparison of methods for detection of plasmid-mediated and chromosomally encoded colistin resistance in Enterobacteriaceae. *Clin Microbiol Infect* 2018;24:175–9.
- Jayol A, Poirel L, Brink A et al. Resistance to colistin associated with a single amino acid change in protein PmrB among *Klebsiella pneumoniae* isolates of worldwide origin. *Antimicrob Agents Chemother* 2014;58:4762–6.
- Jayol A, Poirel L, Dortet L et al. National survey of colistin resistance among carbapenemase-producing Enterobacteriaceae and outbreak caused by colistin-resistant OXA-48-producing *Klebsiella pneumoniae*, France, 2014. *Eurosurveillance* 2016;21:18–25.
- Jean SS, Lee WS, Yu KW et al. Rates of susceptibility of carbapenems, ceftobiprole, and colistin against clinically important bacteria collected from intensive care units in 2007: results from the Surveillance of Multicenter Antimicrobial Resistance in Taiwan (SMART). *J Microbiol Immunol Infect* 2018;51:423–4.
- Joosten P, Ceccarelli D, Odent E et al. Antimicrobial usage and resistance in companion animals: a cross-sectional study in three European countries. *Antibiotics* 2020;9:87.
- Jovcic B, Novovic K, Filipic B et al. Genomic characteristics of colistin-resistant *Salmonella enterica* subsp. *enterica* Serovar infantis from poultry farms in the Republic of Serbia. *Antibiotics* 2020;9:886.
- Kathayat D, Antony L, Deblais L et al. Small molecule adjuvants potentiate colistin activity and attenuate resistance development in *Escherichia coli* by affecting *pmrAB* system. *Infect Drug Resist* 2020;13:2205–22.
- Kato A, Groisman EA. Connecting two-component regulatory systems by a protein that protects a response regulator from dephosphorylation by its cognate sensor. *Genes Dev* 2004;18:2302–13.
- Kato A, Latifi T, Groisman EA. Closing the loop: the PmrA/PmrB two-component system negatively controls expression of its posttranscriptional activator PmrD. *Proc Natl Acad Sci* 2003;100:4706–11.
- Kawahara R, Fujiya Y, Yamaguchi T et al. Most domestic livestock possess colistin-resistant commensal *Escherichia coli* harboring *mcr* in a rural community in Vietnam. *Antimicrob Agents Chemother* 2019;63:e00594–19.
- Khan SA, Everest P, Servos S et al. A lethal role for lipid A in *Salmonella* infections. *Mol Microbiol* 1998;29:571–9.
- Kim S, Woo JH, Kim N et al. Characterization of chromosome-mediated colistin resistance in *Escherichia coli* isolates from livestock in Korea. *Infect Drug Resist* 2019;12:3291–9.
- Koch-Weser J, Sidel VW, Federman EB et al. Adverse effects of sodium colistimethate. Manifestations and specific reaction rates during 317 courses of therapy. *Ann Intern Med* 1970;72:857–68.
- Kocsis B, Kadar B, Toth A et al. MgrB variants in colistin-susceptible and colistin-resistant *Klebsiella pneumoniae* ST258. *J Microbiol Immunol Infect* 2017;50:735–6.
- Kopotsa K, Mbelle NM, Sekyere JO. Epigenomics, genomics, resistome, mobilome, virulome and evolutionary phylogenomics of carbapenem-resistant *Klebsiella pneumoniae* clinical strains. *Microb Genomics* 2020;6:mgen000474.
- Koppe U, von Laer A, Kroll LE et al. Carbapenem non-susceptibility of *Klebsiella pneumoniae* isolates in hospitals from 2011 to 2016, data from the German Antimicrobial Resistance Surveillance (ARS). *Antimicrob Resist Infect Control* 2018;7:71.
- Kox LFF, Wosten MMSM, Groisman EA. A small protein that mediates the activation of a two-component system by another two-component system. *EMBO J* 2000;19:1861–72.
- Kuang Q, He D, Sun H et al. R<sup>93</sup>P Substitution in the PmrB HAMP Domain Contributes to Colistin Heteroresistance in *Escherichia coli* Isolates from Swine. *Antimicrob Agents Chemother* 2020;64:e01509–20.
- Lagace-Wiens PRS, Adam HJ, Poutanen S et al. Trends in antimicrobial resistance over 10 years among key bacterial pathogens from Canadian hospitals: results of the CANWARD study 2007–16. *J Antimicrob Chemother* 2019;74:22–31.
- Landman D, Salamera J, Quale J. Irreproducible and uninterpretable polymyxin B MICs for *Enterobacter cloacae* and *Enterobacter aerogenes*. *J Clin Microbiol* 2013;51:4106–11.
- Laxminarayan R, Matsoso P, Pant S et al. Access to effective antimicrobials: a worldwide challenge. *Lancet North Am Ed* 2016;387:168–75.
- Leavitt A, Navon-Venezia S, Chmelnitsky I et al. Emergence of KPC-2 and KPC-3 in carbapenem-resistant *Klebsiella pneumoniae* strains in an Israeli hospital. *Antimicrob Agents Chemother* 2007;51:3026–9.
- Ledson MJ, Gallagher MJ, Cowperthwaite C et al. Four years' experience of intravenous colomycin in an adult cystic fibrosis unit. *Eur Respir J* 1998;12:592–4.
- Levy SB, Fitzgerald GB, Macone B. Changes in intestinal flora of farm personnel after introduction of a tetracycline-supplemented feed on a farm. *N Engl J Med* 1976;295:583–8.
- Li J, Nation RL, Kaye KS. Polymyxin antibiotics: from laboratory bench to bedside preface. *Adv Exp Med Biol*, 2019;1145:V–Vi.
- Lippa AM, Goulian M. Feedback inhibition in the PhoQ/PhoP signaling system by a membrane peptide. *PLoS Genet* 2009;5:e1000788.
- Liu YY, Wang Y, Walsh TR et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis* 2016;16:161–8.
- Llobet E, Tomas JM, Bengoechea JA. Capsule polysaccharide is a bacterial decoy for antimicrobial peptides. *Microbiology* 2008;154:3877–86.
- Loayza-Villa F, Salinas L, Tijet N et al. Diverse *Escherichia coli* lineages from domestic animals carrying colistin resistance gene *mcr*-1 in an Ecuadorian household. *J Glob Antimicrob Resist* 2020;22:63–7.

- Luo QX, Wang Y, Fu H et al. Serotype is associated with high rate of colistin resistance among clinical isolates of *Salmonella*. *Front Microbiol* 2020;11:592146.
- Luo QX, Yu W, Zhou K et al. Molecular epidemiology and colistin resistant mechanism of *mcr*-positive and *mcr*-negative clinical isolated *Escherichia coli*. *Front Microbiol* 2017;8:2262.
- Maalej SM, Meziou MR, Mahjoubi F et al. Epidemiological study of *Enterobacteriaceae* resistance to colistin in Sfax (Tunisia). *Med Mal Infect* 2012;42:256–63.
- Macesic N, Nelson B, Mcconville TH et al. Emergence of polymyxin resistance in clinical *Klebsiella pneumoniae* through diverse genetic adaptations: a genomic, retrospective cohort study. *Clin Infect Dis* 2020;70:2084–91.
- Magiorakos AP, Srinivasan A, Carey RB et al., Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012;18:268–81.
- Mahalakshmi S, Sunayana MR, SaiSree L et al. *yciM* is an essential gene required for regulation of lipopolysaccharide synthesis in *Escherichia coli*. *Mol Microbiol* 2014;91:145–57.
- Majewski P, Wiczorek P, Sacha PT et al. Emergence of OXA-48 carbapenemase-producing *Enterobacter cloacae* ST89 infection in Poland. *Int J Infect Dis* 2014;25:107–9.
- Malchione MD, Torres LM, Hartley DM et al. Carbapenem and colistin resistance in *Enterobacteriaceae* in Southeast Asia: review and mapping of emerging and overlapping challenges. *Int J Antimicrob Agents* 2019;54:381–99.
- Mammaia C, Bonura C, Di Bernardo F et al. Ongoing spread of colistin-resistant *Klebsiella pneumoniae* in different wards of an acute general hospital, Italy, June to December 2011. *Eurosurveillance* 2012;17:9–14.
- Marin C, Lorenzo-Rebenaque L, Laso O et al. Pet reptiles: a potential source of transmission of multidrug-resistant *Salmonella*. *Front Vet Sci* 2021;7:613718.
- Marshall BM, Levy SB. Food animals and antimicrobials: impacts on human health. *Clin Microbiol Rev* 2011;24:718.
- Martin RM, Bachman MA. Colonization, infection, and the accessory genome of *Klebsiella pneumoniae*. *Front Cell Infect Microbiol*, 2018;8:4.
- Massella E, Reid CJ, Cummins ML et al. Snapshot study of whole genome sequences of *Escherichia coli* from healthy companion animals, livestock, wildlife, humans and food in Italy. *Antibiotics* 2020;9:782.
- Mathur P, Veerarahavan B, Devanga Ragupathi NK et al. Multiple mutations in lipid-A modification pathway & novel *fosA* variants in colistin-resistant *Klebsiella pneumoniae* *Fut Sci OA* 2018;4:FSO319.
- McConville TH, Annavajhala MK, Giddins MJ et al. *CrrB* positively regulates high-level polymyxin resistance and virulence in *Klebsiella pneumoniae*. *Cell Rep* 2020;33:108313.
- Meinersmann RJ, Ladely SR, Plumlee JR et al. Prevalence of *mcr-1* in the cecal contents of food animals in the United States. *Antimicrob Agents Chemother* 2017;61:e02244–16.
- Meletis G, Oustas E, Botziori C et al. Containment of carbapenem resistance rates of *Klebsiella pneumoniae* and *Acinetobacter baumannii* in a Greek hospital with a concomitant increase in colistin, gentamicin and tigecycline resistance. *New Microbiol* 2015;38:417–21.
- Meletis G, Tzampaz E, Sianou E et al. Colistin heteroresistance in carbapenemase-producing *Klebsiella pneumoniae*. *J Antimicrob Chemother* 2011;66:946–7.
- Merida-Vieyra J, De Colsa-Ranero A, Arzate-Barbosa P et al. First clinical isolate of *Escherichia coli* harboring *mcr-1* gene in Mexico. *PLoS ONE* 2019;14:e0214648.
- Mezzatesta ML, Gona F, Caio C et al. Outbreak of KPC-3-producing, and colistin-resistant, *Klebsiella pneumoniae* infections in two Sicilian hospitals. *Clin Microbiol Infect* 2011;17:1444–7.
- Min B, Allen-Scott LK, Buntain B. Transdisciplinary research for complex One Health issues: a scoping review of key concepts. *Prev Vet Med* 2013;112:222–9.
- Mitrophanov AY, Jewett MW, Hadley TJ et al. Evolution and dynamics of regulatory architectures controlling polymyxin B resistance in enteric bacteria. *PLoS Genet* 2008;4:e1000233.
- Moffatt JH, Harper M, Boyce JD. Mechanisms of polymyxin resistance. In: *Polymyxin Antibiotics: From Laboratory Bench to Bedside*, Vol. 1145, Springer, 2019,55–71.
- Monaco M, Giani T, Raffone M et al. Colistin resistance superimposed to endemic carbapenem-resistant *Klebsiella pneumoniae*: a rapidly evolving problem in Italy, November 2013 to April 2014 *Eurosurveillance* 2014;19:14–8.
- Moon K, Gottesman S. A PhoQ/P-regulated small RNA regulates sensitivity of *Escherichia coli* to antimicrobial peptides. *Mol Microbiol* 2009;74:1314–30.
- Morales AS, de Araujo JF, Gomes VTD et al. Colistin resistance in *Escherichia coli* and *Salmonella enterica* strains isolated from swine in Brazil. *Sci World J* 2012;2012:109795.
- Moubareck CA, Mouftah SF, Pal T et al. Clonal emergence of *Klebsiella pneumoniae* ST14 co-producing OXA-48-type and NDM carbapenemases with high rate of colistin resistance in Dubai, United Arab Emirates. *Int J Antimicrob Agents* 2018;52:90–5.
- Mushtaq S, Reynolds R, Gilmore MC et al. Inherent colistin resistance in genogroups of the *Enterobacter cloacae* complex: epidemiological, genetic and biochemical analysis from the BSAC resistance surveillance programme. *J Antimicrob Chemother* 2020;75:2452–61.
- Muyembe T, Vandepitte J, Desmyter J. Natural colistin resistance in *Edwardsiella tarda*. *Antimicrob Agents Chemother* 1973;4:521–4.
- Naha S, Sands K, Mukherjee S et al. KPC-2-producing *Klebsiella pneumoniae* ST147 in a neonatal unit: clonal isolates with differences in colistin susceptibility attributed to AcrAB-TolC pump. *Int J Antimicrob Agents* 2020;55:105903.
- National Antimicrobial Resistance Monitoring System (NARMS). NARMS Update: Integrated Report Summary. 2018. <https://www.fda.gov/animal-veterinary/national-antimicrobial-resistance-monitoring-system/2018-narms-update-in-integrated-report-summary-interactive-version>
- Navon-Venezia S, Kondratyeva K, Carattoli A. *Klebsiella pneumoniae*: a major worldwide source and shuttle for antibiotic resistance. *FEMS Microbiol Rev* 2017;41:252–75.
- Nicolaes V, El Hajjaji H, Davis RM et al. Insights into the function of *YciM*, a heat shock membrane protein required to maintain envelope integrity in *Escherichia coli*. *J Bacteriol* 2014;196:300–9.
- Nordmann P, Jayol A, Poirel L. Rapid detection of polymyxin resistance in *Enterobacteriaceae*. *Emerg Infect Dis* 2016;22:1038–43.
- Okdah L, Leangapichart T, Hadjadj L et al. First report of colistin-resistant *Klebsiella pneumoniae* clinical isolates in Lebanon. *J Glob Antimicrob Resist* 2017;9:15–6.
- Olaitan AO, Dandachi I, Baron SA et al. Banning colistin in feed additives: a small step in the right direction. *Lancet Infect Dis* 2021;21:29–30.

- Olaitan AO, Dia NM, Gautret P et al. Acquisition of extended-spectrum cephalosporin- and colistin-resistant *Salmonella enterica* subsp *enterica* serotype Newport by pilgrims during Hajj. *Int J Antimicrob Agents* 2015;**45**:600–4.
- Olaitan AO, Diene SM, Kempf M et al. Worldwide emergence of colistin resistance in *Klebsiella pneumoniae* from healthy humans and patients in Lao PDR, Thailand, Israel, Nigeria and France owing to inactivation of the PhoP/PhoQ regulator *mgrB*: an epidemiological and molecular study. *Int J Antimicrob Agents* 2014;**44**:500–7.
- Olaitan AO, Thongmalayvong B, Akkhavong K et al. Clonal transmission of a colistin-resistant *Escherichia coli* from a domesticated pig to a human in Laos. *J Antimicrob Chemother* 2015;**70**:3402–4.
- Oliveira J, Reygaert WC. Gram negative bacteria, In: *StatPearls*. Treasure Island, FL: StatPearls Publishing. Copyright © 2021, StatPearls Publishing LLC. 2021.
- Olowo-okere A, Yacouba A. Molecular mechanisms of colistin resistance in Africa: a systematic review of literature. *Germes* 2020;**10**:367–79.
- Ortega-Paredes D, Barba P, Zurita J. Colistin-resistant *Escherichia coli* clinical isolate harbouring the *mcr-1* gene in Ecuador. *Epidemiol Infect* 2016;**144**:2967–70.
- Oteo J, Perez-Vazquez M, Bautista V et al. The spread of KPC-producing Enterobacteriaceae in Spain: WGS analysis of the emerging high-risk clones of *Klebsiella pneumoniae* ST11/KPC-2, ST101/KPC-2 and ST512/KPC-3. *J Antimicrob Chemother* 2016;**71**:3392–9.
- Otter JA, Doumith M, Davies F et al. Emergence and clonal spread of colistin resistance due to multiple mutational mechanisms in carbapenemase-producing *Klebsiella pneumoniae* in London. *Sci Rep* 2017;**7**:12711.
- Padilla E, Llobet E, Domenech-Sanchez A et al. *Klebsiella pneumoniae* AcrAB efflux pump contributes to antimicrobial resistance and virulence. *Antimicrob Agents Chemother* 2010;**54**:177–83.
- Papa-Ezdra R, Grill Diaz F, Vieytes M et al. First three *Escherichia coli* isolates harbouring *mcr-1* in Uruguay. *J Glob Antimicrob Resist* 2020;**20**:187–90.
- Pardon B, Catry B, Dewulf J et al. Prospective study on quantitative and qualitative antimicrobial and anti-inflammatory drug use in white veal calves. *J Antimicrob Chemother* 2012;**67**:1027–38.
- Parisi SG, Bartolini A, Santacatterina E et al. Prevalence of *Klebsiella pneumoniae* strains producing carbapenemases and increase of resistance to colistin in an Italian teaching hospital from January 2012 To December 2014. *BMC Infect Dis* 2015;**15**:1–10.
- Park H, Inouye M. Mutational analysis of the linker region of EnvZ, an osmosensor in *Escherichia coli*. *J Bacteriol* 1997;**179**:4382–90.
- Peleg AY, Hooper DC. CURRENT CONCEPTS hospital-acquired infections due to Gram-negative bacteria. *N Engl J Med* 2010;**362**:1804–13.
- Pena I, Picazo JJ, Rodriguez-Avial C et al. Carbapenemase-producing Enterobacteriaceae in a tertiary hospital in Madrid, Spain: high percentage of colistin resistance among VIM-1-producing *Klebsiella pneumoniae* ST11 isolates. *Int J Antimicrob Agents* 2014;**43**:460–4.
- Petrosillo N, Giannella M, Lewis R et al. Treatment of carbapenem-resistant *Klebsiella pneumoniae*: the state of the art. *Expert Rev Anti Infect Ther* 2013;**11**:159–77.
- Petrosillo N, Taglietti F, Granata G. Treatment options for colistin resistant *Klebsiella pneumoniae*: present and future. *J Clin Med* 2019;**8**:934.
- Pishnian Z, Haeili M, Feizi A. Prevalence and molecular determinants of colistin resistance among commensal Enterobacteriaceae isolated from poultry in northwest of Iran. *Gut Pathog* 2019;**11**:2.
- Pogue JM, Jones RN, Bradley JS et al. Polymyxin susceptibility testing and interpretive breakpoints: recommendations from the United States Committee on Antimicrobial Susceptibility Testing (USCAST). *Antimicrob Agents Chemother* 2020;**64**:e01495–19.
- Pogue JM, Marchaim D, Kaye D et al. Revisiting “Older” Antimicrobials in the era of multidrug resistance. *Pharmacotherapy* 2011;**31**:912–21.
- Poirel L, Jayol A, Nordmann P. Polymyxins: Antibacterial Activity, Susceptibility Testing, and Resistance Mechanisms Encoded by Plasmids or Chromosomes. *Clin Microbiol Rev* 2017;**30**:557–96.
- Prim N, Turbau M, Rivera A et al. Prevalence of colistin resistance in clinical isolates of Enterobacteriaceae: a four-year cross-sectional study. *J Infect* 2017;**75**:493–8.
- Public Health Agency of Canada. *Canadian antimicrobial resistance surveillance system*. 2018.
- Quesada A, Porrero MC, Tellez S et al. Polymorphism of genes encoding PmrAB in colistin-resistant strains of *Escherichia coli* and *Salmonella enterica* isolated from poultry and swine. *J Antimicrob Chemother* 2015;**70**:71–4.
- Raetz CRH, Reynolds CM, Trent MS et al. Lipid a modification systems in Gram-negative bacteria. *Annu Rev Biochem* 2007;**76**:295–329.
- Rahmatallah N, El Rhaffouli H, Amine IL et al. Consumption of antibacterial molecules in broiler production in Morocco. *Vet Med Sci* 2018;**4**:80–90.
- Rebelo AR, Bortolaia V, Kjeldgaard JS et al. Multiplex PCR for detection of plasmid-mediated colistin resistance determinants, *mcr-1*, *mcr-2*, *mcr-3*, *mcr-4* and *mcr-5* for surveillance purposes. *Eurosurveillance* 2018;**23**:29–39.
- Rocha IV, Silva ND, Andrade CAD et al. Diverse and emerging molecular mechanisms award polymyxins resistance to Enterobacteriaceae clinical isolates from a tertiary hospital of Recife. *Infect Genet Evol* 2020;**85**:104584.
- Rossi F, Girardello R, Cury AP et al. Emergence of colistin resistance in the largest university hospital complex of Sao Paulo, Brazil, over five years. *Braz J Infect Dis* 2017;**21**:98–101.
- Rubin EJ, Herrera CM, Crofts AA et al. PmrD is required for modifications to *Escherichia coli* endotoxin that promote antimicrobial resistance. *Antimicrob Agents Chemother* 2015;**59**:2051–61.
- Rumi MV, Mas J, Elena A et al. Co-occurrence of clinically relevant beta-lactamases and MCR-1 encoding genes in *Escherichia coli* from companion animals in Argentina. *Vet Microbiol* 2019;**230**:228–34.
- Ryan KJ, Schainuck LI, Hickman RO et al. Colistimethate toxicity. Report of a fatal case in a previously healthy child. *JAMA* 1969;**207**:2099–101.
- Saavedra SY, Diaz L, Wiesner M et al. Genomic and molecular characterization of clinical isolates of Enterobacteriaceae harboring *mcr-1* in Colombia, 2002 to 2016. *Antimicrob Agents Chemother* 2017;**61**:e00841–17.
- Saini V, McClure JT, Leger D et al. Antimicrobial use on Canadian dairy farms. *J Dairy Sci* 2012;**95**:1209–21.
- Sampaio JL, Gales AC. Antimicrobial resistance in Enterobacteriaceae in Brazil: focus on beta-lactams and polymyxins. *Braz J Microbiol* 2016;**47**:31–7.

- Samra Z, Ofir O, Lishtzinsky Y et al. Outbreak of carbapenem-resistant *Klebsiella pneumoniae* producing KPC-3 in a tertiary medical centre in Israel. *Int J Antimicrob Agents* 2007;**30**:525–9.
- Samuelsen O, Overballe-Petersen S, Bjornholt JV et al. Molecular and epidemiological characterization of carbapenemase-producing *Enterobacteriaceae* in Norway, 2007 to 2014. *PLoS ONE* 2017;**12**:e0187832.
- Sanders WE, Sanders CC. *Enterobacter* spp.: pathogens poised to flourish at the turn of the century. *Clin Microbiol Rev* 1997;**10**:220.
- Satlin MJ, Jenkins SG, Walsh TJ. The global challenge of carbapenem-resistant *Enterobacteriaceae* in transplant recipients and patients with hematologic malignancies. *Clin Infect Dis* 2014;**58**:1274–83.
- Satlin MJ, Walsh TJ. Multidrug-resistant *Enterobacteriaceae*, *Pseudomonas aeruginosa*, and vancomycin-resistant *Enterococcus*: three major threats to hematopoietic stem cell transplant recipients. *Transplant Infect Dis* 2017;**19**:10.
- Sato T, Fukuda A, Suzuki Y et al. Pathogenic lineage of *mcr*-negative colistin-resistant *Escherichia coli*, Japan, 2008–2015. *Emerg Infect Dis* 2016;**22**:2223–5.
- Sato T, Shiraiishi T, Hiyama Y et al. Contribution of novel amino acid alterations in *PmrA* or *PmrB* to colistin resistance in *mcr*-negative *Escherichia coli* clinical isolates, including major multidrug-resistant lineages O25b:H4-ST131-H30Rx and Non-x. *Antimicrob Agents Chemother* 2018;**62**:e00864–18.
- Savin M, Bierbaum G, Hammerl JA et al. ESKAPE bacteria and extended-spectrum-beta-lactamase-producing *Escherichia coli* isolated from wastewater and process water from German poultry slaughterhouses. *Appl Environ Microbiol* 2020;**86**:e02748–19.
- Schmidt JS, Kuster SP, Nigg A et al. Poor infection prevention and control standards are associated with environmental contamination with carbapenemase-producing *Enterobacterales* and other multidrug-resistant bacteria in Swiss companion animal clinics. *Antimicrob Resist Infect Control* 2020;**9**:93.
- Schnaitman CA. Effect of Ethylenediaminetetraacetic Acid, Triton X-100, and Lysozyme on Morphology and Chemical Composition of Isolated Cell Walls of *Escherichia-Coli* *J Bacteriol* 1971;**108**:553.
- Seo KW, Kim JJ, Mo IP et al. Molecular characteristic of antimicrobial resistance of *Salmonella gallinarum* isolates from chickens in Korea, 2014 to 2018. *Poult Sci* 2019;**98**:5416–23.
- Shankar C, Jacob JJ, Sugumar SG et al. Distinctive mobile genetic elements observed in the clonal expansion of carbapenem-resistant *Klebsiella pneumoniae* in India. *Microb Drug Resist* 2021;**27**:1096–104.
- Shen ZQ, Wang Y, Shen YB et al. Early emergence of *mcr*-1 in *Escherichia coli* from food-producing animals. *Lancet Infect Dis* 2016;**16**:293.
- Shimizu S, Iyobe S, Mitsuhashi S. Inducible high resistance to colistin in proteus strains. *Antimicrob Agents Chemother* 1977;**12**:1–3.
- Skarzynska M, Zajac M, Kaminska E et al. *Salmonella* and antimicrobial resistance in wild rodents—true or false threat? *Pathogens* 2020;**9**:771.
- Smet A, Martel A, Persoons D et al. Broad-spectrum beta-lactamases among *Enterobacteriaceae* of animal origin: molecular aspects, mobility and impact on public health. *FEMS Microbiol Rev* 2010;**34**:295–316.
- Snyman Y, Whitelaw AC, Reuter S et al. Colistin resistance mechanisms in clinical *Escherichia coli* and *Klebsiella* spp. isolates from the Western Cape of South Africa. *Microb Drug Resist* 2021;**27**:1249–58.
- Somerville JE, Cassiano L, Bainbridge B et al. A novel *Escherichia coli* lipid a mutant that produces an anti-inflammatory lipopolysaccharide. *J Clin Invest* 1996;**97**:359–65.
- Srinivasan VB, Rajamohan G. KpnEF, a new member of the *Klebsiella pneumoniae* cell envelope stress response regulon, is an SMR-type efflux pump involved in broad-spectrum antimicrobial resistance. *Antimicrob Agents Chemother* 2013;**57**:4449–62.
- Stone GG, Ponce-de-Leon A. In vitro activity of cef-tazidime/avibactam and comparators against Gram-negative bacterial isolates collected from Latin American centres between 2015 and 2017. *J Antimicrob Chemother* 2020;**75**:1859–73.
- Storm DR, Rosenthal KS, Swanson PE. Polymyxin and related peptide antibiotics. *Annu Rev Biochem* 1977;**46**:723–63.
- Sun S, Negrea A, Rhen M et al. Genetic analysis of colistin resistance in *Salmonella enterica* serovar Typhimurium. *Antimicrob Agents Chemother* 2009;**53**:2298–305.
- Taconelli E, Sifakis F, Harbarth S et al. Surveillance for control of antimicrobial resistance. *Lancet Infect Dis* 2018;**18**:E99–106.
- Teo JQM, Chang CWT, Leck H et al. Risk factors and outcomes associated with the isolation of polymyxin B and carbapenem-resistant *Enterobacteriaceae* spp.: a case-control study. *Int J Antimicrob Agents* 2019;**53**:657–62.
- Theobald S, Etter EMC, Gerber D et al. Antimicrobial resistance trends in *Escherichia coli* in South African poultry: 2009–2015. *Foodborne Pathog Dis* 2019;**16**:652–60.
- Timmerman T, Dewulf J, Catry B et al. Quantification and evaluation of antimicrobial drug use in group treatments for fattening pigs in Belgium. *Prev Vet Med* 2006;**74**:251–63.
- Trung NV, Matamoros S, Carrique-Mas JJ et al. Zoonotic transmission of *mcr*-1 colistin resistance gene from small-scale poultry farms, Vietnam. *Emerg Infect Dis* 2017;**23**:529–32.
- U.S. Food and Drug Administration (FDA). *Summary report on antimicrobials sold and distributed for use in food-producing animals*. 2016.
- Uechi K, Tada T, Shimada K et al. Emergence of a carbapenem-resistant and colistin-heteroresistant *Enterobacter cloacae* clinical isolate in Japan. *J Infect Chemother* 2019;**25**:285–8.
- Van TTH, Yidana Z, Smooker PM et al. Antibiotic use in food animals worldwide, with a focus on Africa: pluses and minuses. *J Glob Antimicrob Resist* 2020;**20**:170–7.
- Velkov T, Thompson PE, Nation RL et al., Structure-activity relationships of polymyxin antibiotics. *J Med Chem* 2010;**53**:1898–916.
- Veltman T, Jordan D, McDevitt CA et al. Absence of high priority critically important antimicrobial resistance in *Salmonella* sp. isolated from Australian commercial egg layer environments. *Int J Food Microbiol* 2021;**340**:109042.
- Vila J, Marti S, Sanchez-Cespedes J. Porins, efflux pumps and multidrug resistance in *Acinetobacter baumannii*. *J Antimicrob Chemother* 2007;**59**:1210–5.
- Vounba P, Rhouma M, Arsenaault J et al. Prevalence of colistin resistance and *mcr*-1/*mcr*-2 genes in extended-spectrum beta-lactamase/AmpC-producing *Escherichia coli* isolated from chickens in Canada, Senegal and Vietnam *J Glob Antimicrob Resist* 2019;**19**:222–7.
- Walia K, Madhumathi J, Veeraraghavan B et al. Establishing antimicrobial resistance surveillance and research network in India: journey so far. *Indian J Med Res* 2019;**149**:164–79.
- Walsh TR, Wu YN. China bans colistin as a feed additive for animals. *Lancet Infect Dis* 2016;**16**:1102–3.
- Wand ME, Sutton JM. Mutations in the two component regulator systems *PmrAB* and *PhoPQ* give rise to increased colistin

- resistance in *Citrobacter* and *Enterobacter* spp. *J Med Microbiol* 2020;**69**:521–9.
- Wang Y, Hou NX, Johnston J et al. Low prevalence of mobile colistin-resistance in US meat, catfish, poultry and genomic characterization of a *mcr-1* positive *Escherichia coli* strain. *Food Control* 2020;**118**:107434.
- Wang Y, Xu CY, Zhang R et al. Changes in colistin resistance and *mcr-1* abundance in *Escherichia coli* of animal and human origins following the ban of colistin-positive additives in China: an epidemiological comparative study. *Lancet Infect Dis* 2020;**20**:1161–71.
- Wasył D, Zajac M, Lalak A et al. Antimicrobial resistance in *Escherichia coli* isolated from wild animals in Poland. *Microb Drug Resist* 2018;**24**:807–15.
- Webb HE, Angulo FJ, Granier SA et al. Illustrative examples of probable transfer of resistance determinants from food animals to humans: streptothricins, glycopeptides, and colistin. *F1000Res* 2017;**6**:1805.
- Weterings V, Zhou K, Rossen JW et al. An outbreak of colistin-resistant *Klebsiella pneumoniae* carbapenemase-producing *Klebsiella pneumoniae* in the Netherlands (July to December 2013), with inter-institutional spread. *Eur J Clin Microbiol Infect Dis* 2015;**34**:1647–55.
- World Health Organization (WHO). *Critically important antimicrobials for human medicine*, 6th revision. Geneva: World Health Organization, 2019, 45.
- World Health Organization (WHO). *Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics*. World Health Organization (WHO).2017, 7.
- World Organisation for Animal Health (OIE). *OIE annual report on antimicrobial agents intended for use in animals*, 5th report. World Organisation for Animal Health (OIE). 2021.
- World Organisation for Animal Health (OIE). *OIE list of antimicrobial agents of veterinary importance*. World Organisation for Animal Health (OIE). 2018.
- Wright MS, Suzuki Y, Jones MB et al. Genomic and transcriptomic analyses of colistin-resistant clinical isolates of *Klebsiella pneumoniae* reveal multiple pathways of resistance. *Antimicrob Agents Chemother* 2015;**59**:536–43.
- Yeh JC, Chen CL, Chiou CS et al. Comparison of prevalence, phenotype, and antimicrobial resistance of *Salmonella serovars* isolated from turkeys in Taiwan. *Poult Sci* 2018;**97**:279–88.
- Zakaria AS, Edward EA, Mohamed NM. Genomic insights into a colistin-resistant uropathogenic *Escherichia coli* strain of O23:H4-ST641 lineage harboring *mcr-1.1* on a conjugative IncHI2 plasmid from Egypt. *Microorganisms* 2021;**9**:799.
- Zanardi G, Iemmi T, Spadini C et al. Wild micromammals as bioindicators of antibiotic resistance in ecopathology in Northern Italy. *Animals* 2020;**10**:1184.
- Zhanell GG, Adam HJ, Baxter MR et al. 42936 pathogens from Canadian hospitals: 10 years of results (2007-16) from the CANWARD surveillance study. *J Antimicrob Chemother* 2019;**74**:5–21.
- Zhu Y, Inouye M. The role of the G2 box, a conserved motif in the histidine kinase superfamily, in modulating the function of EnvZ. *Mol Microbiol* 2002;**45**:653–63.