




Pesticide poisoning: A response to Eddleston

Leslie London¹, Erik Jørs^{2,3} and Dinesh Neupane^{4,5} 

¹University of Cape Town, School of Public Health and Family Medicine, Cape Town, South Africa.

²University of Southern Denmark, Department of Clinical Research. ³Clinic of Occupational Medicine, Odense University Hospital, Odense, Denmark. ⁴Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, USA. ⁵Nepal Development Society, Bharatpur-10, Chitwan, Nepal.

We thank Michael Eddleston¹ for his comment on our Editorial in the special issue on Pesticide Poisoning in Low- and Middle-Income Countries² and note the consistent and substantial contribution made by his group on this topic, particularly related to pesticide poisonings and self-harm.^{3–7}

We agree that serious pesticide poisonings are most often due to acts of self-harm, although they are outnumbered by less-severe poisonings in occupational settings and accidental poisonings.^{8–12} Eddleston's argument that suicide should be recognized as an occupational disease is consistent with the idea that categories of work-relatedness of illness should take account of easy access to a work-related hazard.¹³ Prevention of all forms of pesticide poisoning deserves priority attention.

Reducing access to pesticides can lower the number of poisonings, be they due to self-harm, accidents, or occupation. Evidence, first identified in Sri Lanka,⁵ and confirmed in a global systematic review,⁶ suggests that banning highly hazardous pesticides, which are commonly ingested in acts of self-poisoning, can lower the number of fatal self-harm cases. Similarly, farming with reduced use of pesticides (integrated pesticide management [IPM]), can lower the number symptoms of occupational pesticide poisonings.^{6–8}

We note Eddleston's comments confirming the attention paid to providing information to households on safe storage in their trial of ground-installed containers.⁵ Despite this attention, their rigorous study was not effective in reducing suicide, illustrating the limits of interventions requiring individual behaviour change and reinforcing arguments for upstream legislative intervention to ban highly hazardous pesticides. However, not all behaviour change interventions are the same – those that empower workers and farmers to make informed choices may have substantial beneficial impacts.^{14–16}

Preventive measures should then combine multiple approaches for which there is evidence of effectiveness, and we need a hybrid approach advocating for removing highly hazardous pesticides from poor rural communities as suggested by Eddleston et al, as well as educating farmers on IPM and safer storage of pesticide. It may not be realistic to assume that all highly hazardous

pesticides would be removed immediately after adopting policies and the effects of legislation to remove the most hazardous pesticides could be reinforced by IPM promotion.

Such efforts are urgently needed in low- and middle-income countries where pesticide use is increasing rapidly without any concomitant increase in user or consumer knowledge, and without measures to improve agency on the part of those most vulnerable to prevent the harmful effects of pesticide exposure.

Author Contributions

EJ and LL drafted the article. LL, EJ and DN revised the article for important intellectual content. All authors approved the article.

ORCID iD

Dinesh Neupane  <https://orcid.org/0000-0002-1501-2990>

REFERENCES

- Eddleston M. Response to Jors et al, Environmental Health Insights [published online ahead of print July 20, 2018]. *Environ Health Insights*. doi:10.1177/1178630218788554.
- Jørs E, Neupane D, London L. Pesticide poisonings in low- and middle-income countries [published online ahead of print January 3, 2018]. *Environ Health Insights*. doi:10.1177/1178630217750876.
- Konradsen F, Dawson AD, Eddleston M, Gunnell D. Pesticide self-poisoning: thinking outside the box. *Lancet*. 2007;369:169–170. doi:10.1016/S0140-6736(07)60085-3.
- Mohamed F, Manuweera G, Gunnell D, et al. Pattern of pesticide storage before pesticide self-poisoning in rural Sri Lanka. *BMC Pub Health*. 2009;9:405. doi:10.1186/1471-2458-9-405.
- Knipe DW, Chang SS, Dawson AD, et al. Suicide prevention through means restriction: impact of the 2008–2011 pesticide restrictions on suicide in Sri Lanka. *PLoS ONE*. 2017;12:e0172893. doi:10.1371/journal.pone.0172893.
- Gunnell D, Knipe D, Chang SS, et al. Prevention of suicide with regulations aimed at restricting access to highly hazardous pesticides: a systematic review of the international evidence. *Lancet Glob Health*. 2017;5:e1026–e1037. doi:10.1016/S2214-109X(17)30299-1.
- Pearson M, Metcalfe C, Jayamanne S, et al. Effectiveness of household lockable pesticide storage to reduce pesticide self-poisoning in rural Asia: a community-based, cluster-randomised controlled trial. *Lancet*. 2017;390:1863–1872. doi:10.1016/S0140-6736(17)31961-X.
- Lekei EE, Ngowi AV, London L. Under-reporting of acute pesticide poisoning in Tanzania: modelling results from two cross-sectional studies. *Environ Health*. 2016;15:118.
- Jeyaratnam J. Acute pesticide poisoning: a major global health problem. *World Health Stat Q*. 1990;43:139–144.
- Litchfield MH. Estimates of acute pesticide poisoning in agricultural workers in less developed countries. *Toxicol Rev*. 2005;24:271–278.



11. Gunnell D, Eddleston M, Phillips MR, Konradsen F. The global distribution of fatal pesticide self-poisoning: systematic review. *BMC Pub Health*. 2007;7:357.
12. Thundiyil JG, Stober J, Besbelli N, Pronczuk J. Acute pesticide poisoning: a proposed classification tool. *Bull World Health Organ*. 2008;86:205–209.
13. Schilling RSF. Health protection and promotion at work. *British J Ind Med*. 1989;46:683–688.
14. Mancini F, Jiggins JL, O'Malley M. Reducing the incidence of acute pesticide poisoning by educating farmers on integrated pest management in South India. *Int J Occup Med Environ Health*. 2009;15:143–151. doi:10.1179/och.2009.15.2.143.
15. Braun A, Jiggins J, Röling N, Van Den Berg H, Snijders P. A global survey and review of farmer field school experiences. <https://www.share4dev.info/kb/documents/1880.pdf>. Up-dated 2010.
16. Jørs E, Lander F, Huici O, et al. Do Bolivian small holder farmers improve and retain knowledge to reduce occupational pesticide poisonings after training on Integrated Pest Management? *Environ Health*. 2014;13:75. doi:10.1186/1476-069X-13-75.

RECEIVED: December 14, 2018. **ACCEPTED:** December 14, 2018.

TYPE: Letter to the Editor

FUNDING: The author(s) received no financial support for the research, authorship, and/or publication of this article.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Erik Jørs, Department of Clinical Research, University of Southern Denmark; Clinic of Occupational Medicine, Odense University Hospital Sdr. Boulevard 29, 5000 Odense C, Denmark. Email: erik.joers@rsyd.dk