

Psychological, sociodemographic, and clinicopathological predictors of breast cancer patients' decision to undergo breast reconstruction after mastectomy

Azmi Marouf, Medical Student, Hatan Mortada, MBBS, Mahmoud G. Fakiha, MD, DESC.

ABSTRACT

الأهداف: الهدف من البحث هو دراسة أثر العوامل الاجتماعية و النفسية و السمات الشخصية عند مريضات سرطان الثدي على قرار الخضوع لإعادة ترميم الثدي بعد الاستئصال.

المنهجية: في هذه الدراسة المقطعية، تم إجراء مقابلات شخصية مع مريضات سرطان الثدي الاتي خضعن لعملية استئصال الثدي بين يناير 2017م إلى 2019م باستخدام استبيان من 46 بنداً في مستشفى جامعة الملك عبد العزيز في جدة، المملكة العربية السعودية. انقسم الاستبيان إلى 5 أقسام: البيانات الشخصية، السمات الشخصية، مؤشر النرجسية، مقياس القلق والاكتئاب. تم منح الموافقة الأخلاقية من قبل لجنة الأخلاقيات البحثية. أجريت التحليل الاحصائية باستخدام SPSS الإصدار 23.0.

النتائج: حجم العينة المشاركة في الدراسة 196 مرضية، متوسط العمر 48.44 (±9.87) سنة، 60.2% مؤيدتين لعملية ترميم الثدي، 36.7% لديهن أمراض مزمنة، 43.9% من الجنسية السعودية. متوسط مؤشر النرجسية، مقياس القلق و الاكتئاب هو 4.17 (±2.81)، 7.72 (±4.88) و 5.78 (±4.65) على التوالي. ظهرت اختلافات المجموعة في العمر، والاكتئاب، تدخل العقدة الليمفاوية، و اثنين من سمات الشخصية.

الخلاصة: في هذه الدراسة، أثر العمر والاكتئاب بشكل كبير على قرار الحصول على عملية ترميم الثدي. السيدات اللاتي اخترن عملية ترميم الثدي كانوا اصغر سناً، وأقل اكتئاباً، وأقل تدخل للعقد الليمفاوية. أثرت بعض السمات الشخصية أيضاً على قرار الخضوع لعملية ترميم الثدي. يعتبر تشخيص وعلاج مرض الاكتئاب عنصر مهم في رعاية المصابات بسرطان الثدي. أخيراً، يجب ألا يمنع عمر المريضة من تقديم المشورة لها بخصوص عملية ترميم الثدي.

Objectives: To examine the influence of socioeconomic factors, anxiety, depression, and personality traits on the decision to undergo breast reconstruction (BR) post mastectomy.

Methods: In this cross-sectional study, adult female breast cancer (BC) patients who had undergone a mastectomy between January 2017 and 2019 were interviewed using a 46-item questionnaire at King Abdulaziz University Hospital, Jeddah, Saudi Arabia. The questionnaire was divided into 5 sections: demographic data, personality traits, the Narcissism Index (NI), and the Hospital Anxiety and Depression Scale (HADS).

Results: The sample (N=196; mean age: 48.44 [± 9.87] years, opted for BR: 60.2%, with chronic diseases: 36.7%, Saudi nationals: 43.9%) obtained means of 4.17 (± 2.81) on NI score, 7.72 (± 4.88) on anxiety score, and 5.78 (± 4.65) on depression score. Group differences in age, depression, lymph node involvement, and 2 personality traits emerged.

Conclusion: Age and depression significantly influenced the decision to undergo BR. Women who had opted for BR tended to be younger, be less depressed, and be characterized by lesser nodal involvement. Personality traits also influenced the decision to undergo BR. Diagnosing and treating depressed patients is an important component. A patient's age should not deter physicians from counselling them about BR.

Keywords: anxiety, breast cancer, breast reconstruction, depression, mastectomy

*Saudi Med J 2020; Vol. 41 (3): 267-274
doi: 10.15537/smj.2020.3.24946*

From the Faculty of Medicine (Marouf), King Abdulaziz University; from the Department of Plastic Surgery & Burn Unit (Mortada), King Saud Medical City; and from the Division of Plastic & Reconstructive Surgery (Fakiha), Department of Surgery, University of Jeddah, Jeddah, Kingdom of Saudi Arabia.

Received 1st November 2019. Accepted 16th January 2020.

*Address correspondence and reprint request to: Dr. Hatan Mortada, Department of Plastic Surgery & Burn Unit, King Saud Medical City Riyadh, Kingdom of Saudi Arabia. E-mail: hatanmortada@gmail.com
ORCID ID: <https://orcid.org/0000-0003-1283-4136>*

Breast cancer is the most commonly occurring cancer among females, and it affects 2.1 million women worldwide each year.¹ Among Saudi women, it is the most common cancer.² The incidence of breast cancer in the United States (US) is 124 per 100000 women.³ In many cases of breast cancer, a mastectomy is indicated. Mastectomy significantly deprives women of their sense

of femininity and makes them feel less attractive, and this in turn adversely influences their quality of life (QOL).⁴⁻⁷ Postmastectomy breast reconstruction (BR) improves QOL.^{8,9} There are primarily 2 types of BR, namely, implant and autologous (flap) reconstruction, and each of these can be implemented through various approaches. For example, implants include saline and silicone. A flap of the latissimus dorsi or transverse rectus abdominis (TRAM) is used for autologous BR. The latissimus dorsi flap is swung from the back. It provides less fullness than TRAM, thus an implant or a fat graft is often used coupled with it.¹⁰ A TRAM flap is performed by rotating the rectus abdominis muscle or by removing a smaller part of the muscle with its skin (free flap). Free TRAM flaps provide more flexibility and lower risk of tissue necrosis when compared to pedicle TRAM flaps.¹¹ It has been shown that autologous BR yields superior outcomes, when compared to implant BR.¹²⁻¹⁴ Nowadays, most BRs involve implants; in the late twentieth century, autologous implants were more popular.¹⁵ This change is partially attributable to surgeons' efforts to provide the option of reconstruction to the high-risk postmastectomy population (namely, patients with other comorbidities and/or those older than 60 years).¹⁶

Breast cancer patients experience psychological morbidities, primarily mood disorders, after receiving and when living with the diagnosis. The untreated anxiety and depression are important factors that render the QOL of these patients low, not merely the physical aspect of the patients' presentation.^{17,18} These morbidities in women with breast cancer are often overlooked and left untreated because they can have unclear presentations.¹⁷

According to the findings of a Polish study, deciding to undergo BR after breast removal is correlated with personality traits among women. Specifically, introverts rarely opted to undergo BR.¹⁹ Another important personality type is the narcissistic personality. Specifically, narcissists tend to demand perfectionism as it is a sentinel feature of their personality type; hence, they seek to have an extremely attractive body.²⁰⁻²²

Despite recommendations to undergo BR, the rates at which the procedure is performed are low, ranging from 11% to less than 40% (in some tertiary care centers in the US).^{23,24} The decision to undergo BR

is complicated by a variety of procedures.¹² Women are asked to choose from an array of procedures, and the problem has become more strenuous with the development of the perforator flap technique, which minimizes the risk of hernia albeit requires longer operation time.¹² Additionally, the burden that the surgical procedure places on the patient and her relatives makes the decision-making process even more difficult.²⁵ Two studies that were conducted in Europe found that the patient's age influences her decision to undergo BR and the possession of information about the procedure.^{26,27}

Only a small proportion of treated breast cancer patients decide to undergo BR.^{15,28} Therefore, it is important to examine which mood states and social aspects influence this decision-making process. In this study, we aimed to examine the influence of socioeconomic factors, anxiety, depression, and personality traits on the decision to undergo BR after a mastectomy in one center in Jeddah, Saudi Arabia.

Methods. This cross-sectional study was conducted between March and July 2019, using an interview-based questionnaire. This interview was conducted among female patients who were diagnosed with breast cancer at King Abdulaziz University Hospital (KAUH), Jeddah, Saudi Arabia.

Women aged 18 years and older who had undergone mastectomy for breast cancer between January 2017 and 2019 were included. The exclusion criteria were as follows: a) undergone bilateral mastectomy, contralateral mastectomy for prophylactic purposes, or a reoperation and b) breast cancer recurrence. All contributors were knowledgeable before participation, and those who agreed to contribute were registered. Approximately 400 randomly chosen patients were identified from the hospital records, and 241 of them who met the study's inclusion criteria were contacted; only 196 patients agreed to participate (**Figure 1**). Patients were divided into 2 subgroups: Group A, which consisted of women who had opted for BR and Group B, which consisted of women who chose to not undergo BR.

Study variables. We reviewed the literature and developed a valid questionnaire based on the findings of several published studies.^{19,29-32} The questionnaire consisted of 46 questions that were divided into 5 sections: demographic data, personality traits,¹⁹ the Narcissism Index (NI),^{29,30} and the Hospital Anxiety and Depression Scale (HADS).^{31,32} The NI is a reliable tool for predicting the degree of narcissism.^{29,30} The reliability and validity of HADS on similar populations (breast cancer patients) has been studied, and it has been

Disclosure. Authors have no conflict of interests, and the work was not supported or funded by any drug company.




Figure 1 - Flowchart depicting the inclusion and exclusion of the study participants.

shown to have a good predictive value. Furthermore, the Arabic translation used in this study has been shown to be both valid and reliable.³¹⁻³³ The study was conducted after ethical approval was granted by the Research Ethics Committee at the King Abdulaziz University Hospital. Patients participated in this study on an entirely voluntary basis.

Statistical analysis. Data were checked for completeness, and errors were corrected. The normality of the distributions of continuous variables were checked by conducting the Shapiro-Wilk and Kolmogorov Smirnov tests. Their results indicated that all variables were none normally distributed. The sociodemographic variables and clinicopathological features (namely, tumor size, lymph node involvement, and metastasis-TNM) of the 2 participant groups were compared using the Chi-squared test. Respondents' scores on the NI and HADS were calculated by counting the responses. The mean scores of the 2 participant groups were compared using the Mann-Whitney U test. The relationship between TNM and anxiety and depression scores were examined using the Kruskal-Wallis test. The extroversion/introversion personality continuum of the

2 participant groups were compared using Chi-squared and Fisher exact tests. Binary logistic regression analysis was conducted to identify the predictors of the decision to undergo BR. The results have been presented using tables and figures.

All analyses were conducted using the Statistical Package for Social Sciences version 23.0 (IBM, Armonk, NY, USA), and 95% confidence intervals were computed to interpret the results.

Results. A total of 196 patients were interviewed, and 118 (60.2%) of them had opted for BR; the remaining 78 (39.8%) patients had not opted for BR. The mean age of the participants was 48.44 (\pm 9.87) years. Further, 86 (43.9%) patients were Saudi, the monthly income of 118 (60.2%) patients was <5000 Saudi Riyals (SR). One hundred and thirty (66.3%) patients were unemployed/housewives, 72 (36.7%) patients had chronic illnesses, and 76 (38.8%) patients had a family history of breast cancer (BC). Out of the 76 participants with a family history of BC, 71.1% had a relative who had BC and had undergone BR. The

Table 1 -Sociodemographic characteristics and clinicopathological features of the respondents split by participant group.

Characteristics	Attributes	Breast reconstruction (n=118)	No breast reconstruction (n=78)	Odds ratio (95% CI)	P-value
Age (years)	Mean ± SD	45.44 ± 8.66	52.97 ± 9.90		<0.001
	Median	45.0	53.0		
Nationality	Saudi	62 (52.5)	24 (30.8)	2.491 (1.365 - 4.546)	0.003
	Non-Saudi	56 (47.5)	54 (69.2)		
Marital status	Single	9 (7.6)	9 (11.5)		0.779
	Married	93 (78.8)	57 (73.1)		
	Divorced	8 (6.8)	6 (7.7)		
	Widowed	8 (6.8)	6 (7.7)		
Educational level	Below high school	29 (24.6)	19 (24.4)		0.721
	High school	37 (31.4)	29 (37.2)		
	Bachelor's degree	41 (34.7)	25 (32.1)		
	Master	6 (5.1)	4 (5.1)		
	PHD	5 (4.2)	1 (1.3)		
Monthly income (Saudi Riyals)	<5000	68 (57.6)	50 (64.1)		0.506
	5000-10,000	18 (15.3)	14 (17.9)		
	10,000-20,000	24 (20.3)	10 (12.8)		
	>20,000	8 (6.8)	4 (5.1)		
Occupation	Works in Jeddah	26 (22.0)	24 (30.8)		0.041
	Works outside Jeddah	14 (11.9)	2 (2.6)		
	Unemployed	78 (66.1)	52 (66.7)		
Smoking	No	104 (88.1)	76 (97.4)	0.195 (0.043 - 0.886)	0.015
	Yes	14 (11.9)	2 (2.6)		
Chronic illness	No	88 (74.6)	36 (46.2)	3.422 (1.863 - 6.287)	<0.001
	Yes	30 (25.4)	42 (53.8)		
Family history of breast cancer	No	70 (59.3)	50 (64.1)	0.817 (0.452 - 1.474)	0.501
	Yes	48 (40.7)	28 (35.9)		
Family member underwent BR postmastectomy	No	4 (8.3)	8 (28.6)		0.046
	Yes	36 (75)	18 (64.3)		
	Don't know	8 (16.7)	2 (7.1)		
Tumor size	Ductal carcinoma in situ	27 (22.9)	10 (12.8)		0.241
	Breast cancer T1a-c (≤2cm)	33 (28.0)	31 (39.7)		
	Breast cancer T2	34 (28.8)	23 (29.5)		
	Breast cancer T3	10 (8.5)	8 (10.3)		
	Breast cancer T4	14 (11.9)	6 (7.7)		
Lymph node involved	N0	56 (47.5)	35 (44.9)		0.013
	N1	25 (21.1)	30 (38.5)		
	N2	20 (16.8)	10 (12.8)		
	N3	17 (14.4)	3 (3.8)		
Distant metastasis	No	110 (93.2)	70 (89.7)	0.636 (0.228 - 1.773)	0.430
	Yes	8 (6.8)	8 (10.3)		
Adjuvant therapy	None	29 (24.6)	7 (9.0)		0.291
	Chemotherapy	54 (45.8)	47 (60.3)		
	Chemo + endocrine therapy	6 (5.1)	8 (10.3)		
	Chemo + radio therapy	16 (13.6)	8 (10.3)		
	Chemo + radio + endocrine	13 (11.0)	8 (10.3)		

*odds ratio were presented for dichotomous sociodemographic variables. BR: breast reconstruction. 95% CI: 95% conference interval

Table 2 - Distribution of respondents' mean NI score, anxiety score and depression score split by participant group.

Scores	Breast reconstruction (n=118)	No breast reconstruction (n=78)	P-value
Narcissism Index Score	4.53 ± 2.84	3.64 ± 2.70	0.030
Anxiety score	6.44 ± 4.31	9.67 ± 5.07	<0.001
Depression score	4.49 ± 3.97	7.74 ± 4.90	<0.001

Values are presented as mean±SD.
*Mann-Whitney U test was carried out

sociodemographic characteristics for each participant group are presented in **Table 1**.

The group of women who had opted for BR tended to have younger women (mean difference between the 2 groups was 7.53 years, $p<0.001$), fewer nonsmokers ($p=0.015$), and more women without chronic illnesses ($p<0.001$). Unemployed participants were less likely to have opted for BR ($p=0.041$). With regard to the TNM classification, a lymph node involvement was

less prominent among those who had opted for BR ($p=0.13$) (Table 1).

Personality traits and mood status. The mean depression score for all participants was $5.78 (\pm 4.65)$, anxiety score $7.72 (\pm 4.88)$, and NI $4.17 (\pm 2.81)$. Table 2 shows that anxiety (9.67 ± 5.07 versus 6.44 ± 4.31 , $p<0.001$) and depression scores (7.74 ± 4.90 versus 4.49 ± 3.97 , $p<0.001$) were significantly higher among participant who had opted to not undergo BR. However, scores on the NI were higher among those who had opted for BR ($p=0.030$) (Table 2).

The responses that the 2 participant groups provided to the extroversion/introversion personality questions (total: 5 questions) are summarized in Table 3. Group comparisons revealed that there was a statistically

significant difference in the responses that were provided to questions 4 ($p=0.0002$) and 5 ($p<0.001$) (Table 3).

Disease stage and mood status. There was no statistically significant relationship between TNM classification and anxiety and depression scores ($p>0.050$).

Binary logistic regression. Binary logistic regression analysis was used to identify the predictors of the decision to undergo BR. Anxiety and depression scores, sociodemographic variables, TNM classification, and adjuvant treatment were included in the regression analysis. The results revealed that depression scores, age, and lymph node involvement were statistically significant predictors of BR decision. Other included variables were not statistically significant ($p>0.050$) (Table 4).

Table 3 - Extroversion/introversion personality continuum in the 2 study groups.

Personality questions/ answers	Breast reconstruction (n=118)	No breast reconstruction (n=78)	P-value
1) Do you seek success in life?			0.254 [‡]
Definitely yes	100 (84.7)	68 (87.2)	
Rather yes	12 (10.2)	4 (5.1)	
Hard to say	4 (3.4)	6 (7.7)	
Rather no	2 (1.7)	0 (0.0)	
Definitely no	0 (0.0)	0 (0.0)	
2) Do you have the ability to motivate other people?			0.216 [‡]
Definitely yes	80 (67.8)	50 (64.1)	
Rather yes	26 (22.0)	20 (25.6)	
Hard to say	8 (6.8)	6 (7.7)	
Rather no	4 (3.4)	0 (0.0)	
Definitely no	0 (0.0)	2 (2.6)	
3) Are you open to other people?			0.119 [‡]
Definitely yes	74 (62.7)	50 (64.1)	
Rather yes	30 (25.4)	12 (15.4)	
Hard to say	10 (8.5)	8 (10.3)	
Rather no	4 (3.4)	8 (10.3)	
Definitely no	0 (0.0)	0 (0.0)	
4) Does belonging to a social group have an impact of your feeling of safety?			0.002*
Definitely yes	52 (44.1)	40 (51.3)	
Rather yes	22 (18.6)	16 (20.5)	
Hard to say	4 (3.4)	14 (17.9)	
Rather no	20 (16.9)	6 (7.7)	
Definitely no	20 (16.9)	2 (2.6)	
5) Is the opinion of other people about you important for you?			<0.001*
Definitely yes	38 (32.2)	36 (46.2)	
Rather yes	30 (25.4)	20 (25.6)	
Hard to say	8 (6.8)	8 (10.3)	
Rather no	12 (10.2)	12 (15.4)	
Definitely no	30 (25.4)	2 (2.6)	

[‡]Fisher-Freemen-Halton test. *Chi-square test

Discussion. Among breast cancer patients, undergoing mastectomy frequently has a negative impact on the QOL and psychological well-being. Undoubtedly, this body part entails important psychological implications. Consequently, patients who have undergone a mastectomy perceive a decline their sense of femininity and confidence levels. A mastectomy without BR can alter the mood and behaviors of patients in their workspaces.³⁴ Further, it can strongly impact their roles as a family member. Therefore, BR after breast removal can tremendously improve QOL and perceived body image. Accordingly, we aimed to examine the relationship that the decision to undergo BR shares with different personality traits, depression, anxiety and sociodemographic variables among breast cancer patients.

The existing literature suggests that age is a very important determinant on BR.^{24,26,27} A similar result emerged in this study. This finding is attributable to surgeons' fear of postoperative complications, especially in a patient with other chronic illnesses.²⁷ There was a significant difference in BR status between Saudis and non-Saudis. This finding may be attributable to group differences in access to healthcare or the relatively higher mean depression (6.47 ± 4.96 versus 4.91 ± 4.04 , $p=0.035$) and anxiety scores (8.93 ± 5.18 versus 6.19 ± 4.00 , $p<0.001$) that non-Saudi participants obtained (the regression analysis presented in Table 4 yielded no significant predictor).

Another interesting finding was that the patients who were willing to undergo BR were healthier and did not have any chronic illnesses or comorbidities, when compared to the other group. Being healthy is closely related to happiness, and deciding to undergo such a procedure can tremendously improves a patient's

Table 4 - Logistic regression of the predictors of the decision to undergo breast reconstruction postmastectomy.

Variables	Odds ratio	95% CI	P-value
Anxiety score	0.897	0.786 - 1.024	0.108
Depression score	0.794	0.695 - 0.907	0.001
Age (years)	0.870	0.813 - 0.930	0.001
Nationality			
Saudi	Ref.		
Non-Saudi	2.082	0.722 - 6.005	0.175
Occupation			
Unemployed	Ref.		
Works outside Jeddah	0.006	0.000032 - 1.138	0.056
Works in Jeddah	0.101	0.000435 - 23.453	0.410
Monthly income (SR)			
<5000	Ref.		
5000-10000	0.514	0.057 - 4.607	0.552
10000-20000	0.355	0.033 - 3.816	0.392
>20000	1.030	0.104 - 10.243	0.980
Educational level			
PHD	Ref.		
Master	0.643	0.076 - 5.491	0.687
Bachelor degree	0.880	0.104 - 7.450	0.907
High school	1.567	0.044 - 55.619	0.805
Below high school	0.520	0.058 - 4.676	0.560
Marital status			
Married	Ref.		
Single	0.731	0.068 - 7.903	0.797
Divorced	0.918	0.166 - 5.084	0.922
Widowed	0.349	0.041 - 2.929	0.332
Smoking			
No	Ref.		
Yes	7.045	0.085 - 582.244	0.386
Chronic illness			
No	Ref.		
Yes	0.983	0.342 - 2.826	0.975
Tumor size			
Ductal carcinoma in situ	Ref.		
Breast cancer T1a-c	1.776	0.298 - 10.595	0.528
Breast cancer T2	0.632	0.121 - 3.304	0.587
Breast cancer T3	2.461	0.421 - 14.399	0.318
Breast cancer T4	1.309	0.156 - 10.955	0.804
Lymph node involved			
N0	Ref.		
N1	0.217	0.021 - 2.255	0.201
N2	0.041	0.003 - 0.540	0.015
N3	0.090	0.007 - 1.156	0.064
Distal metastasis			
No	Ref.		
Yes	0.756	0.165 - 3.464	0.719
Adjuvant treatment			
None	Ref.		
Chemotherapy	2.475	0.221 - 27.750	0.462
Chemo+endocrine therapy	0.657	0.086 - 5.008	0.685
Chemo+radiotherapy	2.711	0.149 - 49.222	0.500
Chemo+radio+endocrine	1.424	0.129 - 15.732	0.773
Family history of BR			
Yes	Ref.		
No	0.194	0.034 - 1.095	0.063

Ref.: reference, SR: Saudi Riyals, BR: breast reconstruction

QOL.^{4-7,35} This is similar to the findings of 2 studies that were conducted in the US³⁶ and Australia,³⁷ whereby patients with fewer comorbidities were found to be more likely to undergo BR.

Past findings suggest that patients with early-stage BC are more likely to undergo BR. Similarly, in the present study, the percentage of node negative patients was higher in the BR group. It is noteworthy that many of these early stage BC patients do not require postmastectomy chemotherapy and radiotherapy.²⁴

It is noteworthy that, in the present study, patients with a family member who had undergone BR were more likely to have opted for BR than their counterparts. This may be the case because BC survivors prefer to receive information about BR from their friends and peers. Further, they feel less isolated after they receive such information from patients who have undergone BR.³⁸

In this study, the HADS was used to assess the affective states of the participants. The Center for Epidemiological Studies Depression (CES-D) Scale was not used owing to 2 primary reasons: a) it does not screen for anxiety and b) its performance and reliability has been examined primarily mostly on young population.^{39,40} In comparison, the HADS is a simpler assessment, and its reliability has been examined across different contexts (including among BC patients and cancer patients in general) and age groups.³¹⁻³³ Our results suggest that women with depression and anxiety are not prepared to undergo another nonvital procedure as most of the focus is on monitoring BC recurrence with anxiety and depression being vaguely present.¹⁷ Satisfactory doctor-patient communication regarding patient concerns and the provision of crucial information to the patient are important factors that can improve the detection of psychological problems and mitigate depressive symptoms and anxiety among BC patients.¹⁸

The responses that the 2 groups provided to the 5 personality questions, which were adapted from the assessment that was used in a Polish study, were compared using Chi-squared and Fisher exact tests (Table 3). Interestingly, 2 questions for which significant group differences had emerged in the original study had also evidenced significant group differences in the current study.¹⁹ In both the studies, patients who had opted for BR were more likely to be extroverted, as extrovert individuals are mainly outgoing and like to be with others. In the present study, the NI scores were higher among those who had opted for BR.

Study limitations. Although our sample size was moderate, and the study's objectives were addressed, the study has a few limitations. First, our article was

cross-sectional in nature and covered a short time period. Therefore, the participants may not have been representative of the entire population of Saudi Arabia. Second, this study was conducted in a single center, thereby limiting the generalizability of the results. Third, the reliability of the 5 personality questions has not been studied. In the future, studies must include a larger number of hospitals that represent different regions within the country. This study offers insights into the prevalence of depression and anxiety among breast cancer patients after mastectomy. It also uncovers the variability in personality traits within this population. More importantly, it indicates that depression single handedly plays a significant role in the decision to not undergo BR, after potential confounds are controlled for. It is important to screen and detect undiagnosed depression and anxiety among patients who have undergone a mastectomy. This is necessary to effectively treat these conditions through prompt referral to a psychiatrist, improve their QOL, restore at least a moderate level of functioning, and provide assistance during this difficult postmastectomy decision-making period. Patients of all ages should receive counseling about BR after a mastectomy.

In conclusion, undergoing BR is a difficult, yet important decision for a woman who survived BC to make. Age and depression significantly influenced this decision among our study participants. Women who opted to undergo BR tend to be younger, report lower levels of depression, and were characterized by less nodal involvement. Certain personality traits can also influence the decision to undergo BR. Diagnosing and treating depressed patients is an important aspect of caring for BC patients. A patient's age should not deter physicians from counseling patients about BR.

Acknowledgment. *The authors would like to thank Randa Abushanab and Renad Alnemari for their help in data collection. Also, we would like to acknowledge www.Editage.com for English language editing.*

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; 68: 394-424.
- Saudi Health Council. Cancer Incidence Report in Saudi Arabia 2014. Available from: <https://nhic.gov.sa/eServices/Documents/2014.pdf>
- US Cancer Statistics Working Group. US Cancer Statistics Data Visualizations Tool, based on November 2017 submission data (1999-2015): US Department of Health and Human Services, Centers for Disease Control and Prevention and National Cancer Institute. Center for Disease Control and Prevention and National Cancer Institute. [Cited June 2018.] Available from URL: <https://www.cdc.gov/cancer/uscs/dataviz/index.htm>
- Noyan MA, Sertoz OO, Elbi H, Kayar R, Yilmaz R. Variables affecting patient satisfaction in breast surgery: a cross-sectional sample of Turkish women with breast cancer. *Int J Psychiatry Med* 2006; 36: 299-313.
- Ganz PA, Desmond KA, Leedham B, Rowland JH, Meyerowitz BE, Belin TR. Quality of life in long-term, disease-free survivors of breast cancer: a follow-up study. *J Natl Cancer Inst* 2002; 94: 39-49.
- Kenny P, King MT, Shiell A, Seymour J, Hall J, Langlands A, Boyages J. Early stage breast cancer: costs and quality of life one year after treatment by mastectomy or conservative surgery and radiation therapy. *Breast* 2000; 9: 37-44.
- Manganiello A, Hoga LA, Reberte LM, Miranda CM, Rocha CA. Sexuality and quality of life of breast cancer patients post mastectomy. *Eur J Oncol Nurs* 2011; 15: 167-172.
- Cordeiro PG. Breast reconstruction after surgery for breast cancer. *N Engl J Med* 2008; 359: 1590-1601.
- Eltahir Y, Werners LL, Dreise MM, van Emmichoven IA, Jansen L, Werker PM, de Bock GH. Quality-of-life outcomes between mastectomy alone and breast reconstruction: comparison of patient-reported BREAST-Q and other health-related quality-of-life measures. *Plast Reconstr Surg* 2013; 132: 201e-209e.
- di Pompeo FS, Laporta R, Sorotos M, Pagnoni M, Falesiedi F, Longo B. Latissimus dorsi flap for total autologous immediate reconstruction without implants. *Plast Reconstr Surg* 2014; 134: 871e-879e.
- Kroll SS, Gherardini G, Martin JE, Reece GP, Miller MJ, Evans GR, Robb GL, Wang BG. Fat necrosis in free and pedicled TRAM flaps. *Plast Reconstr Surg* 1998; 102: 1502-1507.
- Yueh JH, Slavin SA, Adesiyun T, et al. Patient satisfaction in postmastectomy breast reconstruction: a comparative evaluation of DIEP, TRAM, latissimus flap, and implant techniques. *Plast Reconstr Surg* 2010; 125: 1585-1595.
- Pusic AL, Matros E, Fine N, Buchel E, Gordillo GM, Hamill JB, Kim HM, Qi J, Albornoz C, Klassen AF, Wilkins EG. Patient-reported outcomes 1 year after immediate breast reconstruction: results of the mastectomy reconstruction outcomes consortium study. *J Clin Oncol* 2017; 35: 2499-2506.
- Saulis AS, Mustoe TA, Fine NA. A retrospective analysis of patient satisfaction with immediate postmastectomy breast reconstruction: comparison of three common procedures. *Plast Reconstr Surg* 2007; 119: 1669-1676.
- Albornoz CR, Bach PB, Mehrara BJ, Disa JJ, Pusic AL, McCarthy CM, Cordeiro PG, Matros E. A paradigm shift in US breast reconstruction: increasing implant rates. *Plast Reconstr Surg* 2013; 131: 15-23.
- Farhangkhoe H, Matros E, Disa J. Trends and concepts in post-mastectomy breast reconstruction. *J Surg Oncol* 2016; 113: 891-894.
- Maass SW, Roorda C, Berendsen AJ, Verhaak PF, de Bock GH. The prevalence of long-term symptoms of depression and anxiety after breast cancer treatment: a systematic review. *Maturitas* 2015; 82: 100-108.

18. Turner J, Kelly B, Swanson C, Allison R, Wetzig N. Psychosocial impact of newly diagnosed advanced breast cancer. *Psychooncology* 2005; 14: 396-407.
19. Miśkiewicz H, Antoszewski B, Iljin A. Personality traits and decision on breast reconstruction in women after mastectomy *Polish Journal of Surgery* 2016; 88: 209-214.
20. Siomopoulos V. Narcissistic personality disorder: Clinical features. *American Journal of Psychotherapy* 1988; 42: 240-253.
21. Sorotzkin B. Understanding and treating perfectionism in religious adolescents. *Psychotherapy: Theory, Research, Practice, Training* 1998; 35: 87.
22. Smith MM, Sherry SB, Chen S, Saklofske DH, Flett GL, Hewitt PL. Perfectionism and narcissism: A meta-analytic review. *J Res Pers* 2016; 64: 90-101.
23. Oh DD, Flitcroft K, Brennan ME, Spillane AJ. Patterns and outcomes of breast reconstruction in older women—a systematic review of the literature. *Eur J Surg Oncol* 2016; 42: 604-615.
24. Brennan ME, Spillane AJ. Uptake and predictors of post-mastectomy reconstruction in women with breast malignancy—systematic review. *Eur J Surg Oncol* 2013; 39: 527-541.
25. Somogyi RB, Webb A, Baghdikian N, Stephenson J, Edward KL, Morrison W. Understanding the factors that influence breast reconstruction decision making in Australian women. *Breast* 2015; 24: 124-130.
26. Lardi AM, Myrick ME, Haug M, Schaefer DJ, Bitzer J, Simmen U, Güth U. The option of delayed reconstructive surgery following mastectomy for invasive breast cancer: Why do so few patients embrace this offer?. *Eur J Surg Oncol* 2013; 39: 36-43.
27. Héquet D, Zarca K, Dolbeault S, Couturaud B, Ngô C, Fourchette V, et al. Reasons of not having breast reconstruction: A historical cohort of 1937 breast cancer patients undergoing mastectomy. *Springerplus* 2013; 2: 325.
28. Harcourt D, Rumsey N. Psychological aspects of breast reconstruction: a review of the literature. *J Adv Nurs* 2001; 35: 477-487.
29. Raskin R, Terry H. A principal-components analysis of the Narcissistic Personality Inventory and further evidence of its construct validity. *J Pers Soc Psychol* 1988; 54: 890.
30. Ames DR, Rose P, Anderson CP. The NPI-16 as a short measure of narcissism. *J Pers Soc Psychol* 2006; 40: 440-450.
31. Zigmund AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983; 67: 361-370.
32. Terkawi AS, Tsang S, AlKahtani GJ, Al-Mousa SH, Al Musaed S, AlZoraigi US, Alasfar EM, Doais KS, Abdulrahman A, Altirkawi KA. Development and validation of Arabic version of the Hospital Anxiety and Depression Scale. *Saudi J Anaesth* 2017; 11 (Suppl 1): S11.
33. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale: an updated literature review. *J Psychosom Res* 2002; 52: 69-77.
34. Keesing S, Rosenwax L, McNamara B. The implications of women's activity limitations and role disruptions during breast cancer survivorship. *Women's Health* 2018; 14: 1745505718756381.
35. Offodile AC, Wenger J, Guo L. Relationship between comorbid conditions and utilization patterns of immediate breast reconstruction subtypes post-mastectomy. *The Breast Journal* 2016; 22: 310-315.
36. Reuben BC, Manwaring J, Neumayer LA. Recent trends and predictors in immediate breast reconstruction after mastectomy in the United States. *Am J Surg* 2009; 198: 237-243.
37. Hall SE, Holman CD. Inequalities in breast cancer reconstructive surgery according to social and locational status in Western Australia. *Eur J Surg Oncol* 2003; 29: 519-525.
38. Webb C, Sharma V, Temple-Oberle C. Delivering breast reconstruction information to patients: Women report on preferred information delivery styles and options. *Plast Surg (Oaku)* 2018; 26: 26-32.
39. Pretorius TB. Cross-cultural application of the Center for Epidemiological Studies Depression Scale: A study of black South African students. *Psychol Rep* 1991; 69 (3 Pt 2): 1179-1185.
40. Iwata N, Buka S. Race/ethnicity and depressive symptoms: a cross-cultural/ethnic comparison among university students in East Asia, North and South America. *Soc Sci Med* 2002; 55: 2243-2252.