



CD4+/CD8+ Ratio and Growth Differentiation Factor 8 Levels in Peripheral Blood of Large Canine Males Are Useful Parameters to Build an Age Prediction Model

Han-Jun Lee^{1,*}, Seok-Jin Hong^{1,*}, Seung-Soo Kim^{2,*}, Young-Yon Kwon^{1,*}, Bong-Hwan Choi^{3,*}, Kyung-Mi Choi², Seo-Hyeong Sheen¹, Myung-Jin Lee¹, Sun-Young Hwang¹, Kyoungwan Park⁴, Younghun Joo⁴, Hwayoung Song⁴, Cheol-Koo Lee^{1,2}

¹Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, ²Institute of Animal Molecular Biotechnology, Korea University, Seoul, ³Animal Genomics and Bioinformatics Division, National Institute of Animal Science, Rural Development Administration, Wanju, ⁴Military Working Dog Training Center, Chuncheon, Korea

Purpose: To build an age prediction model, we measured CD4+ and CD8+ cells, and humoral components in canine peripheral blood.

Materials and Methods: Large Belgian Malinois (BGM) and German Shepherd Dog (GSD) breeds (n=27), aged from 1 to 12 years, were used for this study. Peripheral bloods were obtained by venepuncture, then plasma and peripheral blood mononuclear cells (PBMCs) were separated immediately. Six myokines, including interleukin (IL)-6, IL-8, IL-15, leukemia inhibitory factor (LIF), growth differentiation factor 8 (GDF8), and GDF11 were measured from plasma and CD4+/CD8+ T-lymphocytes ratio were measured from PBMC. These parameters were then tested with age prediction models to find the best fit model.

Results: we found that the T-lymphocyte ratio (CD4+/CD8+) was significantly correlated with age (r=0.46, p=0.016). Among the six myokines, only GDF8 showed a significant correlation with age (r=0.52, p=0.005). Interestingly, these two markers showed better correlations in male dogs than females, and BGM breed than GSD. Using these two age biomarkers, we could obtain the best fit in a quadratic linear mixed model (r=0.77, p=3×10⁻⁶).

Conclusions: Age prediction is a challenging task because of complication with biological age. Our quadratic linear mixed model using CD4+/CD8+ ratio and GDF8 level showed a meaningful age prediction.

Keywords: Age prediction model; Aging; CD4-CD8 ratio; Dogs; Myostatin

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

A typical phenomenon of aging is a time-dependent

functional decline. Like other species, dogs show a decline in their physical ability with advanced age. In canines, aging-related multiple functional declines, such

Received: Jan 19, 2021 **Revised:** Sep 14, 2021 **Accepted:** Oct 8, 2021 **Published online** Jan 6, 2022

Correspondence to: Cheol-Koo Lee <https://orcid.org/0000-0003-3927-9195>

Department of Biotechnology, College of Life Sciences and Biotechnology, Korea University, 145 Anam-ro, Seongbuk-gu, Seoul 02841, Korea.

Tel: +82-2-3290-3008, **Fax:** +82-2-3290-3008, **E-mail:** ckleee2005@korea.ac.kr

*These authors contributed equally to this work as co-first authors.

as muscle weakness, and changes in blood composition have been observed [1,2]. Body size shows a negative correlation with lifespan when large and small dogs were compared [3]. But, this negative correlation can be related to a higher death rate caused by cancer in large dogs compared to that in small dogs [4]. Indeed, there is no relationship between lifespan and body size in the same breed [5].

Gradual deterioration of the immune system is a well-known phenomenon of mammalian aging. Consequently, the senescent immune system fails to protect the body against disease, and makes unable to distinguish between self and non-self antigens in the body [6]. Therefore, age-associated changes of the immune system including innate and adaptive immunity have been studied as an important risk factor, and a hallmark of aging and age-related diseases in humans [7]. Also, several anti-aging agents such as rapamycin, aspirin, curcumin are well known to affect immune function [8]. Hematopoietic stem cells (HSCs) give rise to multipotent progenitor cells including lymphoid and myeloid cells that involve in both innate and adaptive immunity [9]. The decline of proliferative potential of these HSCs was observed in conjunction with the advanced age of cell donors [10]. Moreover, total lymphocyte numbers are decreased with age, and the proportion of lymphocyte subset changes in the peripheral blood with age in humans [11]. In particular, the ratio of CD4+ T-lymphocytes to CD8+ T-lymphocytes (CD4+/CD8+ ratio) has been used as a marker of age-associated disease in several studies, including human immunodeficiency virus (HIV)-infected patients, Singapore Longitudinal Ageing Study cohort, and elderly with frailty syndrome [12-14]. In elderly dogs, the decline of immune functionality and responsiveness to vaccination was clearly observed [15]. Consequently, old dogs have poorer defense against infection and are susceptible to autoimmune and immunodeficiency diseases. In 2001, Greeley et al [16] tracked 23 Labrador Retriever dogs from birth to death to identify age-dependent changes in various lymphocyte subsets of the peripheral blood. They found that the absolute numbers of total lymphocytes, including CD3+, CD4+ and CD8+ cells, decreased over time [16]. In particular, the proportion of CD4+ cells was decreased while the proportion of CD8+ cells was increased with age, along with maintaining the proportion of CD3+ cells [17]. To date, many studies have shown a decrease in the

CD4+/CD8+ ratio [15,17].

All living organisms suffer from frailty due to a loss of muscle strength and endurance as they get older. Therefore, aging and muscle seem to be closely related to each other. The skeletal muscle cells produce and release cytokines, termed as myokines, which play active autocrine, paracrine, or endocrine roles [18]. Five representative myokines play important roles in muscles and whole body [18,19]; these include interleukin (IL)-6, leukemia inhibitory factor (LIF), IL-8, IL-15, and growth differentiation factor 8 (GDF8). IL-6 is an exercise mediator with immunoregulatory, anti-inflammatory, metabolic, and hypertrophic effects [19]. IL-6 is released into the blood circulatory system after muscle contraction [20]. LIF, which belongs to the IL-6 cytokine superfamily, has diverse biological functions including platelet formation, bone formation, adipocyte lipid transport, neuronal formation, and muscle satellite cell proliferation [21]. IL-8, primarily produced by the macrophage and endothelial cell, exerts strong chemotactic activity towards neutrophils [20] and acts as an angiogenic factor [22]. IL-15 is known as an anabolic cytokine involving in the resistance exercise for muscle mass in humans [23]. IL-15 is also involved in lipid metabolism and fat oxidation [24]. GDF8, known as myostatin, plays a regulatory role in skeletal muscle growth [25]. When the GDF8 was overexpressed in the transgenic mice, muscle mass was decreased [26]. GDF8 is also involved in regulation of adipose tissue growth [25].

Recently, GDF11, a transforming growth factor β superfamily member, has been suggested as a putative anti-aging factor in mice [27]. GDF11 is a circulating factor that prevents cardiovascular disease, cardiac hypertrophy, and cardiovascular death. Indeed, it was confirmed that GDF11 concentration was lower in older mice [27]. Conversely, in a recent study in humans, GDF11 was found to be involved in the risk of comorbidity, frailty, and surgery in the elderly with cardiovascular disease [28].

Aging causes physical frailty, and frailty is associated with aged muscles [29]. Aged muscles show structural distortions [30] as well as alterations of plasma factors, including the aforementioned myokines, which might induce systemic aging [31]. Immunosenescence, including alterations of T-lymphocytes, has been suggested as a major hallmark of aging [32]. Based on these strong implication of plasma markers related to

the aging process, we measured myokine levels and the proportion of T-lymphocytes in healthy large dogs to test development of an age prediction model.

MATERIALS AND METHODS

1. Large military dogs

While we were not able to prospectively calculate an optimal sample size for this study, we utilized existing manuscripts to justify a target number: 10 dogs in [33], 25 dogs in [34], and 23 dogs in [16]. Thus, we were able to enroll 27 dogs, which were identically managed under the rules of the Military Working Dog Training Center (Chuncheon, Korea). The age range of these dogs was from 1 to 12 years old. Our dog population was maintained in the identical environment, especially in terms of regularity, especially under the military conditions. Their diet contained all essential nutrients requirements, and the proportion of nutrients contained more than 25% crude protein, 10% crude fat, 0.6% Ca, 0.6% P, and less than 4% crude fiber, 10% crude ash, and 10% moisture. All dogs were fed under an identical schedule. Among the 27 dogs, 18 belonged to the breed German Shepherd Dog (GSD) (11 males and 7 females), and 9 belonged to the breed Belgian Malinois (BGM) (5 males and 4 females).

2. Ethical statement

All blood samples were obtained in an ethical manner following guidelines for animal health and welfare at a vet clinic. Advance approval was acquired from the Institutional Animal Care and Use Committee of the National Institute of Animal Science, of the Rural Development Administration, of South Korea.

3. Body size measurement

For abdominal circumference (AC) measurement, the circumference of the narrowest region of the abdomen between the lumbar vertebrae and the pelvis was measured when the dog was in a standing position. For thoracic circumference (TC) measurements, the circumference of the widest region of the thorax between the scapula and thoracic vertebrae was measured.

4. Purification of peripheral blood mononuclear cells (PBMCs) and plasma

Peripheral blood was obtained by venipuncture and transfer into a K₂EDTA coated vacutainer tube (Bec-

ton Dickinson, Plymouth, UK) immediately. Whole blood was diluted 1:1 with phosphate buffered saline (PBS; pH 7.0–7.2) (v:v) and carefully layered into a 4-mL Ficoll-paque PLUS (GE Healthcare Life Science, Uppsala, Sweden) containing conical tube. Tubes were centrifuged at 400×g for 30 minutes at room temperature without brake. The supernatant and white layer were transferred into cryotubes as plasma and PBMCs, respectively. Collected PBMCs were fixed with 2% paraformaldehyde for 10 minutes. Plasma and fixed PBMCs were stored in liquid nitrogen and in a -80°C deep freezer, respectively. All the processes were done in 30 minutes from the blood samples to storage.

5. T-lymphocyte analysis

Fixed and frozen PBMCs were quickly thawed at 37°C, and cells were collected by centrifugation at 400×g for 5 minutes. Cells were washed twice with PBS (supplemented with 0.7% bovine serum albumin). Washed cells were then stained with anti-dog CD3:FITC/CD4:RPE/CD8:Alexa Fluor647[®] (Bio-Rad, Munich, Germany) for 15 minutes [35]. Stained cells were analyzed using FACSVerse (BD, NJ, USA).

6. Enzyme-linked immunosorbent assay (ELISA) for humoral factors

We purchased and used canine-specific ELISA kits for IL-6 and IL-8 (RayBiotech, Peachtree Corners, GA, USA), IL-15 and LIF (USBiological, Salem, MA, USA), GDF11 (Blue Gene, Shanghai, China) and GDF8 (EIAab, Wuhan, China). We measured these humoral factors in accordance with the manufacturer's application notes. For standard curves, we used a four parameter logistic (4-PL) equation:

$$y = B + \frac{(A - B)}{1 + (X/C)^D} \quad (1)$$

In this equation, the dependent variable y represents optical density (450 nm), independent variable x represents the concentration of the detected protein, A is the minimum value that can be obtained, B is the maximum value that can be obtained, C is the point of inflection, and D is the Hill's slope of the standard curve. For measuring the absorbance, we used a multi-detection microplate reader (Hidex, Turku, Finland). We applied samples randomly at different positions of a 96-well plate and measured the average values of the two independent technical replicates.

7. Statistical analysis and age prediction models

For correlation analysis, Pearson's correlation coefficients and p-values were calculated using the `cor.test` function in the R stats package.

Simple and multiple linear regression models were constructed to test the fitness of the factors for predicting age using the `lm` function in the R stats package. We used the following formula:

$$y_i = \alpha + \beta x_i + \varepsilon_i \quad (2)$$

$$y_i = \alpha + \beta_1 x_{1i} + \beta_2 x_{2i} + \varepsilon_i \quad (3)$$

$$y_i = \alpha + \beta_1 x_{1i}^2 + \beta_2 x_{1i} + \beta_3 x_{2i}^2 + \beta_4 x_{2i} + \varepsilon_i \quad (4)$$

In the simple linear regression model (formula 2), each subject (i -th subject, $1 \leq i \leq n=27$), the dependent variable y represents age, α is the intercept, β represents the coefficient value, and the explanatory variable x_i is the measured parameter (either the CD4+/CD8+ ratio or the concentration of GDF8) that is considered as a fixed effect. Finally, general error (ε) was added to the equation. For the multiple linear regression model (formula 3) of each subject, β_1 and β_2 are coefficient values, and phenotypic values are x_1 =CD4+/CD8+ ratio and x_2 =GDF8. Moreover, the quadratic equation was constructed to determine the nonlinear relationship between the explanatory and dependent variables, as indicated in formula 4. The logistic parameters used in formula 4 are identical to those employed in formula 3.

To apply idiosyncratic variation of a random effect and nonlinear relationship to a multiple linear regression model, a quadratic linear mixed-effect model was formed using the `lme` function with the restricted maximum likelihood option in the R `nlme` package. The basic formula of this model is as follows:

$$y_i = \alpha + \beta_1 x_{1i}^2 + \beta_2 x_{1i} + \beta_3 x_{2i}^2 + \beta_4 x_{2i} + Z_i u_i + \varepsilon_i \quad (5)$$

$u \sim N_p(0, D), \quad \varepsilon \sim N_n(0, R_i)$

where the age of each subject (i -th subject, $1 \leq i \leq n$; $\max n=27$) is substituted for y . Explanatory variables, x_1 =CD4+/CD8+ ratio and x_2 =GDF8 were added as fixed effects; $\beta_1, \beta_2, \beta_3,$ and β_4 represent the coefficient values. Z is the $n \times p$ design matrix for covariates of random effects, and u is the corresponding $p \times 1$ vector

of the unobservable random effect coefficients. In this study, breeds and sex were replaced as random effects with nested classification as male and female within the GSD and BGM, respectively ($p=4$; breed/sex in R syntax). D is the $p \times p$ covariance matrix for random effects, and ε was added as the random error. R is an $n \times n$ covariance matrix for the errors.

We used two artificial intelligence models. The first one is a supporting vector regression model using a support vector machine (SVM) with SVM function using the `e1071` package in R. In this model, the CD4+/CD8+ ratio and GDF8 were set as fixed effects, using default options, including radial basis kernel, and the parameters were optimized to cost (cost for false classification)=16, gamma (kernel width)=0.1, and epsilon (tolerance of termination criterion)=0.9 by grid-based parameter tuning function. To calculate the predicted age, the `predict` function in the R stats package was used.

The second model we used was an artificial neural network (ANN) model. We applied this model using IBM SPSS modeler 18.0 (SPSS Inc., Chicago, IL, USA). The CD4+/CD8+ ratio and GDF8 data were used in this model, running multi-layer perceptron with back-propagation and radial basis function (RBF) strategies with the default setting.

RESULTS

1. Body weight has a strong correlation with abdominal circumference and thoracic circumference, but not with age

Body composition of dogs are known to change through aging [36], and one recognized way to estimate body composition is through measuring AC and TC of the dog [37]. To identify the relationship between estimated body composition and age, we measured weight, AC, and TC. We found that AC and TC showed a strong and significant correlation with weight (AC: $r=0.83, p=6 \times 10^{-8}$; TC: $r=0.84, p=3 \times 10^{-8}$) (Fig. 1A, 1B). Next, we analyzed the correlation between the AT ratio and weight. The AT ratio also showed a significant correlation ($r=0.60, p=9 \times 10^{-4}$), but was less robustly than either AC or TC alone (Fig. 1C). These results showed that AC and TC could independently serve as a useful indicator of weight. To investigate the effect of age on weight, we analyzed the relationship between weight and age (Fig. 1D) and found that weight had no correlation




Fig. 1. Weight, abdominal circumference (AC), thoracic circumference (TC), and age. Body size, including AC and TC, showed a clear positive correlation with body weight (A, B). However, their ratio (AC/TC) showed a less positive correlation than the individual body weight parameters (C). Overall, body weight did not correlate with age (D).

with age ($r=-0.07$, $p=0.712$) in this study group.

2. CD4+/CD8+ ratio shows a significant correlation with age

To identify whether the lymphocyte subset is a suitable marker for aging in a canine model, we measured the proportion of helper (CD4+) and cytotoxic (CD8+) T-lymphocytes in the peripheral blood of 27 healthy dogs. We found that CD4+ and CD8+ cells insignificantly decreased ($r=-0.29$, $p=0.144$) and increased ($r=0.30$, $p=0.127$) in conjunction with dog age, respectively (Fig. 2A). On the other hand, the CD4+/CD8+ ratio was significantly decreased with age ($r=-0.46$, $p=0.017$) (Fig. 2A). Sex difference is well known to be associated with immunosenescence and longevity [38]. We examined whether the CD4+/CD8+ ratio is applicable to both males and females. The CD4+/CD8+ ratio was significantly decreased with age in male dogs ($r=-0.56$, $p=0.023$), but not in female dogs ($r=-0.27$, $p=0.415$) (Fig. 2B). The results indicate that the correlation between the CD4+/CD8+ ratio and age could be influenced by sex. We also observed that the lymphocyte subset was differently distributed in the peripheral blood of dogs by breed. The CD4+/CD8+ ratio showed a strong correlation with age in the breed BGM ($r=-0.80$, $p=0.009$), but not for GSD ($r=-0.27$, $p=0.273$) (Fig. 2C).

3. GDF8 levels in plasma significantly correlate with age

To discover humoral markers for age, we measured the plasma levels of five representative myokines, GDF8, IL-8, LIF, IL-15 and IL-6 levels, and GDF11. The level of GDF8 showed a significant correlation with age ($r=0.52$, $p=0.005$) (Fig. 3A). We found that this correlation was dependent on sex. The correlation was significant in males ($r=0.78$, $p=3\times 10^{-4}$), but not significant in females ($r=-0.07$, $p=0.848$) (Fig. 3B). In addition, the level of GDF8 was also dependent on breed and showed a more robust correlation significance in BGM ($r=0.71$, $p=0.032$) but no significant correlation in GSD ($r=0.25$, $p=0.314$) (Fig. 3C). With the exception of GDF8, other humoral factors did not show any correlation with age (Fig. 4). In the case of IL-6 and GDF11, we could only detect them in a few plasma samples, including 10 samples for IL-6 and 11 samples for GDF11 (Table 1, Fig. 4B, 4E).

4. Age prediction model using CD4+/CD8+ ratio and GDF8 level

To develop an age prediction model, we applied various linear models using forward selection of factors as explanatory variables including the CD4+/CD8+ ratio and GDF8 level. First, we tested how these two markers individually fit in a prediction model. We




Fig. 2. Effect of age on T-lymphocyte subset in canine peripheral blood. Immunophenotyping of canine peripheral blood mononuclear cells was analyzed using a flow cytometry. Correlation of anti-CD4 positive cells (helper T cells), anti-CD8 positive cells (cytotoxic T cells), and the CD4+/CD8+ ratio with age (A). Correlation of CD4+/CD8+ ratio with age by sex (B) and by breed (C). GSD: German Shepherd Dog, BGM: Belgian Malinois.

used a simple linear model using each of these factors. Although these two markers showed a significant correlation with chronological age, simple linear model parameters poorly predicted the precise age ($r=0.46$, $p=0.016$ for CD4+/CD8+ ratio; $r=0.52$, $p=0.005$ for GDF8) (Fig. 5A, 5B). Based on these simple linear models as a reference, we tested advanced multiple linear models using two fixed effects, the CD4+/CD8+ ratio and the GDF8 level. This trial showed improved age prediction ($r=0.62$, $p=6 \times 10^{-4}$) compared to reference models (Fig. 5C). Also, we tested several multiple linear models based on various combinations of measurement (Fig. 5D-5G). Finally, we tried a quadratic linear model for

age prediction using the two fixed effect parameters, CD4+/CD8+ ratio and GDF8 level (Fig. 5H). In our data, we already found that sex and breed influenced the age prediction power of the CD4+/CD8+ ratio (Fig. 2B, 2C) and GDF8 level (Fig. 3). To implement the effects of these variables, we set sex and breed as random effects in our quadratic linear model. These random effects were structured as a hierarchical order of male and female within each breed in a quadratic linear mixed model (see details in MATERIALS AND METHODS). Interestingly, by using this age prediction model, we identified the best fit ($r=0.77$, $p=3 \times 10^{-6}$) for chronological age in our study group (Fig. 5I). Recently,




Fig. 3. Plasma level of GDF8. Among the six humoral factors, only GDF8 showed a significant correlation with age (A). Scatter plot of GDF8 with sex (B) and breed (C) effects. GDF: growth differentiation factor, GSD: German Shepherd Dog, BGM: Belgian Malinois.




Fig. 4. Plasma level of leukemia inhibitory factor (LIF), and interleukin (IL)-6, IL-8, IL-15, and growth differentiation factor 11 (GDF11). Scatter plots of LIF (A), IL-6 (B), IL-8 (C), IL-15 (D), and GDF11 (E) showed no significant correlation with age.

machine learning algorithms have been heavily applied in biological data analysis. We also implemented some popular machine learning algorithms including

ANN and SVR for age prediction model. We obtained equivalent results using ANN-RBF ($r=0.72$, $p=2 \times 10^{-5}$) (Fig. 6A) and SVR ($r=0.76$, $p=3 \times 10^{-6}$) (Fig. 6B) compared

Table 1. Measurements of six humoral factors using ELISA

Breed	Sex	Age (y)	GDF8 1× (ng/mL)	IL-15 1× (pg/mL)	LIF 1× (pg/mL)	IL-8 1× (pg/mL)	IL-6 1× (pg/mL)	GDF11 1× (pg/mL)
BGM	Male	2.5	2.02	169	52.2	1,480	68	13.8
GSD	Male	2.9	1.15	379	NA	134	86	33.4
GSD	Male	3.3	1.92	355	43.4	62	214	NA
BGM	Male	2.8	1.93	136	129.0	2,400	214	NA
BGM	Male	2.9	1.05	147	NA	60	656	NA
GSD	Male	6.4	4.52	431	62.2	220	684	33.0
BGM	Male	2.8	2.28	177	NA	68	1,332	NA
GSD	Male	3.2	2.48	321	NA	50	2,028	NA
GSD	Male	6.4	1.43	235	68.2	340	11,634	NA
GSD	Male	3.6	2.30	395	414.2	78	NA	46.4
GSD	Male	5.1	1.52	218	196.4	100	NA	15.2
GSD	Male	5.6	2.10	209	39.2	132	NA	NA
GSD	Male	5.5	1.72	318	36.2	500	NA	NA
BGM	Male	12.1	6.22	204	157.4	540	NA	NA
GSD	Male	7.9	3.70	492	126.0	620	NA	93.0
GSD	Male	5.7	2.04	323	98.4	820	NA	52.2
GSD	Female	7.8	2.86	240	55.0	480	70	27.6
GSD	Female	3.7	1.82	229	682.0	11	NA	10.6
GSD	Female	4.9	2.66	219	55.4	52	NA	NA
BGM	Female	8.0	0.85	347	NA	52	NA	NA
BGM	Female	2.0	1.38	686	NA	84	NA	NA
GSD	Female	5.6	2.44	183	40.6	88	NA	NA
GSD	Female	4.2	1.60	211	49.0	132	NA	NA
GSD	Female	8.3	1.20	216	NA	154	NA	NA
GSD	Female	8.4	1.86	320	36.0	162	NA	12.8
BGM	Female	1.9	1.90	163	52.4	840	NA	10.4
BGM	Female	4.6	2.38	287	53.8	NA	NA	NA

GDF: growth differentiation factor, IL: interleukin, LIF: leukemia inhibitory factor, BGM: Belgian Malinois, GSD: German Shepherd Dog, NA: not available.

to that obtained using the quadratic linear mixed model.

DISCUSSION

In this study, to identify aging-associated factors in dogs, we measured body size (weight, AC, TC, and AT ratio), the proportion of T-lymphocytes (CD4+, CD8+, and the CD4+/CD8+ ratio), the concentration of myokines (IL-8, LIF, IL-15, IL-6, and GDF8), and the concentration of an anti-aging marker, GDF11 (Table 2). In previous studies, body fat showed a correlation in the ratio of abdominal width to thoracic width [37]. In human, elderly people showed more tendency of having visceral fat than young people [39]. In addition, elderly people prone to develop obesity because of their weakened muscles, collectively called sarcopenic obesity [40].

According to these observations, we also would like to see whether there are any relationships between body composition and chronological ages in our dog study. Therefore, we measured AC, TC, and weight of the dogs to estimate body composition. We found that both AC and TC significantly correlated with weight. Also, we found a significant correlation between the AT ratio and weight. However, there was no correlation between AT ratio with age. Therefore, we could not find relationship between estimated body composition (AT ratio) with age in our study.

Helper (CD4+) and cytotoxic T (CD8+) cells play key roles in white blood cells as part of the adaptive immune system. Cytotoxic T cells recognize a specific antigen through the T-cell receptor and directly destroy infected target cells, tumor cells, and damaged cells. On the other hand, helper T cells activate other immune




Fig. 5. Age prediction using linear regression models. Based on the chronological ages, age prediction was performed using given parameters. The black solid lines indicate second-order polynomial fits for the age prediction, and the broken lines represent the isometric reference for perfect prediction of age. Simple linear models were performed using the CD4+/CD8+ ratio (A) and GDF8 (B). Multiple linear models were performed based on the various combinations of fixed effects without considering their interaction including CD4/CD8+GDF8 (C), CD4/CD8+sex+weight (D), CD4/CD8+sex+weight+AC+TC (E), GDF8+sex+weight+AC+TC (F), CD4/CD8+GDF8+sex+weight+AC+TC (G). A quadratic multiple linear model was performed using the CD4/CD8 ratio and GDF8 (H). A quadratic mixed model was performed using the same two parameters with the breed and sex as random effects (I). Symbols: circle for BGM female; triangle for BGM male; square for GSD female; plus for GSD male. GDF: growth differentiation factor, AC: abdominal circumference, TC: thoracic circumference, BGM: Belgian Malinois, GSD: German Shepherd Dog.

cells such as B cells, cytotoxic T cells, and macrophages by releasing cytokines. Furthermore, inversion of the CD4+/CD8+ ratio ($CD4+/CD8+ < 1$) is suggested as a hallmark of immunosenescence and age-related disease in several studies including HIV-infected patients, Singapore Longitudinal Ageing Study cohort [13] and elderly with frailty syndrome [12,14]. In our study, we also examined CD4+ and CD8+ T cells in the peripheral blood of dogs at various ages. From these results, we found that the CD4+/CD8+ ratio was significantly

correlated with dog age (Fig. 2A). However, in the case of the BGM alone (Fig. 2C), there was no correlation between CD4+/CD8+ ratio and the dog age. Although the usefulness of the lymphocyte subset was influenced by sex and breed, the CD4+/CD8+ ratio could be a useful prediction marker of age in certain cases. Several studies on lymphocyte subset analysis gave controversial experimental results in dog models. Some results showed that the levels of CD4+ and CD8+ as well as the CD4+/CD8+ ratio was a significantly correlated




Fig. 6. Age prediction using machine learning algorithms. Age was predicted by machine learning algorithms. Predicted ages were calculated by artificial neural network (ANN) (A) and supporting vector regression (SVR) (B) algorithms. Symbols: circle for BGM female; triangle for BGM male; square for GSD female; plus for GSD male. GDF: growth differentiation factor, AC: abdominal circumference, TC: thoracic circumference, BGM: Belgian Malinois, GSD: German Shepherd Dog.

with dog age [17]. And there were some inconsistent reports including a decreased correlation of CD4+/CD8+ ratio to age [41] and no correlation between them [16] from other research groups. However, our findings showed that the lymphocyte subset distribution was changed in association with dog age (Fig. 2).

The level of GDF8 (myostatin) in the plasma of male canines significantly increased in conjunction with age (Fig. 3B). Moreover, in a study of aging in humans, GDF8 concentration was found to increase as a result of age in 70 men [28]. In both species, GDF8 level could be a reasonable marker of age in males. In contrast, in 70 women [28] and female canines (our study), GDF8 levels were not significantly changed as a result of age (Fig. 3B). GDF8 gene mutation in mice is related to a hypermuscular phenotype, which suggests that GDF8 inhibits skeletal muscle growth [26]. These findings suggest the possibility of muscle weakness when male canines get older. In fact, several studies have already shown that increased GDF8 levels are correlated with loss of muscle mass leading to age-associated sarcopenia [42], glucocorticoid-induced muscle atrophy [43], Duchenne muscular dystrophy [44], and indoxyl sulfate-induced muscle atrophy [45]. All together, these studies suggest that regulation of GDF8 concentration might be important for maintaining and recovering muscle function of individuals not only in old age but also at all ages.

IL-8 and IL-15 locally direct autocrine or paracrine activity in muscles; however, these myokines are rarely released from muscles to the circulatory system [24]. Thus, a change in IL-8 and IL-15 concentration in the plasma might not be detected. However, in our analysis of GSDs, plasma IL-8 levels were clearly detected and

increased in association with aging. Thus, we speculate that IL-8 can be released into the circulatory system in at least some dog breeds, such as the GSD. LIF has a wide variety of functions, included platelet formation, bone formation, adipocyte lipid transport, neuronal formation, and muscle satellite cell proliferation [21]. Therefore, LIF concentration in the plasma may be affected by several factors, not simply by age. We could not observe a change in the plasma LIF concentration as a result of age.

IL-6 is mainly produced in contracting muscles, and its levels are reduced at post-exercise [19,20]. We could not detect IL-6 in our experimental subjects. GDF11 is suggested to be an anti-aging humoral factor [27,28]. In addition, higher levels of GDF11 are associated with decreased risk of cardiovascular events and death in a mice study [27]. However, in humans, GDF11 is involved in sex-specific pathological weakness in older men with cardiovascular disease [28]. In our results, we could not easily detect GDF11 in healthy canine subjects.

To identify potential markers for certain phenotypes, statistical approaches including regression models have been applied [46]. In addition, statistical models for aging have been constructed based on gene expression [47], microRNA [47], proteins [47], metabolites [48], epigenetic markers [49], verbal intelligence [50], and facial patterns [51]. Although various humoral factors have been suggested for age prediction in humans [52], no studies have examined this in canines to the best of our knowledge. Here, we measured specific plasma factors from the peripheral blood of canines, which is easier to non-invasively obtain than other tissue samples. We found that the T-lymphocytes ratio (CD4+/

Table 2. Summary of measurements between male and female

Sex	Statistics	Age (y)	GDF8 1x (ng/mL)	IL-15 1x (pg/mL)	LIF 1x (pg/mL)	IL-8 1x (pg/mL)	IL-6 1x (pg/mL)	GDF11 1x (pg/mL)	Abdominal circumference (cm)	Thoracic circumference (cm)	AC/TC (ratio)	CD8+ (%)	CD4+ (%)	CD4/CD8 (ratio)
Male	Mean	4.9	2.4	282	89	475	1,057	17.9	57	77	0.74	24	44	2.17
	Standard deviation	2.5	1.3	109	105	641	2,879	27.1	5.3	3.9	0.03	11.4	10.0	0.94
Female	Mean	5.4	1.9	282	93	187	6	5.6	51	72	0.71	23	48	2.26
	Standard deviation	2.4	0.6	145	197	253	21	9.0	5.3	5.5	0.05	6.5	6.5	0.69

GDF: growth differentiation factor, IL: interleukin, LIF: leukemia inhibitory factor, AC: abdominal circumference, TC: thoracic circumference.

CD8+) as well as myokine, GDF8, and IL-8 levels were significantly correlated with canine age. Moreover, the significance of the correlation was influenced by both sex and breed difference. Consequently, we could form various models (*e.g.*, linear models, quadratic linear models, ANN, and SVR) of age prediction that were composed of two variables, the CD4+/CD8+ ratio and GDF8 concentration, as fixed effects. In addition, both breed and sex were substituted as random effects to form linear mixed models. We confirmed the fitness of age prediction by established models using a correlation test between chronological age and model-predicted age (Fig. 5, 6). The best model to fit the observed ages was the quadratic linear mixed model (Fig. 5I).

Dogs offer an opportunity to serve as a translational animal model of people, particularly given that they share similar chronic conditions such as obesity, arthritis, diabetes, and age-specific mortality patterns related to metabolic dysfunction and cancer [53]. Relevant to the current work, immunosenescence is one of the major aging traits occurring in both dog and human [6,15,16]. Levels of total lymphocytes and their proportions also seem to be important indicators of age in both species [11,12,15-17]. Furthermore, some myokines, IL-6, TNF- α and NCP-1, have been identified as common players in inflammation of both human and dog [24,54,55]. Given this, the chronological age prediction model established in this work may hold value in studying the influence of age-related treatment attempts on shifting parameters relevant to biological aging. To increase our sample size and include additional breeds in our work, we are currently repeating this work in small dogs and developing more sensitive ELISA methodology for use with smaller sample volumes. Additionally, we recognize that longitudinal *versus* single peripheral blood samples may be needed to determine whether circulating myokines levels fluctuate throughout the day. Finally, future work can be done to characterize T lymphocytes beyond CD4+/CD8+.

CONCLUSIONS

In conclusion, the CD4+/CD8+ ratio and GDF8 concentration in canine blood were useful factors in developing an age prediction model. However, genetic background and sex increased the complexity and limited the ability to predict age (the best $r=0.77$). Although

the aging process varies among individual canines having a diverse distribution of biological age, our age prediction model could be very promising to estimate both biological age and chronological age.

Conflict of Interest

The authors have nothing to disclose.

Funding

This work was supported by the Cooperative Research Program for the Agriculture Science & Technology Development under Grant [Project No. PJ01195001], Rural Development Administration, Republic of Korea, Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2020R1F1A1075785), and a Korea University Grant.

Author Contribution

Conceptualization: CKL, BHC. Investigation: HJL, SJH, SSK, YYK. Methodology: HJL, SJH, SSK, YYK, KMC, SHS, MJL, SYH. Project administration: CKL, BHC. Resources: BHC, KP, YJ, HS. Supervision: CKL. Visualization: HJL, SJH, SSK, YYK. Writing – original draft: HJL, SJH, SSK, YYK, CKL. Writing – review & editing: CKL, BHC, KMC.

Data Sharing Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

REFERENCES

- Pagano TB, Wojcik S, Costagliola A, De Biase D, Iovino S, Iovane V, et al. Age related skeletal muscle atrophy and up-regulation of autophagy in dogs. *Vet J* 2015;206:54-60.
- Radakovich LB, Pannone SC, Truelove MP, Olver CS, Santangelo KS. Hematology and biochemistry of aging-evidence of "anemia of the elderly" in old dogs. *Vet Clin Pathol* 2017;46:34-45.
- Li Y, Deeb B, Pendergrass W, Wolf N. Cellular proliferative capacity and life span in small and large dogs. *J Gerontol A Biol Sci Med Sci* 1996;51:B403-8.
- Fleming JM, Creevy KE, Promislow DE. Mortality in North American dogs from 1984 to 2004: an investigation into age-, size-, and breed-related causes of death. *J Vet Intern Med* 2011;25:187-98.
- Galis F, Van der Sluijs I, Van Dooren TJ, Metz JA, Nussbaumer M. Do large dogs die young? *J Exp Zool B Mol Dev Evol* 2007;308:119-26.
- Castelo-Branco C, Soveral I. The immune system and aging: a review. *Gynecol Endocrinol* 2014;30:16-22.
- Fülöp T, Dupuis G, Witkowski JM, Larbi A. The role of immunosenescence in the development of age-related diseases. *Rev Invest Clin* 2016;68:84-91.
- Yalcin G, Lee CK. The discovery of druggable anti-aging agents. *Ann Geriatr Med Res* 2020;24:232-42.
- Kondo M. Lymphoid and myeloid lineage commitment in multipotent hematopoietic progenitors. *Immunol Rev* 2010;238:37-46.
- Gruver AL, Hudson LL, Sempowski GD. Immunosenescence of ageing. *J Pathol* 2007;211:144-56.
- Huppert FA, Solomou W, O'Connor S, Morgan K, Sussams P, Brayne C. Aging and lymphocyte subpopulations: whole-blood analysis of immune markers in a large population sample of healthy elderly individuals. *Exp Gerontol* 1998;33:593-600.
- Serrano-Villar S, Moreno S, Fuentes-Ferrer M, Sánchez-Marcos C, Avila M, Sainz T, et al. The CD4:CD8 ratio is associated with markers of age-associated disease in virally suppressed HIV-infected patients with immunological recovery. *HIV Med* 2014;15:40-9.
- Ng TP, Camous X, Nyunt MSZ, Vasudev A, Tan CTY, Feng L, et al. Markers of T-cell senescence and physical frailty: insights from Singapore longitudinal ageing studies. *NPJ Aging Mech Dis* 2015;1:15005.
- Rizka A, Setiati S, Harimurti K, Sadikin M, Mansur IG. Effect of alfalcidol on inflammatory markers and T cell subsets in elderly with frailty syndrome: a double blind randomized controlled trial. *Acta Med Indones* 2018;50:215-21.
- HogenEsch H, Thompson S, Dunham A, Ceddia M, Hayek M. Effect of age on immune parameters and the immune response of dogs to vaccines: a cross-sectional study. *Vet Immunol Immunopathol* 2004;97:77-85.
- Greeley EH, Ballam JM, Harrison JM, Kealy RD, Lawler DE, Segre M. The influence of age and gender on the immune system: a longitudinal study in Labrador Retriever dogs. *Vet Immunol Immunopathol* 2001;82:57-71.
- Watabe A, Fukumoto S, Komatsu T, Endo Y, Kadosawa T. Alterations of lymphocyte subpopulations in healthy dogs with aging and in dogs with cancer. *Vet Immunol Immunopathol* 2011;142:189-200.
- Pedersen BK, Febbraio MA. Muscles, exercise and obesity: skeletal muscle as a secretory organ. *Nat Rev Endocrinol*

- 2012;8:457-65.
19. Nielsen S, Pedersen BK. Skeletal muscle as an immunogenic organ. *Curr Opin Pharmacol* 2008;8:346-51.
 20. Pedersen BK, Febbraio MA. Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiol Rev* 2008;88:1379-406.
 21. Metcalf D. The unsolved enigmas of leukemia inhibitory factor. *Stem Cells* 2003;21:5-14.
 22. Baggiolini M. Chemokines in pathology and medicine. *J Intern Med* 2001;250:91-104.
 23. Riechman SE, Balasekaran G, Roth SM, Ferrell RE. Association of interleukin-15 protein and interleukin-15 receptor genetic variation with resistance exercise training responses. *J Appl Physiol* (1985) 2004;97:2214-9.
 24. Nielsen AR, Pedersen BK. The biological roles of exercise-induced cytokines: IL-6, IL-8, and IL-15. *Appl Physiol Nutr Metab* 2007;32:833-9.
 25. Allen DL, Cleary AS, Speaker KJ, Lindsay SF, Uyenishi J, Reed JM, et al. Myostatin, activin receptor IIb, and follistatin-like-3 gene expression are altered in adipose tissue and skeletal muscle of obese mice. *Am J Physiol Endocrinol Metab* 2008;294:E918-27.
 26. Reisz-Porszasz S, Bhasin S, Artaza JN, Shen R, Sinha-Hikim I, Hogue A, et al. Lower skeletal muscle mass in male transgenic mice with muscle-specific overexpression of myostatin. *Am J Physiol Endocrinol Metab* 2003;285:E876-88.
 27. Olson KA, Beatty AL, Heidecker B, Regan MC, Brody EN, Foreman T, et al. Association of growth differentiation factor 11/8, putative anti-ageing factor, with cardiovascular outcomes and overall mortality in humans: analysis of the Heart and Soul and HUNT3 cohorts. *Eur Heart J* 2015;36:3426-34.
 28. Schafer MJ, Atkinson EJ, Vanderboom PM, Kotajarvi B, White TA, Moore MM, et al. Quantification of GDF11 and myostatin in human aging and cardiovascular disease. *Cell Metab* 2016;23:1207-15.
 29. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al.; Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146-56.
 30. Doran P, Donoghue P, O'Connell K, Gannon J, Ohlendieck K. Proteomics of skeletal muscle aging. *Proteomics* 2009;9:989-1003.
 31. Demontis F, Piccirillo R, Goldberg AL, Perrimon N. The influence of skeletal muscle on systemic aging and lifespan. *Aging Cell* 2013;12:943-9.
 32. Chalan P, van den Berg A, Kroesen BJ, Brouwer L, Boots A. Rheumatoid arthritis, immunosenescence and the hallmarks of aging. *Curr Aging Sci* 2015;8:131-46.
 33. Judge SJ, Yanagisawa M, Sturgill IR, Bateni SB, Gingrich AA, Foltz JA, et al. Blood and tissue biomarker analysis in dogs with osteosarcoma treated with palliative radiation and intratumoral autologous natural killer cell transfer. *PLoS One* 2020;15:e0224775.
 34. da C Monini A, Gandini LG Jr, Vianna AP, Martins RP, Jacob HB. Tooth movement rate and anchorage lost during canine retraction: a maxillary and mandibular comparison. *Angle Orthod* 2019;89:559-65.
 35. Pinto LA, Trivett MT, Wallace D, Higgins J, Baseler M, Terabe M, et al. Fixation and cryopreservation of whole blood and isolated mononuclear cells: influence of different procedures on lymphocyte subset analysis by flow cytometry. *Cytometry B Clin Cytom* 2005;63:47-55.
 36. Speakman JR, van Acker A, Harper EJ. Age-related changes in the metabolism and body composition of three dog breeds and their relationship to life expectancy. *Aging Cell* 2003;2:265-75.
 37. Gant P, Holden SL, Biourge V, German AJ. Can you estimate body composition in dogs from photographs? *BMC Vet Res* 2016;12:18.
 38. Caruso C, Accardi G, Virruso C, Candore G. Sex, gender and immunosenescence: a key to understand the different lifespan between men and women? *Immun Ageing* 2013;10:20.
 39. Demerath EW, Sun SS, Rogers N, Lee M, Reed D, Choh AC, et al. Anatomical patterning of visceral adipose tissue: race, sex, and age variation. *Obesity (Silver Spring)* 2007;15:2984-93.
 40. Stenholm S, Harris TB, Rantanen T, Visser M, Kritchevsky SB, Ferrucci L. Sarcopenic obesity: definition, cause and consequences. *Curr Opin Clin Nutr Metab Care* 2008;11:693-700.
 41. Faldyna M, Levá L, Knötigová P, Toman M. Lymphocyte subsets in peripheral blood of dogs--a flow cytometric study. *Vet Immunol Immunopathol* 2001;82:23-37.
 42. Yarasheski KE, Bhasin S, Sinha-Hikim I, Pak-Loduca J, Gonzalez-Cadavid NE. Serum myostatin-immunoreactive protein is increased in 60-92 year old women and men with muscle wasting. *J Nutr Health Aging* 2002;6:343-8.
 43. Ma K, Mallidis C, Bhasin S, Mahabadi V, Artaza J, Gonzalez-Cadavid N, et al. Glucocorticoid-induced skeletal muscle atrophy is associated with upregulation of myostatin gene expression. *Am J Physiol Endocrinol Metab* 2003;285:E363-71.
 44. Pasteuning-Vuhman S, Boertje-van der Meulen JW, van Putten M, Overzier M, Ten Dijke P, Kielbasa SM, et al. New function of the myostatin/activin type I receptor (ALK4) as a mediator of muscle atrophy and muscle regeneration. *FASEB J* 2017;31:238-55.

45. Enoki Y, Watanabe H, Arake R, Sugimoto R, Imafuku T, Tominaga Y, et al. Indoxyl sulfate potentiates skeletal muscle atrophy by inducing the oxidative stress-mediated expression of myostatin and atrogen-1. *Sci Rep* 2016;6:32084.
46. VanRaden PM. Efficient methods to compute genomic predictions. *J Dairy Sci* 2008;91:4414-23.
47. Somel M, Guo S, Fu N, Yan Z, Hu HY, Xu Y, et al. MicroRNA, mRNA, and protein expression link development and aging in human and macaque brain. *Genome Res* 2010;20:1207-18.
48. Yoshida R, Tamura T, Takaoka C, Harada K, Kobayashi A, Mukai Y, et al. Metabolomics-based systematic prediction of yeast lifespan and its application for semi-rational screening of ageing-related mutants. *Aging Cell* 2010;9:616-25.
49. Xu C, Qu H, Wang G, Xie B, Shi Y, Yang Y, et al. A novel strategy for forensic age prediction by DNA methylation and support vector regression model. *Sci Rep* 2015;5:17788.
50. Multani N, Rudzicz F, Wong WY, Namasivayam AK, van Lieshout P. Random item generation is affected by age. *J Speech Lang Hear Res* 2016;59:1172-8.
51. Geng X, Zhou ZH, Smith-Miles K. Automatic age estimation based on facial aging patterns. *IEEE Trans Pattern Anal Mach Intell* 2007;29:2234-40.
52. Frøyshov HM, Bjørnerem Å, Engstad T, Halvorsen DS. Elevated inflammatory markers predict mortality in long-term ischemic stroke-survivors: a population-based prospective study. *Aging Clin Exp Res* 2017;29:379-85.
53. Hoffman JM, Creevy KE, Franks A, O'Neill DG, Promislow DEL. The companion dog as a model for human aging and mortality. *Aging Cell* 2018;17:e12737.
54. Tidball JG. Inflammatory processes in muscle injury and repair. *Am J Physiol Regul Integr Comp Physiol* 2005;288:R345-53.
55. Ceron JJ, Eckersall PD, Martýnez-Subiela S. Acute phase proteins in dogs and cats: current knowledge and future perspectives. *Vet Clin Pathol* 2005;34:85-99.