



Activated Scavenger Receptor A Promotes Glial Internalization of A β

He Zhang^{1,2,3}, Ya-jing Su³, Wei-wei Zhou¹, Shao-wei Wang¹, Peng-xin Xu¹, Xiao-lin Yu^{1*}, Rui-tian Liu^{1*}

1 National Key Laboratory of Biochemical Engineering, Institute of Process Engineering, Chinese Academy of Sciences, Beijing, China, **2** Tsinghua University School of Medicine, Haidian District, Beijing, China, **3** School of Life Science, Ningxia University, Yinchuan, China

Abstract

Beta-amyloid (A β) aggregates have a pivotal role in pathological processing of Alzheimer's disease (AD). The clearance of A β monomer or aggregates is a causal strategy for AD treatment. Microglia and astrocytes are the main macrophages that exert critical neuroprotective roles in the brain. They may effectively clear the toxic accumulation of A β at the initial stage of AD, however, their functions are attenuated because of glial overactivation. In this study, we first showed that heptapeptide XD4 activates the class A scavenger receptor (SR-A) on the glia by increasing the binding of A β to SR-A, thereby promoting glial phagocytosis of A β oligomer in microglia and astrocytes and triggering intracellular mitogen-activated protein kinase (MAPK) signaling cascades. Moreover, XD4 enhances the internalization of A β monomers to microglia and astrocytes through macropinocytosis or SR-A-mediated phagocytosis. Furthermore, XD4 significantly inhibits A β oligomer-induced cytotoxicity to glial cells and decreases the production of proinflammatory cytokines, such as TNF- α and IL-1 β , in vitro and in vivo. Our findings may provide a novel strategy for AD treatment by activating SR-A.

Citation: Zhang H, Su Y-j, Zhou W-w, Wang S-w, Xu P-x, et al. (2014) Activated Scavenger Receptor A Promotes Glial Internalization of A β . PLoS ONE 9(4): e94197. doi:10.1371/journal.pone.0094197

Editor: Joseph El Khoury, Massachusetts General Hospital and Harvard Medical School, United States of America

Received: January 7, 2014; **Accepted:** March 12, 2014; **Published:** April 9, 2014

Copyright: © 2014 Zhang et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from the National Natural Science Foundation of China (30971012 and 81171014) and the National Science and Technology Major Projects of New Drugs (2012ZX09103301-001). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: rtlui@home.ipe.ac.cn (RTL); yuxiaolin@home.ipe.ac.cn (XLY)

† These authors contributed equally to this work.

Introduction

Alzheimer's disease (AD), the most prominent cause of senile dementia, is characterized by two major neuropathological hallmarks as extracellular deposits of fibrillar beta-amyloid (A β) and intracellular neurofibrillary tangles composed of hyperphosphorylated tau [1]. A β peptides are generated through sequential proteolytic processing of the amyloid precursor protein. Overproduction or insufficient clearance of A β breaks A β homeostasis and leads to overall elevation of its steady-state levels, resulting in A β aggregation and deposition, which contribute to AD pathogenesis [2]. Soluble A β oligomers rather than monomers or fibrils are highly neurotoxic species that may trigger the synapse loss and neuronal damage prior to the formation of senile plaques in the early stages of AD [3]. A β oligomers may induce membrane-related toxicity by perturbation of the membrane fluidity or structure, release of lipids from the neuronal plasma membrane, or formation of ion channels [4]. Moreover, A β aggregates may induce the production of reactive oxygen species (ROS), cytokines, and chemokines, such as TNF- α , IL-1 β , IL-6, MCP-1, IL-8 and NO, leading to an initial neurodegenerative characteristic of AD [5]. Therefore, inhibiting A β production or promoting A β clearance is a rational strategy for the prevention or treatment of AD.

Several methods can be used to clear A β from the brain, such as degrading enzymes, LRP receptor or phagocytosis. Glial phagocytosis and degradation of A β are believed to be key mechanisms of the initial defense of the brain against the toxic accumulation of

A β . With the advancement of AD pathology, microglia are overactivated and lose their intrinsic beneficial function of uptake and degradation of A β . These phenomena result in A β accumulation, release of ROS, cytokines, and chemokines, as well as chronic neuroinflammation, and thus, neurodegeneration in AD [6]. Therefore, modulating microglia activation may be a powerful strategy to clear A β and prevent A β -associated pathological events and neuronal loss.

Previous studies have demonstrated that microglia take up and degrade soluble and fibrillar A β in different ways. Soluble A β is internalized into microglia through constitutive, nonsaturable, and fluid phase macropinocytosis, and the internalized soluble A β is rapidly delivered to the lysosomes by the late endolytic pathway [7]. Fibrillar A β interacts with the cell surface innate immune receptor complex, initiates intracellular signaling cascades, and then stimulates phagocytosis [8,9]. Multiple cell surface receptors, including class A scavenger receptor (SR-A), class B scavenger receptor type I (SR-BI), CD36, α 6 β 1 integrin, CD14, CD47 and toll-like receptors (TLRs), reportedly mediate A β uptake and degradation by microglia [10–13].

The roles of members of SRs in AD are diverse, but SR-A and CD36 are the main receptors responsible for A β clearance. SR-A-mediated interactions with A β promote A β phagocytosis and clearance and downregulate inflammatory gene expression in dendritic cells by suppressing TLR4-induced activation of the transcription factor NF- κ B [14]. Meanwhile, the interactions of CD36 with A β initiate a tyrosine kinase-based signaling cascade,

induce the production of neurotoxins and proinflammatory molecules, and promote the recruitment of microglia to A β deposits in the brain [15,16]. The activation of SR-A can induce ligand internalization and direct the traffic of the endocytic signaling transduction pathways, such as PI3K/NF κ B and mitogen-activated protein kinases (MAPKs)/JNK/p38 [17,18]. Moreover, astrocytes, the main glial cells in the brain, can also engulf and degrade A β , which is mediated by the receptors, such as CD36, CD47 and RAGE [19,20]. Recently, a study on the SR-A-deficient AD transgenic mouse model showed that SR-A deficiency markedly accelerated A β accumulation and led to increased mortality, while pharmacological upregulation of SR-A expression increased A β clearance [21]. Therefore, the activation of some glial receptors such as SR-A may be beneficial for A β clearance and AD treatment.

Materials and Methods

Materials

A β _{1–42} was purchased from American Peptide Company (Sunnyvale, CA, USA). XD4 was synthesized by GL Biochem Co., Ltd. (Shanghai, China). The A β _{1–42} and oA β _{1–42} kits for A β measurement were purchased from IBL Co., Ltd. (Gunma, Japan). Anti-A β antibody 6E10 (monoclonal raised against A β N-terminal) and 4G8 (monoclonal raised against A β _{17–24}) were purchased from Signet Laboratories (Covance, Dedham, MA, USA). Antibodies against SR-A and CD36 were purchased from AbD Serotec (Oxford, UK) and Abcam (Cambridge, UK), respectively. Rabbit anti-His-tag polyclonal antibody was from 4bio Co. (Beijing, China). Fucoidan was obtained from Sigma-Aldrich (St. Louis, MO, USA). Poly-L-lysine was obtained from ScienCell (CA, USA). Transmembrane protein extraction kit was purchased from Merck Millipore (MA, USA). Cell-counting kit (CCK)-8 was purchased from Dojindo (Kumamoto, Japan). Phosphotracer ELISA kit was purchased from Abcam (Cambridge, UK).

Monomer A β _{1–42} (mA β _{1–42}) and Oligomer A β _{1–42} (oA β _{1–42}) Preparation

A β _{1–42} was dissolved in 100% 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP) to a concentration of 1 mg/mL, sonicated in a water bath for 10 min, aliquoted into microcentrifuge tubes, dried under vacuum, and stored at –20°C. To prepare the A β _{1–42} monomer, aliquoted A β _{1–42} films were reconstituted by dimethyl sulfoxide (DMSO) at a concentration of 1 mg/ml, sonicated for 1 min, and immediately added to the serum-free cell media. oA β _{1–42} was prepared as previously described, with a slight modification [22]. Briefly, the stored peptide film was resuspended in DMSO to a concentration of 5 mM and diluted by the cell media or PBS to a final concentration of 100 μ M. The media was then rotated at 4°C for 24 h to form the oligomer, and the oligomer-containing media was centrifuged at 14,000 g at 4°C for 10 min to remove the fibrillar and insoluble aggregated A β _{1–42}. Prior to incubation with XD4, the medium was filtered through 0.22 μ m filters (Millipore, MA, USA) to eliminate the residual fibrillar A β _{1–42}.

Cell Culture

Human microglia and rat astrocytes were purchased from ScienCell Research Laboratories (San Diego, CA, USA). They were isolated from human brain tissue and from day 2 rat cerebral cortex, respectively. Cells were delivered frozen at secondary culture. Upon initiating the sub-culture from cryopreserved cells, the astrocyte medium (ScienCell, No. 1801) and microglia medium (ScienCell, No. 1901) was used. Cells were passaged upon

reaching 90% confluence to new poly-L-lysine coated plates, the media was changed every other day before 50% confluence or every day until 90% confluence. The cells used for A β _{1–42} uptake assay were at passage numbers 2 to 5. The immortalized murine BV-2 microglial cells were maintained in Dulbecco's modified Eagle medium (DMEM) supplemented with 10% FBS (Hyclone, USA) and 1% penicillin–streptomycin (Hyclone, USA). Mouse primary microglia was provided by Pricell Corp. (Wuhan, China). The cells were derived from postnatal day 0–2 pups and isolated from mixed cultures at 14 days in vitro. Upon receipt, primary microglia was directly seeded onto a poly-L-lysine pre-coated six-well plate and cultured overnight with the microglia medium provided by the manufacturer (Pricell, MED-0014). Human astrogloma (U251) cells were maintained in low-glucose DMEM (Hyclone, USA) supplemented with 10% FBS and 1% penicillin–streptomycin (Hyclone, USA), and then seeded at a six-well-plate at passage numbers 5 to 6. All the cells were kept at 37°C in 5% CO₂ atmosphere.

Intracellular A β _{1–42} Quantification

Cells were pretreated with or without fucoidan at the indicated concentrations for 20 min, washed twice by PBS, followed by incubation with 1 μ g/ml mA β _{1–42} for 3 h or 1 μ M oA β _{1–42} for 1.5 h in the presence or absence of the indicated concentration of XD4. For antibody blockade assays, antibodies against SR-A (10 μ g/ml) or CD36 (10 μ g/ml) were incubated with BV-2 microglia for 30 min in serum-free media, and then 1 μ M oA β _{1–42} or preincubated oA β _{1–42} with XD4 was added to the cells and incubated for 1.5 h. The cells were washed extensively with PBS and digested with trypsin (Hyclone, USA) for 3 min. Then the cells were lysed with WIP lysis buffer (Bioss Co., China). The A β _{1–42} levels in the lysates were quantified by ELISA using the A β _{1–42} immunoassay kits. Protein amounts in each experimental group were determined using a bicinchoninic acid protein assay (BCA) kit (Pierce, Rockford, IL, USA). Both mA β _{1–42} and oA β _{1–42} levels internalized by cells were standardized to cell protein concentration and normalized as a percentage of control (treatment of A β _{1–42} alone).

Immunocytochemistry

A confocal microscope was used to localize A β _{1–42} and XD4 in BV-2 cells. Cells were seeded into a 12-well plate at a concentration of 5×10^6 per well on coverslips in phenol red-free DMEM (Hyclone, USA) and incubated for 1 d. The cells were treated with 0.25 μ M mA β _{1–42} or mA β _{1–42} mixed with His-XD4 (molar ratio 1:50 and 1:100) in serum-free media for 1.5 h. After washing twice with PBS, cells were fixed with 4% paraformaldehyde at 37°C, washed by PBS and blocked with 5% goat serum in PBS at room temperature for 2 h, and then incubated with primary anti-A β _{1–42} antibody 4G8 (1:500) overnight at 4°C. Consequently, the cells were washed five times with PBS, incubated with goat-anti mouse FITC (1:1000) or rabbit-anti his TIRTC (1:1000) secondary antibody simultaneously in 10% goat serum for 1.5 h, and then incubated with 4',6-diamidino-2-phenylindole (DAPI) in anti-fade mounting medium (Beyotime, Beijing, China) at room temperature for 2 h away from light. The coverslips were kept at 4°C until examination by a confocal laser scanning microscope (Zeiss LSM780). The control experiments were performed according to the same protocol without primary antibodies against A β _{1–42} and His-XD4.

Membrane Protein Extraction

Membrane protein of BV-2 cells was prepared by transmembrane protein extraction kit (Merck Millipore, MA, USA)

according to the manufacturer's instruction. Briefly, 1×10^7 BV-2 cells were collected with a cell scraper and treated with a cytosolic extraction buffer. After centrifugation, the supernatants were collected as cytosolic (soluble) protein fraction. The remaining pellets were resuspended with detergent-free membrane protein extraction buffer with the addition of 5 μ l protein inhibitor cocktail. After incubation and centrifugation, the supernatants enriched with integral membrane proteins were collected in a fresh tube, and the total protein concentration of the cytosolic and the membrane fraction were calculated by BCA assay kit.

Immunoprecipitation

Approximately 30 μ g of 6E10 antibody was cross-linked to 50 μ l slurry of aminolink coupling resin with sodium cyanoborohydride according to the instructions of the manufacturer. The antibody-coupled resins were incubated with oA β_{1-42} , which were pre-incubated with or without XD4 for 2 h and added to homogenized membrane extracts of BV-2 cells. The mixture was incubated for 4 h at room temperature with gentle agitation at 30 min intervals. After the resin was washed four times with PBS, the immune-complex was collected in elution buffer and subjected to subsequent western blot procedures. The control experiment was performed with the same procedure in the absence of oA β_{1-42} .

Western Blot

To identify the abundance of SR-A receptor in the immune-complexes generated in immunoprecipitation, the sample was resolved on 12% SDS-PAGE and transferred onto nitrocellulose membranes. The membranes were washed, blocked by 5% non-fat dried milk for 1 h at room temperature, and then incubated with rat-anti SR-A primary antibody (1:1000) overnight at 4°C. The membranes were washed three times in 1% PBST and incubated with mouse anti-rat HRP-conjugated secondary antibody (1:3000) at room temperature for 2 h. The blots were washed with PBS before being exposed to ECL reagent (Beyotime, Beijing, China).

Cell Viability Assay

Colorimetric CCK-8 assay was used to determine the effect of XD4 on oA β_{1-42} -induced cytotoxicity on glia cells. Glia cells were seeded in 96-well plates at a density of 1×10^4 per well. oA β_{1-42} was pre-incubated with or without different concentrations of XD4 for 1 h before adding to the cells. The final concentration of oA β_{1-42} in each culture well was 5 μ M, and the molar ratios of oA β_{1-42} to XD4 were 1:2 and 1:10, respectively. The plates were then incubated for 48 h at 37°C. Cell viability was monitored by adding 10 μ l of CCK-8 solution reagent to each well and incubated at 37°C for 2 h. The absorbance of the wells was read at 450 nm using an MD SpectraMax M5 microplate reader. Averages from three replicate wells were used for each sample, and each experiment was repeated three times. Cell viability was calculated by dividing the absorbance of the wells containing samples by the absorbance of the wells containing medium alone.

Kinase Phosphorylation Assay

To investigate the effect of XD4 on SR-related protein phosphorylation within the MAPK signaling pathway, the relative amount of phosphorylated p38 MAPK and JNK 1/2/3 was determined by Phosphotracer ELISA kits (Abcam, ab119674) in a semi-quantitative method. Cells were seeded at 96-well plates, and then 1 μ M oA β_{1-42} was mixed with or without XD4 (molar ratio 1:25) added to the cell culture. The amount of the phosphorylated

protein was determined by the fluorescence intensity at 590 nm according to the instructions of the manufacturer.

Measurement of TNF- α and IL-1 β

BV-2 cells were seeded in 12-well culture plates at a density of 3×10^5 per well. oA β_{1-42} was pre-incubated with or without different concentrations of XD4 for 1 h before adding to the cells. The final concentration of oA β_{1-42} was 5 μ M, and the molar ratios of oA β_{1-42} to XD4 were 1:2 and 1:10. After 12 h of incubation, the supernatants were collected to determine the levels of TNF- α and IL-1 β released from the activated BV-2 cells using the TNF- α and IL-1 β ELISA kits according to the protocols of the manufacturer.

Statistical Analysis

Data were obtained from at least three separate experiments for each experimental condition. Data were presented as mean \pm standard deviation, and their statistical significance was analyzed by one-way ANOVA. Multiple comparisons between the groups were performed using Student–Newman–Keuls (SNK) method.

Results

Peptide XD4 Increases Uptake of mA β_{1-42} in Microglia

Mouse primary microglia, human microglia, and BV-2 cells were treated with mA β_{1-42} incubated with or without XD4 for 3 h. The intracellular mA β_{1-42} was measured using immunoassay kits. The results showed that the intracellular A β_{1-42} levels were significantly increased in a concentration-dependent manner in all three types of microglial cells with the addition of XD4 (Figure 1). To investigate the mechanism, fucoidan, which is a general ligand and antagonist of scavenger receptors, was added to three types of microglia and incubated for 20 min. Then, mA β_{1-42} pre-incubated with or without XD4 was added to the cells. The results showed that fucoidan failed to block the mA β_{1-42} engulfment induced by XD4, suggesting that mA β_{1-42} may be taken up through a nonsaturable fluid phase macropinocytosis rather than phagocytosis or receptor-mediated endocytosis, which is consistent with a previous report by Mandrekar *et al.* [7]. Moreover, when XD4 and mA β_{1-42} were added to the cells separately, no increased uptake of mA β_{1-42} was observed (data not shown), indicating that a complex formed from mA β_{1-42} and XD4 may be a prerequisite for enhanced internalization of A β_{1-42} .

mA β_{1-42} and XD4 Colocalize in BV-2 Cells

To validate the internalization of mA β_{1-42} and XD4 in the microglia, confocal microscopy was used to visualize the distribution pattern of the two peptides within BV-2 cells. The results showed that internalized mA β_{1-42} and XD4 colocalized in the cells (Figure 2), further suggesting that mA β_{1-42} may be internalized as mA β_{1-42} -XD4 complex (Figure 2A). Z-stack confocal microscopy was employed to detect the distribution pattern of mA β_{1-42} and XD4. Z-stack images, taken at different depths within the cells, indicated that both A β_{1-42} and XD4 are located in the cytoplasm, but not in the nuclear region (Figure 2B), which is partly consistent with the previous report that the internalized A β_{1-42} is located in the cytoplasm and nuclear region [7]. Moreover, BV-2 cells treated with mA β_{1-42} -XD4 complex displayed a stronger fluorescence signal of A β_{1-42} than those treated by A β_{1-42} alone (Figure 2C and 2D), also suggesting that XD4 promoted the uptake of A β_{1-42} by microglia.




Figure 1. XD4 enhanced A β_{1-42} monomer (mA β_{1-42}) uptake in microglia. mA β_{1-42} was incubated with or without XD4 at the indicated molar ratios for 1 h, and then added to various microglia pretreated with or without fucoidan. After 3 h of incubation, mouse primary microglia (A), human microglia (B), and BV-2 cells (C) were lysed and intracellular mA β_{1-42} was measured by ELISA. The data showed the uptake value as a percentage of control (compared with mA β_{1-42} alone, *, $p < 0.05$, **, $p < 0.01$). doi:10.1371/journal.pone.0094197.g001

Scavenger Receptors are Involved in XD4-mediated oA β_{1-42} Phagocytosis in Microglia

To investigate the effect of XD4 on glial internalization of oA β_{1-42} , the intracellular oA β_{1-42} levels in the three types of microglia in the presence or absence of XD4 were measured. The results showed that XD4 increased oA β_{1-42} phagocytosis by mouse primary microglia, human microglia, and BV-2 cells in a concentration-dependent manner (Figure 3). However, the uptake of oA β_{1-42} was drastically hampered by fucoidan in these microglia, suggesting that increased phagocytosis of oA β_{1-42} induced by XD4 may be mediated by the scavenger receptor, which is consistent with the previous report that internalization of mA β_{1-42} and oA β_{1-42} applies distinct mechanisms.

The phagocytosis of A β_{1-42} by microglia may be orchestrated by receptor ensembles and dysfunction of any component within

the receptor complex [12]. The scavenger receptor-neutralizing antibody attenuates phagocytosis in the microglia and astrocytes [20,23]. SR-A and CD36 receptors have critical roles in A β_{1-42} binding, uptake, and concomitant degradation. To identify which subtype of SRs is involved in XD4-mediated oA β_{1-42} internalization, BV-2 cells were incubated with the neutralizing antibodies against SR-A or CD36 before the addition of oA β_{1-42} or oA β_{1-42} -XD4 complex. The results showed that the anti-SR-A antibody reduced the oA β_{1-42} internalization by 50% in BV-2 cells (Figure 3D), whereas the anti-CD36 antibody did not interfere with oA β_{1-42} internalization. These data suggest that SR-A receptor has an important role in XD4-mediated oA β_{1-42} phagocytosis in the microglia.




Figure 2. Intracellular localization of mA β ₁₋₄₂ and XD4 in BV-2 cells. Microglia were incubated with mA β ₁₋₄₂ (green) or mA β ₁₋₄₂ pre-incubated with XD4-His-tag (red) for 1.5 h at 37°C and immunostained with 4G8 antibody and anti-His antibody, respectively. Nuclei were stained with DAPI (blue). The cells were visualized by confocal microscope to detect the colocalization of mA β ₁₋₄₂ and XD4 in BV-2 cells (A), internalization of mA β ₁₋₄₂ and XD4 in BV-2 cells by apical-to-distal Z-scan (B), and effect of XD4 on uptake of A β ₁₋₄₂ (C). Scale bars = 10 μ m. (D) The densitometric analysis of 2C with IpWin5 (compared with oA β ₁₋₄₂ alone, **, $p < 0.01$). doi:10.1371/journal.pone.0094197.g002

XD4 Strengthens the Interaction between SR-A and oA β ₁₋₄₂

To unravel the mechanism through which XD4 increased oA β ₁₋₄₂ phagocytosis in the microglia, the membrane protein fraction of BV-2 cells incubated with oA β ₁₋₄₂ or oA β ₁₋₄₂-XD4 complex was immunoprecipitated using 6E10-coupled resins. The eluted samples were detected by Western blot using anti-SR-A antibody. Consistent with previous report, the samples immunoprecipitated by 6E10 showed the SR-A bands at 58 KD (Figure 4A). When BV-2 cells were incubated with oA β ₁₋₄₂-XD4 complex, the immunoprecipitated samples showed higher amount of SR-A (Figure 4B). Considering that the expression of SR-A protein cannot be induced to express within 2 h during experimental processing, the greater number of SR-A protein observed in the samples with the addition of oA β ₁₋₄₂-XD4 may result from the strengthened interaction of oA β ₁₋₄₂ with SR-A by XD4. This reinforced binding of oA β ₁₋₄₂ to SR-A may increase the phagocytosis of A β . In addition, CD36 was not observed in the

immunoprecipitated samples, suggesting that XD4 did not affect the interaction of CD36 and A β (data not shown).

To measure the affinity of SR-A for oA β ₁₋₄₂ with or without XD4, high binding 96-well ELISA plates were coated with 1 μ g oA β ₁₋₄₂ in the presence or absence of XD4 at 4°C overnight. The prepared membrane protein extracts of BV-2 cells were added and then the rat-anti SR-A primary antibody, HRP-conjugated anti-rat secondary antibody was added. The ELISA results indicated that XD4 significantly increased the binding affinity of SR-A for oA β ₁₋₄₂ (Figure 4C), which is consistent with above results.

XD4 Increases Uptake of mA β ₁₋₄₂ and oA β ₁₋₄₂ in Astrocytes

To test the effect of XD4 on phagocytosis of A β ₁₋₄₂ in astrocytes, the mA β ₁₋₄₂ and oA β ₁₋₄₂ levels in rat astrocytes and human astrogloma U251 cells pre-incubated with A β ₁₋₄₂ in the presence or absence of XD4 were detected. Data demonstrated that XD4 significantly increased the uptake of both mA β ₁₋₄₂ and




Figure 3. XD4 enhanced oA β_{1-42} phagocytosis in microglia. oA β_{1-42} was incubated with or without XD4 for 1 h, and then added to microglia cell lines pretreated with or without fucoidan. After 1.5 h of incubation, mouse primary microglia (A), human microglia (B), and BV-2 cells (C) were lysed, and intracellular oA β_{1-42} was determined by oligomer-specific ELISA. The data showed the uptake value as a percentage of control. (D) BV-2 cells were incubated with anti-SR-A antibody and anti-CD36 antibody for 30 min, and then exposed to oA β_{1-42} or oA β_{1-42} -XD4 for 1.5 h. The cells were lysed and the intracellular oA β_{1-42} was measured by ELISA. All data were standardized as percentage of control (compared with oA β_{1-42} alone, *, $p < 0.05$, **, $p < 0.01$; compared with oA β_{1-42} +50 μ M XD4, ##, $p < 0.01$). doi:10.1371/journal.pone.0094197.g003

oA β_{1-42} in rat astrocytes and U251 cells, however, these effects were inhibited by 100 μ g/ml fucoidan (Figure 5), indicating that XD4 increased the uptake of mA β_{1-42} and oA β_{1-42} through SR, and microglia and astrocytes applied distinct mechanisms to internalize mA β_{1-42} .

XD4 Attenuates oA β_{1-42} Cytotoxicity in Glia Cells

A β is believed to be a cytotoxic factor that causes neuronal damage in AD progressing. To test whether the enhanced uptake of A β_{1-42} induced by XD4 exerts toxic effects on the glia, human microglia, rat astrocytes, and U251 cells were incubated with

oA β_{1-42} in the presence or absence of XD4. The results indicated that the cell viability of all three glial cells incubated with oA β_{1-42} alone remarkably decreased (Figure 6). However, oA β_{1-42} -induced cytotoxicity can be drastically attenuated by the addition of XD4 in a concentration-dependent manner. Our finding indicated that XD4 can protect glial cells from A β_{1-42} -induced cytotoxicity, although it obviously enhanced intracellular A β levels.

To investigate what happens to the internalized oA β_{1-42} in the glial cells, the BV-2 cells were incubated with oA β_{1-42} for 1.5 h in the presence or absence of 50 μ M XD4, the intracellular oA β_{1-42} levels in BV-2 cells were measured by ELISA at 0 h, 6 h and 12 h, respectively. The results showed that the intracellular oA β_{1-42}




Figure 4. XD4 promoted the interaction between SR-A and oA β ₁₋₄₂. (A) Equal volumes of homogenized membrane fraction of BV-2 cells were added to oA β ₁₋₄₂ or oA β ₁₋₄₂-XD4 preincubated with 6E10-conjugated resin. After 4 h of incubation with gentle agitation, the binding protein from the co-IP process was separated by SDS-PAGE (12% gels) and subjected to immune-blot analysis using anti-SR-A antibody. The control experiment was performed through the same procedure in the absence of oA β ₁₋₄₂. (B) The densitometric analysis was carried out by IpWin5. (C) The affinity of SR-A for oA β ₁₋₄₂ with or without XD4 was tested by ELISA. High binding 96-well ELISA plates were coated with 1 μ g oA β ₁₋₄₂ in the presence or absence of XD4 at 4°C overnight. The prepared membrane protein extracts of BV-2 cells were added and then the rat-anti SR-A primary antibody, HRP-conjugated anti-rat secondary antibody was added followed by TMB substrate (compared with oA β ₁₋₄₂ alone, *, $p < 0.05$, **, $p < 0.01$). doi:10.1371/journal.pone.0094197.g004

levels decreased in a time-dependent manner (Figure 6D). The mechanism of oA β ₁₋₄₂ clearance in microglia may be that the internalized oA β ₁₋₄₂ was trafficked toward lysosomes and degraded by cysteine proteases, including cathepsin B [23].

oA β ₁₋₄₂-XD4 Complex Activates SR-A

JNK and p38 MAPK are involved in the SR ligand-regulated signaling cascade in macrophages and glia. The levels of phosphorylated JNK and p38 MAPK were determined to verify the effect of XD4 on oA β ₁₋₄₂ phagocytosis and to explore the mechanism through which XD4 enhances A β internalization. The results showed that oA β ₁₋₄₂ treatment did not significantly increase JNK and p38 MAPK phosphorylation in BV-2 cells. However, oA β ₁₋₄₂-XD4 complex enhanced this phosphorylation (Figure 7). These results illustrated that the oA β ₁₋₄₂ - XD4 complex can activate SR-A, thus enhancing the phosphorylation of JNK and p38 MAPK.

XD4 Reduces oA β ₁₋₄₂-induced Production of TNF- α and IL-1 β

To assess the effect of XD4 on the production of proinflammatory cytokines, BV-2 cells were incubated with oA β ₁₋₄₂ in the presence or absence of XD4 for 12 h. The supernatants were collected for TNF- α and IL-1 β measurements by ELISA. Our results showed that oA β ₁₋₄₂ alone significantly increased the generation of TNF- α and IL-1 β , whereas XD4 significantly reduced TNF- α and IL-1 β levels in the supernatants in a concentration-dependent manner (Figure 8). These findings indicated that XD4 can attenuate oA β ₁₋₄₂-induced proinflammatory cytokine production, although it enhances A β uptake.

Discussion

AD is the most common neurodegenerative dementia with progressive deposition of A β within the brain. The soluble or small oligomeric forms of A β induce synaptic dysfunction, behavioral deficits, and neuronal degeneration [24]. Therefore, the elimination of toxic A β species in the early stages of AD seems to be a promising strategy for rescuing synaptic damage and relieving




Figure 5. XD4 promoted uptake of A β_{1-42} in astrocytes. mAb β_{1-42} or oAb β_{1-42} was incubated with or without XD4 at the indicated molar ratio for 1 h, and then added to astrocyte cells pretreated with or without fucoidan. After 1.5 h of incubation, rat astrocytes (A, B) or U251 cells (C, D) were lysed and intracellular mAb β_{1-42} (A, C) or oAb β_{1-42} (B, D) was determined by ELISA. The data showed the uptake value as a percentage of control (compared with A β_{1-42} alone, *, $p < 0.05$, **, $p < 0.01$; compared with A β_{1-42} +50 μ M XD4, #, $p < 0.05$, ##, $p < 0.01$). doi:10.1371/journal.pone.0094197.g005

cognitive impairment [23]. Microglia are major phagocytic cells in the brain. Upon injury, microglia migrate to the lesion sites, recognize the pathogen, and then ramify and mount an immune response in response to the stimulus [25]. The activation and accumulation of microglial cells around A β plaques have been reported in the brains of both AD patients and mouse models [26]. Microglia can engulf A β and degrade A β , but this action is a dual sword. Although microglia exert a protective role, possibly by internalization and degradation of A β at initial stages of AD, they may also lose A β phagocytosis and induce neuronal damage by releasing pro-inflammatory and toxic agents, especially at the late stages of AD. Therefore, turning the deleterious state of microglia to beneficial state is a reasonable strategy for AD treatment. To the best of our knowledge, no agent that promotes A β clearance by

SR-A in microglia has been reported. In this study, we showed that glia phagocytosis can be enhanced by peptide XD4.

Aside from microglia, astrocytes are also considered main phagocytes that can engulf A β [20]. Glia can engulf A β in several ways, such as nonsaturable, fluid phase macropinocytosis and phagocytosis [7,11,23]. Our results demonstrated that XD4 increased the uptake of A β monomers by microglia and astrocytes, however, this effect can be blocked by fucoidan in astrocytes rather than in microglia, suggesting that different glial cells apply different mechanisms to engulf A β monomers. Microglia engulf mAb β_{1-42} through nonsaturable, fluid phase macropinocytosis, whereas astrocytes engulf mAb β_{1-42} through phagocytosis by SR-A, which consistent with previous reports [19,27].

Previous reports indicated that SR was also involved in phagocytosis of fibrillar A β by microglia [11,28]. Our present




Figure 6. Protective effect of XD4 on oA β ₁₋₄₂-induced cytotoxicity in glia cells. oA β ₁₋₄₂ was pre-incubated with or without different molar ratios of XD4 for 1 h before adding to the human microglia (A) rat astrocytes (B) and U251 cells (C), respectively. After two days of incubation, cell viability was measured using CCK-8 assay. The absorbance of the wells was read at 450 nm using an MD SpectraMax M5 microplate reader. Cell viability was calculated by dividing the absorbance of the wells containing the samples (corrected for background) by the absorbance of wells containing medium alone (corrected for background) (compared with oA β ₁₋₄₂ alone, *, $p < 0.05$, **, $p < 0.01$). (D) The internalized oA β ₁₋₄₂ was cleared in a time-dependent manner. After BV-2 cells were incubated with oA β ₁₋₄₂ for 1.5 h in the presence or absence of 50 μ M XD4, the intracellular oA β ₁₋₄₂ levels in BV-2 cells were measured by ELISA at 0 h, 6 h and 12 h, respectively. The data showed the value as a percentage of control (compared with oA β ₁₋₄₂ alone at 0 h, **, $p < 0.01$; compared with oA β ₁₋₄₂+50 μ M XD4 at 0 h, ##, $p < 0.01$). doi:10.1371/journal.pone.0094197.g006

results showed that SR-A was involved in the uptake of A β oligomers by microglia, and XD4 enhanced the phagocytosis of A β (Figure 3), which was confirmed by the blockage of phagocytosis with fucoidan and anti-SR-A antibody. Moreover, the phagocytosis of A β oligomers by astrocytes was also enhanced by XD4, and this enhanced phagocytosis was significantly decreased by fucoidan, indicating that XD4 enhanced the internalization of oA β ₁₋₄₂ in astrocytes also via the SR-A pathway. Our immunoprecipitation and Western blot results suggested that the enhanced binding of oA β ₁₋₄₂ to SR-A by XD4 contributed to the enhanced phagocytosis of oA β ₁₋₄₂.

Our previous study showed that XD4 did not inhibit A β aggregation, but attenuated A β ₁₋₄₂-induced cytotoxicity in SH-

SY5Y cells [29]. In this study, we showed that XD4 also inhibited oA β ₁₋₄₂-induced cytotoxicity in glial cells and presented a neuroprotective effect, although more oA β ₁₋₄₂ was internalized into the cells, which may be attributed to the fact that the interaction between XD4 and A β aggregates forms a harmless structure of A β . The activation of SRs by the binding of ligand such as fucoidan can trigger downstream signal transduction pathways in glial cells, increasing the phosphorylation of diverse kinase, such as JNK and p38 [17,30]. Our finding indicated that oA β ₁₋₄₂-XD4 complex can significantly increase the phosphorylation of JNK and p38 MAPK, further confirming that SR-A is involved in XD4-mediated A β phagocytosis.




Figure 7. XD4 enhanced JNK and p38 phosphorylation. oA β_{1-42} or oA β_{1-42} -XD4 was added to the BV-2 cells. After 2 h of incubation, the cells were lysed and the phosphorylated JNK (A) and p38 (B) levels were determined by ELISA (compared with oA β_{1-42} alone, *, $p < 0.05$). doi:10.1371/journal.pone.0094197.g007

Increasing evidence has shown high levels of inflammatory mediators in the brain of AD patients. Activated microglia cells by A β accumulation release high levels of proinflammatory cytokines, such as TNF- α and IL-1 β , which may further cause neuronal dysfunction [31,32]. Although XD4 activated SR-A, it did not increase the generation of TNF- α and IL-1 β in BV-2 cells induced by oA β_{1-42} in vitro and in vivo [28]. This phenomenon may be attributed to the fact that the enhanced phagocytosis lowered the levels of extracellular A β and reduced the A β -mediated stimulation to the cells, decreasing the release of TNF- α and IL-1 β . Moreover, previous reports have demonstrated that the binding of ligands to SR-A attenuated the TLR4-mediated inflammation responses [33], which may partly explain the reduced production of proinflammatory cytokines induced by XD4.

Therefore, XD4 enhanced the glial uptake of A β monomers and oligomers through phagocytosis or macropinocytosis by strengthening the binding of A β to SR-A, inhibited oA β_{1-42} -induced cytotoxicity, and reduced the production of proinflammatory cytokines, including TNF- α and IL-1 β . As a short peptide, XD4 may have lower immunogenicity than antibodies, and is suitable for further engineering to improve specificity and stability. In conjunction with our previous findings that XD4 attenuated memory deficits in AD transgenic mice and reduced amyloid plaque burden and A β 40/42 levels, XD4 may be a promising therapeutic agent for AD treatment.




Figure 8. XD4 reduced production of TNF- α and IL-1 β . BV-2 cells were incubated with oA β_{1-42} in the presence or absence of XD4 for 12 h. The supernatants were collected and the levels of TNF- α (A) and IL-1 β (B) were determined using ELISA kits (compared with oA β_{1-42} alone, *, $p < 0.05$, **, $p < 0.01$). doi:10.1371/journal.pone.0094197.g008

Author Contributions

Conceived and designed the experiments: RTL HZ XLY. Performed the experiments: HZ YJS SWW WWZ PXX. Analyzed the data: HZ YJS RTL XLY. Wrote the paper: HZ RTL XLY.

References

- Lee CY, Landreth GE (2010) The role of microglia in amyloid clearance from the AD brain. *J Neural Transm* 117: 949–960.
- Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297: 353–356.
- Broersen K, Rousseau F, Schymkowitz J (2010) The culprit behind amyloid beta peptide related neurotoxicity in Alzheimer's disease: oligomer size or conformation? *Alzheimers Res Ther* 2: 12.
- Verdier Y, Zarandi M, Penke B (2004) Amyloid beta-peptide interactions with neuronal and glial cell plasma membrane: binding sites and implications for Alzheimer's disease. *J Pept Sci* 10: 229–248.
- Luc LF, Walker DG, Rogers J (2001) Modeling microglial activation in Alzheimer's disease with human postmortem microglial cultures. *Neurobiol Aging* 22: 945–956.
- Prokop S, Miller KR, Heppner FL (2013) Microglia actions in Alzheimer's disease. *Acta Neuropathol* 126: 461–477.
- Mandrekar S, Jiang Q, Lee CY, Koenigsnecht-Talboo J, Holtzman DM, et al. (2009) Microglia mediate the clearance of soluble A β through fluid phase macropinocytosis. *J Neurosci* 29: 4252–4262.
- D'Andrea MR, Cole GM, Ard MD (2004) The microglial phagocytic role with specific plaque types in the Alzheimer disease brain. *Neurobiol Aging* 25: 675–683.
- Mohamed A, Posse de Chaves E (2011) A β internalization by neurons and glia. *Int J Alzheimers Dis* 2011: 127984.
- Husemann J, Loike JD, Anankov R, Febbraio M, Silverstein SC (2002) Scavenger receptors in neurobiology and neuropathology: their role on microglia and other cells of the nervous system. *Glia* 40: 195–205.
- Paresce DM, Ghosh RN, Maxfield FR (1996) Microglial cells internalize aggregates of the Alzheimer's disease amyloid beta-protein via a scavenger receptor. *Neuron* 17: 553–565.
- Bamberger ME, Harris ME, McDonald DR, Husemann J, Landreth GE (2003) A cell surface receptor complex for fibrillar beta-amyloid mediates microglial activation. *J Neurosci* 23: 2665–2674.
- Reed-Geaghan EG, Savage JC, Hise AG, Landreth GE (2009) CD14 and toll-like receptors 2 and 4 are required for fibrillar A β -stimulated microglial activation. *J Neurosci* 29: 11982–11992.
- Yu X, Yi H, Guo C, Zuo D, Wang Y, et al. (2011) Pattern recognition scavenger receptor CD204 attenuates Toll-like receptor 4-induced NF-kappaB activation by directly inhibiting ubiquitination of tumor necrosis factor (TNF) receptor-associated factor 6. *J Biol Chem* 286: 18795–18806.
- Stuart LM, Bell SA, Stewart GR, Silver JM, Richard J, et al. (2007) CD36 signals to the actin cytoskeleton and regulates microglial migration via a p130Cas complex. *J Biol Chem* 282: 27392–27401.
- Wilkinson K, El Khoury J (2012) Microglial scavenger receptors and their roles in the pathogenesis of Alzheimer's disease. *Int J Alzheimers Dis* 2012: 489456.
- Hsu HY, Chiu SL, Wen MH, Chen KY, Hua KF (2001) Ligands of macrophage scavenger receptor induce cytokine expression via differential modulation of protein kinase signaling pathways. *J Biol Chem* 276: 28719–28730.
- Cornejo F, von Bernhardi R (2013) Role of scavenger receptors in glia-mediated neuroinflammatory response associated with Alzheimer's disease. *Mediators Inflamm* 2013: 895651.
- Alarcon R, Fuenzalida C, Santibanez M, von Bernhardi R (2005) Expression of scavenger receptors in glial cells. Comparing the adhesion of astrocytes and microglia from neonatal rats to surface-bound beta-amyloid. *J Biol Chem* 280: 30406–30415.
- Jones RS, Minogue AM, Connor TJ, Lynch MA (2013) Amyloid-beta-induced astrocytic phagocytosis is mediated by CD36, CD47 and RAGE. *J Neuroimmune Pharmacol* 8: 301–311.
- Frenkel DN, Wilkinson K, Zhao L, Hickman SE, Means TK, et al. (2013) Scara1 deficiency impairs clearance of soluble amyloid-beta by mononuclear phagocytes and accelerates Alzheimer's-like disease progression. *Nat Commun* 4: 2030.
- Dahlgren KN, Manelli AM, Stine WB Jr, Baker LK, Krafft GA, et al. (2002) Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *J Biol Chem* 277: 32046–32053.
- Yang CN, Shiao YJ, Shie FS, Guo BS, Chen PH, et al. (2011) Mechanism mediating oligomeric A β clearance by naive primary microglia. *Neurobiol Dis* 42: 221–230.
- Lacor PN, Buniel MC, Furlow PW, Clemente AS, Velasco PT, et al. (2007) A β oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci* 27: 796–807.
- Ransohoff RM, Perry VH (2009) Microglial physiology: unique stimuli, specialized responses. *Annu Rev Immunol* 27: 119–145.
- Frautschy SA, Yang F, Irizarry M, Hyman B, Saido TC, et al. (1998) Microglial response to amyloid plaques in APPsw transgenic mice. *Am J Pathol* 152: 307–317.
- Allaman I, Gavillet M, Belanger M, Laroche T, Viertl D, et al. (2010) Amyloid-beta aggregates cause alterations of astrocytic metabolic phenotype: impact on neuronal viability. *J Neurosci* 30: 3326–3338.
- El Khoury J, Hickman SE, Thomas CA, Cao L, Silverstein SC, et al. (1996) Scavenger receptor-mediated adhesion of microglia to beta-amyloid fibrils. *Nature* 382: 716–719.
- Xue D, Zhao M, Wang YJ, Wang L, Yang Y, et al. (2012) A multifunctional peptide rescues memory deficits in Alzheimer's disease transgenic mice by inhibiting A β 42-induced cytotoxicity and increasing microglial phagocytosis. *Neurobiol Dis* 46: 701–709.
- Coller SP, Paulnock DM (2001) Signaling pathways initiated in macrophages after engagement of type A scavenger receptors. *J Leukoc Biol* 70: 142–148.
- Rogers J, Luc LF (2001) Microglial chemotaxis, activation, and phagocytosis of amyloid beta-peptide as linked phenomena in Alzheimer's disease. *Neurochem Int* 39: 333–340.
- Tuppo EE, Arias HR (2005) The role of inflammation in Alzheimer's disease. *Int J Biochem Cell Biol* 37: 289–305.
- Mukhopadhyay S, Varin A, Chen Y, Liu B, Tryggvason K, et al. (2011) SR-A/MARCO-mediated ligand delivery enhances intracellular TLR and NLR function, but ligand scavenging from cell surface limits TLR4 response to pathogens. *Blood* 117: 1319–1328.