



# Structural Basis of Type 2 Secretion System Engagement between the Inner and Outer Bacterial Membranes

Iain D. Hay, Matthew J. Belousoff, Trevor Lithgow

Infection and Immunity Program, Biomedicine Discovery Institute and Department of Microbiology, Monash University, Clayton, Australia

**ABSTRACT** Sophisticated nanomachines are used by bacteria for protein secretion. In Gram-negative bacteria, the type 2 secretion system (T2SS) is composed of a pseudopilus assembly platform in the inner membrane and a secretin complex in the outer membrane. The engagement of these two megadalton-sized complexes is required in order to secrete toxins, effectors, and hydrolytic enzymes. *Pseudomonas aeruginosa* has at least two T2SSs, with the ancestral nanomachine having a secretin complex composed of XcpQ. Until now, no high-resolution structural information was available to distinguish the features of this *Pseudomonas*-type secretin, which varies greatly in sequence from the well-characterized *Klebsiella*-type and *Vibrio*-type secretins. We have purified the ~1-MDa secretin complex and analyzed it by cryo-electron microscopy. Structural comparisons with the *Klebsiella*-type secretin complex revealed a striking structural homology despite the differences in their sequence characteristics. At 3.6-Å resolution, the secretin complex was found to have 15-fold symmetry throughout the membrane-embedded region and through most of the domains in the periplasm. However, the N1 domain and N0 domain were not well ordered into this 15-fold symmetry. We suggest a model wherein this disordering of the subunit symmetry for the periplasmic N domains provides a means to engage with the 6-fold symmetry in the inner membrane platform, with a metastable engagement that can be disrupted by substrate proteins binding to the region between XcpP, in the assembly platform, and the XcpQ secretin.

**IMPORTANCE** How the outer membrane and inner membrane components of the T2SS engage each other and yet can allow for substrate uptake into the secretin chamber has challenged the protein transport field for some time. This vexing question is of significance because the T2SS collects folded protein substrates in the periplasm for transport out of the bacterium and yet must discriminate these few substrate proteins from all the other hundred or so folded proteins in the periplasm. The structural analysis here supports a model wherein substrates must compete against a metastable interaction between XcpP in the assembly platform and the XcpQ secretin, wherein only structurally encoded features in the T2SS substrates compete well enough to disrupt XcpQ-XcpP for entry into the XcpQ chamber, for secretion across the outer membrane.

**KEYWORDS** *Pseudomonas aeruginosa*, T2SS, protein secretion, protein secretion system, secretin

Bacterial pathogens use a selection of sophisticated protein secretion nanomachines to deliver toxins, effectors, and hydrolytic enzymes into their environments (1, 2). One of these, the type 2 secretion system (T2SS), evolved from an ancestral nanomachine that also gave rise to the type 4 fimbrial system for locomotion through the secretion—and retraction—of long, fibrous pili (3). These pili are projected from an

**Received** 28 July 2017 **Accepted** 12 September 2017 **Published** 17 October 2017  
**Citation** Hay ID, Belousoff MJ, Lithgow T. 2017. Structural basis of type 2 secretion system engagement between the inner and outer bacterial membranes. *mBio* 8:e01344-17. <https://doi.org/10.1128/mBio.01344-17>.

**Editor** Lotte Søgaard-Andersen, Max Planck Institute for Terrestrial Microbiology

**Copyright** © 2017 Hay et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Trevor Lithgow, [trevor.lithgow@monash.edu](mailto:trevor.lithgow@monash.edu).

For a companion article on this topic, see <https://doi.org/10.1128/mBio.01185-17>.


inner membrane protein complex that functions as an assembly platform and ATP-dependent motor, and from this platform, the pili are driven into a cavernous secretin complex embedded in the outer membrane that spans the periplasm to engage the inner membrane. The T2SS itself encompasses a pilus-like structure, referred to as a pseudopilus since it is not driven outside the cell for locomotion but rather driven in a short motor stroke. The pseudopilus is driven into the T2SS secretin complex by either a piston or a screw-like mechanism, thereby promoting secretion of substrate proteins (toxins, effectors, and hydrolytic enzymes) across the outer membrane. Recent high-resolution structures of the *Vibrio*-type and *Klebsiella*-type secretins showed them to be of 15-fold symmetry (4). These structural breakthroughs highlighted a perplexing problem, given that the secretins interact intimately with 6-fold symmetric structures in the T2SS inner membrane platform (5–7).

*Pseudomonas aeruginosa* is a bacterial pathogen of particular concern to patients with burns, wounds, or cystic fibrosis (8). Species of *Pseudomonas* have two characteristic T2SSs, defined by distinct secretins, XcpQ and HxcQ (9–11). Phylogenetic analysis has suggested that while the T2SS containing HxcQ was acquired more recently by lateral gene transfer from betaproteobacteria, the XcpQ secretin defines the ancestral *Pseudomonas*-type T2SS (10). The T2SS containing XcpQ is responsible for the secretion of at least 19 substrates, most of them hydrolases ranging from proteases to hemolytic lipases that are critical for pathogenesis (11–13). After crossing the inner membrane via the SecYEG or TAT translocases, substrates of the T2SS enter the secretin chamber via an unknown mechanism (5, 6, 9, 14). In order to address how the outer membrane and inner membrane components engage together and yet can allow for substrate uptake, high-resolution structural data are required.

Previous detailed sequence analysis has demonstrated the sequence-based features that distinguish the *Klebsiella*-type secretins from the *Vibrio*-type secretins (15). Cluster Analysis of Sequence data (CLANS) is a sensitive tool to depict sequence relationships (16): in a CLANS plot, each protein is represented as a dot clustering with other proteins of similar sequence. A core set of T2SS secretin sequences was extracted from the InterPro “GspD/PilQ family” (IPR001775) (see Text S1 and Table S1 in the supplemental material), and a CLANS plot of these sequences from diverse bacterial lineages showed the evolutionary distances separating the XcpQ and HxcQ secretins of *Pseudomonas* from other functionally characterized secretins (Fig. 1A). Great sequence drift separates the XcpQ and HxcQ secretins from each other and from the previously characterized *Klebsiella*-type and *Vibrio*-type secretins, while the secretin from the recently discovered T2SS of *Acinetobacter baumannii* (17) clusters with the XcpQ secretin from species of *Pseudomonas* and a grouping of marine bacteria (Fig. 1A). Despite their very close sequence-based clustering, the *Klebsiella*-type and *Vibrio*-type secretins are structurally distinct. Thus, with the XcpQ secretin being even more diverse in sequence and representing an archetype for the *Pseudomonas*-type secretins in general, the secretin complex of *P. aeruginosa* PAO1 was purified for structural analysis by single-particle cryo-electron microscopy.


XcpQ oligomers were detergent solubilized and purified by size exclusion chromatography, and the ~1-MDa species was shown to be a pure XcpQ multimer when characterized by SDS-PAGE (Fig. S1). XcpQ samples were flash frozen on vitreous ice and were imaged on a Titan Krios transmission electron microscope equipped with a Gatan K2 Summit direct electron detector (Fig. 1B). After analysis of 18,000+ particles, class averages revealed that XcpQ is a pentadecameric secretin complex (Fig. 1B and S2) with no other symmetries observed in the two-dimensional (2D) class averages.

In the T2SS secretins, each subunit consists of four N domains followed by the highly conserved secretin domain (Pfam PF00263), followed by an S domain. These features, as defined in the secretin structures of the *Klebsiella*-type secretins, are present in the structure of XcpQ (Fig. 2A and B). In the *Klebsiella*-type secretins and *Vibrio*-type secretins, the S domain is necessary for interactions with the pilotin that mediates secretin trafficking to the outer membrane and assembly (15, 18–20). The disposition of the S domain in the XcpQ complex structure is equivalent to that of the other



**FIG 1** Sequence and structural relationships of XcpQ and other secretins. (A) CLANS analysis (16) graphically depicts homology in large data sets of proteins, utilizing all-against-all pairwise BLAST to cluster representations (colored dots) of individual protein sequences in three-dimensional space. Darker lines between sequences represent similar sequences, and lighter lines represent less similar sequences with an E value cutoff of  $1e-10$ . The analysis of T2SS secretin sequences (see Table S1 in the supplemental material) shows that proteins from diverse species cluster into six types. The *Klebsiella* type and *Vibrio* type (red) are barely resolved at this scale, and a diverse group of gammaproteobacterial sequences (yellow) are clustered between the *Klebsiella*-type/*Vibrio*-type group and the other groups. The *Pseudomonas*-type HxcQ secretin clusters among a group of betaproteobacterial T2SS sequences from the *Burkholderiales* (orange). *Pseudomonas aeruginosa* XcpQ clusters among three well-resolved groups: one comprised of various XcpQs from various *Pseudomonas* species, one comprised of marine bacteria, and one comprised of the recently described T2SS secretins from species of *Acinetobacter*. Insets show the distinct structures of the *Klebsiella*-type (PDB accession no. 5WQ7) and *Vibrio*-type (PDB accession no. 5WQ8) secretins. (B) XcpQ was purified by size exclusion chromatography, the fraction containing the ~1-MDa complex was frozen, and individual XcpQ particles were visualized by electron microscopy. 2D class averages of top and side views are shown. (C) The 3.6-Å model of XcpQ is shown intact (left) and sliced open sagittally (right) with color coding to highlight the domain architecture. The top-down view illustrates the 15-fold symmetry of XcpQ.

pilotin-dependent secretins (Fig. 2C). Features previously named the “upper chamber,” “inner gate,” and “cap” were resolved (Fig. 2A and B). Comparison of the XcpQ structure to that of the *Klebsiella*-type secretin complex from *Escherichia coli* K-12 strain DH5 $\alpha$  (Protein Data Bank [PDB] accession no. 5WQ7) revealed a striking structural homology (Fig. 2C) despite the differences in their sequence characteristics (Fig. 1A).



**FIG 2** High-resolution structural analysis of XcpQ. (A) XcpQ molecular model (drawn as a cartoon) built into the electron density. Positions where residues could not be assigned are indicated with dotted circles. (B) XcpQ protomer: domain structure showing the secretin domain, the hinge region (residues 239 to 246), and the periplasmic N domains. (C) XcpQ secretin complex and, by way of comparison, the *Klebsiella*-type secretin (PDB accession no. [5WQ7](#)) shown to scale. The similar positioning of the S domains is indicated. OM, outer membrane. (D) The electron densities of XcpQ (green) and the *Klebsiella*-type secretin (red) are overlaid, showing the relatively conserved structure and the progressive deterioration in resolution through the N2, N1, and N0 domains of the structures. (E) Model representing the C15 symmetry observed in the secretin and N3 domain and a model of the pseudo-6-fold symmetry of a hexamer of dimers for the N0-N2 domain.

The N domains of XcpQ contribute to the chamber that needs to dock onto the XcpP subunits of the inner membrane platform (5, 14). At 3.6-Å resolution, individual  $\alpha$ -helices and  $\beta$ -strands were resolved throughout the upper part of the secretin structure and within the N3 domain, and the majority of the protein could be visualized (Fig. 2C). There is a relatively unstructured linker or hinge between the N3 and N2 domains (corresponding to residues 239 to 246 in the mature form of XcpQ), where the resolution of the structure starts to decline (Fig. 2D, S2D, and S3). Less well resolved density was present for the N1 domains, and no density for the N0 domains could be observed (Fig. 2D, S2D, and S3). The failure to identify these N domains raises the prospect that this region of the secretin complex, proximal to XcpP in the inner membrane, is not arranged in a uniform 15-fold symmetry.

The T2SS has been an enigma to structural and molecular biologists. In order to drive protein secretion, a secretin like XcpQ must dock to an integral inner membrane complex with 6-fold symmetry (XcpP) (5–7). These intimate interactions raise a further structural dilemma, in that substrates access the T2SS in the periplasm and must therefore interpose between the seals formed by the N domains of the secretin (5, 7). Given the recent findings of Douzi et al. (21), in combination with our data, we suggest a plausible scenario that would resolve this conundrum (Fig. 2E).

The three-dimensional (3D) reconstruction of the XcpQ oligomer was performed under a strict cyclic-15-fold (C15) symmetry. This symmetry restriction was judged to be appropriate by unambiguous 2D class averages of “top-down” projections, clearly and only showing a C15 symmetry (Fig. 1B and C). As can be seen from the 2D class averages (Fig. S2), the best-resolved density is that of the secretin ring and N3 domain, leaving the proximal N domains poorly resolved. This holds true for our 3D model, which during the Fourier reconstruction favored the high-contrast features of the secretin ring and N3 domains. Our data suggest that the N0–N2 domains may not fully conform to the C15 symmetry, and attempts to resolve this part of the structure were not successful. We propose that a pseudo-6-fold symmetry exists in the proximal region of the N domains (Fig. 2E). This might be acquired by the XcpQ dimer-based structures defined by Douzi et al. (21) via a radially symmetric array wherein three N domains are displaced away from the XcpP structure to create a metastable arrangement. The flexibility with which the features pack in order to transition from 15-fold to pseudo-6-fold symmetry provides insight into the mechanism for substrate entry into the T2SS, with the substrate-XcpP interactions competing for the XcpQ-XcpP interactions in order to gain entry into the XcpQ chamber; success in this competition would see the substrate be secreted via pseudopilus movements.

Detailed descriptions of methods are available in Text S1 in the supplemental material, with the map and model parameters for the structures presented as Fig. S2 and Table S2, respectively. T2SS secretin sequence accession numbers and annotations used for the CLANS analysis are presented in Table S1. Electron microscopy data were collected at the Ramaciotti Centre for Cryo-Electron Microscopy, Monash University. All figures were generated with either PyMOL (22) or UCSF Chimera (23).

**Accession number(s).** The electron density map has been deposited in the EMDB (EMD-8860). The model of XcpQ has been deposited in the PDB (5WLN).

## SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/mBio.01344-17>.

**TEXT S1**, DOCX file, 0.02 MB.

**FIG S1**, PDF file, 0.1 MB.

**FIG S2**, PDF file, 1.2 MB.

**FIG S3**, PDF file, 0.1 MB.

**TABLE S1**, PDF file, 0.3 MB.

**TABLE S2**, PDF file, 0.05 MB.

## ACKNOWLEDGMENTS

We thank Rebecca Bamert for expert assistance in sample preparation and Christopher Stubenrauch and Rhys Dunstan for critical comments on the manuscript.

I.D.H. was the recipient of an NVIDIA Hardware Grant, and we gratefully acknowledge the support of NVIDIA Corporation for the donation of the TitanX GPU used for this research. Our work in this area is supported by NHMRC program grant 1092262 and the Australian Research Council (ARC; FL130100038). I.D.H. is an ARC Laureate Postdoctoral Fellow, and T.L. is an ARC Australian Laureate Fellow. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

## REFERENCES

- Costa TR, Felisberto-Rodrigues C, Meir A, Prevost MS, Redzej A, Trokter M, Waksman G. 2015. Secretion systems in Gram-negative bacteria: structural and mechanistic insights. *Nat Rev Microbiol* 13:343–359. <https://doi.org/10.1038/nrmicro3456>.
- Gunasinghe SD, Webb CT, Elgass KD, Hay ID, Lithgow T. 2017. Super-resolution imaging of protein secretion systems and the cell surface of gram-negative bacteria. *Front Cell Infect Microbiol* 7:220. <https://doi.org/10.3389/fcimb.2017.00220>.
- Hospenthal MK, Costa TRD, Waksman G. 2017. A comprehensive guide to pilus biogenesis in Gram-negative bacteria. *Nat Rev Microbiol* 15:365–379. <https://doi.org/10.1038/nrmicro.2017.40>.
- Yan Z, Yin M, Xu D, Zhu Y, Li X. 2017. Structural insights into the secretin translocation channel in the type II secretion system. *Nat Struct Mol Biol* 24:177–183. <https://doi.org/10.1038/nsmb.3350>.
- Douzi B, Ball G, Cambillau C, Tegoni M, Voulhoux R. 2011. Deciphering the Xcp *Pseudomonas aeruginosa* type II secretion machinery through multiple interactions with substrates. *J Biol Chem* 286:40792–40801. <https://doi.org/10.1074/jbc.M111.294843>.
- Korotkov KV, Sandkvist M, Hol WG. 2012. The type II secretion system: biogenesis, molecular architecture and mechanism. *Nat Rev Microbiol* 10:336–351. <https://doi.org/10.1038/nrmicro2762>.
- Wang X, Pineau C, Gu S, Guschinskaya N, Pickersgill RW, Shevchik VE. 2012. Cysteine scanning mutagenesis and disulfide mapping analysis of arrangement of GspC and GspD protomers within the type 2 secretion system. *J Biol Chem* 287:19082–19093. <https://doi.org/10.1074/jbc.M112.346338>.
- Breidenstein EB, de la Fuente-Núñez C, Hancock RE. 2011. *Pseudomonas aeruginosa*: all roads lead to resistance. *Trends Microbiol* 19:419–426. <https://doi.org/10.1016/j.tim.2011.04.005>.
- Douzi B, Filloux A, Voulhoux R. 2012. On the path to uncover the bacterial type II secretion system. *Philos Trans R Soc Lond B Biol Sci* 367:1059–1072. <https://doi.org/10.1098/rstb.2011.0204>.
- Durand E, Alphonse S, Brochier-Armanet C, Ball G, Douzi B, Filloux A, Bernard C, Voulhoux R. 2011. The assembly mode of the pseudopilus: a hallmark to distinguish a novel secretion system subtype. *J Biol Chem* 286:24407–24416. <https://doi.org/10.1074/jbc.M111.234278>.
- Ball G, Antelmann H, Imbert PR, Gimenez MR, Voulhoux R, Ize B. 2016. Contribution of the twin arginine translocation system to the exoprotoome of *Pseudomonas aeruginosa*. *Sci Rep* 6:27675. <https://doi.org/10.1038/srep27675>.
- Filloux A. 2011. Protein secretion systems in *Pseudomonas aeruginosa*: an essay on diversity, evolution, and function. *Front Microbiol* 2:155. <https://doi.org/10.3389/fmicb.2011.00155>.
- Passmore IJ, Nishikawa K, Lilley KS, Bowden SD, Chung JC, Welch M. 2015. Mep72, a metzincin protease that is preferentially secreted by biofilms of *Pseudomonas aeruginosa*. *J Bacteriol* 197:762–773. <https://doi.org/10.1128/JB.02404-14>.
- Thomassin JL, Santos Moreno J, Guilvout I, Tran Van Nhieu G, Francetic O. 2017. The trans-envelope architecture and function of the type 2 secretion system: new insights raising new questions. *Mol Microbiol* 105:211–226. <https://doi.org/10.1111/mmi.13704>.
- Dunstan RA, Heinz E, Wijeyewickrema LC, Pike RN, Purcell AW, Evans TJ, Praszkiar J, Robins-Browne RM, Strugnell RA, Korotkov KV, Lithgow T. 2013. Assembly of the type II secretion system such as found in *Vibrio cholerae* depends on the novel pilotin AspS. *PLoS Pathog* 9:e1003117. <https://doi.org/10.1371/journal.ppat.1003117>.
- Frickey T, Lupas A. 2004. CLANS: a Java application for visualizing protein families based on pairwise similarity. *Bioinformatics* 20:3702–3704. <https://doi.org/10.1093/bioinformatics/bth444>.
- Weber BS, Kinsella RL, Harding CM, Feldman MF. 2017. The secrets of *Acinetobacter* secretion. *Trends Microbiol* 25:532–545. <https://doi.org/10.1016/j.tim.2017.01.005>.
- Dunstan RA, Hay ID, Wilksch JJ, Schittenhelm RB, Purcell AW, Clark J, Costin A, Ramm G, Strugnell RA, Lithgow T. 2015. Assembly of the secretion pores GspD, Wza and CsgG into bacterial outer membranes does not require the Omp85 proteins BamA or TamA. *Mol Microbiol* 97:616–629. <https://doi.org/10.1111/mmi.13055>.
- Nickerson NN, Tosi T, Dessen A, Baron B, Raynal B, England P, Pugsley AP. 2011. Outer membrane targeting of secretin PulD protein relies on disordered domain recognition by a dedicated chaperone. *J Biol Chem* 286:38833–38843. <https://doi.org/10.1074/jbc.M111.279851>.
- Tosi T, Nickerson NN, Mollica L, Jensen MR, Blackledge M, Baron B, England P, Pugsley AP, Dessen A. 2011. Pilotin-secretin recognition in the type II secretion system of *Klebsiella oxytoca*. *Mol Microbiol* 82:1422–1432. <https://doi.org/10.1111/j.1365-2958.2011.07896.x>.
- Douzi B, Trinh NTT, Michel-Souzy S, Desmyter A, Ball G, Barbier P, Kosta A, Durand E, Forest KT, Cambillau C, Rousset A, Voulhoux R. 2017. Unraveling the self-assembly of the *Pseudomonas aeruginosa* XcpQ secretin periplasmic domain provides new molecular insights into type II secretion system secretin architecture and dynamics. *mBio* 8:e01185-17. <https://doi.org/10.1128/mBio.01185-17>.
- Schrödinger LLC. The PyMOL molecular graphics system, version 1.8. Schrödinger LLC, Cambridge, MA.
- Petersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. 2004. UCSF Chimera—a visualization system for exploratory research and analysis. *J Comput Chem* 25:1605–1612. <https://doi.org/10.1002/jcc.20084>.