



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Interleukin-1 $\beta$ expression by a recombinant porcine reproductive and respiratory syndrome virus

Steven R. Lawson<sup>a</sup>, Yanhua Li<sup>a</sup>, John B. Patton<sup>b</sup>, Robert J. Langenhorst<sup>a</sup>, Zhi Sun<sup>a</sup>, Zhiyong Jiang<sup>a</sup>, Jane Christopher-Hennings<sup>a</sup>, Eric A. Nelson<sup>a</sup>, David Knudsen<sup>a</sup>, Ying Fang<sup>a,\*</sup>, Kyeong-Ok Chang<sup>b,\*\*</sup>

<sup>a</sup> Department of Veterinary and Biomedical Sciences, South Dakota State University, Brookings, SD 57007, USA

<sup>b</sup> Department of Diagnostic Medicine and Pathobiology, Kansas State University, Manhattan, KS 66506, USA

### ARTICLE INFO

#### Article history:

Received 28 July 2011

Received in revised form 3 November 2011

Accepted 11 November 2011

Available online 20 November 2011

#### Keywords:

PRRSV

Reverse genetics

Cytokine

Interleukin-1 beta

### ABSTRACT

The cytokine interleukin-1 beta (IL-1 $\beta$ ) is a potent inflammatory mediator in response to infection, and can be used as an immunological adjuvant. In this study, we constructed a recombinant porcine reproductive and respiratory syndrome virus (vP129/swIL1 $\beta$ ) expressing swine IL-1 $\beta$  from the separate subgenomic mRNA inserted between the ORF1b and ORF2 genome region. MARC-145 cells infected with vP129/swIL1 $\beta$  secreted 1947 pg of IL-1 $\beta$  per  $2 \times 10^5$  cells at 36 h post-infection. *In vitro* growth kinetics analysis in MARC-145 cells showed that the vP129/swIL1 $\beta$  virus had a similar replication rate as that of parental virus. We further performed *in vivo* characterization of the vP129/swIL1 $\beta$  virus in a nursery pig disease model. The vP129/swIL1 $\beta$  infected pigs did not show visible clinical signs, while respiratory distress and lethargy were evident in pigs infected with the parental virus. The expression of various cytokines from peripheral blood mononuclear cells measured by fluorescent microsphere immunoassay showed that IL-1 $\beta$ , IL-4 and IFN- $\gamma$  expression levels were up-regulated in pigs infected with vP129/swIL1 $\beta$  at 7 and 14 days post-infection. However, no detectable level of IL-1 $\beta$  was found in serum samples from pigs infected with either vP129/swIL1 $\beta$  or parental virus. In summary, this study demonstrated a recombinant PRRSV as a useful tool to study the role of different cytokines in disease progression and immune responses, which represents a new strategy for future therapeutic application and vaccine development.

© 2011 Elsevier B.V. All rights reserved.

### 1. Introduction

Cytokines are signaling molecules for intercellular communication such as the induction of immune responses by recruiting immune cells and regulating cell proliferation and differentiation. During viral infection, cytokines play a central role in the innate and adaptive immunity. For example, both Type I and II interferons (IFNs), and various interleukins (ILs) play crucial roles in the control of viral replication at the early stage of virus infection. The innate cytokines are also important in the induction of adaptive immunity specific to the infecting virus by inducing inflammation responses. The excessive inflammation due to viral infection-mediated cytokine production may also lead to immunopathic conditions and pathogenicity of the viral infections. Because

of the important role that cytokines play in the immune response, cytokines have been studied for their application in therapeutic treatment and in vaccine formulations (Andersson and Tracey, 2011; Heegaard et al., 2011; Leong et al., 1994). However, since cytokines have a short half-life and require large dose administration systemically in order to be fully functional, new methods to deliver cytokines to the site of infection have been developed, including the creation of chimeric molecules, delivery vehicles, and the use of viral vectors to express cytokines (Ahlers et al., 2003; Yilma et al., 2010).

Porcine reproductive and respiratory syndrome virus (PRRSV) is a small-enveloped virus containing a single positive-stranded RNA genome. PRRSV is classified in the Order *Nidovirales*, family *Arteriviridae* (Snijder and Meulenber, 1998). Nucleotide sequence comparisons show that PRRSV can be divided into distinct European (Type I) and North American (Type II) genotypes, possessing only about 63% nucleotide identity at the genomic level (Allende et al., 1999; Nelsen et al., 1999). PRRSV causes mild to severe respiratory disease in infected newborn and growing swine, and reproductive failure in pregnant sows (Rossow et al., 1995; Rossow, 1998). PRRSV infection stimulates a minimal level of innate cytokine production. The inflammatory cytokines, including IL-1 $\beta$ ,

\* Corresponding author at: Department of Veterinary and Biomedical Sciences, South Dakota State University Brookings, SD 57007-1396, USA. Tel.: +1 605 688 6647; fax: +1 605 688 6003.

\*\* Corresponding author. Tel.: +1 785 532 3849; fax: +1 785 532 4039.

E-mail addresses: [ying.fang@sdstate.edu](mailto:ying.fang@sdstate.edu) (Y. Fang), [kchang@vet.ksu.edu](mailto:kchang@vet.ksu.edu) (K.-O. Chang).

IL-6 and IL-8, have been detected in virus-infected pigs (Aasted et al., 2002; Suradhat et al., 2003; Suradhat and Thanawongnuwech, 2003; Thanawongnuwech and Thacker, 2003; Bassaganya-Riera et al., 2004; Thanawongnuwech et al., 2004; Lunney et al., 2010). However, in comparison to the other strong immunogenic swine pathogens, such as swine influenza virus and pseudorabies virus, the quantities of secreted cytokines appeared to be much lower. It is suggested that such poor innate anti-viral IFN and cytokine responses result in a weak adaptive immune response.

There are still no therapeutic drugs or completely effective vaccines to control PRRSV. Currently available modified live vaccines have shown the most promising efficacy but have only been partially successful in protection against a wide array of heterologous PRRSV isolates in the field (Bøtner et al., 1997; Mengeling, 2005). In the effort of new vaccine development, cytokines have been evaluated as adjuvants during the immunization of animals (Zuckermann et al., 1998, 1999; Somasundaram et al., 1999; Foss et al., 2002; Murtaugh et al., 2002; Boyaka et al., 2003; Petrovsky and Aguilar, 2004; Royae et al., 2004). Interleukin-1 (IL-1) is a potent inflammatory mediator in response to infection. IL-1 $\beta$  has been tested as an immunological adjuvant because of its ability to influence a broad spectrum of biological activities within the immune system. It was reported that co-administration of recombinant bovine IL-1 $\beta$  and a modified live BHV-1 vaccine was able to enhance both the humoral and cellular immune response against BHV-1 (Reddy et al., 1990, 1993). A recombinant BHV-1 expressing bovine IL-1 $\beta$  showed that both cellular and secreted forms of recombinant IL-1 $\beta$  possessed biological activity, and the expression of IL-1 $\beta$  protein did not affect the *in vitro* growth efficiency of the virus (Raggio et al., 1996). In this study, we explored the strategy of manipulating the immunogenic properties of a virus by expression of a swine IL-1 $\beta$  (swIL1 $\beta$ ) protein using a PRRSV reverse genetic system. A recombinant PRRSV, designated vP129/swIL1 $\beta$ , was generated. This recombinant virus was further tested in a nursery pig model to determine whether swine IL-1 $\beta$  could be successfully delivered *in vivo*. The effect of *in vivo* expression of swine IL-1 $\beta$  on the pathogenic property of the virus as well as the PRRSV specific immune response was further examined.

## 2. Materials and methods

### 2.1. Cells and viruses

BHK-21 cells were used for initial transfection and recovery of recombinant virus. MARC-145 cells were used for recombinant virus rescue and subsequent experiments. These cells were maintained in Eagle's minimum essential medium (Invitrogen) supplemented with 10% heat-inactivated fetal bovine serum and antibiotics at 37 °C with 5% CO<sub>2</sub>. Porcine alveolar macrophages (PAMs) were obtained by lung lavage of 6-week-old PRRSV-naïve piglets using a method described previously (Zeman et al., 1993). Peripheral blood mononuclear cells (PBMCs) were isolated from infected and control pigs as described previously (Meier et al., 2003) and plated in a 24-well plate ( $4 \times 10^6$  cells/ml). PBMCs were stimulated with 50  $\mu$ l UV-inactivated PRRSV ( $2 \times 10^5$  TCID<sub>50</sub>); parallel cultures were stimulated with the mitogen phytohemagglutinin (PHA), or medium, and served as positive and negative controls, respectively. Stimulated cultures were incubated at 37 °C for 24 h. Cell culture supernatants were collected from stimulated cells and stored at -70 °C for cytokine analysis.

The IL-1 $\beta$  expressing recombinant PRRSV (vP129/swIL1 $\beta$ ) was generated from a North American Type II PRRSV full-length cDNA infectious clone, P129 (Calvert et al., 2002, 2003). The third passage of the MARC-145 cell-recovered parental (vP129) or recombinant virus was used for *in vitro* and *in vivo* experiments. Viral titers were

determined by titration on MARC-145 cells and quantified as 50% tissue culture infective dose (TCID<sub>50</sub>)/ml.

### 2.2. Construction of the plasmid (pCMV-S-P129-1bswIL1 $\beta$ ) for generation of the IL1 $\beta$ expressing recombinant PRRSV (vP129/swIL1 $\beta$ )

The plasmid pCMV-S-P129-1bswIL1 $\beta$  was constructed using pCMV-S-P129-1bGFP2 (Kermit) (Calvert et al., 2009) in which GFP was inserted in the region between ORF1b and ORF2 with *AflIII* and *MluI* enzyme sites (Fig. 1A). The expression of GFP in the construct was derived from the leader/junction (L/J) sequence for ORF2, and the ORF6 L/J was inserted downstream of the GFP gene to drive the expression of ORF2, so that GFP is expressed from the independent subgenomic RNA. The GFP gene was replaced with swine IL-1 $\beta$  gene that was amplified from PAM by RT-PCR using primer pairs of IL-1b-Afl-F (5'-aattcttaagatggccatagctactgaacc) and IL-1b-Mlu-R (5'-aattacgcgtttaggagagaggacttc). The amplicon was digested with *AflIII* and *MluI* enzymes and cloned into the plasmid after it was digested with the same enzyme and removed the GFP gene.

The plasmid pCMV-S-P129-1bswIL1 $\beta$  was transfected into BHK-21 cells as described previously (Calvert et al., 2002, 2003). To rescue the virus, cell culture supernatant obtained at 48 h post-transfection was passaged on MARC-145 cells. Rescue of infectious virus was determined by an indirect immunofluorescent assay (IFA) using PRRSV-specific monoclonal antibody (mAb) SDOW17 as we described previously (Fang et al., 2006, 2008). The expression of swIL1 $\beta$  recombinant protein from the vP129/swIL1 $\beta$  in MARC-145 cells and PAMs was confirmed by Western blot using the polyclonal antibody to the protein (R&D systems).

### 2.3. *In vitro* characterization of vP129/swIL1 $\beta$ recombinant virus


The growth kinetics of the recombinant and parental viruses were examined by infecting MARC-145 cells with viruses at a multiplicity of infection (MOI) of 0.5. Infected cells were collected at 0, 6, 12, 24, 36, 48, 60 and 72 h post infection (hpi), and viral titers were determined by immunofluorescent assay (IFA) on MARC-145 cells and quantified as fluorescent focus units (FFU)/ml. Plaque morphology of each virus was determined by plaque assay on MARC-145 cells as described previously (Fang et al., 2006). To determine the expression of swine IL-1 $\beta$ , the culture supernatant collected at different time points was analyzed using a fluorescent microsphere immunoassay (FMIA) as we described previously (Lawson et al., 2010).

### 2.4. Sequencing of deletion regions

To determine the stability of the IL-1 $\beta$  gene in the recombinant virus, cell lysate from virus-infected cells was harvested, and RNA was extracted using a QIAamp viral RNA kit (Qiagen) following the manufacturer's instruction. The corresponding insertion region was amplified by RT-PCR using primers p129-11931F (5'-gggaggattacaatgatcgctttgc) and p129-12159R (5'-gatggtgaagccaacaataatggcc). PCR products were sequenced at the Iowa State University DNA sequencing facility (Ames, IA, USA).

### 2.5. Animals and infection groups

Eighteen 3-week-old, PRRSV-naïve pigs obtained from a certified PRRSV-negative herd were divided randomly into three groups and housed separately in animal isolation facilities at South Dakota State University (SDSU). After a four-day acclimation period, pigs from each group ( $n = 6$ ) were inoculated with 2 ml ( $1 \times 10^6$  TCID<sub>50</sub>,



**Fig. 1.** Expression of swIL1 $\beta$  by vP129/swIL1 $\beta$  recombinant virus. (A) Construction of the plasmid, pCMV-S-P129-1bswIL1 $\beta$ . The GFP gene in pCMV-S-P129-1bGFP2 (Kermit) backbone was replaced by swine IL-1 $\beta$  gene. The IL-1 $\beta$  is expressed as an independent subgenomic RNA, which was derived from the leader/junction (L/J) sequence of ORF2. The ORF6 L/J was inserted downstream of the IL-1 $\beta$  gene to drive the expression of ORF2; (B) Western blot analysis of the detection of swIL1 $\beta$  in the supernatant fraction (sup) or cell lysates (cell lys) of MARC-145 cells infected with vP129/swIL1 $\beta$  or parental P129 virus. Cells were infected with the MOI of 1 and incubated for 24 h before sample preparation for Western blot analysis, and (C) kinetics of accumulation of swine IL-1 $\beta$  in culture supernatant of MARC-145 cells infected with vP129/swIL1 $\beta$  or parental virus. MARC-145 cells were infected in parallel at MOI of 0.5, and culture supernatant was collected at indicated time points. The swine IL-1 $\beta$  expression levels were determined by Luminex fluorescent microsphere immunoassay (FMIA).

given as 2 ml intranasally) of parental virus (group 1), recombinant virus (group 2) or mock-infected with cell culture medium (group 3).

## 2.6. Clinical signs and serum and tissue sampling

Clinical signs and rectal temperatures were monitored every day during the first five days post-inoculation. Clinical signs were recorded and quantified using a clinical scoring system adapted from Martínez-Lobo et al. (2011). The scoring system was designed to evaluate clinical respiratory signs including sneezing, nasal secretions, coughing and labored breathing. In addition, systemic clinical signs including hair roughness, lethargy and anorexia were also recorded and scored. Severity of clinical symptoms was evaluated on a scale of 0–5 whereby a score of 5 was considered to be the most severe. Blood samples were obtained from all pigs on days 0, 3, 7, 14, 21 and 28 dpi. Pigs were euthanized at 28 dpi to evaluate macroscopic and microscopic lung lesions. Lung lesions of the study animals were evaluated using a previously developed system (Halbur et al., 1995) based on the approximate volume that each lobe contributes to the entire lung: the left and right apical lobes, the left and right cardiac lobes, and the intermediate lobe each contribute 10% of the total lung volume, and the left and right caudal lobes each contribute 25%. These scores were then used to calculate the total lung lesion score based on the relative percentage contributions of each lobe. Blood and tissue [lungs, tonsils and lymph nodes (submandibular lymph node, lateral retropharyngeal lymph node, mesenteric lymph nodes)] samples were collected at the time of euthanasia. For tissue sampling, a representative tissue was collected from each lobe of the lung, while the entire tonsil and lymph node were collected from each pig.

All animal experiments were approved and guided by the Institutional Animal Care and Use Committee (IACUC) at South Dakota State University in accordance with Animal Welfare Assurance Guidelines.

## 2.7. Real-time RT-PCR quantification of viral load in infected animals

For the detection of viral RNA and determination of viral load, serum and tissue samples were examined using a real-time quantitative RT-PCR (Tetracore VetAlert PRRS), which is routinely performed at the South Dakota Animal Disease Research and Diagnostic Laboratory (SD-ADRDL) under strict quality assurance guidelines. The detailed method has been described in our previous publication (Wasilk et al., 2004).

## 2.8. Determination of humoral immunity

To compare the humoral antibody response of the treatment groups, all serum samples were evaluated using an IDEXX Herd-Chek PRRS 2XR ELISA following the manufacturer's instruction. The virus neutralization assay was performed as described previously (Yoon et al., 1994).

## 2.9. FMIA for cytokine expression measurement

Serum samples and culture supernatant from PBMCs were evaluated for expression of selected swine cytokine proteins targeting immune markers for innate immunity: IL-1 $\beta$ , IL-8, IFN- $\alpha$ , TNF- $\alpha$ , IL-12; Th1 immunity: IFN- $\gamma$ ; Th2 immunity: IL-4; and regulatory T cell response: IL-10. Detailed FMIA procedure for cytokine protein detection was described in our previous publication (Lawson et al., 2010). Relative quantification of target cytokine expression was evaluated using mean fluorescent intensity (MFI) values, and each group's results were compared with the mean values for control pigs (or control cultures).

### 2.10. Statistical analysis

Statistical analysis was performed using GraphPad InStat version 3.06 (GraphPad Software). Comparison was performed by one-way analysis of variance with Tukey's multiple comparison tests to determine mean significance. Differences between groups of pigs at different times post-inoculation were considered statistically significant at  $P < 0.05$  for all analyses.

## 3. Results

### 3.1. Expression of swIL1 $\beta$ by recombinant virus


The transfection of pCMV-S-P129-1bswIL1 $\beta$  plasmid DNA into BHK cells yielded the recombinant virus vP129/swIL1 $\beta$ . The vP129/swIL1 $\beta$  viruses were rescued by passage of the cell culture supernatant onto MARC-145 cells, which was evident by IFA staining with anti-nucleocapsid protein monoclonal antibody SDOW17. To detect the IL-1 $\beta$  protein expression in infected cells, Western blot was conducted. As shown in Fig. 1B, a protein band with apparent molecular weight between 17 and 20 kDa was recognized by anti-swIL1 $\beta$  antibody in the cellular and supernatant fractions of vP129/swIL1 $\beta$  infected cells, but not in the wild-type virus infected cells. To measure the amount of swIL1 $\beta$  protein secreted from infected cells, cell culture supernatant collected at 6, 12, 18, 24 and 36 hpi was analyzed (Fig. 1C). The concentration of swIL1 $\beta$  was 400 pg at 18 hpi and reached the maximum concentration of 1947 pg at 36 h, which corresponds to about 2 ng per  $2 \times 10^5$  cells. We further assessed the stability of swine IL-1 $\beta$  gene in recombinant virus by serial passage of the vP129/swIL1 $\beta$  virus at least 10 times on MARC-145 cells. The IL-1 $\beta$  insertion region was analyzed. Total cellular RNA was isolated from cells infected with the passage 3 and 10 of the recombinant virus, and RNA was used as the template for RT-PCR amplification of the insertion region. The PCR product was sequenced, and results showed that the IL-1 $\beta$  gene remained intact as a full-length gene, and no mutation or deletion was detected.

### 3.2. *In vitro* growth characteristics of vP129/swIL1 $\beta$ recombinant virus

To determine whether the insertion/expression of swIL1 $\beta$  has an effect on viral replication, we compared the *in vitro* growth properties of the vP129/swIL1 $\beta$  with that of parental viruses. MARC-145 cells were infected with each of the viruses and harvested at 6, 12, 24, 36, 48, 60, and 72 hpi. Growth kinetics analysis showed that the vP129/swIL1 $\beta$  virus possessed a similar growth rate to that of the parental virus (Fig. 2A). Titers peaked at 48 hpi for both viruses. The peak titer of the vP129/swIL1 $\beta$  virus was  $3.7 \times 10^5$  FFU/ml, compared to  $3.0 \times 10^5$  FFU/ml for the parental virus. Plaque morphology of these viruses was also determined, and the plaque size produced by the vP129/swIL1 $\beta$  virus was similar to the parental virus (Fig. 2B). These results indicate that the insertion of swine IL-1 $\beta$  into the viral genome does not affect the viral replication *in vitro*.

### 3.3. *In vivo* growth and pathogenic property of vP129/swIL1 $\beta$ virus

A total of 18 three-week-old pigs, divided into three groups, were infected with vP129/swIL1 $\beta$  virus, parental virus or cell culture medium (mock infected). The clinical signs and body temperature were monitored during the first five days post-inoculation. As shown in Table 1, all pigs inoculated by parental virus developed clinical signs, showing respiratory distress (coughing, serious nasal discharge) and lethargy, and four pigs had rectal



**Fig. 2.** *In vitro* growth characterization of vP129/swIL1 $\beta$  recombinant virus. (A) Growth kinetics comparison between vP129/swIL1 $\beta$  and parental virus in MARC-145 cells. MARC-145 cells were infected in parallel at MOI of 0.1 with the passage three of chimeric or parental virus. At 6, 12, 24, 36, 48, 60, and 72 h post-infection, cells were harvested, and the virus titers were determined by IFA on MARC-145 cells. The results were mean values from three replications of the experiment, and viral titers were expressed as fluorescence focus units per milliliter (FFU/ml), and (B) Plaque morphology of vP129/swIL1 $\beta$  and parental virus. Confluent cell culture monolayers were infected with viruses at a MOI of 0.1. After 2 h infection, cell culture supernatant was removed and an agar overlay was applied. Plaques were detected after five days incubation at 37 °C and stained by 0.1% crystal violet.

temperatures over 40 °C during the second and third day post-inoculation. In contrast, no apparent clinical signs were observed in those pigs inoculated by vP129/swIL1 $\beta$  virus, and rectal temperature was normal for this group of pigs through the time course of the study. All pigs given virus became infected, which was evident by *in vitro* virus isolation and positive real-time PCR results for the presence of viral RNA in serum. Viral load in serum peaked at 7 dpi (Fig. 3A) for both vP129/swIL1 $\beta$  and parental virus infected groups. Statistically, there was no significant difference in viral load between these two groups, although a slightly lower level of viral load in serum samples from those pigs infected with wild type virus was observed at 14, 21 and 28 dpi. Interestingly, in comparison to the group of pigs infected with wild type virus, a higher level of viral load was observed in the lymph nodes, tonsil and lung tissues from those pigs infected with vP129/swIL1 $\beta$  at 28 dpi (Fig. 3B). At necropsy (28 dpi), minimal lung lesions (Table 1) were observed in vP129 and vP129/swIL1 $\beta$  infected groups. Enlarged mandibular lymph nodes were observed in three of the pigs from the vP129/swIL1 $\beta$  infected group, and in two of the pigs from the vP129 infected group.

To determine the *in vivo* stability of the swIL1 $\beta$  gene in virus, serum samples were used for virus isolation in MARC-145 cells. Virus was recovered from serum samples collected on days 7, 14 and 21 dpi. For each dpi, we sequenced viruses isolated from six pigs, and sequencing was performed using both forward and reverse primers, resulting in a total of 12 sequences for each day analyzed. The results confirmed that the swIL1 $\beta$  gene remained intact as a full-length gene. However, sequencing results revealed that an A299G nucleotide mutation was consistently identified in two of the pigs, which resulted in the amino acid (aa) mutation of Glu to Gly at position 100 of IL-1 $\beta$  protein. Another T553C nucleotide (Cys to Arg) mutation was also found in the virus isolated from a pig at 7 dpi. These results suggest that selection occurred to generate mutations during the viral replication *in vivo*.

**Table 1**  
Evaluation of clinical symptoms and lung lesions for each experimental group.

Clinical symptom <sup>a</sup> and lung lesion <sup>b</sup>	Group scores (mean ± SEM)		
	IL-1β	Parental	Mock
Rectal temp during 5 days post inoculation	39.6 (±0.1)	40.6 (±0.5)	39.7 (±0.3)
Clinical respiratory signs			
Sneezing	0	1.0 (±0.0)	0
Nasal secretions	0.3 (±0.1)	2.0 (±0.0)	0
Coughing	0	2.0 (±0.3)	0
Labored breathing	0	0	0
Systemic clinical signs			
Roughness of hair	0	0	0
Lethargy	0	3.0 (±0.0)	0
Anorexia	0	0	0
Gross lung lesion score	2.0 (±0.5)	2.2 (±0.5)	0

<sup>a</sup> Clinical signs were recorded and quantified using a clinical scoring system adapted from Martínez-Lobo et al. (2011). Severity of clinical symptoms was evaluated on a scale of 0–5 whereby a score of 5 was considered to be the most severe.

<sup>b</sup> Gross lung pathology was assessed by using a gross pig lung lesion scoring system (Halbur et al., 1995) where each lobe of the lung was evaluated for percent pneumonia and the percent pneumonia of each lobe was added for the entire lung.

### 3.4. *In vivo* swIL1β protein expression and immune response in pigs


We further investigated whether *in vivo* expression of swIL1β protein could modulate the viral specific host immune response. In an earlier study, swIL1β was detected in sera or peripheral blood mononuclear cells (PBMCs) (by the fluorescent microsphere immunoassay, FMIA) from PRRSV-infected pigs (Lawson et al., 2010). To compare with these earlier findings, samples were collected from control and virus-infected pigs at 3, 7 and 14 dpi. PBMCs were stimulated *in vitro* with parental virus, PHA, or medium for 24 h. The FMIA results showed that higher levels of swIL1β protein were expressed in PBMCs from pigs infected by PRRSV, either vP129/swIL1β or parental vP129 compared to control pigs at 3, 7 and 14 dpi (Fig. 4). The comparison between vP129/swIL1β and parental vP129 showed that a higher level of swIL1β protein was expressed in pigs infected by vP129/swIL1β at 14 dpi (Fig. 4). No significant difference in swIL1β protein expression was detected from both groups of pigs infected by vP129/swIL1β and parental

virus at 3 and 7 dpi by the FMIA. Interestingly, no detectable level of swIL1β was observed in serum samples from all groups of pigs at 3, 7 and 14 dpi (data not shown). We further monitored the temporal expression of selected innate and cellular immune proteins in PBMCs from virus-infected pigs using FMIA. As shown in Fig. 4, besides IL-1β, IFN-γ expression levels were also significantly higher in pigs infected with vP129/swIL1β compared to pigs with vP129 at 14 dpi. A significantly higher level of IL-4 and TNF-α was also observed in vP129/swIL1β infected pigs compared to pigs with vP129 at 14 dpi. IL-8 and IL-12 expression were higher at an early time point (3 dpi) but lower at 7 dpi in vP129/swIL1β infected pigs compared to pigs infected with vP129. The expression of IL-10 was lower at an early time point (3 dpi) but was higher at 7 dpi in vP129/swIL1β infected pigs compared to pigs infected with vP129. The comparison between virus infected groups (either vP129/swIL1β or parental vP129) and the control group showed that the expression of IL-12 or IL-10 in PBMCs was either up- or down-regulated by virus infection at 3 and 14 dpi.


Further analysis of host humoral immune response showed that both vP129/swIL1β and parental virus infected groups had seroconverted by 14 dpi (Fig. 5A). Interestingly, a higher level of neutralizing antibody response was observed in the group of pigs infected with vP129/swIL1β at 28 dpi (Fig. 5B). Three of the six pigs (50%) in the vP129/swIL1β virus infected group had detectable neutralizing antibody titers by 28 dpi, while two of the six pigs (33%) in the parental virus infected group developed neutralizing antibodies by 28 dpi. Taken together, these results suggested that *in vivo* expression of swIL1β significantly modulated the PRRSV specific immune response.

## 4. Discussion

PRRSV infection appears to elicit poor innate IFN and cytokine responses unlike the other strongly immunogenic viral pathogens. Its effects result in a weak adaptive immune response as demonstrated by prolonged viremia, short duration of cell-mediated immune responses and slow development of virus-specific IFN-γ secreting cells. Anti-PRRSV neutralizing antibodies are generated slowly and their titers remain low. There is an extensive body of literature describing the innate interferon and cytokine effect on PRRSV replication and cytokine-mediated events that occur during PRRSV infection. In an earlier study, inoculation of pigs with porcine respiratory coronavirus, a potent inducer of innate cytokines, provided protection from a subsequent PRRSV infection (Buddaert et al., 1998). Overend et al. (2007) showed that recombinant swine IFN-β protects swine alveolar macrophages from infection with PRRSV. Royae et al. (2004) showed that using an expression




**Fig. 3.** Viral load in pigs infected with vP129/swIL1β or parental virus. Viral load was quantified by real-time RT-PCR, and the result was interpreted as RNA copy numbers per ml. (A) Viral load in serum samples, and (B) Viral load in tissue samples. Different letters ("a" and "b") represent mean viral load of different groups differs significantly ( $P < 0.05$ ).



**Fig. 4.** Comparison of cytokine expression levels in activated PBMCs from pigs infected with vP129/swIL1β with that of parental virus. PBMCs were harvested at 3, 7 and 14 dpi and stimulated in culture with PHA or medium. Cell culture supernatants and pellets were harvested at 24 h post-stimulation. Cytokine protein expression was determined by Luminex fluorescent microsphere immunoassay. Each data point was interpolated using a five-parameter logistic regression curve and displayed as a mean fluorescent unit from three animals with the standard deviation. Different letters (“a”, “b” and “c”) represent mean cytokine level of different groups differs significantly ( $P < 0.05$ ).

plasmid encoding porcine IFN-α as an adjuvant resulted in a temporary increase in the frequency of PRRSV specific IFN-γ secreting cells in the vaccinated animals. These results all support an important role of IFNs and cytokines in the initial innate immune

response and the subsequent activation of adaptive immunity. However, with PRRSV infection, innate immune genes and proteins expression is suppressed and delayed (Van Reeth et al., 1999, 2002; Murtaugh et al., 2002). This initial suppression of the innate



**Fig. 5.** Antibody response in pigs infected with vP129/swIL1β or parental virus. (A) PRRSV-specific serum antibodies measured by IDEXX HerdChek® PRRSV ELISA 2XR kit. S/P ratios of greater than 0.4 are considered positive. (B) Serum neutralizing antibody response determined by fluorescent focus neutralization assay. Results were interpreted as a 90% reduction of the viral infection, and the neutralizing antibody titers were presented as mean value ( $n = 6$ ) and expressed on a  $\log_2$  scale. The parental virus was used for the viral neutralizing assay. Different letters (“a”, “b” and “c”) represent mean neutralizing antibody titer of different groups differs significantly ( $P < 0.05$ ).

immune response, leading to the delayed induction of protective cellular and humoral immunity, provides a window of time that allows PRRSV to replicate, shed and transmit to other contact naïve animals. Therefore, current strategies for vaccine development are directed at constructing a PRRS vaccine capable of inducing a high level of innate and cellular immune responses and robust neutralizing antibody activity within a short period of time after vaccination.

The use of cytokines as adjuvants for enhancing the viral specific immune response has been investigated in both subunit and modified live vaccines (Heath and Playfair, 1992; Gao et al., 1995; Godson et al., 1995; Ahlers et al., 2003; Yilma et al., 2010; Heegaard et al., 2011). In particular, IL-1 $\beta$  has been tested as a potent immune modulator in vaccine development. The recombinant bovine IL-1 $\beta$  has been tested as an adjuvant to co-administrate with a modified live BHV-1 vaccine, and an *in vivo* study showed enhanced humoral and cellular immune response. Although the study demonstrated the effect of IL-1 $\beta$  on the enhancement of host immunity, the systemic administration of IL-1 $\beta$  to develop an adjuvant effect requires a large dose of recombinant IL-1 $\beta$  and multiple injections at a local site. This could result in severe side effects and is not cost-effective in veterinary medicine. The use of recombinant virus as a vector to express the cytokines, such as IL-1 $\beta$ , has the potential to overcome these drawbacks. The viral vector system allows delivery of the cytokine to the specific site of infection. It reduces the dose requirement and thereby decreases the cost and side effects. A number of cytokines have been tested by inserting into recombinant viruses (Bembridge et al., 1998; Karaca et al., 1998; Giavedoni et al., 1997; Leong et al., 1994; Flexner et al., 1990). For example, expression of IFN- $\gamma$  by recombinant respiratory syncytial virus attenuated the virus *in vivo* and protected animals from subsequent challenge with wild type virus (Bukreyev et al., 1999). In vaccinia virus, expression of IL-2 attenuated the virus while maintained the immunogenicity *in vivo* (Flexner et al., 1990). In our study, a recombinant PRRSV (vP129/swIL1 $\beta$ ) was constructed, which expresses the IL-1 $\beta$  gene as a separate mRNA from an additional transcriptional product placed between ORF1b and ORF2 genome region. The recombinant virus produced high levels of IL-1 $\beta$  protein in cell culture. *In vitro* growth kinetics analysis indicated that the production of swIL1 $\beta$  was not toxic to the cells and did not affect viral replication. In comparison to the parental virus, the vP129/swIL1 $\beta$  virus grew to a viral titer similar to that of the parental virus in MARC-145 cells. This property is important for recombinant vaccine development. Genetic manipulation/attenuation of the live virus are typically accompanied by reduced viral titer. However, to achieve a satisfactory level of immunogenicity, the virus titer must be adequately maintained. The ability to produce high titers of vP129/swIL1 $\beta$  virus *in vitro* makes it easy to produce sufficient quantities for large-scale vaccine production. In addition, the large amount of recombinant IL-1 $\beta$  protein produced by the recombinant virus could be an alternative method for the *in vitro* production of recombinant cytokines.

We further demonstrated that the recombinant PRRSV could be used as an *in vivo* cytokine delivery system to study the mechanism of different cytokine effects on disease progression. First, the expression of IL-1 $\beta$  by vP129/swIL1 $\beta$  did not show a difference in viral load in the sera compared to those by the parental virus in animals. Also, our data showed that the IL-1 $\beta$  gene is maintained in the virus isolated from serum samples of the vP129/swIL1 $\beta$  infected pigs. Most importantly, the *in vivo* co-expression of swIL1 $\beta$  attenuated the pathogenic property of the virus. The parental virus P129 is known to be a highly pathogenic virus (Calvert et al., 2002, 2003). Results from our study demonstrated that parental P129 virus infected pigs showed significant symptoms of lethargy and respiratory distress. In contrast, the swIL1 $\beta$  expressing vP129/swIL1 $\beta$  virus infected pigs did not show apparent clinical symptoms. It is not clear why the expression of IL-1 $\beta$  by vP129/swIL1 $\beta$  resulted

in attenuated symptoms in the animals. Initially, we hypothesized that vP129/swIL1 $\beta$  might be more virulent than the parental virus because of the expression of the proinflammatory cytokine IL-1 $\beta$ . It is possible that the attenuation may be related to the site (alveolar region) of virus replication and the expression of the cytokine where the anti-inflammatory mechanism is strong (Spight et al., 2005). Another interesting phenomenon related to this is that although the viral load in blood was similar, a higher level of viral load in tissues (tonsil, lymph nodes, and lung) was observed in pigs infected with vP129/swIL1 $\beta$  at 28 dpi. In this case, the viral load at the local tissues did not correlate with the disease progression, since the vP129/swIL1 $\beta$  infected pigs did not develop apparent clinical symptoms and no significant lung lesion was observed. It is possible that the continued expression of IL-1 $\beta$  in virus infected cells may promote virus replication by recruiting monocytes to differentiate into macrophages in the local tissues. Currently, we are investigating the potential mechanism of the attenuation by IL-1 $\beta$  in virus-infected animals.

Although no detectable level of swIL1 $\beta$  was observed in sera from animals infected with vP129/swIL1 $\beta$  or parental vP129, there were significantly higher levels of IL-1 $\beta$  detected in vP129/swIL1 $\beta$  infected pigs in PBMCs (Fig. 4). Further analysis of a time course pattern of cytokine proteins expression using a FMIA suggested that the co-expression of IL-1 $\beta$  could lead to a strong T cell response. In comparison to the parental virus, infection with vP129/swIL1 $\beta$  resulted in a significant increase of both Th1 marker (IFN- $\gamma$ ) and Th2 marker (IL-4) in PBMCs at 14 dpi. The result of the humoral immune response showed that both vP129/swIL1 $\beta$  and parental virus infected groups had seroconverted by 14 dpi at the same levels (Fig. 5A). However, a higher level of neutralizing antibody response was observed in the group of pigs infected with vP129/swIL1 $\beta$  compared to that with parental virus at 28 dpi (Fig. 5B), which may be due to the strong T cell response by vP129/swIL1 $\beta$ .

In summary, our study demonstrated that the recombinant PRRSV with the reverse genetics system can be used as a potential *in vivo* cytokine delivery system to study the role of different cytokines in disease progression and immune responses. This study represents a new approach for future *in vivo* application of therapeutic cytokines and development of genetic engineered vaccines for controlling PRRS.

## Acknowledgements

We thank Jay G. Calvert (Pfizer Animal Health, MI, USA) for providing pCMV-S-P129-1bGFP2 (Kermit) plasmid. Special thanks to Russell Ransburgh, Brianna Sandager, Travis Hawkins, Lyle Brown, Elizabeth Brown and Diane Baker for excellent technical assistance in animal study. This project was supported by the National Research Initiative of the USDA Cooperative State Research, Education and Extension Service (to Y. Fang, grant# 2007-01745).

## References

- Aasted, B., Bach, P., Nielsen, J., Lind, P., 2002. Cytokine profiles in peripheral blood mononuclear cells and lymph node cells from piglets infected in utero with porcine reproductive and respiratory syndrome virus. *Clin. Diagn. Lab. Immunol.* 9 (6), 1229–1234.
- Ahlers, J.D., Belyakov, I.M., Berzofsky, J.A., 2003. Cytokine, chemokine, and costimulatory molecule modulation to enhance efficacy of HIV vaccines. *Curr. Mol. Med.* 3, 285–301.
- Allende, R., Lewis, T.L., Lu, Z., Rock, D.L., Kutish, G.F., Ali, A., Doster, A.R., Osorio, F.A., 1999. North American and European porcine reproductive and respiratory syndrome viruses differ in non-structural protein coding regions. *J. Gen. Virol.* 80, 307–315.
- Andersson, U., Tracey, K.J., 2011. Molecular basis of applied biological therapeutics. *J. Intern. Med.* 269, 2–7.
- Bassaganya-Riera, J., Thacker, B.J., Yu, S., Strait, E., Wannemuehler, M.J., Thacker, E.L., 2004. Impact of immunizations with porcine reproductive and respiratory syndrome virus on lymphoproliferative recall responses of CD8<sup>+</sup> T cells. *Viral Immunol.* 17, 25–37.



- Bembridge, G.P., Lopez, J.A., Cook, R., Melero, J.A., Taylor, G., 1998. Recombinant vaccinia virus coexpressing the F protein of respiratory syncytial virus (RSV) and interleukin-4 (IL-4) does not inhibit the development of RSV-specific memory cytotoxic T lymphocytes, whereas priming is diminished in the presence of high levels of IL-2 or gamma interferon. *J. Virol.* 72, 4080–4087.
- Bøtner, A., Strandbygaard, B., Sorensen, K.J., Have, P., Madsen, K.G., Madsen, E.S., Alexandersen, S., 1997. Appearance of acute PRRS-like symptoms in sow herds after vaccination with a modified live PRRS vaccine. *Vet. Rec.* 141, 497–499.
- Boyaka, P.N., Tafaro, A., Fischer, R., Fujihashi, K., Jirillo, E., McGhee, J.R., 2003. Therapeutic manipulation of the immune system: enhancement of innate and adaptive mucosal immunity. *Curr. Pharm. Des.* 9, 1965–1972 (Review).
- Buddaert, W., Van Reeth, K., Pensaert, M., 1998. In vivo and in vitro interferon (IFN) studies with the porcine reproductive and respiratory syndrome virus (PRRSV). *Adv. Exp. Med. Biol.* 440, 461–467.
- Bukreyev, A., Whitehead, S.S., Bukreyeva, N., Murphy, B.R., Collins, P.L., 1999. Interferon  $\gamma$  expressed by a recombinant respiratory syncytial virus attenuates virus replication in mice without compromising immunogenicity. *Proc. Natl. Acad. Sci. U.S.A.* 96, 2367–2372.
- Calvert, J.G., Sheppard, M.G., Welch, S.K.W., 2002. Infectious cDNA clone of North American porcine reproductive and respiratory syndrome (PRRS) virus and use thereof. US Patent 6,500,662.
- Calvert, J.G., Sheppard, M.G., Welch, S.K.W., 2003. Infectious cDNA clone of North American porcine reproductive and respiratory syndrome (PRRS) virus and use thereof. US Patent 20,030,157,689.4.
- Calvert, J., Sheppard, G., Michael, G., Welch, S.K.W., 2009. Infectious cDNA clone of North American porcine reproductive and respiratory syndrome (PRRS) virus and uses thereof. US Patent 7618797.
- Fang, Y., Rowland, R.R., Roof, M., Lunney, J.K., Christopher-Hennings, J., Nelson, E.A., 2006. A full-length cDNA infectious clone of North American type 1 porcine reproductive and respiratory syndrome virus: expression of green fluorescent protein in the nsp2 region. *J. Virol.* 80, 11447–11455.
- Fang, Y., Christopher-Hennings, J., Brown, E., Liu, H., Chen, Z., Lawson, S., Breen, R., Clement, T., Gao, X., Bao, J., Knudsen, D., Daly, R., Nelson, E., 2008. Development of genetic markers in the nonstructural protein 2 region of a US type 1 porcine reproductive and respiratory syndrome virus: implications for future recombinant marker vaccine development. *J. Gen. Virol.* 89, 3086–3096.
- Flexner, C., Moss, B., London, W.T., Murphy, B.R., 1990. Attenuation and immunogenicity in primates of vaccinia virus recombinants expressing human interleukin-2. *Vaccine* 8, 17–21.
- Foss, D.L., Zilliox, M.J., Meier, W., Zuckermann, F., Murtaugh, M.P., 2002. Adjuvant danger signals increase the immune response to porcine reproductive and respiratory syndrome virus. *Viral Immunol.* 15, 557–566.
- Gao, Y., Daley, M.J., Splitter, G.A., 1995. BHV-1 glycoprotein 1 and recombinant interleukin 1 beta efficiently elicit mucosal IgA response. *Vaccine* 13, 871–877.
- Giavedoni, L., Ahmad, S., Jones, L., Yilma, T., 1997. Expression of gamma interferon by simian immunodeficiency virus increases attenuation and reduces postchallenge virus load in vaccinated rhesus macaques. *J. Virol.* 71, 866–872.
- Godson, D.L., Baca-Estrada, M.E., Van Kessel, A.G., Hughes, H.P., Morsy, M.A., Van Donkersgoed, J., Harland, R.J., Shuster, D.E., Daley, M.J., Babiuk, L.A., 1995. Regulation of bovine acute phase responses by recombinant interleukin-1 beta. *Can. J. Vet. Res.* 59, 249–255.
- Halbur, P.G., Paul, P.S., Frey, M.L., Landgraf, J., Eernisse, K., Meng, X.J., Lum, M.A., Andrews, J.J., Rathje, J.A., 1995. Comparison of the pathogenicity of two US porcine reproductive and respiratory syndrome virus isolates with that of the Lelystad virus. *Vet. Pathol.* 32, 648–660.
- Heath, A.W., Playfair, J.H., 1992. Cytokines as immunological adjuvants. *Vaccine* 10, 427–434 (Review. Erratum in: *Vaccine* 10, 808).
- Heegaard, P.M., Dedieu, L., Johnson, N., Le Potier, M.F., Mockey, M., Mutinelli, F., Vahlenkamp, T., Vascellari, M., Sørensen, S.N.S., 2011. Adjuvants and delivery systems in veterinary vaccinology: current state and future developments. *Arch. Virol.* 156, 183–202.
- Karaca, K., Sharma, J.M., Winslow, B.J., Junker, D.E., Reddy, S., Cochran, M., McMillen, J., 1998. Recombinant fowlpox viruses coexpressing chicken type I IFN and Newcastle disease virus HN and F genes: influence of IFN on protective efficacy and humoral responses of chickens following in ovo or post-hatch administration of recombinant viruses. *Vaccine* 16, 1496–1503.
- Lawson, S., Lunney, J., Zuckermann, F., Osorio, F., Nelson, E., Welbon, C., Clement, T., Fang, Y., Wong, S., Kulas, K., Christopher-Hennings, J., 2010. Development of an 8-plex Luminex assay to detect swine cytokines for vaccine development: assessment of immunity after porcine reproductive and respiratory syndrome virus (PRRSV) vaccination. *Vaccine* 28, 5356–5364.
- Leong, K.H., Ramsay, A.J., Boyle, D.B., Ramshaw, I.A., 1994. Selective induction of immune responses by cytokines coexpressed in recombinant fowlpox virus. *J. Virol.* 68, 8125–8130.
- Lunney, J.K., Fritz, E.R., Reedy, J.M., Kuhar, D., Prucnal, E., Molina, R., Christopher-Hennings, J., Zimmerman, J., Rowland, R.R., 2010. Interleukin-8, interleukin-1 $\beta$ , and interferon- $\gamma$  levels are linked to PRRS virus clearance. *Viral Immunol.* 23, 127–134.
- Martínez-Lobo, F.J., Díez-Fuertes, F., Segalés, J., García-Artiga, C., Simarro, I., Castro, J.M., Prieto, C., 2011. Comparative pathogenicity of type 1 and type 2 isolates of porcine reproductive and respiratory syndrome virus (PRRSV) in a young pig infection model. *Vet. Microbiol.* (Epub ahead of print).
- Meier, W.A., Galeota, J., Osorio, F.A., Husmann, R.J., Schnitzlein, W.M., Zuckermann, F.A., 2003. Gradual development of the interferon-gamma response of swine to porcine reproductive and respiratory syndrome virus infection or vaccination. *Virology* 309, 18–31.
- Mengeling, W.L., 2005. The porcine reproductive and respiratory syndrome quary. Part II: vaccines and vaccination strategy. *J. Swine Health Prod.* 13, 152–156.
- Murtaugh, M.P., Xiao, Z., Zuckermann, F., 2002. Immunological responses of swine to porcine reproductive and respiratory syndrome virus infection. *Viral Immunol.* 15, 533–547.
- Nelsen, C.J., Murtaugh, M.P., Faaberg, K.S., 1999. Porcine reproductive and respiratory syndrome virus comparison: divergent evolution on two continents. *J. Virol.* 73, 270–280.
- Overend, C., Mitchell, R., He, D., Rompato, G., Grubman, M.J., Garmendia, A.E., 2007. Recombinant swine beta interferon protects swine alveolar macrophages and MARC-145 cells from infection with porcine reproductive and respiratory syndrome virus. *J. Gen. Virol.* 88, 925–931.
- Petrovsky, N., Aguilar, J.C., 2004. Vaccine adjuvants: current state and future trends. *Immunol. Cell. Biol.* 82, 488–496.
- Raggio, C., Fitzpatrick, D.R., Babiuk, L.A., Liang, X., 1996. Expression of bovine interleukin-1 $\beta$  in a bovine herpesvirus-1 vector: in vitro analysis. *Virology* 221, 78–86.
- Reddy, D.N., Reddy, P.G., Minocha, H.C., Fenwick, B.W., Baker, P.E., Davis, W.C., Blecha, F., 1990. Adjuvanticity of recombinant bovine interleukin-1 beta: influence on immunity, infection, and latency in a bovine herpesvirus-1 infection. *Lymphokine Res.* 9, 295–307.
- Reddy, D.N., Reddy, P.G., Xue, W., Minocha, H.C., Daley, M.J., Blecha, F., 1993. Immuno-potential of bovine respiratory disease virus vaccines by interleukin-1 beta and interleukin-2. *Vet. Immunol. Immunopathol.* 37, 25–38.
- Rossow, K.D., 1998. Porcine reproductive and respiratory syndrome. *Vet. Pathol.* 35, 1–20.
- Rossow, K.D., Collins, J.E., Goyal, S.M., Nelson, E.A., Christopher-Hennings, J., Benfield, D.A., 1995. Pathogenesis of porcine reproductive and respiratory syndrome virus infection in gnotobiotic pigs. *Vet. Pathol.* 32, 361–373.
- Royae, A.R., Husmann, R., Dawson, H.D., Calzada-Nova, G., Schnitzlein, W.M., Zuckermann, F., Lunney, J.K., 2004. Deciphering the involvement of innate immune factors in the development of the host responses to PRRSV vaccination. *Vet. Immunol. Immunopathol.* 102, 199–216.
- Snijder, E.J., Meulenber, J.J., 1998. The molecular biology of arteriviruses. *J. Gen. Virol.* 79, 961–979.
- Somasundaram, C., Takamatsu, H., Andréoni, C., Audonnet, J.C., Fischer, L., Lefèvre, F., Charley, B., 1999. Enhanced protective response and immuno-adjuvant effects of porcine GM-CSF on DNA vaccination of pigs against Aujeszky's disease virus. *Vet. Immunol. Immunopathol.* 70, 277–287.
- Spight, D., Zhao, B., Haas, M., Wert, S., Denenberg, A., Shanley, T.P., 2005. Immunoregulatory effects of regulated, lung-targeted expression of IL-10 in vivo. *Am. J. Physiol. Lung Cell Mol. Physiol.* 288, L251–L265.
- Suradhat, S., Thanawongnuwech, R., 2003. Upregulation of interleukin-10 gene expression in the leukocytes of pigs infected with porcine reproductive and respiratory syndrome virus. *J. Gen. Virol.* 84, 2755–2760.
- Suradhat, S., Thanawongnuwech, R., Poovorawan, Y., 2003. Upregulation of IL-10 gene expression in porcine peripheral blood mononuclear cells by porcine reproductive and respiratory syndrome virus. *J. Gen. Virol.* 84, 453–459.
- Thanawongnuwech, R., Thacker, E.L., 2003. Interleukin-10, interleukin-12, and interferon-gamma levels in the respiratory tract following *Mycoplasma hyopneumoniae* and PRRSV infection in pigs. *Viral Immunol.* 16, 357–367.
- Thanawongnuwech, R., Thacker, B., Halbur, P., Thacker, E.L., 2004. Increased production of proinflammatory cytokines following infection with porcine reproductive and respiratory syndrome virus and *Mycoplasma hyopneumoniae*. *Clin. Diagn. Lab. Immunol.* 11, 901–908.
- van Reeth, K., Labarque, G., Nauwynck, H., Pensaert, M., 1999. Differential production of proinflammatory cytokines in the pig lung during different respiratory virus infections: correlations with pathogenicity. *Res. Vet. Sci.* 67, 47–52.
- van Reeth, K., van Gucht, S., Pensaert, M., 2002. In vivo studies on cytokine involvement during acute viral respiratory disease of swine: troublesome but rewarding. *Vet. Immunol. Immunopathol.* 87, 161–168.
- Wasilk, A., Callahan, J.D., Christopher-Hennings, J., Gay, T.A., Fang, Y., Dammen, M., Reos, M.E., Torremorell, M., Polson, D., Mellencamp, M., Nelson, E., Nelson, W.M., 2004. Detection of US, Lelystad, and European-like porcine reproductive and respiratory syndrome viruses and relative quantitation in boar semen and serum samples by real-time PCR. *J. Clin. Microbiol.* 42, 4453–4461.
- Yilma, T., Verardi, P., Jones, L., 2010. Development of safe and efficacious viral vaccines for animals. *Crit. Rev. Immunol.* 30, 223–237.
- Yoon, I.J., Joo, H.S., Goyal, S.M., Molitor, T.W., 1994. A modified serum neutralization test for the detection of antibody to porcine reproductive and respiratory syndrome virus in swine sera. *J. Vet. Diagn. Invest.* 6, 289–292.
- Zeman, D., Neiger, R., Yaeger, M., Nelson, E., Benfield, D., Leslie-Steen, P., Thomson, J., Miskimins, D., Daly, R., Minehart, M., 1993. Laboratory investigation of PRRS virus infection in three swine herds. *J. Vet. Diagn. Invest.* 5, 522–528.
- Zuckermann, F.A., Husmann, R.J., Schwartz, R., Brandt, J., Mateu de Antonio, E., Martin, S., 1998. Interleukin-12 enhances the virus-specific interferon gamma response of pigs to an inactivated pseudorabies virus vaccine. *Vet. Immunol. Immunopathol.* 63, 57–67.
- Zuckermann, F.A., Martin, S., Husmann, R.J., Brandt, J., 1999. Use of interleukin 12 to enhance the cellular immune response of swine to an inactivated herpesvirus vaccine. *Adv. Vet. Med.* 41, 447–461 (Review).