

7,11-Bis(4-methylphenyl)-2,4,8,10-tetraazaspiro[5.5]undecane-1,3,5,9-tetraone

Ali Mohammad Astaraki,^a Ayoob Bazgir^{a*} and Fereshteh Faraji^b

^aDepartment of Chemistry, Islamic Azad University, Dorood Branch, Dorood 688173551, Iran, and ^bDepartment of Chemistry, Faculty of Science, Karaj Branch, Karaj, Iran
Correspondence e-mail: a_bazgir@yahoo.com


Received 11 July 2008; accepted 15 July 2008

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.005 \text{ \AA}$; R factor = 0.081; wR factor = 0.148; data-to-parameter ratio = 16.0.

In the molecule of the title compound, $\text{C}_{21}\text{H}_{20}\text{N}_4\text{O}_4$, the two methylphenyl rings are oriented at a dihedral angle of $59.32(4)^\circ$. The other two rings have flattened-boat conformations. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules. There are $\text{C}-\text{H}\cdots\pi$ contacts between a methylphenyl ring and methyl and methine groups.

Related literature

For general background, see: Pradhan *et al.* (2006); Useglio *et al.* (2006); Kazmierski *et al.* (2006). For bond-length data, see: Allen *et al.* (1987). For ring conformation puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{21}\text{H}_{20}\text{N}_4\text{O}_4$
 $M_r = 392.41$

Monoclinic, $P2_1/n$
 $a = 8.852(2)$ Å

$b = 12.538(3)$ Å
 $c = 17.259(4)$ Å
 $\beta = 104.483(18)^\circ$
 $V = 1854.6(8)$ Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.1 \text{ mm}^{-1}$
 $T = 298(2)$ K
 $0.15 \times 0.11 \times 0.1$ mm

Data collection

Stoe IPDSII diffractometer
Absorption correction: numerical (*X-SHAPE*; Stoe & Cie, 2005)
 $T_{\min} = 0.979$, $T_{\max} = 0.991$

17766 measured reflections
4456 independent reflections
3107 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.093$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.081$
 $wR(F^2) = 0.148$
 $S = 1.13$
4456 reflections
278 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.44 \text{ e \AA}^{-3}$

Table 1

Hydrogen-bond geometry (Å, °).

$Cg4$ is the centroid of the C11–C14/C16/C17 ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1–H1B···O2 ⁱ	0.80 (5)	2.39 (5)	3.004 (3)	135 (4)
N2–H2B···O3 ⁱⁱ	0.77 (4)	2.32 (3)	3.065 (4)	164 (3)
N3–H3···O1 ⁱ	0.82 (4)	2.54 (4)	3.181 (3)	136 (3)
N4–H4D···O1 ⁱⁱⁱ	0.87 (4)	1.92 (4)	2.785 (3)	174 (3)
C4–H4B···Cg4 ^{iv}	0.96	3.02	3.721 (3)	131
C10–H10···Cg4 ^v	0.98	3.11	3.914 (3)	141

Symmetry codes: (i) $-x + 2, -y + 1, -z + 1$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$; (iii) $x - \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}$; (iv) $-x + \frac{3}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (v) $-x + 1, -y + 1, -z$.

Data collection: *X-AREA* (Stoe & Cie, 2005); cell refinement: *X-AREA*; data reduction: *X-RED*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors are grateful to the Islamic Azad University, Dorood Branch, for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2496).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Kazmierski, W. M., Furfine, E., Spaltenstein, A. & Wright, L. L. (2006). *Bioorg. Med. Chem. Lett.* **16**, 5226–5230.
- Pradhan, R., Patra, M., Behera, A. K., Mishra, B. K. & Behera, R. K. (2006). *Tetrahedron*, **62**, 779–828.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Stoe & Cie (2005). *X-AREA*, *X-RED* and *X-SHAPE*. Stoe & Cie, Darmstadt, Germany.
- Useglio, M., Castellano, P. M., Operto, M. A., Torres, R. & Kaufmen, T. S. (2006). *Bioorg. Med. Chem. Lett.* **16**, 5097–5101.

supplementary materials

Acta Cryst. (2008). E64, o1555 [doi:10.1107/S1600536808022113]

7,11-Bis(4-methylphenyl)-2,4,8,10-tetraazaspiro[5.5]undecane-1,3,5,9-tetraone

A. M. Astaraki, A. Bazgir and F. Faraji

Comment

Spiro compounds having cyclic structures fused at a central carbon are of recent interest, due to their interesting conformational feature and structural implications on biological systems (Pradhan *et al.*, 2006). The asymmetric characteristic of the molecule due to the chiral spiro carbon is one of the important criteria of the biological activities. For example, some spiro compounds show antibacterial and antiviral activities (Usegljo *et al.*, 2006, Kazmierski *et al.*, 2006). We report herein the synthesis and crystal structure of the title compound.

In the molecule of the title compound (Fig. 1), the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Ring A (C1-C3/C5-C7) and C (C11-C14/C16/C17) are, of course, planar and they are oriented at a dihedral angle of A/C = 59.32 (4)°. Rings B (N1/N2/C8-C10/C18) and D (N3/N4/C18-C21) have flattened-boat [$\varphi = -54.22 (2)$ °, $\theta = 129.53 (3)$ ° (for ring B) and $\varphi = 52.72 (3)$ °, $\theta = 21.44 (3)$ ° (for ring D)] conformations, having total puckering amplitudes, Q_T , of 1.186 (3) and 0.174 (3) Å, respectively (Cremer & Pople, 1975).

In the crystal structure, intermolecular N-H···O hydrogen bonds (Table 1) link the molecules (Fig. 2), in which they may be effective in the stabilization of the structure. The C—H···π contacts (Table 1) between the ring C and the methyl and methine groups further stabilize the structure.

Experimental

For the preparation of the title compound, a mixture of 4-methylbenzaldehyde (0.24 g, 2 mmol), barbituric acid (0.13 g, 1 mmol), and urea (0.06 g, 1 mmol) was heated at 373 K. After 2 h, the reaction mixture was washed with water (10 ml). The residue recrystallized from ethanol to afford the pure product (yield; 0.25 g, 65%, m.p. 519–521 K).

Refinement

H1B, H2B, H3 and H4D atoms (for NH) were located in difference syntheses and refined isotropically [N-H = 0.77 (4)–0.87 (4) Å and $U_{\text{iso}}(\text{H}) = 0.032 (8)$ –0.076 (14) Å²]. The remaining H atoms were positioned geometrically, with C-H = 0.93, 0.98 and 0.96 Å for aromatic, methine and methyl H, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

supplementary materials




Fig. 2. A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

7,11-Bis(4-methylphenyl)-2,4,8,10-tetraazaspiro[5.5]undecane-1,3,5,9-tetraone

Crystal data

C ₂₁ H ₂₀ N ₄ O ₄	$F_{000} = 824$
$M_r = 392.41$	$D_x = 1.405 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/n$	Mo $K\alpha$ radiation
	$\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2yn	Cell parameters from 2021 reflections
$a = 8.852 (2) \text{ \AA}$	$\theta = 2.0\text{--}28.1^\circ$
$b = 12.538 (3) \text{ \AA}$	$\mu = 0.1 \text{ mm}^{-1}$
$c = 17.259 (4) \text{ \AA}$	$T = 298 (2) \text{ K}$
$\beta = 104.483 (18)^\circ$	Prism, yellow
$V = 1854.6 (8) \text{ \AA}^3$	$0.15 \times 0.11 \times 0.1 \text{ mm}$
$Z = 4$	

Data collection

Stoe IPDSII diffractometer	$R_{\text{int}} = 0.093$
rotation method scans	$\theta_{\text{max}} = 28.1^\circ$
Absorption correction: Numerical shape of crystal determined optically (X-SHAPE; Stoe & Cie, 2005))	$\theta_{\text{min}} = 2.0^\circ$
$T_{\text{min}} = 0.979, T_{\text{max}} = 0.991$	$h = -11 \rightarrow 11$
17766 measured reflections	$k = -16 \rightarrow 16$
4456 independent reflections	$l = -22 \rightarrow 22$
3107 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	H atoms treated by a mixture of independent and constrained refinement
Least-squares matrix: full	$w = 1/[\sigma^2(F_o)^2 + (0.0701P)^2 + 0.8069P]$ where $P = (F_o^2 + 2F_c^2)/3$
$R[F^2 > 2\sigma(F^2)] = 0.081$	$(\Delta/\sigma)_{\text{max}} = 0.003$
$wR(F^2) = 0.148$	$\Delta\rho_{\text{max}} = 0.43 \text{ e \AA}^{-3}$
$S = 1.13$	$\Delta\rho_{\text{min}} = -0.44 \text{ e \AA}^{-3}$
4456 reflections	Extinction correction: none
278 parameters	

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	1.0239 (3)	0.62989 (17)	0.65204 (12)	0.0446 (5)
O2	1.0451 (3)	0.60240 (17)	0.43863 (13)	0.0463 (6)
O3	0.7614 (3)	0.63921 (19)	0.18421 (13)	0.0522 (6)
O4	0.6359 (3)	0.85414 (18)	0.36650 (13)	0.0520 (6)
N1	0.8229 (3)	0.6195 (2)	0.54163 (15)	0.0417 (6)
H1B	0.809 (5)	0.559 (4)	0.553 (3)	0.076 (14)*
N2	0.9943 (3)	0.7595 (2)	0.55764 (14)	0.0403 (6)
H2B	1.069 (4)	0.787 (3)	0.582 (2)	0.046 (10)*
N3	0.8909 (3)	0.6150 (2)	0.31407 (15)	0.0393 (6)
H3	0.936 (4)	0.563 (3)	0.3020 (19)	0.032 (8)*
N4	0.7093 (3)	0.7516 (2)	0.27656 (14)	0.0381 (6)
H4D	0.646 (4)	0.787 (3)	0.239 (2)	0.045 (9)*
C1	0.4632 (4)	0.5999 (3)	0.3870 (2)	0.0453 (7)
H1	0.4228	0.6665	0.3944	0.054*
C2	0.3664 (4)	0.5232 (3)	0.3421 (2)	0.0516 (8)
H2	0.2622	0.5397	0.3196	0.062*
C3	0.4211 (4)	0.4227 (3)	0.33000 (18)	0.0455 (7)
C4	0.3157 (5)	0.3384 (3)	0.2819 (2)	0.0634 (10)
H4A	0.3123	0.2776	0.3152	0.076*
H4B	0.3556	0.3172	0.2374	0.076*
H4C	0.2125	0.3668	0.2626	0.076*
C5	0.5765 (4)	0.4012 (3)	0.3651 (2)	0.0489 (8)
H5	0.6162	0.334	0.3587	0.059*
C6	0.6746 (4)	0.4773 (2)	0.40958 (19)	0.0448 (7)
H6	0.7787	0.4606	0.432	0.054*
C7	0.6191 (3)	0.5784 (2)	0.42101 (16)	0.0363 (6)
C8	0.7243 (3)	0.6631 (2)	0.46862 (16)	0.0357 (6)
H8	0.6568	0.7166	0.4842	0.043*
C9	0.9505 (4)	0.6683 (2)	0.58658 (16)	0.0361 (6)
C10	0.9253 (3)	0.8080 (2)	0.48001 (16)	0.0349 (6)
H10	0.8511	0.8621	0.4885	0.042*
C11	1.0480 (4)	0.8656 (2)	0.44822 (17)	0.0374 (7)
C12	1.0027 (4)	0.9543 (2)	0.39888 (18)	0.0417 (7)
H12	0.8984	0.9746	0.3844	0.05*
C13	1.1118 (4)	1.0120 (2)	0.3715 (2)	0.0491 (8)
H13	1.0792	1.0695	0.3374	0.059*
C14	1.2688 (4)	0.9861 (2)	0.3938 (2)	0.0474 (8)
C15	1.3894 (5)	1.0521 (3)	0.3674 (3)	0.0694 (11)

supplementary materials

H15A	1.3866	1.124	0.3862	0.083*
H15B	1.3673	1.0521	0.3101	0.083*
H15C	1.4911	1.0224	0.3892	0.083*
C16	1.3123 (4)	0.8966 (3)	0.4421 (2)	0.0511 (8)
H16	1.4168	0.8768	0.4568	0.061*
C17	1.2051 (4)	0.8368 (3)	0.46843 (19)	0.0455 (8)
H17	1.2375	0.7769	0.4999	0.055*
C18	0.8296 (3)	0.7230 (2)	0.42113 (15)	0.0340 (6)
C19	0.9342 (3)	0.6434 (2)	0.39333 (16)	0.0335 (6)
C20	0.7848 (3)	0.6665 (2)	0.25345 (17)	0.0356 (6)
C21	0.7178 (3)	0.7831 (2)	0.35308 (16)	0.0343 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0555 (13)	0.0453 (11)	0.0249 (10)	-0.0030 (10)	-0.0049 (9)	0.0011 (8)
O2	0.0490 (13)	0.0443 (11)	0.0358 (11)	0.0170 (10)	-0.0075 (10)	-0.0017 (9)
O3	0.0634 (15)	0.0587 (14)	0.0279 (11)	0.0076 (12)	-0.0013 (10)	-0.0114 (10)
O4	0.0593 (14)	0.0517 (13)	0.0381 (12)	0.0261 (11)	-0.0007 (10)	-0.0024 (10)
N1	0.0491 (15)	0.0430 (14)	0.0264 (12)	-0.0046 (12)	-0.0030 (11)	0.0014 (10)
N2	0.0495 (15)	0.0402 (13)	0.0231 (11)	-0.0030 (12)	-0.0059 (11)	-0.0012 (10)
N3	0.0442 (14)	0.0381 (13)	0.0312 (13)	0.0117 (11)	0.0012 (10)	-0.0082 (10)
N4	0.0440 (14)	0.0405 (13)	0.0233 (11)	0.0111 (11)	-0.0040 (10)	0.0010 (10)
C1	0.0410 (16)	0.0415 (16)	0.0491 (19)	0.0038 (13)	0.0031 (14)	-0.0023 (13)
C2	0.0364 (16)	0.059 (2)	0.053 (2)	-0.0003 (15)	-0.0014 (14)	0.0024 (16)
C3	0.0520 (19)	0.0473 (17)	0.0335 (15)	-0.0100 (14)	0.0041 (14)	0.0023 (13)
C4	0.070 (2)	0.051 (2)	0.060 (2)	-0.0199 (18)	0.0006 (19)	-0.0005 (17)
C5	0.059 (2)	0.0394 (16)	0.0446 (18)	0.0011 (15)	0.0052 (15)	-0.0016 (13)
C6	0.0437 (17)	0.0436 (16)	0.0422 (17)	0.0069 (13)	0.0012 (14)	-0.0002 (13)
C7	0.0394 (15)	0.0402 (15)	0.0259 (13)	0.0013 (12)	0.0017 (11)	0.0023 (11)
C8	0.0411 (15)	0.0398 (14)	0.0231 (12)	0.0060 (12)	0.0023 (11)	-0.0020 (11)
C9	0.0441 (16)	0.0401 (14)	0.0214 (12)	0.0050 (12)	0.0034 (11)	-0.0023 (11)
C10	0.0435 (15)	0.0327 (13)	0.0233 (13)	0.0067 (12)	-0.0014 (11)	-0.0034 (10)
C11	0.0508 (17)	0.0291 (13)	0.0283 (13)	0.0034 (12)	0.0025 (12)	-0.0022 (11)
C12	0.0504 (18)	0.0349 (14)	0.0375 (16)	0.0116 (13)	0.0067 (13)	0.0021 (12)
C13	0.072 (2)	0.0317 (14)	0.0444 (18)	0.0090 (15)	0.0163 (16)	0.0030 (13)
C14	0.058 (2)	0.0382 (15)	0.0455 (18)	0.0030 (14)	0.0127 (15)	-0.0037 (13)
C15	0.074 (3)	0.054 (2)	0.081 (3)	-0.007 (2)	0.021 (2)	0.009 (2)
C16	0.0467 (18)	0.0494 (18)	0.053 (2)	0.0031 (15)	0.0053 (15)	0.0036 (15)
C17	0.0488 (18)	0.0423 (16)	0.0401 (17)	0.0082 (14)	0.0012 (14)	0.0065 (13)
C18	0.0415 (15)	0.0334 (13)	0.0221 (12)	0.0081 (12)	-0.0018 (11)	-0.0019 (10)
C19	0.0374 (15)	0.0328 (13)	0.0263 (13)	0.0054 (11)	0.0001 (11)	-0.0011 (10)
C20	0.0372 (15)	0.0376 (14)	0.0277 (13)	-0.0025 (12)	0.0002 (11)	-0.0015 (11)
C21	0.0367 (14)	0.0347 (13)	0.0265 (13)	0.0078 (12)	-0.0013 (11)	0.0006 (11)

Geometric parameters (\AA , $^\circ$)

C1—C7	1.384 (4)	C11—C12	1.398 (4)
C1—C2	1.388 (5)	C12—C13	1.381 (5)

C1—H1	0.93	C12—H12	0.93
C2—C3	1.385 (5)	C13—C14	1.384 (5)
C2—H2	0.93	C13—H13	0.93
C3—C5	1.384 (5)	C14—C16	1.393 (5)
C3—C4	1.513 (5)	C14—C15	1.509 (5)
C4—H4A	0.96	C15—H15A	0.96
C4—H4B	0.96	C15—H15B	0.96
C4—H4C	0.96	C15—H15C	0.96
C5—C6	1.385 (4)	C16—C17	1.372 (5)
C5—H5	0.93	C16—H16	0.93
C6—C7	1.390 (4)	C17—H17	0.93
C6—H6	0.93	C18—C19	1.518 (4)
C7—C8	1.512 (4)	C18—C21	1.531 (4)
C8—N1	1.449 (4)	C19—O2	1.207 (3)
C8—C18	1.577 (4)	C19—N3	1.372 (3)
C8—H8	0.98	C20—O3	1.210 (3)
C9—O1	1.250 (3)	C20—N4	1.370 (4)
C9—N2	1.344 (4)	C20—N3	1.379 (4)
C9—N1	1.346 (4)	C21—O4	1.207 (3)
C10—N2	1.459 (3)	C21—N4	1.362 (4)
C10—C11	1.516 (4)	N1—H1B	0.80 (5)
C10—C18	1.568 (4)	N2—H2B	0.77 (4)
C10—H10	0.98	N3—H3	0.82 (3)
C11—C17	1.393 (4)	N4—H4D	0.87 (4)
C7—C1—C2	120.8 (3)	C12—C13—C14	121.4 (3)
C7—C1—H1	119.6	C12—C13—H13	119.3
C2—C1—H1	119.6	C14—C13—H13	119.3
C3—C2—C1	121.6 (3)	C13—C14—C16	117.6 (3)
C3—C2—H2	119.2	C13—C14—C15	121.6 (3)
C1—C2—H2	119.2	C16—C14—C15	120.8 (3)
C5—C3—C2	117.2 (3)	C14—C15—H15A	109.5
C5—C3—C4	120.9 (3)	C14—C15—H15B	109.5
C2—C3—C4	121.9 (3)	H15A—C15—H15B	109.5
C3—C4—H4A	109.5	C14—C15—H15C	109.5
C3—C4—H4B	109.5	H15A—C15—H15C	109.5
H4A—C4—H4B	109.5	H15B—C15—H15C	109.5
C3—C4—H4C	109.5	C17—C16—C14	121.8 (3)
H4A—C4—H4C	109.5	C17—C16—H16	119.1
H4B—C4—H4C	109.5	C14—C16—H16	119.1
C3—C5—C6	121.7 (3)	C16—C17—C11	120.3 (3)
C3—C5—H5	119.2	C16—C17—H17	119.8
C6—C5—H5	119.2	C11—C17—H17	119.8
C5—C6—C7	120.8 (3)	C19—C18—C21	114.1 (2)
C5—C6—H6	119.6	C19—C18—C10	112.2 (2)
C7—C6—H6	119.6	C21—C18—C10	107.7 (2)
C1—C7—C6	117.9 (3)	C19—C18—C8	109.7 (2)
C1—C7—C8	120.4 (3)	C21—C18—C8	106.3 (2)
C6—C7—C8	121.7 (3)	C10—C18—C8	106.3 (2)
N1—C8—C7	111.3 (2)	O2—C19—N3	121.0 (3)

supplementary materials

N1—C8—C18	109.0 (2)	O2—C19—C18	122.5 (2)
C7—C8—C18	114.6 (2)	N3—C19—C18	116.5 (2)
N1—C8—H8	107.2	O3—C20—N4	122.2 (3)
C7—C8—H8	107.2	O3—C20—N3	122.1 (3)
C18—C8—H8	107.2	N4—C20—N3	115.7 (2)
O1—C9—N2	122.1 (3)	O4—C21—N4	120.6 (2)
O1—C9—N1	121.1 (3)	O4—C21—C18	121.3 (3)
N2—C9—N1	116.8 (3)	N4—C21—C18	118.0 (2)
N2—C10—C11	110.9 (2)	C9—N1—C8	124.5 (3)
N2—C10—C18	110.2 (2)	C9—N1—H1B	117 (3)
C11—C10—C18	114.1 (2)	C8—N1—H1B	118 (3)
N2—C10—H10	107	C9—N2—C10	127.1 (3)
C11—C10—H10	107	C9—N2—H2B	118 (3)
C18—C10—H10	107	C10—N2—H2B	114 (3)
C17—C11—C12	118.3 (3)	C19—N3—C20	127.0 (3)
C17—C11—C10	123.1 (3)	C19—N3—H3	115 (2)
C12—C11—C10	118.5 (3)	C20—N3—H3	118 (2)
C13—C12—C11	120.4 (3)	C21—N4—C20	126.5 (2)
C13—C12—H12	119.8	C21—N4—H4D	117 (2)
C11—C12—H12	119.8	C20—N4—H4D	117 (2)
C7—C1—C2—C3	0.4 (5)	N1—C8—C18—C19	65.8 (3)
C1—C2—C3—C5	0.5 (5)	C7—C8—C18—C19	-59.7 (3)
C1—C2—C3—C4	179.3 (4)	N1—C8—C18—C21	-170.3 (2)
C2—C3—C5—C6	-1.0 (5)	C7—C8—C18—C21	64.2 (3)
C4—C3—C5—C6	-179.8 (3)	N1—C8—C18—C10	-55.8 (3)
C3—C5—C6—C7	0.5 (5)	C7—C8—C18—C10	178.7 (2)
C2—C1—C7—C6	-0.8 (5)	C21—C18—C19—O2	167.7 (3)
C2—C1—C7—C8	178.9 (3)	C10—C18—C19—O2	44.8 (4)
C5—C6—C7—C1	0.4 (5)	C8—C18—C19—O2	-73.1 (3)
C5—C6—C7—C8	-179.4 (3)	C21—C18—C19—N3	-16.2 (4)
C1—C7—C8—N1	136.0 (3)	C10—C18—C19—N3	-139.0 (3)
C6—C7—C8—N1	-44.2 (4)	C8—C18—C19—N3	103.0 (3)
C1—C7—C8—C18	-99.7 (3)	C19—C18—C21—O4	-174.6 (3)
C6—C7—C8—C18	80.0 (3)	C10—C18—C21—O4	-49.3 (4)
N2—C10—C11—C17	27.2 (4)	C8—C18—C21—O4	64.3 (3)
C18—C10—C11—C17	-98.0 (3)	C19—C18—C21—N4	7.6 (4)
N2—C10—C11—C12	-150.1 (3)	C10—C18—C21—N4	132.9 (3)
C18—C10—C11—C12	84.6 (3)	C8—C18—C21—N4	-113.5 (3)
C17—C11—C12—C13	-0.4 (4)	O1—C9—N1—C8	174.7 (3)
C10—C11—C12—C13	177.0 (3)	N2—C9—N1—C8	-5.4 (5)
C11—C12—C13—C14	-1.9 (5)	C7—C8—N1—C9	164.5 (3)
C12—C13—C14—C16	2.8 (5)	C18—C8—N1—C9	37.2 (4)
C12—C13—C14—C15	-176.9 (3)	O1—C9—N2—C10	175.3 (3)
C13—C14—C16—C17	-1.4 (5)	N1—C9—N2—C10	-4.6 (5)
C15—C14—C16—C17	178.4 (4)	C11—C10—N2—C9	-146.8 (3)
C14—C16—C17—C11	-0.9 (5)	C18—C10—N2—C9	-19.4 (4)
C12—C11—C17—C16	1.8 (5)	O2—C19—N3—C20	-168.4 (3)
C10—C11—C17—C16	-175.5 (3)	C18—C19—N3—C20	15.4 (4)
N2—C10—C18—C19	-72.4 (3)	O3—C20—N3—C19	175.2 (3)

C11—C10—C18—C19	53.3 (3)	N4—C20—N3—C19	-3.9 (5)
N2—C10—C18—C21	161.2 (2)	O4—C21—N4—C20	-173.8 (3)
C11—C10—C18—C21	-73.2 (3)	C18—C21—N4—C20	4.0 (4)
N2—C10—C18—C8	47.6 (3)	O3—C20—N4—C21	174.2 (3)
C11—C10—C18—C8	173.3 (2)	N3—C20—N4—C21	-6.7 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1B···O2 ⁱ	0.80 (5)	2.39 (5)	3.004 (3)	135 (4)
N2—H2B···O3 ⁱⁱ	0.77 (4)	2.32 (3)	3.065 (4)	164 (3)
N3—H3···O1 ⁱ	0.82 (4)	2.54 (4)	3.181 (3)	136 (3)
N4—H4D···O1 ⁱⁱⁱ	0.87 (4)	1.92 (4)	2.785 (3)	174 (3)
C4—H4B···Cg4 ^{iv}	0.96	3.02	3.721 (3)	131
C10—H10···Cg4 ^v	0.98	3.11	3.914 (3)	141

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $x+1/2, -y+3/2, z+1/2$; (iii) $x-1/2, -y+3/2, z-1/2$; (iv) $-x+3/2, y-1/2, -z+1/2$; (v) $-x+1, -y+1, -z$.

supplementary materials

Fig. 1




Fig. 2

