



The Equidae from Cooper's D, an early Pleistocene fossil locality in Gauteng, South Africa

Shaw Badenhorst^{1,*} and Christine M. Steininger^{1,2,*}

¹ Evolutionary Studies Institute, University of the Witwatersrand, Johannesburg, South Africa

² DST-NRF Centre of Excellence in Palaeosciences, University of the Witwatersrand, Johannesburg, South Africa

* These authors contributed equally to this work.

ABSTRACT

Cooper's D is a fossil locality in the Bloubank Valley close to other important sites such as Sterkfontein and Kromdraai in Gauteng, South Africa. The fossil deposits of Cooper's D date to 1.38 ± 0.11 Ma. Hominins like *Paranthropus robustus* and early *Homo* have been recovered from Cooper's Cave. We report here on the Equidae remains. Our sample contains specimens from the extinct *Equus capensis*, and a specimen which represents an extinct hipparion *Eurygnathohippus cf. cornelianus*. This particular specimen was previously identified as plains zebra (*Equus quagga*). The contribution of Equidae to the total fossil assemblage of Cooper's D is relatively low, and these remains were likely accumulated by various predators such as spotted and brown hyenas and leopards. The Equidae, as well as the other fauna from Cooper's D supports the existence of grassland, wooded and water components in the vicinity of the site.

Subjects Paleontology

Keywords Equidae, *Eurygnathohippus cornelianus*, *Equus capensis*, Cooper's D, Hipparion

INTRODUCTION

The Bloubank Valley in the Cradle of Humankind close to Johannesburg in South Africa is well-known for the range of hominin taxa discovered at fossil localities like Sterkfontein, Kromdraai and Swartkrans dating from the Plio-Pleistocene. The animals found associated with these hominins are of considerable interest, and a large body of research has been produced, investigating aspects such as the palaeo-environments (e.g., [Reed, 1996](#); [Adams, 2006](#); [Kuhn, Werdelin & Steininger, 2017](#); [Sénégal & Thackeray, 2008](#); [Steininger, 2011](#); [Adams et al., 2016](#)), indirect dating of deposits through faunal seriation (e.g., [McKee, Thackeray & Berger, 1995](#); [Badenhorst et al., 2011](#); [Van Zyl, Badenhorst & Brink, 2016](#)), taphonomy (e.g., [Brain, 1981](#); [Adams et al., 2007](#); [Val, Taru & Steininger, 2014](#); [Kruger & Badenhorst, 2018](#)) and taxonomy (e.g., [Churcher, 1970](#); [Churcher, 1974](#); [Vrba, 1976](#)).

Cooper's Cave is located in the Gauteng province of South Africa. The site is over 250 m² in size and contains three distinct localities, Cooper's A, B, and D all located in the Monte Cristo Formation (Malmani Subgroup, Transvaal Supergroup) ([Fig. 1](#); see [De Ruiter et al., 2009](#)). The most fauna from Cooper's Cave are from the D component, including the specimens described in this paper. Cooper's D represents a de-roofed cave. Three facies are recognised at Cooper's D (Facies A, B and C). Previous uranium-lead (U-Pb) analyses

Submitted 2 November 2018

Accepted 2 April 2019

Published 15 May 2019

Corresponding author

Shaw Badenhorst,
shaw.badenhorst@wits.ac.za

Academic editor

Matt Sponheimer

Additional Information and
Declarations can be found on
page 11

DOI 10.7717/peerj.6909

© Copyright

2019 Badenhorst and Steininger

Distributed under

Creative Commons CC-BY 4.0

OPEN ACCESS




Figure 1 (A) Location of Cooper's D in South Africa, and (B) site plan of the site. The location of Cooper's D in the Gauteng province of South Africa. The site map is indicated

Full-size [DOI: 10.7717/peerj.6909/fig-1](https://doi.org/10.7717/peerj.6909/fig-1)

date Cooper's D to between 1.5 and 1.4 million years ago (De Ruiter et al., 2009). Recently refined uranium-lead dates for Cooper's D suggest a 1.38 ± 0.11 Ma date for the site (Pickering et al., 2019). Remains of *Paranthropus robustus* are present at Cooper's D. This taxon has only been recovered at Swartkrans, Kromdraai, Sterkfontein, Gondolin and Drimolen in the Bloubaank Valley and lived between 2 and 1.2 million years ago in South Africa (De Ruiter, Sponheimer & Lee-Thorp, 2008; Steininger, Berger & Kuhn, 2008).

The animal remains from Cooper's D have been well studied. Berger et al. (2003) and De Ruiter et al. (2009) were the first to produce a preliminary list of taxa present in the faunal assemblage from Cooper's D, and these include primates, carnivores, hyraxes, perissodactyls, artiodactyls, rodents and lagomorphs. Val, Taru & Steininger (2014) studied the primate remains from Cooper's D, and argued that at least some primates, including *P. robustus*, were hunted and killed by leopards and hyenas. However, given the abundance of juvenile and sub-adult primates, the geomorphology of the cave, and the low impact

of carnivore modification, Cooper's D likely represented a locality that was occupied by large-bodied cercopithecids who died a natural death in the cave (Val, Taru & Steininger, 2014). Cooper's D also yielded primate remains of *Theropithecus*, *Papio*, and a large papionin that may belong to *Gorgopithecus* (DeSilva, Steininger & Patel, 2013; Folinsbee & Reisz, 2013). O'Regan & Steininger (2017) studied the felid sample from Cooper's D. This sample indicates the presence of four large felid genera, namely *Dinofelis*, *Megantereon*, *Panthera* and *Acinonyx*. This sample may mark the first appearance of the modern cheetah in Africa, as well the first occurrence of the East African species *Dinofelis cf. aronoki* in southern Africa. Moreover, the site also yielded remains of two mustelids, *Propoecilogale bolti* and *Mellivora capensis*, as well as a viverrid, *Civettictis cf. civetta* (O'Regan, Cohen & Steininger, 2013). The Herpestidae from Cooper's D were collected by brown hyena (Cohen, O'Regan & Steininger, 2019). The Hyaenidae from Cooper's D include *Chasmaporthetes nitidula*, *Crocuta ultra*, *Parahyaena brunnea*, *Hyaena hyaena* and *cf. Proteles* sp. (Kuhn, Werdelin & Steininger, 2017). The preliminary studies indicate the presence of the extant *Equus quagga* and extinct *Equus capensis* at Cooper's D, (Berger et al., 2003; De Ruiter et al., 2009). As these Equidae specimens have not been thoroughly considered from the site, they are described for this paper, and their taphonomic and palaeoecological implications discussed.

Brief Overview of the Taxonomic Status of Equidae

The taxonomic status of the Equidae in southern Africa has been debated for several decades. Morphometric analyses suggest that the plains zebra (or Burchell's zebra) *Equus burchellii* and the extinct quagga *E. quagga* are distinct species (e.g., Thackeray, 1988; Klein & Cruz-Uribe, 1999). However, the evidence from cranial morphology, body stripes and molecular data consider *E. quagga* as a subspecies of the plains zebra (e.g., Lowenstein & Ryder, 1985; Rau, 1986; but see Leonard et al., 2005). It is now widely accepted that the two taxa are conspecific with the older name of *E. quagga* retained (Bronner et al., 2003; Skinner & Chimimba, 2005). There are three extant species of zebra in sub-Saharan Africa. They are Grévy's zebra *Equus grevyi* from East Africa, the mountain zebra *Equus zebra* from the western and southern parts of southern Africa and south-western Angola, and *E. quagga* from the north-eastern parts of southern Africa west to Angola (Bronner et al., 2003; Skinner & Chimimba, 2005).

Early in the 21st century, Broom (1909) described a large extinct Equid from South Africa, which he named *Equus capensis*. This taxon, often referred to as the extinct Cape zebra had shorter limbs, was more heavily built than extant horses with a larger cranium (Brain, 1981). Poor understanding of individual variation in teeth morphology and the effects of wear on teeth resulted in a flurry of different extinct zebras described in South Africa (Wells, 1959). However, these are all now considered synonyms of *Equus capensis* (Churcher & Richardson, 1978).

The taxonomic status of *Equus capensis* remains controversial (e.g., Peters et al., 1994), but we retained the taxon following Bernor et al. (2010: 705). Some authors (e.g., Cooke, 1950; Churcher, 1986; Churcher, 2006) suggested that *Equus capensis* is closely related to the extinct *Equus oldowayensis* from East Africa, and its immediate descendant, Grévy's

zebra (*Equus grevyi*). *Equus grevyi* has a large head with short legs (Churcher & Richardson, 1978). According to this view, *Equus capensis* still exists as Grévy's zebra in Eastern Africa today (Churcher, 2006). Recent studies on the DNA from *Equus capensis* indicate they are closer related to *Equus quagga* (Orlando et al., 2009). However, extensive osteological comparisons are still lacking between these forms (Peters et al., 1994:24). Often, in practice, Equid specimens of adult individuals that are larger than *Equus quagga* are regarded as belonging to *Equus capensis* (e.g., Brink, 1987; Peters et al., 1994; Stynder, 1997; Badenhorst et al., 2011; Adams et al., 2016).

Equus capensis had a wide distribution in southern and eastern Africa, and existed from the Late Pliocene to the Early Holocene (ca. 3.6 mya–10,000 ka). Stable isotopes indicate that, on average *Equus capensis* were grazers (Lee-Thorp & Beaumont, 1995; Codron et al., 2008) of coarse grasses like modern plains zebras (Hofmann & Stewart, 1972; Stynder, 1997). In the interior of South Africa, *Equus quagga* and *Equus capensis* occur concurrently in many Pleistocene deposits (Churcher & Richardson, 1978). *Equus quagga* appeared about one million years ago in southern Africa, although they occurred in Africa from the Late Pliocene–Early Pleistocene (Bernor et al., 2010).

Another species of extinct zebra existed in southern Africa, formerly called *Hipparion lybicum steytleri* (Churcher & Richardson, 1978). Bernor et al. (2010) refers to the hipparion material from Ethiopia, Tanzania, Kenya and South Africa as *Eurygnathohippus cornelianus*, which existed between 2.5 and .05 million years ago. Hipparions have been identified at localities such as Swartkrans Member 1 and Kromdraai A (Brain, 1981). This three-toed zebra had a slightly-built, and appreciably smaller than *Equus quagga* (Brain, 1981) with distinct dental morphology (Churcher & Richardson, 1978; Brain, 1981; MacFadden, 1992; Churcher, 2000; Churcher & Watson, 2004; Churcher, 2006).

METHODS

All the available Equidae material from Cooper's D was considered for this paper. Specimens were identified using the collections housed at the University of the Witwatersrand (modern BP and fossil Swartkrans SK collections) and the Ditsong National Museum of Natural History (AZ collection) coupled with published descriptions (Broom, 1909; Cooke, 1950; Churcher, 1974; Churcher & Richardson, 1978; Brain, 1981; Churcher, 2000; Churcher & Watson, 2004; Churcher, 2006). Following common practice (e.g., Brink, 1987; Peters et al., 1994; Stynder, 1997; Badenhorst et al., 2011; Adams et al., 2016), specimens larger than the extant *Equus quagga* were identified as the extinct *Equus capensis*. The specimens are quantified using the Number of Identified Specimens (NISP) and the Minimum Number of Individuals (MNI). Detail about each identified specimen is listed in [Data S1](#).

All measurements for postcranial material follow Von den Driesch (1976). Comparative measurements are included from specimens housed in the collections at the University of the Witwatersrand and the Ditsong National Museum of Natural History. Almost no comparative measurements are available for Equidae from sites in the Bloubank Valley (but see Churcher & Watson, 2004). It is well beyond the scope of this paper to include such a comprehensive osteometrical study. However, we included measurements of *Equus quagga*

Table 1 The Equidae identified in the Cooper's Cave assemblage.

Taxon	NISP	MNI
<i>Equus capensis</i>	13	2(1 adult,1 juvenile)
<i>cf. Equus capensis</i>	3	–
<i>Eurygnathohippus cornelianus</i>	1	1
Equidae indeterminate	18	–
Total	42	

and *Equus capensis* where such data is available for comparative purposes. All measurements are in millimetres to accuracy of 0.1 mm, and when an estimated measurement was taken, it is indicated as 'est', following common practice in osteometrical research (e.g., *Von den Driesch, 1976*). For the II and IV metapodia of Equidae, an anterior-posterior length was taken of the proximal part of the element, as well as a medial-lateral breadth measurement. We calculate the proportion of Equidae to large and very large Bovidae (Bov III's and IV's, following *Brain (1974)* in samples from the Bloubank Valley where remains of *P. robustus* have been discovered. We selected large and very large Bovidae, as these ungulates are similar in size to Equidae, and may have been prey to similar predators.

RESULTS

The extinct *Equus capensis* and *Eurygnathohippus cf. cornelianus* were identified from Cooper's D (Table 1). Many specimens from Cooper's D retained sufficient morphological features to allow identification to species level. *Equus capensis* is represented by at least one adult and one juvenile individual in the sample. No taphonomic modifications were noted on the material. None of the teeth from Cooper's D could be assigned to hipparion (Fig. 2). However, one postcranial specimen is likely from a hipparion (Fig. 3). It is a right metacarpal IV, which is smaller in size than the average *Equus quagga* (Table 2) and larger than a modern donkey. As a result, we postulate that this specimen is a hipparion (*Eurygnathohippus cf. cornelianus*). Teeth and post-crania of *Equus capensis* are larger in size than *Equus quagga* (Tables 3 and 4). Specimen CD 991 is proximal portion of a left IV metatarsal, and specimen CD 3508 is a proximal portion of a left metatarsal II. Both are identified as *Equus capensis* due to their large size (Table 5). Using NISP, most samples contain between 4 and 15% Equidae (Swartkrans Members 1, 2 and 3, Drimolen and Sterkfontein Member 5). At Cooper's D, Equidae are also poorly represented (3% based on MNI). Gondolin 1 and 2 has the highest representation of Equidae of 30%. Kromdraai contains no Equidae remains (Table 6).

DISCUSSION

Equus capensis was identified based on the large size of the specimens compared to the smaller extant *Equus quagga*. The use of size differences to distinguish Pleistocene and extant species remains controversial (*Watson & Plug, 1995*), also in the case of *Equus capensis* (*Peters et al., 1994*). However, extensive morphological comparisons are lacking to determine if *Equus capensis* is related to an extant species of zebra (*Peters et al., 1994*). *Equus*



Figure 2 Example of teeth assigned to *Equus capensis* (Specimen CD 5881, Upper M1). Note that the protocone is not isolated as is the case in hipparions. Photo credit: Ashley Kruger.

Full-size  DOI: [10.7717/peerj.6909/fig-2](https://doi.org/10.7717/peerj.6909/fig-2)

capensis appeared some two million years ago in South Africa, and existed until about 10 000 years ago. In contrast, the plains zebra only appeared about a million years ago in eastern and southern Africa, and still exists today (summary in [Bernor et al. \(2010\):702](#)). This suggests that the two taxa had different evolutionary trajectories as two separate species or subspecies. However, more research is required to investigate the relationship between them. Assuming *Equus capensis* is a distinct species, it is present at Cooper's D. The deposit predates the appearance of *Equus quagga* in the Bloubank Valley by millennia. *Equus capensis* occurs at sites in the Bloubank Valley like Swartkrans (Members 1–3) ([De Ruiter, 2003](#)), Gladysvale (GVED) ([Lacruz et al., 2003](#)), Sterkfontein Member 4, Kromdraai A ([Brain, 1981](#)) and Garage Ravine at Bolt's Farm ([Badenhorst et al., 2011](#)).

The extant *Equus quagga* is present in Africa since 2.33 million years ago ([Geraads, Raynal & Eisenmann, 2004](#); [Adams et al., 2016](#)). Remains have been found at sites in the Bloubank Valley such as Swartkrans (Member 3) ([De Ruiter, 2003](#)), Drimolen Main Quarry ([Adams et al., 2016](#)), Gladysvale (GVED) ([Lacruz et al., 2003](#)), Sterkfontein (Members 5 and 6), Swartkrans Channel Fill and Kromdraai A ([Brain, 1981](#)). A previous, preliminary



Figure 3 (A) a right metacarpal IV of *Equus quagga* (AZ 1131). B: CD 24345, a right metacarpal IV, which is likely from *Eurygnathohippus cornelianus* (see Table 2). Photo credit: Brett Eloff.

Full-size DOI: 10.7717/peerj.6909/fig-3

Table 2 Comparison between metacarpal IV Cooper's D *Eurygnathohippus cf. cornelianus* and modern Equidae. For some specimens, the sex is unknown.

Taxa	Specimen Number	Sex	Length	Breadth
<i>Eurygnathohippus cf. cornelianus</i>	CD 24 345	–	18.65	14.14
<i>Equus quagga</i>	AZ 3252	–	22.45	16.57
<i>Equus quagga</i>	AZ 1283	♀	19.79 (est)	–
<i>Equus quagga</i>	AZ 1131	–	23.81	16.0
<i>Equus caballus</i>	AZ 585	♀	22.52 (est)	18.26 (est)
<i>Equus africanus asinus</i>	AZ 423	–	15.78	12.0

analysis of the Equidae reported *Equus quagga* from Cooper's D (Berger et al., 2003; De Ruiter et al., 2009).

The small size of the specimen, comparable to modern donkey, suggest that the specimen originally designated as *Equus quagga* from Cooper's D is actually *Eurygnathohippus cf.*

Table 3 Comparisons of teeth measurements of *Equus capensis* from Cooper's D and *Equus quagga*. The measurements indicate *Equus capensis* is consistently larger than *Equus quagga*.

Taxa	Specimen Number	Side	Mesiodistal Length	Buccolingual Breadth
Upper M1				
<i>Equus capensis</i>	CD 5881	R	30.28 (est)	30.47
<i>Equus quagga</i>	BP/4/147	R	23.46 (est)	26.38
Upper M3				
<i>Equus capensis</i>	CD 16973	R	31.06	25.98
<i>Equus quagga</i>	BP/4/147	R	21.67 (est)	20.89
Indeterminate Upper Molar				
<i>Equus capensis</i>	CD 9070	R	36.19	26.99
<i>Equus capensis</i>	SK 41515	–	31.80	22.46
Upper Deciduous P3 or P4				
<i>Equus capensis</i>	CA 1159	R	38.55	25.84 (est)
<i>Equus capensis</i>	CD 9293	R	36.20	26.40
<i>Equus capensis</i>	SK 28881	R	39.1	28.2
Indeterminate Lower Molar				
<i>Equus capensis</i>	CD 992	–	27.10	–
<i>Equus quagga</i> (M1)	BP/4/147	–	22.25	–
<i>Equus quagga</i> (M2)	BP/4/147	–	22.60	–
Lower M3				
<i>Equus capensis</i>	CD 11067	L	40.65 (est)	18.48 (est)
<i>Equus quagga</i>	BP/4/147	L	31.20 (est)	15.22 (est)

cornelianus. The latter species is only recognised from teeth and lower leg bones, and no morphological differences necessarily exist to separate the smaller *Eurygnathohippus cornelianus* from *Equus quagga*. *Eurygnathohippus cornelianus* is not abundant in fossil deposits in the interior of South Africa, but specimens have been found at Cornelia-Uitzoek (Brink et al., 2012), Makapansgat Member 3 (Reed, 1996), Kromdraai A (Churcher, 1970), Swartkrans Members 1–3 (Churcher & Watson, 2004), Sterkfontein Member 4 (Reynolds & Kibii, 2011), Gondolin GD 2 (Adams, 2006) and Drimolen Makondo Infill (Rovinsky et al., 2015). *Eurygnathohippus cornelianus* occur mainly in Pliocene deposits in South Africa, but they persist to the Early Pleistocene as is evident from Cornelia-Uitzoek, which dates to between 1.07 and 0.99 million years ago (Brink et al., 2012; Rovinsky et al., 2015).

Previous research at Cooper's D suggests brown hyena and leopard involvement in the accumulation of the deposits (De Ruiter et al., 2009; Cohen, O'Regan & Steininger, 2019). At sites like Swartkrans, Sterkfontein, Kromdraai and Minnaar's Cave, the extinct sabre-toothed cat, *Dinofelis* could have preyed on *Equus capensis* (Lesnik & Thackeray, 2006) although other predators could also have been involved. The main predators of extant Grévy and plains zebras are lions, leopards, spotted hyenas and hunting dogs (Smuts, 1979; Grubb, 1981; Churcher, 1993). Of these predators, lions and hunting dogs do not make use of shelters (e.g., Pitman et al., 2013) although carnivores such as hyenas could scavenge their kills and bring bones into shelters. The low to near absence of Equid remains

Table 4 Comparisons of measurements of post-crania of *Equus capensis* from Cooper's D with other Equidae. *Equus capensis* is consistently larger than *Equus quagga*, and similar in size to the modern horse *Equus caballus*.

Taxa	Specimen Number	Sex	Side	Measurements
Astragalus				
<i>Equus capensis</i>	CD 6747	–	Left	GH: 61.58, LmT: 59.68, GB: 59.12, BFd: 46.26
<i>Equus capensis</i>	SK 6288	–	Right	GH: 63.04, LmT: 63.23
<i>Equus quagga</i>	AZ 1424	♀	Left	GH: 54.56, LmT: 54.80, GB: 59.15, BFp: 46.46
Scapula				
<i>Equus capensis</i>	CD 6061	–	Left	GLP: 103.00, SLC: 72.15, LG: 65.31, BG: 56.11
<i>Equus quagga</i>	AZ 1424	♀	Left	GLP: 77.78, SLC: 51.01, LG: 48.10, BG: 42.60
<i>Equus quagga</i>	BP/4/911	–	–	GLP: 75.73, SLC: 50.30, LG: 49.72, BG: 38.54
<i>Equus caballus</i>	BP/4/929	–	–	GLP: 100.33 (est), SLC: 69.34, LG: 57.19, BG: 49.06
Femur				
<i>Equus capensis</i>	CD 10416	–	Left	DC: 56.94
<i>Equus quagga</i>	AZ 1424	♀	Left	DC: 48.71

Notes.

Astragalus GH, greatest height; LmT, length of the medial part of the trochlea tali; GB, greatest breadth; BFd, breadth of the facies articularis distalis; Scapula GLP, greatest length of the processus articularis; SLC, smallest length of the collum scapulae; LG, length of the glenoid cavity; BG, breadth of the glenoid cavity; Femur DC, greatest depth of the caput femoris (Von den Driesch, 1976).

Table 5 Comparisons of measurements of metatarsals II and IV of *Equus capensis* from Cooper's D and *Equus quagga*. The measurements indicate *Equus capensis* is consistently larger than *Equus quagga*.

Taxa	Specimen Number	Sex	Length of Proximal Articulation	Breadth of Proximal Articulation
Metatarsal IV				
<i>Equus capensis</i>	CD 991	–	32.77	22.26 (est)
<i>Equus quagga</i>	AZ 1424	–	25.81	19.40
<i>Equus quagga</i>	AZ 1131	–	27.61	20.30
<i>Equus quagga</i>	AZ 1132	♂	27.05	18.94
<i>Equus quagga</i>	AZ 3252	–	26.04	20.08
<i>Equus quagga</i>	AZ 1283	♀	27.16	17.84
Metatarsal II				
<i>Equus capensis</i>	CD 3508	–	23.13	16.09
<i>Equus quagga</i>	AZ 1424	–	17.85	11.94
<i>Equus quagga</i>	AZ 1131	–	17.54	11.71
<i>Equus quagga</i>	AZ 1132	♂	18.18	11.68
<i>Equus quagga</i>	AZ 3252	–	16.77	11.90
<i>Equus quagga</i>	AZ 1283	♀	18.40	11.83

at sites in the Bloubank Valley may reflect this. Small sample sizes aside (Rovinsky et al., 2015), Equidae may have also been absent from some portions of the Bloubank Valley and larger region for some times of the year (Mills & Shenk, 1992). In addition, while Equids may have been regularly preyed upon, not all attempts to hunt them would have been

Table 6 Proportions of Equidae and large and very large Bovidae (Bovidae III and IV, see [Brain \(1974\)](#) representation in samples containing *Paranthropus robustus*. The dates are from [Herries, Curnoe & Adams \(2009\)](#).

Site and samples	Relative age (mya)	Equidae (NISP/MNI)	Bovidae III and IV (NISP/MNI)	% Equidae NISP (%MNI)	Reference
Swartkrans Member 1	2	27/11	594/82	4 (12)	Brain, 1981 : 322–323
Swartkrans Member 1	2	30/4	418/12	7 (25)	Watson, 2004 : 40
Kromdraai B	2.11–1.65	-/-	66/12	0 (0)	Brain, 1981 : 339–340
Kromdraai Member 2	2.11–1.65	-/-	84/-	0 (0)	Fourvel et al., 2016 : 83
Gondolin 1 and 2	1.78	19/2	45/8	30 (20)	Adams et al., 2007 : 2533
Sterkfontein Member 5	1.78–0.82	23/3	157/15	13 (17)	Brain, 1981 : 317
Drimolen	2–1.5	3/1	64/23	4 (4)	Adams et al., 2016
Cooper's D	1.5–1.4	-/2	-/67	- (3)	De Ruiter et al., 2009 : 506. This study
Swartkrans Member 2	1.65–1.07	60/16	335/43	15 (27)	Brain, 1981 : 328–329
Swartkrans Member 2	1.65–1.07	22/4	268/15	8 (21)	Watson, 2004 : 40
Swartkrans Member 3	1.04–0.62	74/7	892/34	7 (17)	Watson, 2004 : 40

successful. Plains and Grévy's zebras are formidable prey, being able to kick and bite to defend themselves and run at high speeds (overviews in [Grubb, 1981](#); [Churcher, 1993](#)).

Like the plains and Grevy's zebra ([Rautenbach, 1982](#); [Churcher, 1993](#); [Skinner & Chimimba, 2005](#)), *Equus capensis* ([Lee-Thorp & Beaumont, 1995](#); [Codron et al., 2008](#)) and *Eurygnathohippus* were grazers ([Franz-Odenaal, Kaiser & Bernor, 2003](#); [White et al., 2009](#); [Cerling et al., 2015](#)). The plains zebra must drink water daily, and is always in close proximity to water ([Rautenbach, 1982](#)). Grévy's zebra require less water than the plain zebra. While they prefer to drink daily, they can go without water for three days ([Churcher, 1993](#)). The source of water was presumably the palaeo-Bloubaank River ([De Ruiter et al., 2009](#)). The large mammalian fauna from Cooper's D suggest a mosaic environment with grassland, woodland and water components ([De Ruiter et al., 2009](#); [Steininger, 2011](#); [Cohen, O'Regan & Steininger, 2019](#)).

CONCLUSION

The Equidae sample from Cooper's D contains two extinct species, *Equus capensis* and *Eurygnathohippus cf. cornelianus*. While a previous preliminary report indicate the presence of the extant *Equus quagga*, we do not consider this identification to be correct. Instead, the small size of the specimen, and the deposit predating the arrival of extant *Equus quagga* in South Africa, supports this notion. A variety of carnivores preyed on Equidae including hyenas, leopards and lions and likely contributed bone remains to the deposits, directly or indirectly. The presence of the two extinct species supports the presence of grassland, woodland and water components around Cooper's D at 1.38 ± 0.11 Ma.

ACKNOWLEDGEMENTS

The Ditsong National Museum of Natural History allowed access to their collections. The comments and suggestions of the reviewers are greatly appreciated.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The excavations at Cooper's D were supported financially by the Palaeoanthropological Scientific Trust (PAST), National Research Foundation, and the DST-NRF Centre of Excellence in Palaeosciences (CoE-Palaeo). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Palaeoanthropological Scientific Trust (PAST).

National Research Foundation.

DST-NRF Centre of Excellence in Palaeosciences (CoE-Palaeo).

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Shaw Badenhorst conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Christine M. Steininger conceived and designed the experiments, contributed reagents/materials/analysis tools, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft, excavated the site.

Data Availability

The following information was supplied regarding data availability:

Detail for each identified specimen is listed in [Data S1](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6909#supplemental-information>.

REFERENCES

- Adams JW. 2006.** Taphonomy and paleoecology of the Gondolin Plio-Pleistocene cave site, South Africa. Ph.D. dissertation, Washington University, St. Louis.
- Adams JW, Herries AIR, Kuykendall KL, Conroy GC. 2007.** Taphonomy of a South African cave: geological and hydrological influences on the GD 1 fossil assemblage at Gondolin, a Plio-Pleistocene paleocave system in the Northwest Province, South Africa. *Quaternary Science Reviews* **26**:2526–2543
[DOI 10.1016/j.quascirev.2007.05.006](https://doi.org/10.1016/j.quascirev.2007.05.006).
- Adams JW, Rovinsky DS, Herries AIR, Menter CG. 2016.** Macromammalian faunas, biochronology and palaeoecology of the early Pleistocene Main Quarry hominin-bearing deposits of the Drimolen Palaeocave System, South Africa. *PeerJ* **4**:e1941
[DOI 10.7717/peerj.1941](https://doi.org/10.7717/peerj.1941).

- Badenhorst S, Sénégas F, Gommery D, Potze S, Kgasi L, Thackeray JF. 2011.** Pleistocene animal remains from Garage Ravine Cave on Bolt's Farm in the Cradle of Humankind, South Africa. *Annals of the Ditsong National Museum of Natural History* 1:33–40.
- Berger LR, De Ruiter DJ, Steininger CM, Hancox J. 2003.** Preliminary results of excavations at the newly investigated Cooper's D deposit, Gauteng, South Africa. *South African Journal of Science* 99:276–278.
- Bernor RL, Armour-Chelu MJ, Gilbert H, Kaiser TM, Schulz E. 2010.** Equidae. In: Werdelin L, Sander WJ, eds. *Cenozoic mammals of Africa*. Berkeley: University of California Press, 685–721.
- Brain CK. 1974.** Some suggested procedures in the analysis of bone accumulations from southern African Quaternary sites. *Annals of the Transvaal Museum* 29:1–8.
- Brain CK. 1981.** *The hunters or the hunted? An introduction to African cave taphonomy*. Chicago: University of Chicago Press.
- Brink JS. 1987.** The archaeozoology of Florisbad, Orange Free State. In: *Memoirs van die Nasionale Museum Bloemfontein no. 24*. Bloemfontein: National Museum.
- Brink JS, Herries AIR, Moggi-Cecchi J, Gowlett JAJ, Bousman CB, Hancox JP, Grün R, Eisenmann V, Adams JW, Rossouw L. 2012.** First hominine remains from a ~1.0 million year old bone bed at Cornelia-Uitzoek, Free State Province, South Africa. *Journal of Human Evolution* 63:527–535 DOI [10.1016/j.jhevol.2012.06.004](https://doi.org/10.1016/j.jhevol.2012.06.004).
- Bronner GN, Hoffmann M, Taylor PJ, Chimimba CT, Best PB, Mathee CA, Robinson TJ. 2003.** A revised systematic checklist of the extant mammals of the southern African subregion. *Durban Museum Novitates* 28:56–103.
- Broom R. 1909.** On evidence of a large horse recently extinct in South Africa. *Annals of the South African Museum* 7:281–282.
- Cerling TE, Andanje SA, Blumenthal SA, Brown FH, Chritz KL, Harris JM, Hart JA, Kirera FM, Kaleme P, Leakey LN, Leakey MG, Levin NE, Manthi FK, Passey BH, Uno KT. 2015.** Dietary changes of large herbivores in the Turkana Basin, Kenya from 4 to 1 Ma. *Proceedings of the National Academy of Sciences of the United States of America* 112(37):11467–11472 DOI [10.1073/pnas.1513075112](https://doi.org/10.1073/pnas.1513075112).
- Churcher CS. 1970.** The fossil Equidae from the Krugersdorp caves. *Annals of the Transvaal Museum* 26(6):145–168.
- Churcher CS. 1974.** *Sivatherium maurusium* (Pomel) from the Swartkrans Australopithecine site, Transvaal (Mammalia: Giraffidae). *Annals of the Transvaal Museum* 29(6):65–69.
- Churcher CS. 1986.** The extinct Cape zebra. *Sagittarius* 1(4):4–5.
- Churcher CD. 1993.** *Equus grevyi*. *Mammalian Species* 453:1–9.
- Churcher CS. 2000.** Extinct equids from Limeworks Cave and Cave of Hearths, Makapansgat, Northern Province, and a consideration of variation in the cheek teeth of *Equus capensis* Broom. *Palaeontologia Africana* 36:97–117.
- Churcher CS. 2006.** Distribution and history of the Cape zebra (*Equus capensis*) in the Quaternary of Africa. *Transactions of the Royal Society of South Africa* 61(2):89–95 DOI [10.1080/00359190609519957](https://doi.org/10.1080/00359190609519957).

- Churcher CS, Richardson ML. 1978.** Equidae. In: Maglio J, Cooke HBS, eds. *Evolution of African mammals*. Cambridge: Harvard University Press, 379–422.
- Churcher CS, Watson V. 2004.** Additional fossil Equidae from Swartkrans. In: Brain CK, ed. *Swartkrans. A cave's chronical of early man. Transvaal Museum Monograph no. 8*. Pretoria: Transvaal Museum, 137–150.
- Codron D, Brink JS, Rossouw L, Clasuss M. 2008.** The evolution of ecological specialization in southern African ungulates: competition or physical environmental turnover. *Oikos* **117**:334–353.
- Cohen BF, O'Regan HJ, Steininger CM. 2019.** Mongoose manor: herpestidae remains from the Early Pleistocene Cooper's D locality in the Cradle of Humankind, Gauteng, South Africa. *Palaeontologia Africana* **53**:97–113.
- Cooke HBS. 1950.** A critical revision of the Quaternary Perissodactyla of southern Africa. *Annals of the South African Museum* **31**:393–476.
- De Ruiter DJ. 2003.** Revised faunal list for Members 1-3 for Swartkrans, South Africa. *Annals of the Transvaal Museum* **40**:29–41.
- De Ruiter DJ, Pickering R, Steininger CM, Kramers JD, Hancox PJ, Churchill SE, Berger LR, Backwell L. 2009.** New Australopithecus robustus fossils and associated U-Pb dates from Cooper's Cave (Gauteng, South Africa). *Journal of Human Evolution* **56**(5):497–513 DOI [10.1016/j.jhevol.2009.01.009](https://doi.org/10.1016/j.jhevol.2009.01.009).
- De Ruiter DJ, Sponheimer M, Lee-Thorp JA. 2008.** Indications of habitat association of *Australopithecus robustus* in the Bloubank Valley, South Africa. *Journal of Human Evolution* **55**(6):1015–1030 DOI [10.1016/j.jhevol.2008.06.003](https://doi.org/10.1016/j.jhevol.2008.06.003).
- DeSilva JM, Steininger CM, Patel BA. 2013.** Cercopithecoid primate postcranial fossils from Cooper's D, South Africa. *Geobios* **46**:381–394 DOI [10.1016/j.geobios.2013.07.001](https://doi.org/10.1016/j.geobios.2013.07.001).
- Folinsbee KE, Reisz RR. 2013.** New craniodental fossils of papiodin monkeys from Cooper's D, South Africa. *American Journal of Physical Anthropology* **151**:613–629 DOI [10.1002/ajpa.22317](https://doi.org/10.1002/ajpa.22317).
- Fourvel JB, Brink J, O'Regan H, Beaudet A, Pavia M. 2016.** Some preliminary interpretations of the oldest faunal assemblage from Kromdraai. In: Braga J, Thackeray JF, eds. *Kromdraai. A birthplace of Paranthropus in the Cradle of Humankind. A South African heritage site*. Cape Town: Sun Press, 71–104.
- Franz-Odendaal TA, Kaiser TM, Bernor RL. 2003.** Systematics and dietary evaluation of a fossil Equid from South Africa. *South African Journal of Science* **99**:453–459.
- Geraads D, Raynal J, Eisenmann V. 2004.** The earliest human occupation of North Africa: a reply to Sahnouni et al. (2002). *Journal of Human Evolution* **46**:751–761 DOI [10.1016/j.jhevol.2004.01.008](https://doi.org/10.1016/j.jhevol.2004.01.008).
- Grubb P. 1981.** *Equus burchelli*. *Mammalian Species* **157**:1–9.
- Herries AIR, Curnoe F, Adams JW. 2009.** A multi-disciplinary seriation of early *Homo* and *Paranthropus* bearing palaeocaves in southern Africa. *Quaternary International* **202**:14–28 DOI [10.1016/j.quaint.2008.05.017](https://doi.org/10.1016/j.quaint.2008.05.017).
- Hofmann RR, Stewart DRM. 1972.** Grazer or browser: a classification based on the stomach structure and feeding habits of East African ruminants. *Mammalia* **36**:226–240.

- Klein RG, Cruz-Urbe K. 1999.** Craniometry of the genus *Equus* and the taxonomic affinities of the extinct South African quagga. *South African Journal of Science* 95:81–86.
- Kruger A, Badenhorst S. 2018.** Remains of a barn owl (*Tyto alba*) from the Dinaledi Chamber, Rising Star Cave, South Africa. *South African Journal of Science* 114(11/12):Article #5152 DOI 10.17159/sajs.2018/5152.
- Kuhn BF, Werdelin L, Steininger CM. 2017.** Fossil Hyaenidae from Cooper's Cave, South Africa, and the palaeoenvironmental implications. *Palaeobiodiversity and Palaeoenvironments* 97:355–365 DOI 10.1007/s12549-016-0247-y.
- Lacruz R, Ungar P, Hancox PJ, Brink JS, Berger LR. 2003.** Gladysvale: fossils, strata and GIS analysis. *South African Journal of Science* 99:283–285.
- Lee-Thorp JA, Beaumont PB. 1995.** Vegetation and seasonality shifts during the late Quaternary deduced from $^{13}\text{C}/^{12}\text{C}$ ratios of grazers at Equus Cave, South Africa. *Quaternary Research* 43:426–432 DOI 10.1006/qres.1995.1048.
- Leonard JA, Rohland N, Glaberman S, Fleischer RC, Caccione A, Hofreiter M. 2005.** A rapid loss of stripes: the evolutionary history of the extinct quagga. *Biological Letters* 1:291–295 DOI 10.1098/rsbl.2005.0323.
- Lowenstein JM, Ryder OA. 1985.** Immunological systematics of the extinct quagga (Equidae). *Experientia* 41:1192–1193 DOI 10.1007/BF01951724.
- MacFadden BJ. 1992.** Fossil horses. In: *Systematics, paleobiology, and evolution of the Family Equidae*. Cambridge: Cambridge University Press.
- McKee JK, Thackeray JF, Berger LR. 1995.** Faunal assemblage seriation of southern African Pliocene and Pleistocene fossil deposits. *American Journal of Physical Anthropology* 96:235–250 DOI 10.1002/ajpa.1330960303.
- Mills MGL, Shenk TM. 1992.** Predator-prey relationships: the impact of lion predation on wildebeest and zebra populations. *Journal of Animal Ecology* 61:693–702 DOI 10.2307/5624.
- O'Regan HJ, Cohen BF, Steininger CM. 2013.** Mustelid and viverrid remains from the Pleistocene site of Cooper's D, Gauteng, South Africa. *Palaeontologia Africana* 48:19–23.
- O'Regan HJ, Steininger CM. 2017.** Felidae from Cooper's Cave, South Africa (Mammalia: Carnivora). *Geodiversitas* 39(2):315–332 DOI 10.5252/g2017n2a8.
- Orlando L, Metcalf JL, Alberdi MT, Telles-Antunes M, Bonjean D, Otte M, Martin F, Eisenmann V, Mashkour M, Morello F, Prado JL, Salas-Gismondi R, Shockey BJ, Wrinn PJ, Vasilev SK, Ovodov ND, Cherry MI, Hopwood B, Male D, Austin JJ, Hänni C, Cooper A. 2009.** Revising the recent evolutionary history of Equids using ancient DNA. *Proceedings of the National Academy of Science of the United States of America* 106:21754–21759 DOI 10.1073/pnas.0903672106.
- Peters J, Gautier A, Brink JS, Haenen W. 1994.** Late Quaternary extinction of ungulates in Sub-Saharan Africa: a reductionist's approach. *Journal of Archaeological Science* 21:17–28 DOI 10.1006/jasc.1994.1004.
- Pickering R, Herries AIR, Woodhead JD, Hellstrom JC, Green HE, Paul B, Ritzman T, Strait DS, Schoville BJ, Hancox PJ. 2019.** U–Pb-dated flowstones restrict

- South African early hominin record to dry climate phases. *Nature* **565**:226–229
DOI [10.1038/s41586-018-0711-0](https://doi.org/10.1038/s41586-018-0711-0).
- Pitman RT, Kilian PJ, Ramsay PM, Swanepoel LH. 2013.** Foraging and habitat specialization by female leopards (*Panthera pardus*) in the Waterberg Mountains of South Africa. *South African Journal of Wildlife Research* **43**(2):167–176
DOI [10.3957/056.043.0204](https://doi.org/10.3957/056.043.0204).
- Rau RE. 1986.** The quagga and its kin. *Sagittarius* **1**(2):8–10.
- Rautenbach IL. 1982.** Mammals of the Transvaal. In: *Ecoplan Monograph no. 1*. Pretoria: Ecoplan.
- Reed KE. 1996.** The paleoecology of Makapansgat and other African Plio-Pleistocene hominid localities. Ph.D. dissertation, State University of New York, Stony Brook.
- Reynolds SC, Kibii JM. 2011.** Sterkfontein at 75: review of palaeoenvironments, fauna and archaeology from the hominin site Sterkfontein (Gauteng, South Africa). *Palaeontologia Africana* **46**:59–88.
- Rovinsky DS, Herries AIR, Menter CG, Adams JW. 2015.** First description of *in situ* primate and faunal remains from the Plio-Pleistocene Drimolen Makondo palaeocave infill, Gauteng, South Africa. *Palaeontologia Electronica* **18.2.34A**:1–21.
- Sénégas F, Thackeray JF. 2008.** Temperature indices based on relative abundances of rodent taxa represented in South African Plio-Pleistocene assemblages. *Annals of the Transvaal Museum* **45**:143–144.
- Skinner JD, Chimimba CT. 2005.** *The mammals of the southern African sub-region*. Cambridge: Cambridge University Press.
- Smuts GL. 1979.** Diet of lions and spotted hyaenas assessed from stomach contents. *South African Journal of Wildlife Research* **9**:19–25.
- Steininger CM. 2011.** The dietary behaviour of Early Pleistocene bovids from Cooper's Cave and Swartkrans, South Africa. PhD thesis, University of the Witwatersrand, Johannesburg.
- Steininger C, Berger LR, Kuhn BF. 2008.** A partial skull of *Paranthropus robustus* from Cooper's Cave, South Africa. *South African Journal of Science* **104**:143–146.
- Stynder DD. 1997.** The use of faunal evidence to reconstruct site history at Hoedjiespunt 1 (HDP1), Western Cape. Masters dissertation, Cape Town: University of Cape Town.
- Thackeray JF. 1988.** Zebras from Wonderwerk Cave, Northern Cape Province, South Africa: attempts to distinguish *Equus burchelli* and *E. quagga*. *South African Journal of Science* **84**(2):99–101.
- Val A, Taru P, Steininger CM. 2014.** New taphonomic analysis of large-bodied primate assemblage from Cooper's D, Bloubaank Valley, South Africa. *South African Archaeological Bulletin* **69**(199):49–58.
- Van Zyl WJ, Badenhorst S, Brink JS. 2016.** Pleistocene Bovidae from X Cave on Bolt's Farm in the Cradle of Humankind in South Africa. *Annals of the Ditsong National Museum of Natural History* **6**:39–73.
- Von den Driesch A. 1976.** A guide to the measurement of animal bones from archaeological sites. In: *Peabody Museum Bulletin 1*. Cambridge: Harvard University.

- Vrba ES. 1976.** The fossil Bovidae of Sterkfontein, Swartkrans and Kromdraai. In: *Transvaal Museum Memoir no. 21*. Pretoria: Transvaal Museum.
- Watson V. 2004.** Composition of the Swartkrans bone accumulations, in terms of skeletal parts and animals represented. In: Brain CK, ed. *Swartkrans. A cave's chronical of early man. Transvaal Museum Monograph no. 8*. Pretoria: Transvaal Museum, 35–73.
- Watson V, Plug I. 1995.** *Oreotragus major* Wells and *Oreotragus oreotragus* (Zimmerman) (Mammalia: Bovidae): two species? *Annals of the Transvaal Museum* **36(13)**:183–191.
- Wells LH. 1959.** The nomenclature of South African fossils equids. *South African Journal of Science* **55(3)**:64–66.
- White TD, Ambrose SH, Suwa G, Su DF, DeGusta D, Bernor RL, Boisserie J-R, Brunet M, Delson E, Frost S, Garcia N, Giaourtsakis IX, Haile-Selassie Y, Howell FC, Lehmann T, Likius A, Pehlevan C, Saegusa H, Semprebon G, Teaford M, Vrba E. 2009.** Macrovertebrate paleontology and the Pliocene habitat of *Ardipithecus ramidus*. *Science* **326(5949)**:87–93 DOI [10.1126/science.1175822](https://doi.org/10.1126/science.1175822).