



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Expression of nitric oxide synthase-2 in glia associated with CNS pathology

Angela K. Loihl and Sean Murphy\*

*Department of Pharmacology and Neuroscience Program, University of Iowa College of Medicine, Iowa City, IA 52242, USA*

## Introduction

While the production of nitric oxide (NO) in the nervous system is catalyzed by three, highly homologous isoforms of NO synthase (NOS) (Forstermann et al., 1995), in this chapter we focus exclusively on NOS-2. This dimeric, heme-containing, soluble protein whose activity is independent of a rise in intracellular calcium, is variously termed 'inducible', 'immunologic', and 'macNOS' (Griffith and Stuehr, 1995). These terms are somewhat misleading because NOS-2 has been reported in a wide variety of normal and neoplastic cell types, in some cases constitutively (Xie and Nathan 1994). All CNS parenchymal cells, together with the smooth muscle and endothelium of the vascular wall, can be transcriptionally activated by a variety of agents to express NOS-2, as well as cells (neutrophils and monocytes) infiltrating the parenchyma following changes in blood-brain barrier properties (Grzybicki et al., 1997, 1998). In addition, there are reports of developmentally related NOS-2 expression in particular brain regions (Galea et al., 1995; Arnhold et al., 1997).

Initially from studies with cultured cells (Murphy et al., 1993), but now exemplified by a host of

in vivo conditions (see later), identified astrocytes have been shown to express NOS-2 upon activation. Induction can be triggered in vitro (see Table 1) by endotoxin and/or combinations of such cytokines as interleukin (IL)-1 $\beta$ , tumor necrosis factor (TNF)- $\alpha$  and interferon (IFN)- $\gamma$ ; HIV envelope proteins also activate the astrocyte gene (Koka et al., 1995). The promoter region of the glial NOS-2 gene displays a wide variety of potential response elements for transcription factors, such as AP-1, Hif, NF $\kappa$ B, NF-IL6, TNF, IRF, and GAS (Spitsin et al., 1997). In cultured astrocytes the transcript can be detected within a few hours of exposure to cytokines, is maximally transcribed by 4 h and then declines rapidly, such that it is not detectable at 12 h (Park and Murphy, 1994). In contrast, expression of NOS-2 in the CNS in vivo is an unexpectedly delayed event, which may, as we discuss later, have functional implications.

The observation that HIV envelope proteins can induce NOS-2 is important as it demonstrates that priming cells with IFN- $\gamma$  (a cytokine not produced by resident CNS cells) is not obligatory for gene activation. In addition, cytokine-activated astrocytes, macrophages and cerebrovascular cells release TNF $\alpha$  which, in turn, induces NOS-2 in naive cells (Borgerding and Murphy, 1995; Shafer and Murphy, 1997). This implies that induction of NOS-2 in CNS parenchymal cells could occur

\*Corresponding author. Tel.: 319 335 7958; fax: 319 335 8930;  
e-mail: sean-murphy@uiowa.edu

TABLE 1

Regulators of NOS-2 expression *in vitro*.

Inducers	Blockers
Proinflammatory cytokines (IL-1 $\beta$ , TNF $\alpha$ , IFN $\gamma$ )	Anti-inflammatory cytokines (TGF $\beta$ 1, IFN $\beta$ , IL-4, IL-10)
Endotoxin	Dexamethasone
HIV env. proteins	Chemokines (MCP-1, IL-8)
Hypoxia	Neurotransmitters (norepinephrine, glutamate, ATP, angiotensin II)
$\beta$ -amyloid	Heat shock protein 70 nitric oxide (NO)

*in vivo* following events on the luminal side of an intact blood-brain barrier. Transcriptional activation of the cerebral vasculature by endotoxin or cytokines from adhering hematogenous cells may be sufficient to induce NOS-2 in astrocytes and microglia on the parenchymal side.

The timing and extent of NOS-2 expression associated with various neuropathologies may also be related to the prior history of the cells involved. For example, heat shock protein 70 and the anti-inflammatory cytokines transforming growth factor (TGF)- $\beta$ 1 and IFN $\beta$  block NOS-2 induction at the transcriptional level (Park et al., 1994; Feinstein et al., 1996; Stewart et al., 1997). Activation of glutamate, ATP, angiotensin-II and  $\beta$ -adrenergic receptors on astrocytes blunt the NOS-2 response to endotoxin and pro-inflammatory cytokines (Kopnisky et al., 1997; Lin and Murphy, 1997; Feinstein 1998). It is clear that some of these modulatory effects result from interference with the subsequent activation of a key transcription factor, NF $\kappa$ B, and/or with its binding to the promoter.

Nitric oxide inhibits not only NOS-2 activity but also regulates the level of NOS-2 mRNA expression through a mechanism involving NF- $\kappa$ B. Some reports suggest a blockade of NF- $\kappa$ B activation (Colasanti et al., 1995; Peng et al., 1995), whereas others propose inhibition of binding to its response element on the NOS-2 promoter via nitrosylation of a key cysteine residue on one of

the NF- $\kappa$ B subunits (DelaTorre et al., 1997; Park et al., 1997). This regulation might represent a means to limit or focus NO production in discrete groups of cells in the CNS (Togashi et al., 1997). Such a feedback effect of NO has been confirmed *in vivo* (Luss et al., 1994), pointing to potential rebound effects with the therapeutic use of NOS-2 inhibitors.

The NOS-2 transcript is rapidly degraded, implying there may be factors co-induced in cells which are responsible for causing this instability (Park and Murphy, 1996). The 3'-untranslated region of NOS-2 has AUUUA repeats (Galea et al., 1994) which are suggested to be determinants of mRNA stability and this could represent yet another mechanism by which the amount and duration of NO production is regulated.

With all of these potential regulatory mechanisms for NOS-2 expression in mind, what is the evidence for expression *in vivo*? what are the roles that NO from this source play in CNS function? There are many potential targets for NO and its oxidation products (such as peroxynitrite). These include heme and non-heme iron, thiols and DNA. Such reactions account for the reported biological effects of NOS activation, such as mitochondrial dysfunction (Knowles, 1997), neuromodulation (Christopherson and Bredt, 1997), and cell and organism killing (Nathan, 1997). In addition, it is evident that NO has subtle effects on the regulation of a number of cytokine and chemokine genes (Merrill and Murphy, 1997). However, it is important to distinguish between effects due to NO and those resulting from peroxynitrite exposure. The latter is a strong oxidant with effects that, in general, are biologically deleterious; the production of NO in the absence of superoxide anion produces different effects, many of which may be beneficial (Butler et al., 1995; Nathan, 1997).

### Expression of NOS-2 *in vivo*

Since last we reviewed this literature (Murphy and Grzybicki, 1996) there have been considerable efforts to identify specific neuropathological con-

ditions that result in the induction of NOS-2. The earlier, suggestive reports of NOS-2 mRNA and/or protein expression associated with certain types of CNS injury have now been confirmed by *in situ* hybridization and immunohistochemistry and provide evidence for the cellular source. There is specific evidence for glial expression of NOS-2 associated with neuronal injury and infection of the CNS, and in neurodegenerative and demyelinating diseases (see Table 2). We refer to these most recent reports for completeness, and then focus on ischemia and the evidence for glial NOS-2 expression. Finally, we speculate on the possible roles for NO in the development of pathology, and in recovery from injury.

### Trauma

Direct injury in the CNS results in a reactive gliosis, characterized by induction of the glial fibrillary acidic protein gene and changes in astrocyte morphology (Eddleston and Mucke, 1993). In a freeze-lesion model, we found that NOS-2 was induced within a few hours in cells infiltrating the lesion (Grzybicki et al., 1998). Only after 24 h could we find NOS-2 expression in astrocytes delimiting the lesion edge. In a compression injury to the spinal cord, Hamada et al. (1996) report NOS-2 mRNA, which peaks at 24 h and persists for three days. The expression was associated with motor dysfunction and application

of a NOS inhibitor reduced motor disturbance. This delayed expression of NOS-2 in parenchymal cells is of interest and we speculate on its meaning later in the chapter.

There is also suggestive evidence for NOS-2 expression in glia after indirect neuronal injury, which occurs with axotomy or chemical toxicity, and results in a marked microglial, but lesser astroglial cell response. Following a local injection of kainic acid into the cortex, Lei et al. (1996) describe neuronal degeneration, followed 2–3 days later by expression of NOS-2 in reactive astrocytes and in endothelial cells at the edge of the lesion.

### Infection

How viruses cause lesions, or dysfunction without direct damage to the CNS, is unclear. Diffusible cytokines, together with reactive oxygen and nitrogen species produced as a component of the immune response presumably contribute to the pathology. In many instances, NOS-2 expression is associated with viral infection of the CNS and, in some, the resulting NO appears to be beneficial.

Intranasal inoculation of mice with the neurotropic hepatitis virus leads to expression of NOS-2 in the brain (Grzybicki et al., 1997). The cells responsible for its expression appear to be entirely infiltrating, with no evidence for parenchymal cell expression. However, this infection proves fatal within seven days. Lane et al. (1997) used a neuroattenuated strain of this virus which is cleared and the animals thus survive. They report NOS-2 in astrocytes and macrophages in the olfactory bulb. We have looked also at expression of NOS-2 in persistently infected animals, generated by raising inoculated pups with immunized dams. In these animals, which display marked demyelinating lesions and associated hindlimb paralysis, NOS-2 is expressed very prominently in astrocytes surrounding such lesions (Sun et al., 1995; Grzybicki et al., 1997). A similar expression was reported by Oleszak et al. (1997) in astrocytes surrounding necrotizing lesions in mice infected with Theiler's virus. There is also suggestive evidence that NOS-2 expression in the CNS is

TABLE 2

Role of NOS-2 in neuropathologies.

Pathology	After NOS-2 block
Direct cortical injury	Reduced
Indirect injury	?
Hepatitis, Theiler's, HIV	?
LCM virus	Worsened
Endotoxin	Reduced
ALS	?
Retinal dystrophy	?
Alzheimer	?
EAE/MS	Reduced or worsened
Global ischemia	Reduced
Stroke	Reduced

associated with HIV-1 (Adamson et al., 1996) and SIV infection (Lane et al., 1996).

The study by Lane et al. (1997) suggests that, while NO from NOS-2 can block viral replication *in vitro*, this is not the determining factor for clearing hepatitis virus in the CNS. Campbell (1996), employing intracerebral infection with lymphocytic choriomeningitis virus, found NOS-2 expression only in macrophages. While the viral burden was unaffected by a NOS inhibitor, such treatment exacerbated clinical severity and decreased survival time, suggesting a protective role for NO.

In wild-type mice early in the course of systemic inflammation, resulting from an injection of endotoxin, there is a marked expression of NOS-2 in vascular, glial and neuronal structures adjacent to areas devoid of a blood-brain barrier (Wong et al., 1996; Htain et al., 1997). Boje (1996) found that intracisternal administration of endotoxin resulted in NO production from the meninges and alteration in blood-brain barrier properties, which could be reversed by a NOS inhibitor. It is of interest to note that mice with deletions in the NOS-2 gene are protected from death after high-doses of endotoxin (MacMicking et al., 1995).

### Neurodegenerative diseases

Characteristic of such diseases is the progressive loss of a subset of neurons and there is evidence to implicate NO as causal or, at least, contributory.

In a proportion (20%) of cases of familial amyotrophic lateral sclerosis (ALS) there are multiple missense mutations in cytosolic Cu/Zn superoxide dismutase (SOD) and it is proposed that these result in either a gain of function, or increase an injurious reaction catalyzed by SOD. For example, Crow and Beckman (1995) suggest that there is enhanced nitration by NO of neurofilament protein. Evidence for nitration being causal in ALS comes from the observation that nitration blocks phosphorylation of tyrosine residues and that nitrotyrosine resembles the strongly antigenic dinitrophenol, perhaps provoking an autoimmune reaction.

Inherited retinal dystrophy results from accumulation of cellular debris in the subretinal space and visual cell degeneration. Cotinet et al. (1997) looked at the retinal Muller (astrocyte-like) cell from dystrophic rats and found that they produce abnormal amounts of TNF $\alpha$  and NOS-2 derived NO. As NO can decrease phagocytosis of photoreceptor segments, which underlies this dystrophy, they suggest that NO may disrupt transmission in retinal neurons.

Alzheimer's disease (AD) occurs in 10–15% of individuals over 65 and is characterized by the loss of pyramidal neurons in the hippocampus and neocortex, and of cholinergic neurons in the median forebrain. The hallmark of AD is the deposition of  $\beta$ -amyloid aggregates which can be toxic to neurons and also leads to NO production from glial cells (Rossi and Bianchini, 1996; Barger and Harmon, 1997) and neurons (Vodovotz et al., 1996). Weldon et al. (1998) injected soluble and fibrillar  $\beta$ -amyloid into rat striatum. They noted that microglia and astrocytes nearby showed a marked increase in NOS-2 expression and speculate that the resultant NO may be involved in neuronal killing.

### Demyelinating diseases

Demyelinating diseases are characterized by preferential damage to myelin, with relative preservation of axons. Multiple sclerosis (MS) and animal models such as experimental autoimmune encephalomyelitis (EAE) are inflammatory demyelinating diseases characterized by early damage to the blood-brain barrier. Lymphocyte and monocyte infiltration occurs, as well as activation of indigenous glia. What produces the resulting myelin damage is unknown and both indigenous as well as infiltrating cell types may contribute to myelin destruction, either directly or indirectly by secretion of proinflammatory cytokines or NO (Merrill and Benveniste, 1996).

A number of recent reports confirm a suggested role for NOS-2 in demyelinating disease. Using a spin trap method *in vivo* Hooper et al. (1995) could detect NO in the CNS of EAE animals,

which correlated with the onset of neurological symptoms. Increased NO production occurs in MS as illustrated by elevated levels of nitrite and nitrate levels in CSF (Johnson et al., 1995). Using immunocytochemistry, NOS-2 and nitrotyrosine have been found in diseased brain (Okuda et al., 1995; Hooper et al., 1995; Bagasra et al., 1995; DeGroot et al., 1997; Merrill et al., 1998). Both the protein and the transcript for NOS-2 colocalize with markers for macrophages/microglia (Bagasra et al., 1995; Hooper et al., 1997; van der Veen, 1997) and/or astrocytes (Okuda et al., 1997; Tran et al., 1997; Merrill et al., 1998). Cross et al. (1996) showed that NOS-2 transcript and enzyme activity correlate with active EAE. It is of interest that MS patients have elevated circulating antibodies to S-nitrosocysteine, suggesting nitrosylation of proteins in vivo (Boullerne et al., 1995). Akin to the human situation with MS, Ding et al. (1997) find gender-relatedness in NOS-2 expression and NO production in mice with EAE.

Based on the evidence for NO production in MS and EAE, several studies have assessed roles for NO using NOS inhibitors. The results have been mixed. In one study of acute EAE inhibitors of NOS provided no benefit (Zielasek et al., 1995). Other EAE studies showed a small rise in clinical score (Ruuls et al., 1996) or a decrease in disease incidence, severity and duration (Cross et al., 1994; Zhao et al., 1996). Most recently, Hooper et al. (1997) report significant therapeutic effects on EAE with NOS inhibitors, and scavengers of NO and peroxynitrite.

The current therapeutic available for the treatment of MS is IFN $\beta$ , demonstrated to be a blocker of NOS-2 expression (Stewart et al., 1997). Hall et al. (1997) suggest that it is beneficial through its antagonism of endogenously produced IFN $\gamma$ , hence impairing the activation of macrophages and microglia. Treatment of EAE animals with anti-inflammatory cytokines such as IL10, another NOS-2 modulator, inhibits disease (Willenborg et al., 1995). However, using the EAE model in NOS-2 gene deficient mice, Fenyk-Melody et al. (1998) report that disease is more severe and prolonged. This might suggest either that NO is immunosup-

pressive at the time of immunization or that, without NO, there is a loss of the normal regulation of infiltration of inflammatory cells into the CNS (De Caterina et al., 1995; Hickey et al., 1997).

### **Cerebral ischemia**

Elucidating the sequence of physiological and biochemical events following cerebral ischemia, in an attempt to identify means to reduce the amount of damage that develops, has been the focus of extensive investigation. The early events that occur following blood flow reduction, leading to cell death, have been well characterized in animal models of cerebral ischemia (Siesjo, 1992). Briefly, ischemia induces mitochondrial injury leading to a rapid depletion of ATP levels and a disruption of normal ion homeostasis. Large amounts of glutamate are released and intracellular calcium levels rise because of NMDA-receptor activation. The massive increase in free intracellular calcium following ischemia is a precursor to a variety of events capable of promoting neuronal and glial cell death such as inflammation, gene activation, and free radical production (for a review see Iadecola, 1997). While the pathogenesis of ischemic brain injury is probably the result of the activation of multiple biochemical pathways, it is evident, given the chain of events that occur, that NO has the potential to participate in the mechanisms of cerebral ischemia.

There is sufficient evidence that NO is produced following both global and focal cerebral ischemia. Using techniques such as in vivo spin trapping (Sato et al., 1994), porphyrinic microsensor measurement (Malinski et al., 1993), and measurements of brain nitrite and cGMP levels as a reflection of increased NO production (Kader et al., 1993; Shibata et al., 1996), increased generation of NO in the brain following cerebral ischemia has been demonstrated.

### **NOS-2 and global ischemia**

Transient global ischemia is seen clinically associated with disorders of systemic circulation, such as

heart failure or other cardiac dysfunction. Blood flow to the brain may be reduced below levels of autoregulation or may be completely interrupted for a period of time. Global ischemia induces a gliotic reaction, and delayed neuronal necrosis occurs in selectively vulnerable cells such as pyramidal neurons in the CA1 subfield of the hippocampus. NOS-2 protein expression has been shown to occur in a rat model of transient global ischemia (Endoh et al., 1994). Three days after a 10 min ischemic episode, NOS-2 protein was detected in the CA1 region of the hippocampus by immunohistochemistry, with immunoreactivity progressively increasing to day 30. Double-labeling experiments identified these NOS-2-expressing cells as reactive astrocytes, and ruled out microglial expression.

In our own studies with spontaneously hypertensive rats, we can detect NOS-2 protein by 72 h of reperfusion following 30 min of global ischemia. These cells morphologically resemble astrocytes in the corpus callosum, basal ganglia, and the hippocampus (Fig. 1). In response to global ischemia, NOS-2 induction parallels, spatially and temporally, the gliotic reaction. The induction of NOS-2 appears to be restricted to astrocytes in regions of neuronal damage, as there is no evidence for NOS-2 expression in other cell types or other regions. The mechanism of delayed NOS-2 induction, and the role of NO generated by this isoform following global ischemia remains to be determined. To date, no studies have been performed investigating global ischemic damage in mice deficient in NOS-2 expression. This may be due to the difficulties associated with mice not surviving the ischemic episode (Yang et al., 1997), or it may be due to the complications of strain variability in global ischemic outcome (Fujii et al., 1997).

### **NOS-2 and focal ischemia**

Focal cerebral ischemia, or stroke, occurs as a regional injury and results from the prolonged interruption of blood flow due to the occlusion of a single CNS artery. A focal vascular occlusion









Fig. 1. Expression of NOS-2 in astrocytes following global ischemia. In tissue sections stained with hematoxylin and eosin, neuronal cell death can be seen in the CA1 region of the rat hippocampus three days following 30 min of transient global ischemia (A, B). Astrogliosis, indicated by an increase in GFAP immunoreactivity, is also evident three days after the insult (C, D). Immunohistochemistry reveals NOS-2 protein expressed after three days in cells morphologically resembling astrocytes (E, F). Monoclonal GFAP antibody (Sigma, 1:200) and polyclonal NOS-2 antibody (Transduction Labs., 1:100) labelling was detected using standard immunoperoxidase techniques with 3, 3'-diaminobenzidine as substrate. All sections were counterstained with hematoxylin and eosin. Magnification: A, 40x; B, 200x; C, 400x; D-F, 600x.

produces a localized, destructive lesion or infarct, with necrosis of all cellular elements in the territory normally perfused by the occluded vessel. Ischemic stroke occurs most often in humans as the result of a thromboembolic vascular occlusion, secondary to atherosclerosis, in which there is a focal deposition of plasma-derived lipids that inhibits normal flow. In rodent models, focal ischemia can be induced by either permanently or transiently occluding the middle cerebral artery (MCA).

Several studies have provided biochemical, molecular, and immuno-histochemical evidence of NOS-2 induction following focal cerebral ischemia. Two separate reports demonstrated the induction of NOS-2 activity following permanent MCA occlusion in rats (Yoshida et al., 1995; Iadecola 1995a). Both studies found an increase in NOS-2 activity at 2-3 days post-occlusion, with a return to baseline by seven days. The time course of NOS-2 induction suggests that NO generated from this isoform does not contribute to the early stages of ischemic injury. Iadecola et al. (1995c; 1996) have also investigated NOS-2 mRNA and protein expression in response to focal ischemia. Following permanent MCA occlusion in rats, NOS-2 mRNA was detected by 12 h, peaked at 48 h and returned to baseline by seven days post-occlusion. Immunoreactivity for NOS-2, observed at 48-96 hours after ischemia, was detected only in neutrophils invading the infarct and was not seen in resident cells (Iadecola et al., 1996). Transient





focal ischemia in rats also leads to NOS-2 induction, though with different spatial and temporal patterns of expression than seen following permanent occlusion. Two hours of MCA occlusion, followed by reperfusion, resulted in maximal expression of NOS2 mRNA at an earlier time point, and detection of protein predominantly in vascular cells, rather than in neutrophils (Iadecola et al., 1996).

In mice, permanent MCA occlusion using the filament model of ZeaLonga et al. (1989) leads to the induction of NOS-2 mRNA and protein expression. Using *in situ* hybridization, we were able to detect NOS-2 message at 24, 48 and 72 h (latest point studied), but not at 6 h post-occlusion (Fig. 2). At 24 hours after ischemia, NOS-2 immunoreactivity was observed primarily in what appeared, morphologically, to be inflammatory cells within the infarct (Figure 2). However, at 48 and 72 h NOS-2 immunoreactivity was also observed in cells morphologically resembling astrocytes, as well as by GFAP staining on adjacent sections (Fig. 2). These NOS-2 positive cells were located primarily in regions surrounding the infarct. In contrast to previously published reports, these data indicate that resident glial cells, in addition to infiltrating cells, express NOS-2 in the ischemic brain. The differences in our findings compared to previous reports may reflect the differences in total infarct volumes produced as a result of using different surgical procedures (cauterization versus filament occlusion). The infarct volume generated with the filament model of MCA occlusion was typically greater than 80 mm<sup>3</sup> (Loihl and Murphy, 1997) while infarcts produced by cauterization were generally smaller (< 40 mm<sup>3</sup>), and predominantly cortical (Iadecola et al., 1997).

### Role of NOS-2 in cerebral ischemia

While it is clear that NO participates in the mechanisms of ischemia, much of the early work investigating the precise role of NO yielded contradictory results. This controversy stemmed largely from the fact that much of the research being










Fig. 2. Expression of NOS-2 revealed by *in situ* hybridization and immunohistochemistry at 48 and 72 h following middle cerebral artery occlusion in mice. NOS-2 mRNAs, detected with a 35S-labelled cRNA probe (Grzybicki et al., 1997), was observed at 48 and 72 h following permanent MCA occlusion (A, B). NOS-2 immunoreactivity was evident at 48 h post-occlusion, in what appear to be infiltrating inflammatory cells within the infarct (C). NOS-2 immunoreactivity at 72 h post-occlusion was also seen in cells morphologically resembling astrocytes located at the border of the infarct (D). GFAP immunoreactivity reveals astrocytes located at the infarct border at 48 h post-occlusion (E). NOS-2 immunoreactivity at 48 h post-occlusion, in an adjacent section to that shown in E, in cells resembling astrocytes (F). See legend to Fig. 1 for details of methods. Magnification: A, B, 200x; C-F, 400x.

done involved the use of pharmacological inhibitors that, while somewhat selective, are not actually specific for a particular isoform of NOS. Experiments *in vivo* have shown that inhibition of NO synthesis ameliorates (Nowicki et al., 1991; Buisson et al., 1992; Nagafugi et al., 1992; Ashwal et al., 1993; Iadecola et al., 1995b, 1996; Zhang et al., 1996) and worsens (Yamamoto et al., 1992; Dawson et al., 1992; Kuluz et al., 1993) tissue damage in animal models of cerebral ischemia. Given the scope of potential biological actions of NO, and the different spatial and temporal patterns of expression of the NOS isoforms, it is likely that NO generated by the different isoforms is subserving different roles at different times following ischemic insult. Therefore, non-specifically blocking NOS activity overshadows the potential for multiple roles of NO in ischemia. To circumvent the limitations of the currently available pharmacological inhibitors, and to elucidate the contributions of the different isoforms of NOS, ischemic brain damage has been investigated in genetically altered mice, which are deficient in a particular isoform of NOS.

Evidence provided in the last few years using gene targeting strategies has resolved the debate over the roles of the constitutive isoforms of NOS in the pathophysiology of focal cerebral ischemia (Iadecola, 1997). Infarct volume, resulting from MCA occlusion, in mice deficient for NOS-1 is reduced compared to wild type mice. Infarct



volume, however, in these mutant mice increased following administration of nitro-L-arginine, a NOS inhibitor, presumably due to inhibition of NOS-3 (Huang et al., 1994). In mice deficient for NOS-3, infarct volumes were larger than in wild type littermates. These results indicate that NO derived from NOS-1 acts early following cerebral ischemia and contributes to the development of tissue damage, while NO generated from NOS-3 serves a protective function, most likely by regulating blood flow to the ischemic penumbra.

More recently, evidence has been provided that begins to elucidate the role of the NOS-2 isoform in focal cerebral ischemia. A report by Iadecola et al. (1997) demonstrated that 24 h following MCA occlusion, infarct volumes in mice deficient for NOS-2 did not vary from infarct volumes in wildtype animals. We have found similar results in our own studies (Loihl and Murphy, 1997). This is consistent with the idea, suggested by enzyme activity studies, that NOS-2 does not contribute to the early stages of ischemic damage, and is not particularly surprising given the time course of NOS-2 induction. Iadecola et al. (1997) did find, however, that 96 h after MCA occlusion infarct volumes in NOS-2 deficient mice were 28% smaller than in wild-type controls. This smaller infarct at 96 h post-occlusion was accompanied by less severe motor deficits in mutant mice, indicating that the lack of NOS-2 correlates with improved functional outcome.

### **Regulation and role of astroglial NOS-2 in neuropathologies**

Perhaps the only conclusions that can be drawn from the neuropathological evidence presented above are that NOS-2 is sometimes expressed in astroglial cells and that, when there is expression, its appearance is delayed (>1 day) beyond what we might predict from *in vitro* observations (a few hours). This tardy appearance of NOS-2 might simply reflect development of the appropriate conditions for gene activation, such as infiltration of inflammatory cells either producing the relevant cytokines or initiating their production in

resident cells. Indeed, in some pathologies other stimuli may predominate, such as hypoxia. On the other hand, we could interpret the delay in NOS-2 expression as a reflection of the presence of factors that suppress gene activation. In this case, the timing of NOS-2 expression and the production of NO from this source may represent an adaptive response, and thus hint at a beneficial function.

If NOS-2 gene expression is initially suppressed following damage, then the *in vitro* evidence points to a number of likely candidates. While the *in vivo* data are not definitive, they are at least consistent. Levels of transforming growth factor  $\beta$ 1 rise following ischemia (Wang et al., 1995; Krupinski et al., 1996). Extracellular glutamate and ATP are elevated and persist for a few hours following CNS injury and the activation of selective chemokine genes is an early event. Specific chemokines are elevated following trauma (Ghirnikar et al., 1996; Grzybicki et al., 1998), exposure to  $\beta$ -amyloid (Meda et al., 1996), with bacterial infection (Spanaus et al., 1997), in EAE and MS (Hulkower et al., 1993; Karpus et al., 1995; Miyagishi et al., 1995; Ransohoff et al., 1996) and in ischemia (Kim et al., 1995; Yamasaki et al., 1995, 1997; Takami et al., 1997; Yoshimoto et al., 1997). Not only will chemokines such as IL-8 and MCP-1 block NOS-2 induction (McCall et al., 1992; Rojas et al., 1993) but NO can also inhibit the expression of IL-8 and MCP-1 (Andrew et al., 1995; Zeiher et al., 1995; De Caterina et al., 1995). A delayed appearance of NOS-2 might therefore ensure an environment in which chemokine gene expression can proceed and the entry of hematogenous cells is unhindered by NO effects on endothelial adhesion molecule expression (Khan et al., 1996; Tsao et al., 1996; Hickey et al., 1997). That the later production of NO might then terminate infiltration of inflammatory cells into the parenchyma, or downregulate expression of pro-inflammatory cytokines, are intriguing possibilities. Together with its general role as a vasodilator, thus improving perfusion in compromised tissue and leading to remodelling, NO may therefore prove beneficial by limiting the inflammatory response.

From functional and genetic studies, it seems that the effects of NO from NOS-2 can either worsen or improve recovery from damage. Evidence from the various enzyme inhibition studies is mixed, fraught with problems regarding specificity. Evidence from studies with NOS-2 gene-deficient mice is emerging to suggest that the presence of the enzyme can be beneficial, deleterious or without consequence (Nathan 1997). In cerebral ischemia, while NOS-2 clearly does not contribute to pathology within the first 24 h (Iadecola et al., 1997; Loihl and Murphy, 1997), gene-deficient mice are reported to have smaller infarcts at 4 days (Iadecola et al., 1997). Such NOS-2 deficient animals have a greater leukocyte response in endotoxemia (Hickey et al., 1997) which might explain why the condition of such mice with EAE is made worse (Fenyk-Melody et al., 1998).

The interpretation of results from gene-deficient mice is itself not without problems because of the possibility of developmental compensation and the real potential for expression of a truncated protein, perhaps with novel or unregulated functions. However, and in the absence of truly selective enzyme inhibitors, observing neuropathological outcome in experimental models employing such NOS-2 deficient mice is a powerful indicator of the roles of NOS-2. Verification could then be sought by virally re-introducing the gene in deficient animals. It would also be informative to observe pathological outcomes in transgenic mice in which NOS-2 can be overexpressed. However, there are no reports of the generation of such animals to date.

### Acknowledgements

We recognize the contributions of Sherry Kardos and Hsin Lee Lin to this work, which is supported by the NIH (NS29226) and American Heart Association (Grant-in-Aid).

### References

- Adamson, D.C., Wildemann, B., Sasaki, M., Glass, J.D., McCarthur, J.C., Christov, V.I., Dawson, T.M. and Dawson, V.L. (1996) Immunologic NO synthase: Elevation in severe AIDS dementia and induction by HIV-1 gp41. *Science*, 274: 1917–1921.
- Andrew, P.J., Harant, H. and Lindley, I.J.D. (1995) Nitric oxide regulates IL-8 expression in melanoma cells at the transcriptional level. *Biochem. Biophys. Res. Commun.*, 214: 949–956.
- Arnhold, S., Andressen, C., Bloch, W., Mai, J.K. and Addicks, K. (1997) NO synthase II is transiently expressed in embryonic mouse olfactory receptor neurions. *Neurosci. Lett.*, 229: 165–168.
- Ashwal, S., Cole, D.J., Osborne, T.N. and Pearce, W.J. (1993) Low dose L-NAME reduces infarct volume in the rat MCA/O reperfusion model. *J. Neurosurg. Anesthesiol.*, 5: 241–259.
- Bagasra, O., Michaels, F.H., Zheng, Y.M., Bobroski, L.E., Spitsin, S.V., Fu, Z.F., Tawadros, R. and Koprowski, H. (1995) Activation of the inducible form of nitric oxide synthase in the brains of patients with multiple sclerosis. *Proc. Natl. Acad. Sci. USA*, 92: 12041–12045.
- Barger, S.W. and Harmon, A.D. (1997) Microglial activation by Alzheimer amyloid precursor protein and modulation by apolipoprotein E. *Nature*, 388: 878–881.
- Boje, K.M.K. (1996) Inhibition of nitric oxide synthase attenuates blood–brain barrier disruption during experimental meningitis. *Brain Res.*, 720: 75–83.
- Borgerding, R. and Murphy, S. (1995) Expression of NO synthase in cerebral endothelial cells is regulated by cytokine-activated astrocytes. *J. Neurochem.*, 65: 1342–1347.
- Boullerne, A.I., Petry, K.G., Meynard, M. and Geffard, M. (1995) Indirect evidence for nitric oxide involvement in multiple sclerosis by characterization of circulating antibodies directed against conjugated S-nitrosocysteine. *J. Neuroimmunol.*, 60: 117–124.
- Buisson, A., Plotkine, M. and Boulu, R.G. (1992) The neuroprotective effect of a nitric oxide inhibitor in a rat model of focal cerebral ischemia. *Br. J. Pharmacol.*, 106: 766–767.
- Butler, A.R., Flitney, F.W. and Williams, D.L.H. (1995) NO, nitrosonium ions, nitroxide ions, nitrosothiols and iron-nitrosyls in biology. *Trends Pharmacol. Sci.*, 16: 18–22.
- Campbell, I.L. (1996) Exacerbation of lymphocytic choriomeningitis in mice treated with the inducible nitric oxide synthase inhibitor aminoguanidine. *J. Neuroimmunol.*, 71: 31–36.
- Christopherson, K.S. and Bredt, D.S. (1997) Nitric oxide in excitable tissues. *J. Clin. Invest.*, 100: 2424–2429.
- Colasanti, M., Persichini, T., Menegazzi, M., Mariotto, S., Giodano, E., Calderera, C.M., Sogos, V., Lauro, G.M. and Suzuki, H. (1995) Induction of nitric oxide synthase mRNA expression. *J. Biol. Chem.*, 270: 26731–26733.
- Cotinet, A., Goureau, O., Hicks, D., Thillaye-Goldenberg, B. and de Kozak, Y. (1997) TNF and NO production by retinal Muller glial cells from rats exhibiting inherited retinal dystrophy. *Glia*, 20: 59–69.
- Cross, A.H., Misko, T.P., Lin, R.F., Hickey, W.F., Trotter, J.L. and Tilton, R.G. (1994) Aminoguanidine, an inhibitor of

- inducible nitric oxide synthase, ameliorates experimental autoimmune encephalomyelitis in SJL mice. *J. Clin. Invest.*, 93: 2684–2690.
- Cross, A.H., Keeling, R.M., Goorha, S., San, M., Rodi, C., Wyatt, P.S., Manning, P.T. and Misko, T.P. (1996) Inducible nitric oxide synthase gene expression and enzyme activity correlate with disease activity in murine EAE. *J. Neuroimmunol.*, 71: 145–153.
- Crow, J.P. and Beckman, J.S. (1995) Reactions between nitric oxide, superoxide and peroxynitrite. In: L. Ignarro and F. Murad (Eds.), *Nitric Oxide*, Academic Press, San Diego, pp. 17–43.
- Dawson, D.A., Kusumoto, K., Graham, D.I., McCulloch, J. and Macrae, I.M. (1992) Inhibition of nitric oxide synthesis does not reduce infarct volume in a rat model of focal cerebral ischemia. *Neurosci. Lett.*, 142: 151–154.
- De Caterina, R., Libby, P., Peng, H.-B., Thannickal, V.J., Rajavshisth, T.B., Gimbrone, M.A. Jr., Shin, W.S. and Liao, J.K. (1995) Nitric oxide decreases cytokine-induced endothelial activation. *J. Clin. Invest.*, 96: 60–68.
- De Groot, C.J.A., Ruuls, S.R., Theeuwes, J.W.M., Dijkstra, C.D. and Van der Walk, P. (1997) Immunocytochemical characterization of the expression of inducible and constitutive isoforms of nitric oxide synthase in demyelinating multiple sclerosis lesions. *J. Neuropathol. Exp. Neurol.*, 56: 10–20.
- DelaTorre, A., Schroeder, R.A. and Kuo, P.C. (1997) Alteration of NF- $\kappa$ B p50 DNA binding kinetics by S-nitrosylation. *Biochem. Biophys. Res. Commun.*, 238: 703–706.
- Ding, M., Wong, J.L., Rogers, N.E., Ignarro, L.J. and Voslhuil, R.R. (1997) Gender differences of inducible nitric oxide production in SJL/J mice with EAE. *J. Neuroimmunol.*, 77: 99–106.
- Eddleston, M. and Mucke, L. (1993) Molecular profile of reactive astrocytes-implications for their role in neurologic disease. *Neuroscience*, 54: 15–36.
- Endoh, M., Maiese, K. and Wagner, J.A. (1994) Expression of the inducible form of nitric oxide synthase by reactive astrocytes after transient global ischemia. *Brain Res.*, 651: 92–100.
- Feinstein, D.L. (1998) Suppression of astroglial nitric oxide synthase expression by norepinephrine results from decreased NOS-2 promoter activity. *J. Neurochem.* 70: 1484–1496.
- Feinstein, D.L., Galea, E., Aquino, D.A., Li, G.C., Xu, H. and Reis, D.J. (1996) Heat shock protein 70 suppresses astroglial inducible nitric oxide synthase expression by decreasing NF $\kappa$ B activation. *J. Biol. Chem.*, 271: 17724–17732.
- Fenyk-Melody, J., Garrison, A., Brunnert, S., Weidner, J., Shen, F., Shelton, B. and Mudgett, J.S. (1998) Experimental autoimmune encephalomyelitis is exacerbated in mice lacking the NOS-2 gene. *J. Immunol.* 160: 2940–2949.
- Forstermann, U., Gath, I., Schwarz, P., Closs, E.I. and Kleinert, H. (1995) Isoforms of nitric oxide synthase. *Biochem. Pharmacol.*, 50: 1321–1332.
- Fujii, M., Hara, H., Meng, W., Vonsattel, J.P., Huang, Z. and Moskowitz, M.A. (1997) Strain-related differences in susceptibility to transient forebrain ischemia in SV-129 and C57.BL6 mice. *Stroke*, 28: 1805–1810.
- Galea, E., Reis, D.J. and Feinstein, D.L. (1994) Cloning and expression of inducible nitric oxide synthase from rat astrocytes. *J. Neurosci. Res.*, 37: 406–414.
- Galea, E., Reis, D.J., Xu, H. and Feinstein, D.L. (1995) Transient expression of calcium-independent nitric oxide synthase activity in brain blood vessels during development. *FASEB J.*, 9: 1632–1637.
- Ghirnikar, R.S., Lee, Y.L., He, T.R. and Eng, L.F. (1996) Chemokine expression in rat stab wound brain injury. *J. Neurosci. Res.*, 46: 727–733.
- Griffith, O.W. and Stuehr, D.J. (1995) Nitric oxide synthases. *Annu. Rev. Physiol.*, 57: 707–736.
- Grzybicki, D., Kwack, K.B., Perlman, S. and Murphy, S. (1997) Nitric oxide synthase type II expression by different cell types in MHV-JHM encephalitis suggests distinct roles for nitric oxide in acute vs. persistent virus infection. *J. Neuroimmunol.*, 73: 15–27.
- Grzybicki, D., Moore, S.A., Schelper, R., Glabinski, A., Ransohoff, R.M. and Murphy, S. (1998) Expression of monocyte chemoattractant protein (MCP-1) and nitric oxide synthase-2 following cerebral trauma. *Acta Neuropathol.*, 95: 98–103.
- Hall, G.L., Compston, A. and Scolding, N.J. (1997)  $\beta$ -interferon and multiple sclerosis. *Trends Neurosci.*, 20: 63–67.
- Hamada, Y., Ikata, T., Katoh, S., Tsuchiya, K., Niwa, M., Tsutsumishata, Y. and Fukuzawa, K. (1996) Roles of nitric oxide in compression injury of rat spinal cord. *Free Rad. Biol. Med.*, 20: 1–9.
- Hickey, M.J., Sharkey, K.A., Sihota, E.G., Reinhardt, P.H., MacMicking, J.D., Nathan, C. and Kubes, P. (1997) Inducible nitric oxide synthase-deficient mice have enhanced leukocyte-endothelium interactions in endotoxemia. *FASEB J.*, 11: 955–964.
- Hooper, D.C., Ohnishi, S.T., Kean, R., Numagami, Y., Dietzschold, B. and Koprowski, H. (1995). Local nitric oxide production in viral and autoimmune diseases of the central nervous system. *Proc. Natl. Acad. Sci. USA*, 92: 5312–5316.
- Hooper, D.C., Bagasra, O., Marini, J.C., Zborek, A., Ohnishi, S.T., Kean, R., Champoin, J.M., Sarler, A.B., Bobroski, L., Farber, J.L., Akaike, T., Maeda, H. and Koprowski, H. (1997) Prevention of EAE by targeting nitric oxide and peroxynitrite. *Proc. Natl. Acad. Sci. USA*, 94: 2528–2533.
- Htain, W.W., Leong, S.K. and Ling, E.A. (1997) In vivo expression of inducible nitric oxide synthase in supra-ventricular amoeboid microglial cells in neonatal BALB/c and athymic mice. *Neurosci. Lett.*, 223: 53–56.
- Huang, Z., Huang, P.L., Panahian, N., Dalkara, T., Fishman, M.C. and Moskowitz, M.A. (1994) Effects of cerebral

- ischemia in mice deficient in neuronal nitric oxide synthase. *Science*, 265: 1883–1885.
- Hulkower, K., Brosnan, C.F., Aquino, D.A., Cammer, W., Kulshrestha, S., Guida, M.P., Rapoport D.A. and Berman, J.W. (1993) Expression of CSF-1, c-fms, and MCP-1 in the central nervous system of rats with experimental allergic encephalomyelitis. *J. Immunol.*, 150: 2525–2533.
- Iadecola, C. (1997) Bright and dark sides of nitric oxide in ischemic brain injury. *Trends Neurosci.*, 20: 132–139.
- Iadecola, C., Zhang, F., Xu, S., Casey, R. and Ross, M.E. (1995a) Inducible nitric oxide synthase expression in brain following cerebral ischemia. *J. Cerebral Blood Flow Metab.*, 15: 378–384.
- Iadecola, C., Zhang, F. and Xu, X. (1995b) Inhibition of inducible nitric oxide synthase ameliorates cerebral ischemic damage. *Am. J. Physiol.*, 268: R286–R292.
- Iadecola, C., Xu, X., Zhang, F., El-Fakahany, E.E. and Ross, M.E. (1995c) Marked induction of calcium-independent nitric oxide synthase activity after focal cerebral ischemia. *J. Cereb. Blood Flow Metab.*, 15: 52–59.
- Iadecola, C., Zhang, F., Casey, R., Clark, H.B. and Ross M.E. (1996) Inducible nitric oxide synthase gene expression in vascular cells after transient focal cerebral ischemia. *Stroke*, 27: 1373–1380.
- Iadecola, C. Zhang, F., Casey, R., Nagayama, M. and Ross, M.E. (1997) Delayed reduction of ischemic brain injury and neurological deficits in mice lacking the inducible nitric oxide synthase gene. *J. Neurosci.*, 17: 9157–9164.
- Johnson, A.W., Land, J.M., Thompson, E.J., Bolanos, J.P., Clark, J.B. and Heales, S.J. (1995) Evidence for increased nitric oxide production in multiple sclerosis. *J. Neurol. Neurosur. Psych.*, 58: 107–116.
- Kader, A., Frazzini, V.I., Solomon, R.A. and Trifiletti, R.R. (1993) Nitric oxide production during focal cerebral ischemia in rats. *Stroke*, 24: 1709–1716.
- Karpus, W.J., Lukacs, N.W., McRae, B.L., Strieter, R.M., Kunkel, S.L. and Miller, S.D. (1995) An important role for the chemokine macrophage inflammatory protein-1 $\alpha$  in the pathogenesis of the T cell-mediated autoimmune disease, experimental autoimmune encephalomyelitis. *J. Immunol.*, 155: 5003–5010.
- Khan, B.V., Harrison, D.G., Olbrych, M.T., Alexander, R.W. and Medford, R.M. (1996) Nitric oxide regulates vascular cell adhesion molecule 1 gene expression and redox-sensitive transcriptional events in human vascular cells. *Proc. Natl. Acad. Sci. USA*, 93: 9114–9119.
- Kim, J.S., Gautam, S.C., Chopp, M., Zaloga, C., Jones, M.L., Ward, P.A. and Welch, K.M.A. (1995) Expression of monocyte chemoattractant protein-1 and macrophage inflammatory protein-1 after focal cerebral ischemia in the rat. *J. Neuroimmunol.*, 56: 127–134.
- Knowles, R.G. (1997) Nitric oxide, mitochondria and metabolism. *Trans. Biochem. Soc.*, 25: 895–901.
- Koka, P., He, K., Zack, J.A., Kitchen, S., Peacock, W., Fried, I., Tran, T., Yashar, S.S. and Merrill, J.E. (1995) Human immunodeficiency virus 1 envelope proteins induce interleukin 1, tumor necrosis factor  $\alpha$  and nitric oxide in glial culture from fetal, neonatal, and adult human brain. *J. Exp. Med.*, 182: 941–952.
- Kopnisky, K.L., Sumners, C. and Chandler, L.J. (1997) Cytokine- and endotoxin-induced nitric oxide synthase in rat astroglial cultures. *J. Neurochem.*, 68: 935–944.
- Kossmann, T., Stahel, P.F., Lenzlinger, P.M., Redl, H., Dubs, R.W., Trentz, O., Schalg, G. and Morganti-Kossmann, M.C. (1997) Interleukin 8 released into the CSF after brain injury is associated with blood-brain barrier dysfunction and NGF production. *J. Cereb. Blood Flow Metab.*, 17: 280–289.
- Krupinski, J., Kumar, P., Kumar, S. and Kaluza, J. (1996) Increased expression of TGF- $\beta$ 1 in brain tissue after ischemic stroke in humans. *Stroke*, 27: 852–857.
- Kuluz, J.W., Prado, R.J., Dietrich, W.D., Schleine, C.L. and Watson, B.D. (1993) The effect of nitric oxide synthase inhibition on infarct volume after reversible focal cerebral ischemia in conscious rats. *Stroke*, 24: 2023–2029.
- Lane, T.E., Buchmeier, M.J., Watry, D.D. and Fox, H.S. (1996) Expression of inflammatory cytokines and inducible nitric oxide synthase in brains of SIV-infected Rhesus monkeys: Applications to HIV-induced central nervous system disease. *Molec. Med.*, 2: 27–37.
- Lane, T.E., Paoletti, A.D. and Buchmeier, M.J. (1997) Disassociation between the invitro and in vivo effects of nitric oxide on a neurotropic murine coronavirus. *J. Virol.*, 71: 2202–2210.
- Lei, D.L., Yang, D.L. and Liu, H.M. (1996) Local injection of kainic acid causes widespread degeneration of NADPH-d neurons and induction of NADPH-d in neurons, endothelial cells and reactive astrocytes. *Brain Res.*, 730: 199–206.
- Lin, H.L. and Murphy, S. (1997) Regulation of astrocyte nitric oxide synthase type II expression by ATP and glutamate involves loss of transcription factor binding to DNA. *J. Neurochem.*, 69: 612–616.
- Loibl, A.K. and Murphy, S. (1997) Expression of nitric oxide synthase-2 in various cell types following focal cerebral ischemia in mice. *Abst. Soc. Neurosci.*, 23: 2437.
- Luss, H., DiDilvio, M., Litton, A.L., Molin y Vedia, L., Nussler, A.K. and Billiar, T.R. (1994) Inhibition of nitric oxide synthesis enhances the expression of inducible nitric oxide synthase mRNA and protein in a model of chronic liver inflammation. *Biochem. Biophys. Res. Commun.*, 204: 635–640.
- MacMicking, J.D., Nathan, C., Hom, G., Chartrain, N., Fletcher, D.S., Trumbauer, M., Stevens, K., Xie, Q., Sokol, K., Hutchinson, N., Chen, H. and Mudgett, J.S. (1995) Altered responses to bacterial infection and endotoxin shock in mice lacking inducible nitric oxide synthase. *Cell*, 81: 641–650.

- Malinski, T., Bailey, F., Zhang, Z.G. and Chopp, M. (1993) Nitric oxide measured by a porphyrinic microsensor in rat brain after transient middle cerebral artery occlusion. *J. Cereb. Blood Flow Metab.*, 13: 355-358.
- McCall, T.B., Palmer, R.M.J. and Moncada, S. (1992) Interleukin-8 inhibits the induction of nitric oxide synthase in rat peritoneal neutrophils. *Biochem. Biophys. Res. Commun.*, 186: 680-685.
- Meda, L., Bernasconi, S., Bonaiuto, C., Sozzani, S., Zhou, D., Otvos, L., Mantovani, A., Rossi, F. and Cassatella, M.A. (1996)  $\beta$ -amyloid (25-35) peptide and IFN- $\gamma$  synergistically induce the production of the chemotactic cytokine MCP-1/JE in monocytes and microglial cells. *J. Immunol.*, 157: 1213-1218.
- Merrill, J.E., Genain, C.P., Parkinson, J.F., Medberry, P., Halks-Miller, M., DelVecchio, V., Kardos, S. and Murphy, S. (1998) iNOS and nitrotyrosine in macrophages and glia in demyelinating lesions. Submitted
- Merrill, J.E. and Murphy, S. (1997) Regulation of gene expression in the nervous system by reactive oxygen and nitrogen species. *Metab. Brain Disease*, 12: 97-112.
- Merrill, J.E. and Benveniste, E.N. (1996) Cytokines in inflammatory brain lesions: Helpful and harmful. *Trends Neurosci.*, 19: 331-338.
- Miyagishi, R., Kikuchi, S., Fukazawa, T. and Tashiro, K. (1995) Macrophage inflammatory protein-1 $\alpha$  in the cerebrospinal fluid of patients with multiple sclerosis and other inflammatory neurological diseases. *J. Neurol. Sci.*, 129: 223-227.
- Murphy, S. and Grzybicki, D. (1996) Glial NO: Normal and pathological roles. *Neuroscientist*, 2: 91-100.
- Murphy, S., Simmons, M.S., Agullo, L., Garcia, A., Feinstein, D.L., Gallea, E., Reis, D.L., Minc-Golomb, D. and Schwartz, J.P. (1993) Synthesis of nitric oxide in CNS glial cells. *Trends Neurosci.*, 16: 323-328.
- Nagafugi, T., Matsui, T., Koide, T. and Asano, T. (1992) Blockade of nitric oxide formation by N<sup>w</sup>-nitro-L-arginine mitigates ischemic brain edema and subsequent cerebral infarction in rats. *Neurosci. Lett.*, 147: 159-162.
- Nathan, C. (1997) Inducible nitric oxide synthase. *J. Clin. Invest.*, 100: 2417-2423.
- Nowicki, J.P., Duval, D., Poignet, H. and Scatton, B. (1991) Nitric oxide mediates neuronal death after focal cerebral ischemia in the mouse. *Eur. J. Pharmacol.*, 204: 339-340.
- Okuda, Y., Nakatsuji, Y., Fujimura, H., Esumi, H., Ogura, T., Yanagihara, T. and Sakoda, S. (1995) Expression of the inducible isoform of nitric oxide synthase in the central nervous system of mice correlates with the severity of actively induced experimental allergic encephalomyelitis. *J. Neuroimmunol.*, 62: 103-112.
- Okuda, Y., Sakoda, S., Fujimura, H. and Yanagihara, T. (1997) Nitric oxide via an inducible isoform of nitric oxide synthase is a possible factor to eliminate inflammatory cells from the CNS of mice with EAE. *J. Neuroimmunol.*, 73: 107-116.
- Oleszak, E.L., Katsetos, C.D., Kuzmak, J. and Varadhachary, A. (1997) Inducible nitric oxide synthase in Theiler's murine encephalomyelitis virus infection. *J. Virol.*, 71: 3228-3235.
- Park, S.K. and Murphy, S. (1994) Duration of expression of inducible nitric oxide synthase in glial cells. *J. Neurosci. Res.*, 39: 405-411.
- Park, S.K. and Murphy, S. (1996) Nitric oxide synthase type II mRNA stability is translation- and transcription-dependent. *J. Neurochem.*, 67: 1766-1769.
- Park, S.K., Lin, H.L. and Murphy, S. (1994) Nitric oxide limits transcriptional induction of nitric oxide synthase in CNS glial cells. *Biochem. Biophys. Res. Commun.*, 201: 762-768.
- Park, S.K., Lin, H.L. and Murphy, S. (1997) Nitric oxide regulates nitric oxide synthase-2 gene expression by inhibiting NF- $\kappa$ B binding to DNA. *Biochem. J.*, 322: 609-613.
- Peng, H., Libby, P. and Liao, J.K. (1995) Induction and stabilization of I $\kappa$ B $\alpha$  by nitric oxide mediates inhibition of NF- $\kappa$ B. *J. Biol. Chem.*, 270: 14214-14219.
- Ransohoff, R.M., Glabinski, A. and Tani, M. (1996) Chemokines in immune-mediated inflammation of the central nervous system. *Cytokine and Growth Factor Rev.*, 7: 35-46.
- Rojas, A., Delgado, R., Glaria, L. and Palacios, M. (1993) MCP-1 inhibits the induction of nitric oxide synthase in J774 cells. *Biochem. Biophys. Res. Commun.*, 196: 274-279.
- Rossi, F. and Bianchini, E. (1996) Synergistic induction of nitric oxide by  $\beta$ -amyloid and cytokines in astrocytes. *Biochem. Biophys. Res. Commun.*, 225: 474-478.
- Ruuls, S.R., Van Der Linden, S., Sontrop, K., Huitinga, I. and Dijkstra, C.D. (1996) Aggravation of experimental allergic encephalomyelitis by administration of nitric oxide synthase inhibitors. *Clin. Exp. Immunol.*, 103: 467-476.
- Sato, S., Tominaga, T., Ohnishi, T. and Ohnishi, S.T. (1994) Role of nitric oxide in brain ischemia. *Ann. NY Acad. Sci.*, 738: 369-373.
- Shafer, R. and Murphy, S. (1997) Activated astrocytes induce nitric oxide synthase-2 in cerebral endothelium via TNF $\alpha$ . *GLIA*, 21: 370-379.
- Shibata, M., Araki, N., Hamada, J., Sasaki, T., Shimazu, K. and Fukuuchi, Y. (1996) Brain nitrite production during global ischemia and reperfusion. *Brain Res.*, 735: 86-90.
- Siesjö, B.K. (1992) Pathophysiology and treatment of focal cerebral ischemia. *J. Neurosurg.*, 77: 169-184.
- Spanaus, K-S., Nadal, D., Pfister, H-W., Seebach, J., Widmer, U., Frei, K., Gloor, S. and Fontana, A. (1997) C-X-C and C-C chemokines are expressed in the CSF in bacterial meningitis and mediate chemotactic activity on peripheral blood-derived polymorphonuclear and mononuclear cells in vitro. *J. Immunol.*, 158: 1956-1964.
- Spitsin, S.V., Koprowski, H. and Michaels, F.H. (1996) Characterization and functional analysis of the human inducible nitric oxide synthase gene promoter. *Molec. Med.*, 2: 226-235.

- Stewart, V.C., Giovannoni, G., Land, J.M., McDonald, W.I., Clark, J.B. and Heales, S.J.R. (1997) Pretreatment of astrocytes with interferon  $\alpha/\beta$  impairs interferon  $\gamma$  induction of nitric oxide synthase. *J. Neurochem.*, 68: 2547-2551.
- Sun, N., Grzybicki D., Castro, R.F., Murphy, S. and Perlman, S. (1995) Activation of astrocytes in the spinal cord of mice chronically infected with a neurotropic coronavirus. *Virology*, 213: 482-493.
- Takami, S., Nishikawa, H., Minami, M., Nishiyori, A., Sato, M., Akaike, A. and Satoh, M. (1997) Induction of MIP-1 $\alpha$  mRNA on glial cells after focal ischemia in the rat. *Neurosci. Lett.*, 227: 173-176.
- Togashi, H., Sasaki, M., Frohman, E., Taira, E., Ratan, R.R., Dawson, T.M. and Dawson, V.L. (1997) Neuronal (type I) nitric oxide synthase regulates nuclear factor  $\kappa$ B activity and immunologic (type II) nitric oxide synthase expression. *Proc. Natl. Acad. Sci. USA*, 94: 2676-2680.
- Tran, E.H., Hardin-Pouzet, H., Verge, G. and Owens, T. (1997) Astrocytes and microglia express inducible nitric oxide synthase in mice with EAE. *J. Neuroimmunol.*, 74: 121-129.
- Tsao, P.S., Buitrago, R., Chan, J.R. and Cooke, J.P. (1996) Fluid flow inhibits endothelial adhesiveness. *Circulation*, 94: 1682-1689.
- van der Veen, R.C., Hinton, D.R., Incardonna, F. and Hofman, F.M. (1997) Extensive peroxynitrite activity during progressive stages of CNS inflammation. *J. Immunol.*, 77: 1-7.
- Vodovotz, Y., Lucia, M.S., Flanders, K.C., Chesler, L., Xie, Q.W., Smith, T.W., Weidner, J., Mumford, R., Webber, R., Nathan, C., Roberts, A.B., Lippa, C.F. and Sporn, M.B. (1996) Inducible nitric oxide synthase in tangle-bearing neurons of patients with Alzheimer's disease. *J. Exp. Med.*, 184: 1425-1433.
- Wang, X., Yue, T., White, R.F., Baroen, F.C. and Feuerstein, G.Z. (1995) Transforming growth factor  $\beta$ 1 exhibits delayed gene expression following focal cerebral ischemia. *Brain Res. Bull.*, 36: 607-609.
- Weldon, D.T., Rogers, S.D., Ghilardi, J.R., Finke, M.P., Cleary, J.P., O'Hare, E., Esler, W.P., Maggio, J.E. and Mantyh, P.W. (1998) Fibrillar  $\beta$ -amyloid induces microglial phagocytosis expression of inducible nitric oxide synthase and loss of a select population of neurons in the rat CNS in vivo. *J. Neurosci.*, 18: 2161-2173.
- Willenborg, D.O., Fordham, S.A., Cowden, W.B. and Ramshaw, I.A. (1995) Cytokines and murine autoimmune encephalomyelitis: Inhibition or enhancement of disease with antibodies to select cytokines, or by delivery of exogenous cytokines using a recombinant vaccinia virus system. *Scand. J. Immunol.*, 41: 31-41.
- Wong, M.L., Rettori, V., Al-Shekhlee, A., Bongiorno, P.B., Canteros, G., McCann, S.M., Gold, P.W. and Licinio, J. (1996) Inducible nitric oxide synthase gene expression in the brain during systemic inflammation. *Nature Med.*, 2: 581-584.
- Xie, Q-W. and Nathan C. (1994) The high output nitric oxide pathway: Role and regulation. *J. Leuk. Biol.*, 56: 576-582.
- Yamamoto, S., Golanov, E.V., Berger, S.B. and Reis, D.J. (1992) Inhibition of nitric oxide synthase increases focal ischemic infarction in rat. *J. Cereb. Blood Flow Metab.*, 12: 717-726.
- Yamasaki, Y., Matsuo, Y., Matsuura, N., Onodera, H., Itoyama, Y. and Kogure, K. (1995) Transient increase of cytokine-induced neutrophil chemoattractant, a member of the interleukin-8 family, in ischemic brain areas after focal ischemia in rats. *Stroke*, 26: 318-323.
- Yamasaki, Y., Matsuo, Y., Zagorski, J., Matsuura, N., Onodera, H., Itoyama, Y. and Kogure, K. (1997) New therapeutic possibility of blocking cytokine-induced neutrophil chemoattractant on transient ischemic brain damage in rats. *Brain Res.*, 759: 103-111.
- Yang, G., Kitagawa, K., Matsushita, K., Mabuchi, T., Yagita, Y., Yanagihara, T. and Matsumoto, I. (1997) C57BL/6 strain is most susceptible to cerebral ischemia following bilateral common carotid occlusion among seven mouse strains. *Brain Res.*, 752: 209-218.
- Yoshida, T., Waechter, C., Huang, Z. and Moskowitz, M.A. (1995) Induction of nitric oxide synthase activity in rodent brain following middle cerebral artery occlusion. *Neurosci. Lett.*, 194: 214-218.
- Yoshimoto, T., Houkin, K., Tada, M. and Abe, H. (1997) Induction of cytokines, chemokines and adhesion molecule mRNA in a rat forebrain reperfusion model. *Acta Neuropathol.*, 93: 154-158.
- Zea Longa, E., Weinstein, P.R., Carlson, S. and Cummins, R. (1989) Reversible middle cerebral artery occlusion without craniectomy in rats. *Stroke*, 20: 84-91.
- Zeiger, A.M., Fisslthaler, B., Schray-Utz, B. and Busse, R. (1995) Nitric oxide modulates the expression of monocyte chemoattractant protein 1 in cultured human endothelial cells. *Circ. Res.*, 76: 980-986.
- Zhang, F., Casey, R., Ross, M.E. and Iadecola, C. (1996) Aminoguanidine ameliorates and L-arginine worsens brain damage from intraluminal middle cerebral artery occlusion. *Stroke*, 27: 317-323.
- Zhao, W., Tilton, R.G., Corbett, J.A., McDaniel, M.L., Misko, T.P., Williamson, J.R., Cross, A.H. and Hickey, W.F. (1996) Experimental allergic encephalomyelitis in the rat is inhibited by aminoguanidine, an inhibitor of nitric oxide synthase. *J. Neuroimmunol.*, 64: 123-133.
- Zielasek, J., Jung, S., Gold, R., Liew, F.Y., Toyka, K.V. and Hartung, H.-P. (1995) Administration of nitric oxide synthase inhibitors in experimental autoimmune neuritis and experimental autoimmune encephalitis. *J. Neuroimmunol.*, 58: 81-88.