

**ERRATUM**    **OPEN**

# Erratum: Treating tobacco dependence: guidance for primary care on life-saving interventions. Position statement of the IPCRG

O. C. P. Van Schayck<sup>1</sup>, S. Williams<sup>2</sup>, V. Barchilon<sup>3</sup>, N. Baxter<sup>2,4</sup>, M. Jawad<sup>5</sup>, P. A. Katsaounou<sup>6</sup>, B. J. Kirenga<sup>7</sup>, C. Panaitescu<sup>8</sup>, I. G. Tsiligianni<sup>9</sup>, N. Zwar<sup>10</sup> and A. Ostrem<sup>11</sup>

*npj Primary Care Respiratory Medicine* (2017)27:52; doi:10.1038/s41533-017-0048-4

**Erratum to:** *npj Primary Care Respiratory Medicine* (2017); doi:10.1038/s41533-017-0039-5; Published 09 June 2017

The original version of this article contained an error in the spelling of the author I. G. Tsiligianni, which was incorrectly given as K. W. I. G. Tsiligianni. This has now been corrected in both the PDF and HTML versions of the article.

adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing,

© The Author(s) 2017

<sup>1</sup>Department of Family Medicine, CAPHRI, Maastricht University, Maastricht, The Netherlands; <sup>2</sup>International Primary Care Respiratory Group, Aberdeen, UK; <sup>3</sup>Andalusian Health Service (SAS), Tobacco group of GRAP (Primary Care Respiratory Group), Andalusia, Spain; <sup>4</sup>Southwark Clinical Commissioning Group, London, UK; <sup>5</sup>Faculty of Medicine, School of Public Health, Imperial College London, London, UK; <sup>6</sup>Pulmonary Medicine, Medical School, National and Kapodistrian University of Athens, Evangelismos Hospital, Athens, Greece; <sup>7</sup>Lung Institute and Division of Pulmonary Medicine, Makerere University College of Health Sciences, Kampala, Uganda; <sup>8</sup>Family Medicine Solo Practice, RespiRo-Romanian Primary Care Respiratory Group, Bucharest, Romania; <sup>9</sup>Clinic of Social and Family Medicine, Faculty of Medicine, University of Crete, Crete, Greece; <sup>10</sup>School of Public Health and Community Medicine, UNSW Australia, Sydney, NSW, Australia and <sup>11</sup>General Practitioner, Gransdalen Legesenter, Oslo, Norway  
Correspondence: O. C. P. Van Schayck (onno.vanschayck@maastrichtuniversity.nl)

Published online: 05 September 2017

Published in partnership with Primary Care Respiratory Society UK