

Journal section: Oral Surgery
 Publication Types: Review

doi:10.4317/medoral.17130
<http://dx.doi.org/doi:10.4317/medoral.17130>

Survival of implants placed with the osteotome technique: An update

Jose Viña-Almunia ¹, Laura Maestre-Ferrín ¹, Teresa Alegre-Domingo ¹, María Peñarrocha-Diago ²

¹ DDS. Master in Oral Surgery and Implantology. Faculty of Medicine and Dentistry. University of Valencia

² Associate Professor of Oral Surgery. Professor of the Master in Oral Surgery and Implantology. Faculty of Medicine and Dentistry. University of Valencia. Valencia (Spain)

Correspondence:
 Clínicas Odontológicas
 Gascó Oliag 1
 46021 - Valencia (Spain)
maria.penarrocha@uv.es

Viña-Almunia J, Maestre-Ferrín L, Alegre-Domingo T, Peñarrocha-Diago MA. Survival of implants placed with the osteotome technique: An update. Med Oral Patol Oral Cir Bucal. 2012 Sep 1;17 (5):e765-8.
<http://www.medicinaoral.com/medoralfree01/v17i5/medoralv17i5p765.pdf>

Received: 03/06/2010
 Accepted: 03/12/2011

Article Number: 17130 <http://www.medicinaoral.com/>
 © Medicina Oral S. L. C.I.F. B 96689336 - pISSN 1698-4447 - eISSN: 1698-6946
 eMail: medicina@medicinaoral.com
Indexed in:
 Science Citation Index Expanded
 Journal Citation Reports
 Index Medicus, MEDLINE, PubMed
 Scopus, Embase and Emcare
 Indice Médico Español

Abstract

A literature review is made to analyze the survival of implants placed with the osteotome technique. A PubMed search was made based on the key words “osteotome AND dental implants”, corresponding to publications between 1998-2008. The inclusion criteria were: a) a minimum of 10 patients; b) a minimum follow-up of 6 months; c) implants placed using the osteotome technique with or without indirect sinus lift; and d) specification of the implant number and survival rate. Sixty-four articles were identified, of which 20 met the inclusion criteria. A total of 2006 implants were placed in 1312 patients using the osteotome technique. The duration of follow-up after prosthetic loading ranged from 6-144 months. Indirect sinus lift was carried out in all but one of the studies. The residual crest height ranged from 2.3-11.7 mm. with a mean gain in bone after sinus lift of 2.5-5.5 mm. The time from implant placement to prosthetic loading varied from 1.5-9 months. The percentage implant survival rate was 90.5-100%. The survival rate of implants placed with the osteotome technique is high and does not differ with respect to implant placement with the conventional technique.

Key words: Osteotomes, dental implants, indirect sinus lift.

Introduction

Summers was the first to describe the osteotome technique to increase bone density in the dental implant bed (1,2) and perform localized maxillary sinus lift (1,3). Benign paroxysmal vertigo has been reported as a complication secondary to tapping of the osteotome with the mallet (4).

The literature offers little information on the predictability of implant placement using the osteotome technique without added sinus lift. In most clinical studies, implant placement using the osteotome technique is carried out in combination with indirect sinus lift (5,6). A literature review is made to analyze the survival of implants placed with the osteotome technique.

Material and Methods

A PubMed search was made based on the key words “osteotome AND dental implants”, limiting the search to human studies published in English in dental journals during the period 1998-2008. The inclusion criteria were: a) a minimum of 10 patients; b) a minimum follow-up of 6 months; c) implants placed using the osteotome technique with or without indirect sinus lift; and d) specification of the implant number and survival rate. The following data were collected from each study: year of publication, inclusion criteria, type of intervention, results obtained and follow-up.

Results and Discussion

Sixty-four articles were identified with the key words “osteotome AND dental implants”. Of these articles, 20 met the inclusion criteria and were subjected to analysis (Table 1).

A total of 2006 implants were placed in 1312 patients using the osteotome technique. The duration of follow-up after prosthetic loading ranged from 6-144 months. Indirect sinus lift was carried out in all but one of the studies (7). Specifically, Strietzel et al. used osteotomes only for alveolar crest expansion, and concluded that

this technique is not indicated in Lekholm and Zarb type I and II bone, because osteotome pressure in such cortical bone adversely affects the vascular supply (7). The residual crest height ranged from 2.3-11.7 mm in the different studies (8-16) (Table 2). Rosen et al. (17), Diserens et al. (18) and Sforza et al. (19) performed indirect sinus lift with a minimum residual crest height of 3, 4 and 5 mm, respectively. The mean gain in bone after sinus lift was 2.5-5.5 mm (10,11,13,15, 20-23) (Table 2). Most of the studies used bone graft material when performing indirect sinus lift: particulate autologous bone (10,24,25), xenograft (Bio-Oss®) (9,16,18), or a combination of both (11,12,15,17,19). Five of the studies used no graft material (8,20,22,23,26). One study (21) made use of platelet-rich fibrin, while Barone et al. (13) used a mixture of collagen gel and porcine bone particles (Gel 40®, Osteobiol, Tecness). The implant survival rate in the sinus lift procedures made with graft material varied from 90.5-98.5%, versus 96-100% when no graft material was added.

The time from implant placement to prosthetic loading varied from 1.5 (15,21,24) to 9 months (10). In no case was immediate loading performed.

Table 1. Data collected from the articles meeting the study inclusion criteria.

Author and year	No. patients	No. implants	No. failures	Implant survival rate (%)	Follow-up (months)
Komarnyckij and London, 1998 (10)	16	43	2	95.3	9-47
Rosen et al., 1999 (17)	101	174	8	95.4	6-66
Deporter et al., 2000 (9)	16	26	0	100	6
Fugazzotto, 2002 (24)	103	116	2	98.3	48
Fugazzotto and De, 2002 (25)	150	167	3	97.8	36
Strietzel et al., 2002 (7)	22	22	2	91	3-12
Toffler, 2004 (11)	167	276	18	93.5	28
Brägger et al., 2004 (12)	19	25	1	96	12
Leblebicioglu et al., 2005 (23)	40	75	2	97.4	25
Li, 2005 (8)	42	61	5	98.1	6
Deporter et al., 2005 (14)	70	104	2	98.1	37.6
Nedir et al., 2006 (20)	17	25	0	100	12
Ferrigno et al., 2006 (15)	323	588	9	90.5	12-144
Diserens et al., 2006 (18)	55	66	2	98.5	6
Fermergard and Astrand; 2008 (22)	36	53	2	96	15-16
Diss et al., 2008 (21)	20	35	1	97.1	12
Kerrmalli et al., 2008 (16)	45	57	3	94.8	33.1
Lai et al., 2008 (26)	32	42	2	95.2	10
Barone et al., 2008 (13)	12	12	1	91.7	18
Sforza et al., 2008 (19)	26	39	1	97.4	12 (minimum)

Table 2. Residual crest height and bone gain in the included articles that specified these parameters.

Study	Mean residual crest height (range)(in mm)	Mean bone gain (range)(in mm)
Komarnyckyj and London, 1998 (10)	7.1 (3-10)	3.8
Bragger et al., 2004 (12)	7.7 (6-9)	4
Toffler, 2004 (11)	7 (2.3-10.3)	5.5
Leblebicioglu et al., 2005 (23)	5.4	3.2
Deporter et al., 2005 (14)	6.5	3.2
Nedir et al., 2006 (20)	5.4	2.5
Ferrigno et al., 2006 (15)	6.3	4.4
Diserens et al., 2006 (18)	9.1 (4.9-11.7)	3.3
Fermegard and Astrand, 2008 (22)	(3-12)	-
Lai et al., 2008 (26)	6.4 (4-8)	-
Kermalli et al., 2008 (16)	-	(0.5-6.5)
Barone et al., 2008 (13)	7.8	4.2
Diss et al., 2008 (21)	4.2	-

Sixty-six implants failed in 58 patients. The percentage implant survival rate with the osteotome technique was 90.5-100%. A recent study (27) observed no differences in the survival of implants placed after direct or indirect sinus lift, or in native bone in the posterior maxilla. Several authors (11,17,22) have pointed to residual bone height as a predictor of the survival of implants placed using the osteotome technique with sinus lift. Toffler et al. (11) recorded a 73.3% survival rate when the residual crest height measured 4 mm or less, versus 93.5% in the case of the total implants. Rosen et al. (17) obtained similar results: the global implant survival rate was 96% and 85.7% in the presence of residual crest heights of 4 mm or less, respectively. Fermegård et al. (22) documented two failures out of 53 implants. In both cases the residual bone height measured 4 mm or less.

Conclusion

The survival rate of implants placed with the osteotome technique is high and does not differ with respect to implant placement with the conventional technique.

References

- Summers RB. A new concept in maxillary implant surgery: the osteotome technique. *Compendium*. 1994;15:152, 154-6.
- Summers RB. The osteotome technique: Part 2-The ridge expansion osteotomy (REO) procedure. *Compendium*. 1994;15:422,424,426.
- Summers RB. The osteotome technique: Part 3-Less invasive methods of elevating the sinus floor. *Compendium*. 1994;15:698,700,702-4.
- Peñarrocha-Diago M, Rambla-Ferrer J, Perez V, Pérez-Garrigues H. Benign paroxysmal vertigo secondary to placement of maxillary implants using the alveolar expansion technique with osteotomes: a study of 4 cases. *Int J Oral Maxillofac Implants*. 2008;23:129-32.

- Shalabi MM, Manders P, Mulder J, Jansen JA, Creugers NH. A meta-analysis of clinical studies to estimate the 4.5-year survival rate of implants placed with the osteotome technique. *Int J Oral Maxillofac Implants*. 2007;22:110-6.
- Calvo-Guirado JL, Gómez-Moreno G, López-Marí L, Ortiz-Ruiz AJ, Guardia-Muñoz J. Atraumatic maxillary sinus elevation using threaded bone dilators for immediate implants. A three-year clinical study. *Med Oral Patol Oral Cir Bucal*. 2010;15:e366-70.
- Strietzel FP, Nowak M, Kuchler I, Friedmann A. Peri-implant alveolar bone loss with respect to bone quality after use of the osteotome technique: results of a retrospective study. *Clin Oral Implants Res*. 2002;13:508-13.
- Li TF. Sinus floor elevation: a revised osteotome technique and its biological concept. *Compend Contin Educ Dent*. 2005;26:619-26.
- Deporter D, Todescan R, Caudry S. Simplifying management of the posterior maxilla using short, porous-surfaced dental implants and simultaneous indirect sinus elevation. *Int J Periodontics Restorative Dent*. 2000;20:476-85.
- Komarnyckyj OG, London RM. Osteotome single-stage dental implant placement with and without sinus elevation: a clinical report. *Int J Oral Maxillofac Implants*. 1998;13:799-804.
- Toffler M. Minimally invasive sinus floor elevation procedures for simultaneous and staged implant placement. *N Y State Dent J*. 2004;70:38-44.
- Brägger U, Gerber C, Joss A, Haenni S, Meier A, Hashorva E, et al. Patterns of tissue remodeling after placement of ITI dental implants using an osteotome technique: a longitudinal radiographic case cohort study. *Clin Oral Implants Res*. 2004;15:158-66.
- Barone A, Cornelini R, Ciaglia R, Covani U. Implant placement in fresh extraction sockets and simultaneous osteotome sinus floor elevation: a case series. *Int J Periodontics Restorative Dent*. 2008;28:283-9.
- Deporter DA, Caudry S, Kermalli J, Adegbenbo A. Further data on the predictability of the indirect sinus elevation procedure used with short, sintered, porous-surfaced dental implants. *Int J Periodontics Restorative Dent*. 2005;25:585-93.
- Ferrigno N, Laureti M, Fanali S. Dental implants placement in conjunction with osteotome sinus floor elevation: a 12-year life-table analysis from a prospective study on 588 ITI implants. *Clin Oral Implants Res*. 2006;17:194-205.

16. Kermalli JY, Deporter DA, Lai JY, Lam E, Atenafu E. Performance of threaded versus sintered porous-surfaced dental implants using open window or indirect osteotome-mediated sinus elevation: a retrospective report. *J Periodontol.* 2008;79:728-36.
17. Rosen PS, Summers R, Mellado JR, Salkin LM, Shanaman RH, Marks MH, et al. The bone-added osteotome sinus floor elevation technique: multicenter retrospective report of consecutively treated patients. *Int J Oral Maxillofac Implants.* 1999;14:853-8.
18. Diserens V, Mericske E, Schäppi P, Mericske-Stern R. Transcrestal sinus floor elevation: report of a case series. *Int J Periodontics Restorative Dent.* 2006;26:151-9.
19. Sforza NM, Marzadori M, Zucchelli G. Simplified osteotome sinus augmentation technique with simultaneous implant placement: a clinical study. *Int J Periodontics Restorative Dent.* 2008;28:291-9.
20. Nedir R, Bischof M, Vazquez L, Szmukler-Moncler S, Bernard JP. Osteotome sinus floor elevation without grafting material: a 1-year prospective pilot study with ITI implants. *Clin Oral Implants Res.* 2006;17:679-86.
21. Diss A, Dohan DM, Mouhyi J, Mahler P. Osteotome sinus floor elevation using Choukroun's platelet-rich fibrin as grafting material: a 1-year prospective pilot study with microthreaded implants. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2008;105:572-9.
22. Fermergård R, Astrand P. Osteotome sinus floor elevation and simultaneous placement of implants--a 1-year retrospective study with Astra Tech implants. *Clin Implant Dent Relat Res.* 2008;10:62-9.
23. Leblebicioglu B, Ersanli S, Karabuda C, Tosun T, Gokdeniz H. Radiographic evaluation of dental implants placed using an osteotome technique. *J Periodontol.* 2005;76:385-90.
24. Fugazzotto PA. Immediate implant placement following a modified trephine/osteotome approach: success rates of 116 implants to 4 years in function. *Int J Oral Maxillofac Implants.* 2002;17:113-20.
25. Fugazzotto PA, De PS. Sinus floor augmentation at the time of maxillary molar extraction: success and failure rates of 137 implants in function for up to 3 years. *J Periodontol.* 2002;73:39-44.
26. Lai HC, Zhang ZY, Wang F, Zhuang LF, Liu X. Resonance frequency analysis of stability on ITI implants with osteotome sinus floor elevation technique without grafting: a 5-month prospective study. *Clin Oral Implants Res.* 2008;19:469-75.
27. Huynh-Ba G, Friedberg JR, Vogiatzi D, Ioannidou E. Implant failure predictors in the posterior maxilla: a retrospective study of 273 consecutive implants. *J Periodontol.* 2008;79:2256-61.